expressions for Titchmarsh-Weyl m-functions

associated with the Hydrogen Atom on the half line

Using a computer algebra system to simplify

by

Cecilia Knoll and Charles Fulton

TECHNICAL REPORT

FLORIDA INSTITUTE OF TECHNOLOGY September 2007

Using a computer algebra system to simplify expressions for Titchmarsh-Weyl m-functions associated with the Hydrogen Atom on the half line^{*}

CECILIA KNOLL Department of Applied Mathematics Florida Institute of Technology Melbourne, Florida 32901-6975

CHARLES FULTON Department of Mathematical Sciences Florida Institute of Technology Melbourne, Florida 32901-6975

Abstract

Abstract: In this paper we give simplified formulas for certain polynomials which arise in some new Titchmarsh-Weyl m-functions for the radial part of the separated Hydrogen atom on the half line $(0, \infty)$ and two independent programs for generating them using the symbolic manipulator Mathematica.

1 Introduction

Recently Fulton[1] and Fulton and Langer [2] considered the following Sturm-Liouville problem for the hydrogen atom on the half line, $x \in (0, \infty)$:

$$-y'' + \left(-\frac{a}{x} + \frac{\ell(\ell+1)}{x^2}\right)y = \lambda y, \quad 0 < x < \infty$$

$$(1.1)$$

$$\lim_{x \to 0} W_x \left(y, x^{\frac{1}{2}} J_{2\ell+1}(\sqrt{4ax}) \right) = 0.$$
(1.2)

In [1] a fundamental system of Frobenius solutions defined at x = 0 was introduced having the forms:

$$\phi(x,\lambda) = x^{\ell+1} \left[1 + \sum_{n=1}^{\infty} a_n(\lambda) x^n \right] = \frac{1}{(-2\sqrt{\lambda})^{\ell+1}} M_{\beta,\ell+\frac{1}{2}}(-2ix\sqrt{\lambda}), \quad \beta := \frac{ia}{2\sqrt{\lambda}}$$
(1.3)

0

^{*}This research partially supported by National Science Foundation Grant DMS-0109022 to Florida Institute of Technology.

and

$$\theta(x,\lambda) = -\frac{1}{2\ell+1} \left[K_{\ell}(\lambda)\phi(x,\lambda)\ln x + x^{-1} + \sum_{n=1}^{\infty} d_n(\lambda)x^{-\ell+n} \right]$$
(1.4)

Where $a_n(\lambda)$ and $d_n(\lambda)$ are polynomials in λ , $M_{\beta,\ell+\frac{1}{2}}$ is the Whittaker function of first kind and

$$K_{\ell}(\lambda) = -\frac{a}{(2\ell+1)!(2\ell)!} \prod_{j=1}^{\ell} \left[4\lambda j^2 + a^2 \right] = \frac{(-2i\sqrt{\lambda})^{2\ell+1}(-\ell-\beta)_{2\ell+1}}{(2\ell)!(2\ell+1)!}.$$
(1.5)

Then a Titchmarsh-Weyl m-function was introduced in [1] by the requirement

$$\theta(x,\lambda) - m_{\ell}(\lambda)\phi(x,\lambda) \in L_2(0,\infty),$$
(1.6)

which gives m_{ℓ} as

$$m_{\ell}(\lambda) = -ak_{\ell}(\lambda) \left\{ \log(-2i\sqrt{\lambda}) + \Psi \left(1 - \frac{ia}{2\sqrt{\lambda}} \right) - H_{2\ell} + 2\gamma \right\}$$
$$- ak_{\ell}(\lambda) \left\{ \sum_{j=1}^{\ell} \frac{1}{\left(j - \frac{ia}{2\sqrt{\lambda}}\right)} \right\} + \frac{(i\sqrt{\lambda})^{2\ell+1}}{(2\ell+1)!} \left[\sum_{k=0}^{2\ell} \frac{2^{k} \left(-\ell - \frac{ia}{2\sqrt{\lambda}} \right)_{k}}{k!(2\ell+1-k)} \right]$$
(1.7)

where $\Psi(z) = \frac{\Gamma'(z)}{\Gamma(z)}$ is the psi or digamma function, $H_{2\ell} = \sum_{j=1}^{2\ell} \frac{1}{j}$, $\gamma =$ Euler's constant, and the Pochhammer symbol is defined for any complex z as $(z)_k = z(z+1)...(z+k-1)$. Here $k_\ell(\lambda)$ is the polynomial of degree ℓ defined by

$$k_{\ell}(\lambda) = -\frac{1}{a(2\ell+1)} K_{\ell}(\lambda) = -\frac{1}{(2\ell+1)!} \prod_{j=1}^{\ell} \left[4\lambda j^2 + a^2 \right].$$
(1.8)

For $\ell = 0$, we define $k_0(\lambda) = 1$. The function $m_\ell(\lambda)$ is an analytic function in the half planes Im $\lambda < 0$ and Im $\lambda > 0$, has poles on the negative x-axis at the eigenvalues of the problem (1.1)-(1.2), and a branch cut, corresponding to the continuous spectrum, on the positive real λ -axis. In [2] a Pick-Nevalinna representation of $m_0(\lambda)$ was obtained for $\ell = 0$ and for $\ell \ge 1$ it was shown that $m_\ell(\lambda)$ has a Q-function representation which puts it in the class N_κ of generalized Nevalinna functions with $\kappa = \left[\frac{\ell+1}{2}\right]$.

Our purpose in this paper is to show that the last two terms in $m_{\ell}(\lambda)$ can be decomposed into real and imaginary parts as

$$-ak_{\ell}(\lambda)\left\{\sum_{j=1}^{\ell}\frac{1}{j-\frac{ia}{2\sqrt{\lambda}}}\right\}+\frac{(i\sqrt{\lambda})^{2\ell+1}}{(2\ell+1)!}\left[\sum_{k=0}^{2\ell}\frac{2^{k}\left(-\ell-\frac{ia}{2\sqrt{\lambda}}\right)_{k}}{k!(2\ell+1-k)}\right]=i\sqrt{\lambda}k_{\ell}(\lambda)+\frac{r_{\ell}(\lambda)}{2\ell+1},$$
(1.9)

where $r_{\ell}(\lambda)$ is a polynomial of degree ℓ in λ .

Effectively, the decomposition (1.9) into real and imaginary parts becomes a defining equation for $r_{\ell}(\lambda)$. Our first *Mathematica* program enables the decomposition to be verified. Next, in Section 3 we separate the left hand side of (1.9) into real and imaginary parts by introducing polynomial representations of $(-\ell - t)_k$ in t and

$$\prod_{\substack{j=1\\j\neq m}}^{\ell} \left(\lambda + \frac{a^2}{4j^2}\right) \text{ in } \lambda,$$

so that the real part can be represented as a polynomial of degree ℓ . This yields a real, somewhat explicit, representation for $r_{\ell}/(2\ell+1)$, and using it a second *Mathematica* program shows that this real representation yields the same result as the first *Mathematica* program.

2 The First Mathematica program

A simple program in *Mathematica* can be used to verify that the polynomial $r_{\ell}(\lambda)$ in (1.9) is real valued. This makes use of the built-in function **Pochhammer** [a, k] which executes the multiplications in the Pochhammer symbol $(a)_k$.

Using the second expression in (1.5) for $K_{\ell}(\lambda)$, we have using (1.8) that

$$b := -ak_{\ell}(\lambda) = \frac{K_{\ell}(\lambda)}{2\ell + 1} = \frac{(-2i\sqrt{\lambda})^{2\ell + 1}(-\ell - \beta)_{2\ell + 1}}{[(2\ell + 1)!]^2}$$

Accordingly, solving (1.9) for $\frac{r_{\ell}(\lambda)}{2\ell+1}$, we have

$$\frac{r_{\ell}(\lambda)}{2\ell+1} = b \left\{ \sum_{j=1}^{\ell} \frac{1}{j-\beta} \right\} + c + \frac{bi\sqrt{\lambda}}{a}, \text{ where } c := \frac{(i\sqrt{\lambda})^{2\ell+1}}{(2\ell+1)!} \left[\sum_{k=0}^{2\ell} \frac{2^k \left(-\ell - \frac{ia}{2\sqrt{\lambda}}\right)_k}{k!(2\ell+1-k)} \right].$$
(2.1)

Following is the *Mathematica* program which implements equation (2.1). The output for $\frac{r_{\ell}(\lambda)}{2\ell+1}$ for $\ell = 1, 2, 3$, and 4 is shown. The program was executed up to $\ell = 30$ showing that $r_{\ell}(\lambda)$ remained real-valued. Observe that the constant *a* multiplies all terms of $\frac{r_{\ell}(\lambda)}{2\ell+1}$ which in the *Mathematica* output is ans1. This is also poved in (3.7) below.

First Program

Program input: ℓ

$$\begin{split} \ell &= 4; \\ \text{While}[\ell < 6, \beta = \frac{Ia}{2\sqrt{\lambda}}; \\ b &= \text{Expand}[(-2I\sqrt{\lambda})^{2\ell+1} \frac{\text{Pochhammer}[-\ell-\beta,2\ell+1]}{(2\ell+1)!^2}; (*\text{This generates } K_{\ell}/(2\ell+1). \text{ See } (1.5).*) \\ c &= \text{Apart}[\text{Simplify}[\frac{(I\sqrt{\lambda})^{2\ell+1}}{(2\ell+1)!} \sum_{k=0}^{2\ell} \frac{\text{Pochhammer}[-\ell-\beta,k]2^k}{k!(2\ell+1-k)}]]; \\ g &= b \left(-\sum_{j=1}^{2\ell} \frac{1}{j}\right); \\ e &= b \sum_{j=1}^{\ell} \frac{1}{j-\beta}; \\ f &= \frac{bI\sqrt{\lambda}}{a}; \\ \text{Print}[\text{"For } \ell = \text{", } \ell]; \\ ans1 &= \text{Simplify}[c + e + f]; \\ \text{Print}[\text{"ans1} = \text{"ans1}]; (*\text{This generates the RHS of } (2.1).*) \\ ans2 &= \text{Series}[ans1, \{\lambda, 0, \ell\}]; \text{Print}[\text{"ans3} = \text{", ans3}]; (*\text{This generates the RHS of } (2.6)*) \\ ans4 &= \text{Series}[ans3, \{\lambda, 0, \ell\}]; \text{Print}[\text{"= ", ans4}]; \\ \ell &= \ell + 1] \end{split}$$

1

Program output:

For $\ell = 1$

$$ans1 = -\frac{a\lambda}{36}$$

$$ans3 = \frac{1}{72}(3a^3 + 10a\lambda) = \frac{a^3}{24} + \frac{5a\lambda}{36}$$
(2.2)

For $\ell = 2$

$$ans1 = -\frac{a\lambda(a^2 + 13\lambda)}{7200} = -\frac{a^3\lambda}{7200} - \frac{13a\lambda^2}{7200}$$

$$ans3 = \frac{25a^5 + 476a^3\lambda + 1288a\lambda^2}{172800} = \frac{a^5}{6912} + \frac{119a^3\lambda}{43200} + \frac{161a\lambda^2}{21600}$$

(2.3)

For $\ell = 3$

$$ans1 = -\frac{a\lambda(a^4 + 46a^2\lambda + 400\lambda^2)}{8467200} = -\frac{a^5\lambda}{8467200} - \frac{23a^3\lambda^2}{4233600} - \frac{a\lambda^3}{21168}$$

$$ans3 = \frac{49a^7 + 2684a^5\lambda + 35656a^3\lambda^2 + 88896a\lambda^3}{508032000}$$

$$= \frac{a^7}{10368000} + \frac{671a^3\lambda}{127008000} + \frac{4457a^3\lambda^2}{63504000} + \frac{463a\lambda^3}{2646000}$$
(2.4)

For $\ell = 4$

$$ans1 = -\frac{a\lambda(a^{6} + 107a^{4}\lambda + 3124a^{2}\lambda^{2} + 22548\lambda^{3})}{32920473600}$$

$$= -\frac{a^{7}\lambda}{32920473600} - \frac{107a^{5}\lambda^{2}}{32920473600} - \frac{781a^{3}\lambda^{3}}{8230118400} - \frac{1879a\lambda^{4}}{2743372800}$$

$$ans3 = \frac{761a^{9} + 90200a^{7}\lambda + 3204208a^{5}\lambda^{2} + 36438400a^{3}\lambda^{3} + 86960256a\lambda^{4}}{36870930432000}$$

$$= \frac{761a^{9}}{36870930432000} + \frac{451a^{7}\lambda}{184354652160} + \frac{4087a^{3}\lambda^{3}}{11522165760} + \frac{226459a\lambda^{4}}{96018048000}.$$

$$(2.5)$$

Output for the polynomials

$$p_{\ell}(\lambda) := \frac{r_{\ell}(\lambda)}{2\ell + 1} + ak_{\ell}(\lambda)H_{2\ell}, \qquad (2.6)$$

which arise in the representation of the m_{ℓ} function from [1, Equation(8.15)],

$$m_{\ell}(\lambda) = k_{\ell}(\lambda) \left\{ -a \log(-2i\sqrt{\lambda}) - a\Psi\left(1 - \frac{ia}{2\sqrt{\lambda}}\right) - 2\gamma a + i\sqrt{\lambda} \right\} + p_{\ell}(\lambda)$$

is also given below for $\ell=1,2,3,4.$

$$p_{1} = \frac{a^{3}}{24} + \frac{5a\lambda}{36},$$

$$p_{2} = \frac{a^{5}}{6912} + \frac{119a^{3}}{43200}\lambda + \frac{161a}{21600}\lambda^{2},$$

$$p_{3} = \frac{a^{7}}{10368000} + \frac{671a^{5}}{127008000}\lambda + \frac{4457a^{3}}{63504000}\lambda^{2} + \frac{463a}{2646000}\lambda^{3}$$

$$p_{4} = \frac{761a^{9}}{36870930432000} + \frac{451a^{7}}{184354652160}\lambda + \frac{4087a^{5}}{47029248000}\lambda^{2} + \frac{11387a^{3}}{11522165760}\lambda^{3} + \frac{226459a}{96018048000}\lambda^{4}.$$

Here p_ℓ is printed as ans3 in the above program.

3 Explicit Representation for r_{ℓ}

In this section we give a method for separating the expression in equation (1.9) into real and imaginary parts, yielding a real representation for the polynomial $r_{\ell}(\lambda)$. The difficulty arises from the complicated product in the Pochhammer symbol $(\ell - \beta)_k$ where $\beta = \frac{ia}{2\sqrt{\lambda}}$. Replacing β by a real variable t, we let the coefficients of the polynomial $(-\ell - t)_k$ be defined by

$$g_k(t) := (-\ell - t)_k = \prod_{j=0}^{k-1} (-\ell + j - t) = \sum_{n=0}^k \alpha(k, n) t^n = \sum_{j=0}^{k_1} \alpha(k, 2j) t^{2j} + \sum_{j=0}^{k_2} \alpha(k, 2j + 1) t^{2j+1}, \quad (3.1)$$

where $k_1 = \left[\frac{k}{2}\right]$ and $k_2 = \left[\frac{k-1}{2}\right]$. Here $\alpha(k, n) = \alpha_\ell(k, n)$, and we are interested for fixed ℓ to have $\alpha_\ell(k, n)$ available for all $0 \le k \le 2\ell$, and all $0 \le n \le k$. For n = 0, the constant term is $\alpha_\ell(k, 0) = \prod_{j=0}^{k-1} (-\ell + j) = (-\ell)_k$, $k = 0, 1, \dots 2\ell$. Formulas for larger values of n become increasingly more complicated and are not known in closed form. We can, however, represent the real and the imaginary parts of (1.9) in terms of $\alpha(k, n)$. Putting $t = \beta = \frac{ia}{2\sqrt{\lambda}}$ in (3.1) gives

$$g_{k}(\beta) = \sum_{n=0}^{k} \alpha(k,n) \left(\frac{a}{2}\right)^{n} i^{n} (\lambda^{-\frac{1}{2}})^{n}$$

$$= \sum_{j=0}^{k_{1}} (-1)^{j} \alpha(k,2j) \left(\frac{a}{2}\right)^{2j} i^{n} (\lambda^{-\frac{1}{2}})^{2j}$$

$$+ i \sum_{j=0}^{k_{2}} (-1)^{j} \alpha(k,2j+1) \left(\frac{a}{2}\right)^{2j+1} (\lambda^{-\frac{1}{2}})^{2j+1}.$$
(3.2)

Now for the $(i\sqrt{\lambda})^{2\ell+1}$ term in (1.9) we have $(i)^{2\ell+1} = (-1)^{\ell}i$, $\ell = 0, 1, 2, ...$ Thus the second sum in (1.9) may be written as

$$i(-1)^{\ell} \sum_{k=0}^{2\ell} \frac{2^{k}}{(2\ell+1)!k!(2\ell+1-k)} g_{k}(\beta) (\lambda^{\frac{1}{2}})^{2\ell+1}$$

$$= -(-1)^{\ell} \sum_{k=1}^{2\ell} \frac{2^{k}}{(2\ell+1)!k!(2\ell+1-k)} \left[\sum_{j=0}^{k_{2}} (-1)^{j} \alpha(k,2j+1) \left(\frac{a}{2}\right)^{2j+1} (\lambda^{\frac{1}{2}})^{2\ell+1-(2j+1)} \right]$$

$$+i(-1)^{\ell} \sum_{k=0}^{2\ell} \frac{2^{k}}{(2\ell+1)!k!(2\ell+1-k)} \left[\sum_{j=0}^{k_{1}} (-1)^{j} \alpha(k,2j) \left(\frac{a}{2}\right)^{2j+1} (\lambda^{\frac{1}{2}})^{2\ell+1-2j} \right]$$

$$= \sum_{k=0}^{2\ell} \frac{2^{k} A_{k}}{(2\ell+1)!k!(2\ell+1-k)} + i \sum_{k=0}^{2\ell} \frac{2^{k} B_{k}}{(2\ell+1)!k!(2\ell+1-k)}$$

$$(3.3)$$

where

$$A_k = (-1)^{\ell+1} \sum_{j=0}^{k_2} (-1)^j \alpha(k, 2j+1) \left(\frac{a}{2}\right)^{2j} \lambda^{\ell-j}$$
(3.4)

and

$$B_k = (-1)^{\ell} \sum_{j=0}^{k_1} (-1)^j \alpha(k, 2j) \left(\frac{a}{2}\right)^{2j} \lambda^{\ell-j} \lambda^{\frac{1}{2}}$$
(3.5)

Similarly, we may decompose the first term in (1.9) as

$$\sum_{m=1}^{\ell} \frac{1}{m - \frac{ia}{2\sqrt{\lambda}}} = \sum_{m=1}^{\ell} \frac{4\lambda m}{4\lambda m^2 + a^2} + i \sum_{m=1}^{\ell} \frac{2a\lambda^{\frac{1}{2}}}{4\lambda m^2 + a^2}.$$
(3.6)

Using (1.8) and cancelling one $4\lambda j^2 + a^2$ factor for each term in the above sums, we find that

$$-ak_{\ell}(\lambda) \left[\sum_{m=1}^{\ell} \frac{1}{m - \frac{ia}{2\sqrt{\lambda}}} \right] = -\frac{a}{[(2\ell+1)!]^2} \left[\sum_{m=1}^{\ell} 4\lambda m \prod_{\substack{j=1\\ j \neq m}}^{\ell} (4\lambda j^2 + a^2) \right] +i \left(-\frac{a}{[(2\ell+1)!]^2} \left[\sum_{m=1}^{\ell} 2a\lambda^{\frac{1}{2}} \prod_{j=1}^{\ell} (4\lambda j^2 + a^2) \right] \right).$$
(3.7)

Combining (3.3) and (3.7) and taking real and imaginary parts of (1.9), we thus obtain:

$$\sqrt{\lambda}k_{\ell}(\lambda) = -\frac{2a^2\sqrt{\lambda}}{[(2\ell+1)!]^2} \left[\sum_{\substack{m=1\\j\neq m}}^{\ell} \prod_{\substack{j=1\\j\neq m}}^{\ell} (4\lambda j^2 + a^2) \right] + \sum_{k=0}^{2\ell} \frac{2^k B_k}{(2\ell+1)!k!(2\ell+1-k)}$$
(3.8)

and

$$\frac{r_{\ell}(\lambda)}{2\ell+1} = -\frac{4\lambda a}{[(2\ell+1)!]^2} \left[\sum_{m=1}^{\ell} m \prod_{\substack{j=1\\j\neq m}}^{\ell} (4\lambda j^2 + a^2) \right] + \sum_{k=1}^{2\ell} \frac{2^k A_k}{(2\ell+1)!k!(2\ell+1-k)}.$$
(3.9)

Since every A_k involves a as a factor, it follows that a factor of a occurs in both terms on the right hand side of (3.9). Hence a = 0 will give $r_{\ell}(\lambda) = 0$.

To bring the expression in (3.9) into a simpler form, we now isolate the coefficients of the powers of λ . For the second term in (3.9) we insert the formula (3.4) for A_k into the sum, change the summation index $(m = \ell - j)$, and then interchange the order of summation:

7

$$\sum_{k=1}^{2\ell} \frac{2^k A_k}{(2\ell+1)!k!(2\ell+1-k)}$$

$$= \sum_{k=1}^{2\ell} \sum_{j=0}^{k_2} \frac{2^k (-1)^{\ell+1+j} \alpha(k, 2j+1) \left(\frac{a}{2}\right)^{2j+1} \lambda^{\ell-j}}{(2\ell+1)!k!(2\ell+1-k)}$$

$$= \sum_{k=1}^{2\ell} \sum_{m=\ell-k_2}^{\ell} \left[\frac{2^k (-1)^{m+1} \alpha(k, 2(\ell-m)+1) \left(\frac{a}{2}\right)^{2(\ell-m)+1}}{(2\ell+1)!k!(2\ell+1-k)} \right] \lambda^m$$

$$= \sum_{m=1}^{\ell} \sum_{k=2(\ell-m)+1}^{2\ell} \left[\frac{2^k (-1)^{m+1} \alpha(k, 2(\ell-m)+1) \left(\frac{a}{2}\right)^{2(\ell-m)+1}}{(2\ell+1)!k!(2\ell+1-k)} \right] \lambda^m$$

$$= \sum_{j=1}^{\ell} d_j \lambda^j$$
(3.10)

where

$$d_j := \frac{a}{2} \sum_{k=2(\ell-j)+1}^{2\ell} \left[\frac{2^k (-1)^{m+1} \left(\frac{a}{2}\right)^{2(\ell-j)} \alpha(k, 2(\ell-j)+1)}{(2\ell+1)! k! (2\ell+1-k)} \right].$$
(3.11)

Similarly, to isolate the powers of λ in the first term in (3.7) we first define the coefficients $\gamma(m, n)$ of the $\ell - 1$ degree polynomial,

$$\prod_{\substack{j=1\\j\neq m}}^{\ell} \left(\lambda + \frac{a^2}{4j^2}\right) = \sum_{n=0}^{\ell-1} \gamma(m, n) \lambda^n.$$
(3.12)

Insertion of this into the first term in (3.9), changing the summation index (j = n + 1), and interchanging the order of summation then yields:

$$\frac{-4\lambda a}{[(2\ell+1)!]^2} \left[\sum_{m=1}^{\ell} m \prod_{\substack{j=1\\ j \neq m}}^{\ell} (4\lambda j^2 + a^2) \right] \\
= \frac{-a4^{\ell}(\ell!)^2}{[(2\ell+1)!]^2} \left[\sum_{m=1}^{\ell} \sum_{n=0}^{\ell-1} \left[\frac{\gamma(m,n)}{m} \right] \lambda^{n+1} \right] \\
= \frac{-a4^{\ell}(\ell!)^2}{[(2\ell+1)!]^2} \left[\sum_{m=1}^{\ell} \sum_{j=1}^{\ell} \left[\frac{\gamma(m,j-1)}{m} \right] \lambda^j \right] \\
= \frac{-a4^{\ell}(\ell!)^2}{[(2\ell+1)!]^2} \left[\sum_{j=1}^{\ell} \sum_{m=1}^{\ell} \left[\frac{\gamma(m,j-1)}{m} \right] \lambda^j \right] \\
= \sum_{j=1}^{\ell} c_j \lambda^j \tag{3.13}$$

where

$$c_j := -\frac{a4^{\ell}(\ell!)^2}{[(2\ell+1)!]^2} \sum_{m=1}^{\ell} \left(\frac{\gamma(m,j-1)}{m}\right).$$
(3.14)

Putting (3.10) and (3.13) in (3.9) we thus have for the polynomial $r_{\ell}(\lambda)/(2\ell+1)$ the representation,

$$\frac{r_{\ell}(\lambda)}{2\ell+1} = \sum_{j=1}^{\ell} (c_j + d_j)\lambda^j.$$
 (3.15)

Here it is clear that a is a common factor in c_j and d_j for all j, and hence for the case a = 0, $r_{\ell} \equiv 0$. The expression (3.15) is somewhat more explicit than the expression (2.1) and (3.9) since the coefficients of the powers of λ are isolated, and there are no complex terms present. On the other hand, further simplifications are certainly desirable; however, this requires closed form formulas for $\alpha(k,n)$ and $\gamma(m,n)$ which remain elusive. However, the formulas for c_j and d_j are easily implemented using a symbolic manipulator.

4 The Second Mathematica Program

As an independent check on the first *Mathematica* program we implemented the formulas (3.11) and (3.14) for d_j and c_j and computed the polynomial of degree ℓ in (3.15). This required computing and storing the

coefficient $\alpha(k,n)$ and $\gamma(m,n)$ in (3.1) and (3.12). Following is the *Mathematica* program that does this to compute $r_{\ell}(\lambda)/(2\ell+1)$. The output for $\ell = 1, 2, 3$ and 4 is shown. The program was executed up to $\ell = 30$ and gave exact agreement with the first *Mathematica* program.

Second Program

Program input: ℓ

$$\begin{split} \ell &= 4; k = 0; \\ &\text{While}[k \leq 2\ell, \\ &p[t_{-}] = \text{Expand}[\text{Pochhammer}[-\ell - t, k]]; n = k; i = 0; \\ &\text{While}[i \leq n, \\ &\alpha[k, i] = p[0]; \\ &p[t_{-}] = \text{Simplify}[\text{Expand}[(p[t] - \alpha[k, i])/t]]; \\ &i + +] \\ &k + +] \\ &\text{Clear}[k]; j = 1; \\ &\text{While}[j \leq \ell, \\ &d[j] = \sum_{k=2(\ell-j)+1}^{2^{\ell}(-1)^{j-1}\alpha[k, 2\ell - 2j + 1]\left(\frac{a}{2}\right)^{2\ell - 2j + 1}} ; j + +] \\ &\text{Clear}[j]; m = 1; \\ &\text{While}[m \leq \ell, \\ &q[t_{-}] = \left(\prod_{j=1}^{\ell} \left(t + \frac{a^2}{4j^2}\right)\right) / \left(t + \frac{a^2}{4m^2}\right); n = \ell - 1; i = 0; \\ &\text{While}[m \leq \ell, \\ &\gamma[m, i] = q[0]; \\ &q[t_{-}] = \text{Simplify}[\text{Expand}[(q[t] - \gamma[m, i])/t]]; \\ &i + +] \\ &m + +] \\ &\text{Clear}[i]; j = 1; \\ &\text{While}[j \leq \ell, \\ &c[j] = \frac{-a4^{\ell}(\ell!)^2}{((2\ell+1)!)^2} \sum_{m=1}^{\ell} \frac{\gamma[m, j - 1]}{m}; j + +; \\ &\text{Print}[\sum_{j=1}^{\ell}(c[j] + d[j])t^j] \end{split}$$

Output of second program

$$\frac{r_\ell}{(2\ell+1)} = -\frac{at}{36}$$

For $\ell = 2$:

For $\ell = 1$:

$$\frac{r_{\ell}}{(2\ell+1)} = -\frac{a^3t}{7200} - \frac{13at^2}{7200}$$

For
$$\ell = 3$$
:

$$\frac{r_{\ell}}{(2\ell+1)} = -\frac{a^5t}{8467200} - \frac{23a^3t^2}{4233600} - \frac{at^3}{21168}$$
For $\ell = 4$:

$$\frac{r_{\ell}}{(2\ell+1)} = -\frac{a^7t}{32920473600} - \frac{107a^5t^2}{32920473600} - \frac{781a^3t^3}{8230118400} - \frac{1879at^4}{2743372800}$$

This output is in agreement with the output of the first program, and was also checked up to $\ell = 30$. The two *Mathematica* programs verify that the imaginary part of the expression (2.1) for $r_{\ell}(\lambda)/(2\ell+1)$ is zero

for all values of ℓ for which the programs were executed. A general proof that this is true for all ℓ , that is, a proof of (3.8) (or, equivalently, a proof of (3.9)) requires that the $\alpha(k, n)$ and $\gamma(m, n)$ coefficients be obtained in a simplified form, and this appears to be a formidable task. It is quite difficult to perform induction on ℓ to prove (3.8) or (3.9) because of the complicated manner in which $\alpha(k, n) = \alpha_{\ell}(k, n)$ and $\gamma(m, n) = \gamma_{\ell}(m, n)$ change with ℓ . Nevertheless, it may be possible to construct rigorous proofs by making use of some combinatorial analysis. For example there are general forms for the coefficients of a polynomial having k known roots x_i , i = 1, 2, ..., k. Sen and Krishnamurthy [3], for example, show that

$$q_k(x) := \prod_{j=1}^k (x - x_j) = x^k + \sum_{m=1}^k a_m x^{k-m}$$
(4.1)

with

$$a_m := (-1)^m S_m := (-1)^m \sum x_1 x_2 \dots x_m, \tag{4.2}$$

where the sum is taken over all the products of x_i taken m at a time. Thus, for the polynomial $g_k(t)$ of (3.1) we have

$$g_k(t) = (-1)^k \prod_{j=0}^{k-1} [t - (j-\ell)] = (-1)^k \prod_{j=1}^k [t - (j-1-\ell)] = (-1)^k \left(t^k + \sum_{m=1}^k a_m t^{k-m} \right),$$
(4.3)

with

$$a_m = a_\ell(k,m) = (-1)^m \sum t_1 t_2 \dots t_m \tag{4.4}$$

where the sum is taken over all the products of $t_j := j - 1 - \ell$ taken m at a time. But, a general solution for a_m as a function of k and ℓ , valid for all ℓ and all $0 \le k \le 2\ell$, remains elusive. Another idea would be to establish (3.8) or (3.9) by induction on ℓ , but it unfortunately appears difficult to employ the induction hypothesis. So, an analytic proof of (3.8) or (3.9) valid for all $\ell \ge 1$ remains an open problem.

5 Conclusion

The simplification of $a_{\ell}(k, n)$ and a rigorous proof of (3.8) and/or (3.9) remain as open problems. We have, however, given in this note two *Mathematica* programs which implicitly establish both of these results.

References

- C. Fulton, Titchmarsh-Weyl m-functions for Second-order Sturm Liouville Problems with two singular endpoints, Math. Nachr. 281 (10) (2008), 1418-1475.
- [2] C. Fulton and H. Langer, Sturm-Liouville operators with singularities and Generalized Nevanlinna functions, Complex Anal. and Oper. Theory, to appear.
- [3] E.V. Krishnamurthy and S.K. Sen, Numerical Algorithms: Computations in Science and Engineering, Affiliated East West Press, New Delhi, 2001.