# ANTI-PLURICANONICAL SYSTEMS ON Q-FANO 3-FOLDS

# MENG CHEN

ABSTRACT. We investigate birationality of the anti-pluricanonical map  $\varphi_{-m}$ , the rational map defined by the anti-pluricanonical system  $|-mK_X|$ , on weak Q-Fano 3-folds.

### 1. Introduction

Throughout a normal projective variety X is called a *terminal weak*  $\mathbb{Q}$ -Fano variety if X has at worst terminal singularities, X is  $\mathbb{Q}$ -factorial and  $-K_X$  is a nef and big  $\mathbb{Q}$ -Cartier divisor where  $K_X$  is the canonical Weil divisor of X. A weak  $\mathbb{Q}$ -Fano variety is said to be  $\mathbb{Q}$ -Fano if  $-K_X$  is  $\mathbb{Q}$ -ample.

According to the Minimal Model Program (MMP), Q-Fano varieties form a fundamental aspect in birational geometry.

Let X be a terminal weak Q-Fano 3-fold. The number  $g(X) := h^0(X, \omega_X^{[-1]}) - 2$  is usually called the genus of X. Denote by  $r_X$  the Cartier index of X. For a positive integer m, we define  $|-mK_X|$  to be the natural closure of the complete linear system  $|-mK_{X^0}|$  where  $X^0 := X \setminus X_{\text{Sing}}$ . Whenever  $P_{-m} := h^0(X, \mathcal{O}_X^{[-m]}) \geq 2$ , the rational map  $\varphi_{-m}$  corresponding to  $|-mK_X|$  gives rise to the so-called "anti-canonical geometry" of X. Therefore a natural interesting question is to find a practical number m such that  $\varphi_{-m}$  is birational onto its image. Such a number m (independent of X) exists due to the boundedness theorem which was proved by Kawamata [9] (for Picard number one case) and by Kollár-Miyaoka-Mori-Takagi [12] (for general case). Even though, it is very interesting to ask the following:

**Problem 1.** Can one find an optimal constant c such that  $\varphi_{-c}$  is birational onto the image for all weak Q-Fano 3-folds?

When X is Gorenstein (i.e.  $r_X = 1$ ), one may take c = 5 according to Ando [2]. In fact, considerable classification has been done for Fano 3-folds (see, for example, [6], [7], [8], [14], [15], [16], [18], [24] and so on). When X is non-Gorenstein, Problem 1 is open in general.

The author was supported by National Outstanding Young Scientist Foundation (#10625103) and National Natural Science Foundation of China (Key Project: #10731030).

In contrast to  $\varphi_m$  or  $|mK_X|$ ,  $\varphi_{-m}$  shows some pathological nature. For instance, the birationality behavior of  $\varphi_{-m}$  is not birationally invariant! This often makes the situation be more complicated.

The aim of this note is to build an effective mechanic to study the birationality of  $\varphi_{-m}$  on weak Q-Fano 3-folds. Our main technical theorem is **Theorem** 3.5, which has direct applications to various situations in Section 4 (see Theorems 4.1, 4.2, 4.5). In fact, since  $P_{-8} > 1$  by [3, Theorem 1.1 (ii)], the rational map  $\varphi_{-8} : X \dashrightarrow \mathbb{P}^N$  can be properly defined. Thus the geometry of X can be detected by studying  $\varphi_{-8}$ . Here is a direct consequence of our general arguments in Section 4:

**Theorem 1.1.** Let X be a terminal weak  $\mathbb{Q}$ -Fano 3-fold with  $r_X > 1$ . Then  $\varphi_{-m}$  is birational onto its image under one of the following conditions:

- (i) dim  $\varphi_{-8}(X) = 3$  and  $m \ge 32$ ;
- (ii) dim  $\overline{\varphi_{-8}(X)} = 2$  and  $m \ge max\{2r_X + 16, 48\};$
- (iii) dim  $\overline{\varphi_{-8}(X)} = 1$  and

$$m \ge \begin{cases} 16, & r_X = 2; \\ 20, & r_X = 3; \\ 3r_X + 10, & r_X \ge 4. \end{cases}$$

**Remark 1.2.** By virtue of the boundedness theorem for Q-Fano 3-folds,  $r_X$  is upper bounded. Therefore Theorem 1.1 has actually obtained a universal constant c with  $\varphi_{-c}$  birational. However such a constant c might still be a little bit far from optimal.

Restricting our interest to Q-Fano 3-folds with Picard number one, what we can prove is slightly favorable. Note that Q-Fano 3-folds with Picard number one form an important class since the 3-dimensional MMP (see, for instance, [11], [13], [21] and [25]) says that any terminal object (after running MMP) of a given smooth projective variety is either a Mori fiber space (whose fibers are Q-Fano with Picard number one) or a minimal 3-fold. Furthermore, many important works relating to the classification of non-Gorenstein Q-Fano 3-folds with Picard number one have been achieved so far (see [1], [20], [27], [28], [31] and others for a sample of references). Here we would especially like to mention a conjecture of Reid: a general member  $D \in |-K_X|$  is a K3 surface for a "general" Q-Fano 3-fold of Picard number one. The conjecture in Fano case was proved by Reid and Shokurov [26], although it is still open in general.

In order to make a concise statement of our result, we say that a  $\mathbb{Q}$ -Fano 3-fold X is *standard* if

- the Picard number  $\rho(X) = 1$ ;
- ·  $g(X) \ge 0$ , a general member  $D \in |-K_X|$  is irreducible and reduced, and a smooth birational model S of D has non-negative Kodaira dimension, i.e.  $\kappa(S) \ge 0$ .

Roughly speaking, if Reid's conjecture holds true, then a "general" terminal Q-Fano 3-fold with Picard number one is standard.

Here is our another result:

**Theorem 1.3.** Let X be a standard Q-Fano 3-fold. Then  $\varphi_{-m}$  is birational for all  $m \geq 6$ .

As mentioned earlier, the linear system  $|-mK_X|$  shows a lot of pathological properties for Q-Fano 3-folds. To overcome the obvious difficulties, we will extend the Q-divisor method that was used to treat 3-folds of general type. Thanks to the self-optimization function of our mechanic and to the boundedness theorem for Q-Fano 3-folds, we have managed to prove the effective birationality of  $\varphi_{-m}$ . Unfortunately, this method is less effective for those Q-Fano 3-folds on which, for some integer l > 0,  $|-lK_X|$  is composed of birationally ruled surfaces, though that kind of "bad" Q-Fano 3-folds can be regarded as "minority".

This paper is organized as follows. Section 2 mainly sets up the rational map  $\varphi_{-m_0}$  and then reduces the problem to that on a nonsingular model. In Section 3, we prove our key theorem on the birationality of  $\varphi_{-m}$ . Various concrete applications will be presented in Section 4. In Appendix, we prove  $P_{-m} > 0$  for all  $m \ge 6$  on any Q-Fano 3-folds, which is applied to prove our main theorems.

Sometimes, we use the notation  $\mathcal{O}_X(mK_X)$  to denote the reflexive sheaf  $\omega_X^{[m]}$ . The symbol ~ always means linear equivalence while  $\equiv$  represents numerical equivalence.

I would like to thank Jungkai A. Chen, Hélène Esnault, Christopher D. Hacon, Yujiro Kawamata, Janos Kollár and De-Qi Zhang either for stimulating discussions or for their generous helps.

# 2. Preliminaries and the main reduction

**2.1. Convention.** Let X be a terminal weak Q-Fano 3-fold. Pick a canonical divisor  $K_X$  on X. Denote by  $r_X$  the Cartier index of X, which is nothing but the minimal positive integer with  $r_X K_X$  being Cartier. For any positive integer m, the number  $P_{-m} := h^0(X, \mathcal{O}_X(-mK_X))$  is called the *anti-plurigenus* of X. By convention,  $g := h^0(X, \mathcal{O}_X(-K_X)) - 2$  is referred to as the *genus* of X. Clearly, since  $-K_X$  is nef and big, Kawamata's vanishing theorem [11, Theorem 1-2-5] implies

$$h^i(\mathcal{O}_X) = h^i(X, K_X - K_X) = 0$$

for all i > 0. Thus  $\chi(\mathcal{O}_X) = 1$ . Furthermore, the positive rational number  $-K_X^3$  is called the *anti-canonical volume*.

By [3, Theorem 1.1 (ii)], one has  $P_{-8} \ge 2$ . Thus, for a given weak  $\mathbb{Q}$ -Fano 3-fold X, there exists a positive integer  $m_0 \le 8$  such that  $P_{-m_0} \ge 2$ . We shall begin from studying the geometry induced by the rational map  $\varphi_{-m_0}$ .

**2.2.** Set up for  $\varphi_{-m_0}$ . We study the  $m_0$ -th anti-canonical map of X:

$$X \xrightarrow{\varphi_{-m_0}} \mathbb{P}^{P_{-m_0}-1}$$

which is not necessarily well-defined everywhere. First of all we fix an effective Weil divisor  $\hat{K}_0 \sim -m_0 K_X$ . By Hironaka's big theorem, we can take successive blow-ups  $\pi : Y \to X$  such that:

- (i) Y is nonsingular projective;
- (ii) the movable part  $|M_{-m_0}|$  of the linear system  $|\lfloor \pi^*(-m_0K_X) \rfloor|$ is base point free and, consequently, the rational map  $\gamma := \varphi_{-m_0} \circ \pi$  is a morphism;
- (iii) the support of the union of  $K_0$  and the exceptional divisors of  $\pi$  is of simple normal crossings (to secure the usage of vanishing theorems).

Let  $Y \xrightarrow{f} \Gamma \xrightarrow{s} Z$  be the Stein factorization of  $\gamma$  with  $Z := \gamma(Y) \subset \mathbb{P}^{P_{-m_0}-1}$ . In summary, we have the following commutative diagram:



Recall that

$$\pi^*(K_X) := K_Y - \frac{1}{r_X} E_\pi$$

with  $E_{\pi}$  effective and Q-Cartier since X has at worst terminal singularities. Equivalently,  $-K_Y = \pi^*(-K_X) - \frac{1}{r_X}E_{\pi}$ .

**Case**  $(f_{np})$ . If dim $(\Gamma) \geq 2$ , a general member S of  $|M_{-m_0}|$  is a nonsingular projective surface by Bertini's theorem. We say that  $|-m_0K_X|$  is not composed with a pencil of surfaces.

**Case**  $(f_{\mathbf{p}})$ . If dim $(\Gamma) = 1$ , then  $\Gamma \cong \mathbb{P}^1$  since  $g(\Gamma) \leq q(Y) = q(X) := h^1(\mathcal{O}_X) = 0$ . Furthermore, a general fiber S of f is an irreducible smooth projective surface by Bertini's theorem again. We may write

$$M_{-m_0} = \sum_{i=1}^{a_{m_0}} S_i \equiv a_{m_0} S$$

where  $S_i$  is a smooth fiber of f for all i and  $a_{m_0} = P_{-m_0} - 1$ . In this case, we say that  $|-m_0K_X|$  is composed with a (rational) pencil of surfaces.

Define

$$\iota := \begin{cases} 1, & \text{Case } (f_{np}) \\ a_{m_0}, & \text{Case } (f_p). \end{cases}$$

Clearly, at any case,  $M_{-m_0} \equiv \iota S$  with  $\iota \geq 1$ .

**Definition 2.3.** For both Case  $(f_{np})$  and Case  $(f_p)$ , we call S a generic irreducible element of  $|M_{-m_0}|$ .

Lemma 2.4. Keep the same notation as above. The number

$$r_X(\pi^*(-K_X)^2 \cdot S)_Y$$

is a positive integer.

Proof. Note that the number  $(\pi^*(-K_X)^2 \cdot S)$  is positive since  $(\pi^*(-K_X)^2 \cdot S)_Y = (\pi^*(-K_X)|_S)_S^2$  and  $\pi^*(-K_X)|_S$  is nef and big on S. It is also independent of the choice of  $\pi$  according to the projection formula of the intersection theory. So we may choose such a modification  $\pi$  that dominates a resolution of singularities  $\tau : \hat{W} \to X$ . Then we see  $(\pi^*(-K_X)^2 \cdot S)_Y = (\tau^*(-K_X)^2 \cdot S_1)_{\hat{W}}$  where  $S_1 = \delta_*(S)$  is a divisor on  $\hat{W}$  and  $\delta : Y \to \hat{W}$  is a birational morphism. Note, however,  $S_1$  is a generic element in an algebraic family though it is not necessarily nonsingular.

We may write  $K_{\hat{W}} = \tau^*(K_X) + \Delta_{\tau}$  where  $\Delta_{\tau}$  is an exceptional effective  $\mathbb{Q}$ -divisor over those isolated terminal singularities on X. Now, by intersection theory, we have  $(r\tau^*(-K_X)\cdot\tau^*(-K_X)\cdot S_1)_{\hat{W}} = (r\tau^*(-K_X)\cdot (-K_Z)\cdot S_1)_{\hat{W}}$  is an integer.  $\Box$ 

The proof for the next lemma was suggested by C. D. Hacon.

**Lemma 2.5.** Let W be a normal projective variety on which there is an integral Weil  $\mathbb{Q}$ -Cartier divisor D. Let  $h: V \longrightarrow W$  be any resolution of singularities. Assume that E is an effective exceptional  $\mathbb{Q}$ -divisor on V with  $h^*(D) + E$  a Cartier divisor on V. Then

$$h_*\mathcal{O}_V(h^*(D) + E) = \mathcal{O}_W(D)$$

where  $\mathcal{O}_W(D)$  is the reflexive sheaf corresponding to the Weil divisor D.

*Proof.* By [17, Lemma 2.11],  $h_*\mathcal{O}_V(\lfloor h^*(D) \rfloor) = \mathcal{O}_W(D)$ . Consequently, one has:

$$\mathcal{O}_W(D) = h_* \mathcal{O}_V(\lfloor h^*(D) \rfloor) \hookrightarrow h_* \mathcal{O}_V(h^*(D) + E) \xrightarrow{\jmath} \mathcal{O}_W(D)$$

where, for any section  $g \in h_*\mathcal{O}_V(h^*(D) + E)(U)$  over a Zariski open subset  $U \subset W$ , g comes from a rational function  $\tilde{g}$  on  $h^{-1}(U)$  with

$$((\tilde{g}) + (h^*(D) + E))|_{h^{-1}(U)} \ge 0.$$

However, since  $\operatorname{Codim}(W_{\operatorname{Sing}}) \geq 2$  and by the valuation theory, one has  $((g) + D)|_U = ((g) + D)|_{U \cap W_{\operatorname{Sing}}} = ((\tilde{g}) + h^*(D) + E)|_{h^{-1}(U \cap W_{\operatorname{Sing}})} \geq 0.$ Thus g can be viewed as a section j(g) in  $\mathcal{O}_W(D)(U)$  and j is defined in this way.

Since j is identical over  $W^0 := W \setminus W_{\text{Sing}}$  and  $h_* \mathcal{O}_V(h^*(D) + E)$  is torsion free, j is an inclusion. On the other hand, pick a rational function  $\xi$  on an open set U of X such that  $(\xi) + D \ge 0$ . Then  $(\xi \circ h) + h^*(D) \ge 0$  and so  $(\xi \circ h) + h^*(D) + E \ge 0$ . So  $\xi$  belongs to  $h_*\mathcal{O}_V(h^*D + E)(U)$  and j is surjective. Hence j is the identity.  $\Box$ 

**2.6. The main reduction.** Let  $\pi : Y \longrightarrow X$  be as in 2.2. Let m > 0 be an integer. One has:

$$K_{Y} + \lceil (m+1)\pi^{*}(-K_{X}) \rceil$$
  
=  $\pi^{*}(K_{X}) + \frac{1}{r}E_{\pi} + \pi^{*}(-(m+1)K_{X}) + E_{m+1}$   
=  $\pi^{*}(-mK_{X}) + (\frac{1}{r}E_{\pi} + E_{m+1})$ 

where  $E_{m+1}$  is an effective  $\mathbb{Q}$ -divisor on Y. Lemma 2.5 implies

$$\pi_*\mathcal{O}_Y(K_Y + \lceil (m+1)\pi^*(-K_X) \rceil) = \mathcal{O}_X(-mK_X).$$

Hence  $\varphi_{-m}$  is birational if and only if  $\Phi_{|K_Y+\lceil (m+1)\pi^*(-K_X)\rceil|}$  is birational. Noting that

$$\begin{array}{rcl} H^0(\mathcal{O}_X(-mK_X)) &\cong& H^0(\mathcal{O}_Y(\lfloor -m\pi^*(K_X) \rfloor) \\ &\cong& H^0(\mathcal{O}_Y(K_Y + \lceil (m+1)\pi^*(-K_X) \rceil) ) \end{array}$$

we denote by  $|M_{-m}|$  the movable part of  $|\lfloor -m\pi^*(K_X)\rfloor|$ . Clearly, one has the equality:

$$-m\pi^*(K_X) = M_{-m} + F_m \tag{2.1}$$

where  $F_m$  is an effective  $\mathbb{Q}$ -divisor.

Another direct consequence is that we may write:

$$K_Y + \lceil (m+1)\pi^*(-K_X) \rceil \sim M_{-m} + N_{-m}$$
 (2.2)

where  $N_{-m}$  is the fixed part.

**Corollary 2.7.** Let X be a terminal weak  $\mathbb{Q}$ -Fano 3-fold. If

 $P_{-m_0} > m_0 r_X (-K_X)^3 + 1$ 

for some integer  $m_0$ , then  $|-m_0K_X|$  is not composed with a pencil.

Proof. Assume  $|-m_0K_X|$  is composed with a pencil. Keep the same notation as in 2.2. Then we have  $m_0\pi^*(-K_X) \ge M_{-m_0} \equiv a_{m_0}S$  with  $a_{m_0} = P_{-m_0} - 1$ . Thus one has  $m_0(-K_X)^3 \ge (P_{-m_0} - 1)(\pi^*(-K_X)^2 \cdot S) \ge \frac{1}{r_X}(P_{-m_0} - 1)$  by Lemma 2.4, a contradiction. We are done.  $\Box$ 

# 3. The key theorem

Let X be a terminal weak Q-Fano 3-fold on which  $P_{-m_0} \geq 2$  for some integer  $m_0 > 0$ . Keep the same notation as in 2.2. Pick a generic irreducible element S of  $|M_{-m_0}|$ . Suppose we have already a base point free linear system |G| on S.

**3.1. Notations.** Denote by *C* a generic irreducible element of |G|. Since  $\pi^*(-K_X)|_S$  is nef and big, there definitely exists a rational number  $\varrho > 0$  such that

$$\pi^*(-K_X)|_S - \varrho C$$

is numerically equivalent to en effective  $\mathbb{Q}$ -divisor on S. Define

$$\zeta := (\pi^*(-K_X) \cdot C)_Y = (\pi^*(-K_X)|_S \cdot C)_S$$

which will be the key quantity accounting for the birationality of  $\varphi_{-m}$ . Clearly, since  $r_X K_X$  is a Cartier divisor, one has

$$\zeta \ge \frac{1}{r_X}.\tag{3.1}$$

We define two more quantities as follows:

$$\varepsilon := (m + 1 - \frac{m_0}{\iota} - \frac{1}{\varrho})\zeta;$$
  
$$\varepsilon_0 := [\varepsilon].$$

**Theorem 3.2.** Let X be a terminal weak Q-Fano 3-fold with  $P_{-m_0} \ge 2$ for some positive integer  $m_0$ . Assume that |G| is a base point free linear system on a generic irreducible element S of  $|M_{-m_0}|$ . Let m > 0 be an integer. Keep the same notation as above. Then the inequality:

$$\zeta \ge \frac{2g(C) - 2 + \varepsilon_0}{m}$$

holds under one of the following conditions:

(i) g(C) > 0 and  $\varepsilon > 1$ ; (ii) g(C) = 0 and  $\varepsilon > 2$ .

*Proof.* Equality (2.1) gives:

$$m_0 \pi^*(-K_X) = M_{-m_0} + F_{m_0}$$

for some effective  $\mathbb{Q}$ -divisor  $F_{m_0}$ . For the given integer m > 0, one has:

$$|K_Y + \lceil (m+1)\pi^*(-K_X) - \frac{1}{\iota}F_{m_0}\rceil| \subset |K_Y + \lceil (m+1)\pi^*(-K_X)\rceil|.$$
(3.2)

Under the assumption  $\varepsilon > 0$ , the Q-divisor

$$(m+1)\pi^*(-K_X) - \frac{1}{\iota}F_{m_0} - S \equiv (m+1 - \frac{m_0}{\iota})\pi^*(-K_X)$$

is nef and big and thus  $H^1(Y, K_Y + \lceil (m+1)\pi^*(-K_X) - \frac{1}{\iota}F_{m_0}\rceil - S) = 0$ by Kawamata-Viehweg vanishing theorem [10, 30]. Hence one has the surjective map:

$$H^{0}(Y, K_{Y} + \lceil (m+1)\pi^{*}(-K_{X}) - \frac{1}{\iota}F_{m_{0}}\rceil) \longrightarrow H^{0}(S, K_{S} + L_{m}) \quad (3.3)$$

where

$$L_m := \left( \left\lceil (m+1)\pi^*(-K_X) - \frac{1}{\iota}F_{m_0} \right\rceil - S \right) |_S \ge \left\lceil \mathcal{L}_m \right\rceil$$

and  $\mathcal{L}_m := ((m+1)\pi^*(-K_X) - \frac{1}{\iota}F_{m_0} - S)|_S$ . On the other hand, we have

$$\pi^*(-K_X)|_S \equiv \varrho C + H$$

for an effective  $\mathbb{Q}$ -divisor H on S. Thus the  $\mathbb{Q}$ -divisor

$$\mathcal{L}_m - \frac{1}{\varrho}H - C \equiv \varepsilon \pi^*(-K_X)|_S$$

is nef and big and, by the vanishing theorem again,

$$H^{1}(S, K_{S} + \lceil \mathcal{L}_{m} - \frac{1}{\varrho}H \rceil - C) = 0.$$

Therefore, one has the following surjective map:

$$H^{0}(S, K_{S} + \lceil \mathcal{L}_{m} - \frac{1}{\varrho} H \rceil) \longrightarrow H^{0}(C, K_{C} + D_{m})$$
(3.4)

where

$$D_m := \left\lceil \mathcal{L}_m - \frac{1}{\varrho} H - C \right\rceil |_C \ge \left\lceil \mathcal{D}_m \right\rceil$$

and  $\mathcal{D}_m := (\mathcal{L}_m - \frac{1}{\varrho}H - C)|_C$  with deg $\lceil \mathcal{D}_m \rceil \ge \varepsilon_0$ .

Now under the situations (i) and (ii),  $|K_C + \lceil \mathcal{D}_m \rceil|$  is base point free with

 $\deg(K_C + \lceil \mathcal{D}_m \rceil) \ge 2g(C) - 2 + \varepsilon_0.$ 

Denote by  $\mathcal{N}_m$  the movable part of  $|K_S + \lceil \mathcal{L}_m - \frac{1}{\varrho}H\rceil|$ . Noticing the relations (3.2), (3.3) and  $|K_S + \lceil \mathcal{L}_m - \frac{1}{\varrho}H\rceil| \subset |K_S + \lceil \mathcal{L}_m\rceil|$  while applying [4, Lemma 2.7], one gets:

$$m\pi^*(-K_X)|_S \ge M_{-m_0}|_S \ge \mathcal{N}_m$$

and  $\mathcal{N}_m|_C = K_C + \lceil \mathcal{D}_m \rceil$  simply because the later one is base point free under either of the conditions.

So one has  $m\zeta \ge \deg(K_C + \lceil \mathcal{D}_m \rceil) \ge 2g(C) - 2 + \varepsilon_0$ . We are done.  $\Box$ 

Riemann-Roch theorem on C directly implies the following:

**Corollary 3.3.** Under the same assumption as that of Theorem 3.2, then one has:

- (1)  $P_{-m} > 0$  whenever g(C) = 0 and  $\varepsilon > 1$ ;
- (2)  $P_{-m} > 0$  whenever g(C) > 0 and  $\varepsilon > 0$ .

From now on, we shall work on the birationality. We shall always require that the linear system  $\Lambda_m := |K_Y + \lceil (m+1)\pi^*(-K_X)\rceil|$  satisfies the following assumption for some integer m > 0.

### **3.4.** Assumption:

- (1) The linear system  $\Lambda_m$  distinguishes different generic irreducible elements of  $|M_{-m_0}|$  (namely,  $\Phi_{\Lambda_m}(S') \neq \Phi_{\Lambda_m}(S'')$  for two different generic irreducible elements S', S'' of  $|M_{-m_0}|$ ).
- (2) The linear system  $\Lambda_{m|S}$  distinguishes different generic irreducible elements of the given linear system |G| on S.

Here is our key theorem:

**Theorem 3.5.** Let X be a terminal weak Q-Fano 3-fold with  $P_{-m_0} \ge 2$ for some positive integer  $m_0$ . Assume that |G| is a base point free linear system on a generic irreducible element S of  $|M_{-m_0}|$ . Let m > 0 be an integer. If Assumption 3.4 is satisfied and  $\varepsilon > 2$ , then  $\varphi_{-m}$  is birational onto its image.

*Proof.* Since Assumption 3.4 (1) is satisfied, the usual birationality principle reduces the birationality of  $\varphi_{-m}$  to that of  $\Phi_{\Lambda_m}|_S$  for a generic irreducible element S. Similarly, due to Assumption 3.4 (2), one only needs to prove the birationality of  $\Phi_{\Lambda_m}|_C$  for a generic irreducible element C of |G|.

Now since one has the surjective maps (3.3) and (3.4), it is sufficient to prove that  $|K_C + \lceil \mathcal{D}_m \rceil|$  gives a birational map. Clearly this is the case whenever  $\varepsilon > 2$ , which in fact implies deg( $\lceil \mathcal{D}_m \rceil$ )  $\geq 3$ . We are done.

While applying Theorem 3.5, one needs to verify Assumption 3.4 in advance. Hereby we are working on this.

**Proposition 3.6.** Let X be a terminal weak  $\mathbb{Q}$ -Fano 3-fold with  $P_{-m_0} \geq 2$  for some positive integer  $m_0$ . Then Assumption 3.4 (1) is satisfied for all

$$m \ge \begin{cases} m_0 + 6, & m_0 \ge 2\\ 2, & m_0 = 1 \end{cases}$$

*Proof.* By Corollary 5.1 in Appendix, one has

$$K_{Y} + \lceil (m+1)\pi^{*}(-K_{X}) \rceil$$
  

$$\geq K_{Y} + \lceil (m-m_{0}+1)\pi^{*}(-K_{X}) + M_{-m_{0}} \rceil$$
  

$$= (K_{Y} + \lceil (m-m_{0}+1)\pi^{*}(-K_{X}) \rceil) + M_{-m_{0}}$$
  

$$\geq M_{-(m-m_{0})} + M_{-m_{0}} \geq M_{-m_{0}}$$

for all  $m - m_0 \geq 6$  whenever  $m_0 \geq 2$  (resp.  $\geq 1$  whenever  $m_0 = 1$ ). When  $f: Y \to \Gamma$  is of type  $(f_{np})$ , [29, Lemma 2] implies that  $\Lambda_m$  can distinguish different generic irreducible elements of  $|M_{-m_0}|$ . When f is of type  $(f_p)$ , since the rational (i.e.  $\Gamma \cong \mathbb{P}^1$ ) pencil  $|M_{-m_0}|$  can already separate different fibers of f,  $\Lambda_m$  can naturally distinguish different generic irreducible elements of  $|M_{-m_0}|$ .

It is slightly more complicated to verify Assumption 3.4 (2).

**Lemma 3.7.** Let T be a nonsingular projective surface on which there is a base point free linear system |G|. Let Q be an arbitrary Q-divisor on T. Then the linear system  $|K_T + \lceil Q \rceil + G|$  can distinguish different generic irreducible elements of |G| under one of the following conditions:

(i)  $K_T + \lceil Q \rceil$  is effective and |G| is not composed with an irrational pencil of curves;

- (ii) Q is nef and big,  $g(C) \ge 0$  and |G| is composed with an irrational pencil of curves;
- (iii) Q is nef and big,  $Q \cdot G > 1$ , g(C) = 0 and |G| is composed with an irrational pencil of curves.

*Proof.* The statement corresponding to (i) follows from [29, Lemma 2] and the fact that a rational pencil can automatically separate its different generic irreducible elements.

For situations (ii) and (iii), we pick a generic irreducible element C of |G|. Then, since  $h^0(S,G) \ge 2$ ,  $G \equiv sC$  for some integer  $s \ge 2$  and  $C^2 = 0$ . Denote by C' another generic irreducible element of |G|. The Kawamata-Viehweg vanishing theorem gives the surjective map:

$$H^0(T, K_T + \lceil Q \rceil + G) \longrightarrow H^0(C, K_C + D) \oplus H^0(C', K_{C'} + D')$$

where  $D := (\lceil Q \rceil + G - C)|_C$  and  $D' := (\lceil Q \rceil + G - C')|_{C'}$  with  $\deg(D) \ge Q \cdot C > 0$ ,  $\deg(D') \ge Q \cdot C' > 0$ .

Whenever  $g(C) \ge 0$ , Riemann-Roch formula gives  $h^0(C, K_C + D) = h^0(C', K_{C'} + D') > 0$ . Thus  $|K_T + \lceil Q \rceil + G|$  can distinguish C and C'. Whenever g(C) = 0 and  $Q \cdot C > 1$ ,  $H^0(C, K_C + D) \ne 0$  and  $H^0(C', K_{C'} + D') \ne 0$ . So  $|K_T + \lceil Q \rceil + G|$  can also distinguish C and C'.

**Proposition 3.8.** Let X be a terminal weak Q-Fano 3-fold with  $P_{-m_0} \geq 2$  for some positive integer  $m_0$ . Keep the same notation as in 2.2. Take  $|G| = |M_{-m_0|S}|$  on a generic irreducible element S of  $|M_{-m_0}|$ . If  $f: Y \to Z$  is of type  $(f_{np})$ , then Assumption 3.4 (2) is satisfied under one of the following conditions:

(1) 
$$g(C) > 0$$
 and  $m \ge \begin{cases} \max\{m_0 + 6, 2m_0\} & m_0 \ge 2\\ 2, & m_0 = 1; \end{cases}$   
(2)  $g(C) = 0, \ \varepsilon > 1 \ and \ m \ge \begin{cases} m_0 + 6, & m_0 \ge 2\\ 2, & m_0 = 1. \end{cases}$ 

Proof. By the relation (3.2) and the surjective map (3.3), we see  $\Lambda_m|_S \supset |K_S + L_m|$ , which will be proved to be able to distinguish different generic irreducible elements of |G|. By definition, we have  $\iota = 1$ . Since  $m_0 \pi^*(-K_X) \ge M_{-m_0}$ , we may take  $\varrho = \frac{1}{m_0}$ .

Now one has the following:

$$K_{S} + L_{m} = (K_{Y})|_{S} + \lceil (m+1)\pi^{*}(-K_{X}) - F_{m_{0}} \rceil|_{S}$$
  

$$\geq (K_{Y} + \lceil (m-m_{0}+1)\pi^{*}(-K_{X}) \rceil)|_{S} + M_{-m_{0}}|_{S}.$$

Thus, if |G| is not composed with an irrational pencil of curves,  $|K_S + L_m|$  can distinguish different generic irreducible elements provided that  $m - m_0 \ge 6$  whenever  $m_0 \ge 2$  (or that  $m \ge 2$  whenever  $m_0 = 1$ ).

Suppose |G| is composed with an irrational pencil of curves. One has:

$$\begin{array}{rcl} K_{S} + L_{m} & \geq & K_{S} + \lceil \mathcal{L}_{m} \rceil \\ & = & K_{S} + \lceil ((m - m_{0} + 1)\pi^{*}(-K_{X}) - F_{m_{0}}) |_{S} \rceil \\ & \geq & K_{S} + \lceil (m - 2m_{0} + 1)\pi^{*}(-K_{X}) |_{S} \rceil + M_{-m_{0}} |_{S}. \end{array}$$

If g(C) > 0, Lemma 3.7 (ii) implies that Assumption 3.4 (2) is satisfied for  $m \ge 2m_0$ . If g(C) = 0, by Lemma 3.7 (iii), we need the condition  $\varepsilon = (m - 2m_0 + 1)\zeta = Q \cdot C > 1$ , where we may take  $Q = (m - 2m_0 + 1)\pi^*(-K_X)|_S$ .

When f is of type  $(f_p)$ , we have the similar treatment.

**Proposition 3.9.** Let X be a terminal weak Q-Fano 3-fold. Assume that  $P_{-m_0} \geq 2$  for some positive integer  $m_0$  and that  $|-m_1K_X|$  is not composed with a pencil of surfaces for another integer  $m_1 \geq m_0$ . Keep the same notation as in 2.2. Take  $|G| = |M_{-m_1|S}|$  on a generic irreducible element S of  $|M_{-m_0}|$ . Let C be a generic irreducible element of |G|. If  $f: Y \to Z$  is of type  $(f_p)$ , then Assumption 3.4 (2) is satisfied under one of the following conditions:

(1) 
$$g(C) > 0, \ m \ge \begin{cases} \max\{m_1 + 6, m_0 + m_1\} & m_0 \ge 2\\ m_1 + 1, & m_0 = 1; \end{cases}$$
  
(2)  $g(C) = 0, \ \varepsilon > 1 \ and \ m \ge \begin{cases} m_1 + 6, & m_0 \ge 2\\ m_1 + 1, & m_0 = 1. \end{cases}$ 

Proof. First, we may re-modify our  $\pi$  in 2.2 such that the movable part  $|M_{-m_1}|$  of  $|\lfloor \pi^*(-m_1K_X) \rfloor|$  is base point free too. Since  $m_1\pi^*(-K_X) \ge M_{-m_1}$ , we may take  $\varrho = \frac{1}{m_1}$ . Similar to the proof of Proposition 3.8, it suffices to prove that  $|K_S + L_m|$  can distinguish different generic irreducible elements of |G|.

For a suitable integer m > 0, one has the following:

$$K_S + L_m = (K_Y)|_S + \lceil (m+1)\pi^*(-K_X) - F_{m_0}\rceil|_S \geq (K_Y + \lceil (m-m_1+1)\pi^*(-K_X)\rceil)|_S + M_{-m_1}|_S.$$

Thus, if |G| is not composed with an irrational pencil of curves,  $|K_S + L_m|$  can distinguish different irreducible elements provided that  $K_Y + \lceil (m - m_1 + 1)\pi^*(-K_X) \rceil$  is effective, i.e.  $m - m_1 \ge 6$  whenever  $m_0 \ge 2$  (or  $m \ge m_1 + 1$  whenever  $m_0 = 1$ ).

Assume |G| is composed with an irrational pencil of curves. One has:

$$\begin{array}{rcl}
K_{S} + L_{m} &\geq & K_{S} + \left\lceil \mathcal{L}_{m} \right\rceil \\
&= & K_{S} + \left\lceil ((m - m_{0} + 1)\pi^{*}(-K_{X}) - F_{m_{0}}) |_{S} \right\rceil \\
&\geq & K_{S} + \left\lceil (m - m_{0} - m_{1} + 1)\pi^{*}(-K_{X}) |_{S} \right\rceil + M_{-m_{1}} |_{S}.
\end{array}$$

If g(C) > 0, Lemma 3.7(ii) implies that Assumption 3.4 (2) is satisfied for  $m \ge m_0 + m_1$ . If g(C) = 0, by Lemma 3.7(iii), we need the

condition  $\varepsilon = (m - m_0 - m_1 + 1)\zeta = Q \cdot C > 1$ , where we may take  $Q = (m - m_0 - m_1 + 1)\pi^*(-K_X)|_S$ . We are done.

# 4. Applications

In this section, we shall present some concrete applications on the birationality of  $\varphi_{-m}$  using Theorem 3.5. In practice, we take the smallest integer  $\hat{m}_0 := \hat{m}_0(X)$  with  $P_{-\hat{m}_0} \ge 2$ . Such a number  $\hat{m}_0$  is uniquely determined by X and  $\hat{m}_0 \le 8$  by [3, Theorem 1.1 (ii)].

**Theorem 4.1.** Let X be a terminal weak Q-Fano 3-fold. Assume  $\dim \overline{\varphi_{-\hat{m}_0}(X)} = 3$ . Then  $\varphi_{-m}$  is birational onto its image for all  $m \ge 4\hat{m}_0$ . In particular,  $|-mK_X|$  gives a birational map for  $m \ge 32$ .

*Proof.* Keep the same notation as in 2.2. We have  $\iota = 1$ . Take  $|G| = |M_{-\hat{m}_0|S}|$  on a generic irreducible element S of  $|M_{-\hat{m}_0}|$ . Clearly, we may take  $\varrho = \frac{1}{\hat{m}_0}$ . Since, by assumption,  $\Phi_{|M_{-\hat{m}_0|S}|}$  maps a generic irreducible element C of |G| onto a curve, Riemann-Roch formula and Clifford's theorem on C gives

$$\hat{m}_0 \pi^*(-K_X) \cdot C \ge M_{-\hat{m}_0|S} \cdot C \ge \begin{cases} 2, & g(C) > 0\\ 1, & g(C) = 0. \end{cases}$$
(4.1)

We prove the theorem according to the value of g(C).

Assume g(C) > 0. We have seen  $\zeta \geq \frac{2}{\hat{m}_0}$  by inequality (4.1). By Proposition 3.6 and Proposition 3.8, Assumption 3.4 is satisfied for

$$m \ge \begin{cases} \max\{\hat{m}_0 + 6, 2\hat{m}_0\} & \hat{m}_0 \ge 2\\ 2, & \hat{m}_0 = 1. \end{cases}$$

Take any  $m \ge 3\hat{m}_0$ . Then  $\varepsilon \ge 2 + \frac{2}{\hat{m}_0} > 2$ . Thus Theorem 3.5 implies that  $\varphi_{-m}$  is birational for all  $m \ge 3\hat{m}_0$  and  $\hat{m}_0 \ne 2$ . For  $\hat{m}_0 = 2$ , we need to set  $m \ge 8$  to satisfy  $m \ge \hat{m}_0 + 6$ .

Assume g(C) = 0. We have already seen  $\zeta \geq \frac{1}{\hat{m}_0}$  by inequality (4.1). Take any  $m \geq 4\hat{m}_0$ . Then  $\varepsilon > 2$ . Again Propositions 3.6, 3.8 imply that  $\varphi_{-m}$  is birational for all  $m \geq 4\hat{m}_0$ .

**Theorem 4.2.** Let X be a terminal weak Q-Fano 3-fold with  $r_X > 1$ . Assume dim  $\overline{\varphi_{-\hat{m}_0}(X)} = 2$ . Denote by C a generic irreducible element of  $|G| := |M_{-\hat{m}_0|S}|$ . Then  $\varphi_{-m}$  is birational onto its image under one of the following conditions:

- (1)  $g(C) > 0, m \ge 6\hat{m}_0;$
- (2)  $g(C) = 0, \ m \ge 2\hat{m}_0 + 2r_X.$

*Proof.* Again we have  $\iota = 1$  and may take  $\varrho = \frac{1}{\hat{m}_0}$ . Clearly |G| is composed with a pencil of curves.

(1) Assume g(C) > 0. If we take a sufficiently large  $m = m' > 4\hat{m}_0$ such that  $\varepsilon = (m' + 1 - 2\hat{m}_0)\zeta > 1$  (thus  $\varepsilon_0 \ge 2$ ), then Theorem 3.3 implies  $\zeta \ge \frac{2}{m'}$ . Take m = m' - 1. Then  $\varepsilon \ge (m' - 2\hat{m}_0)\frac{2}{m'} > 1$ . Theorem 3.3 again gives  $\zeta \geq \frac{2}{m'-1}$ . This means, inductively, that

 $\zeta \geq \frac{2}{4\hat{m}_0} = \frac{1}{2\hat{m}_0}.$ Whenever  $m \geq 6\hat{m}_0$ , then Assumption 3.4 is naturally satisfied by Propositions 3.6 and 3.8. On the other hand,  $\varepsilon \geq (6\hat{m}_0 + 1 - 2\hat{m}_0)\frac{1}{2\hat{m}_0} >$ 2. Theorem 3.5 implies that  $\varphi_m$  is birational.

(2) Assume g(C) = 0. Since we have  $\zeta \geq \frac{1}{r_X}$ , we may take  $m \geq \frac{1}{r_X}$  $2\hat{m}_0 + 2r_X$  and then  $\varepsilon \ge (m+1-2\hat{m}_0)\zeta > 2$ . When  $r_X > 1$ , Theorem 3.5, Propositions 3.6 and 3.8 imply that  $\varphi_m$  is birational for all  $m \geq 1$  $2\hat{m}_0 + 2r_X$ . 

Especially, when  $\hat{m}_0 = 1$ , we have the following:

**Corollary 4.3.** Let X be a standard Q-Fano 3-fold. Then  $\varphi_{-m}$  is birational for all  $m \geq 6$ .

*Proof.* We have  $\hat{m}_0 = 1$ . According to [1, Theorem 2.18],  $|-K_X|$  has no fixed parts and is not composed with a pencil of surfaces. Since Xis standard, S is not fibred by rational curves. Thus we have q(C) > 0. Now the result follows directly from Theorem 4.1 and Theorem 4.2 (1).

From now on within this section, we study the most difficult case:  $\dim \varphi_{-\hat{m}_0}(X) = 1$ . First, we need to look for a number  $m_1 > 0$  such that  $|-m_1K_X|$  is not composed with a pencil of surfaces.

**Proposition 4.4.** Let X be a terminal weak  $\mathbb{Q}$ -Fano 3-fold with  $r_X >$ 1. Then  $|-m_1K_X|$  is not composed with a pencil of surfaces under one of the following situations:

- (1)  $r = 2, m_1 = 4;$ (2)  $r = 3, m_1 = 6;$
- (3)  $r > 4, m_1 = r + 2.$

*Proof.* Let  $\rho: V \to X$  be a resolution of singularities of X. According to Lemma 2.5 and Kawamata-Viehweg Vanishing theorem, for all positive integer m,

$$P_{-m} := P_{-m}(X) = h^0(V, K_V + \lceil (m+1)\rho^*(-K_X) \rceil) \\ = \chi(V, \mathcal{O}_V(K_V + \lceil (m+1)\rho^*(-K_X) \rceil)).$$

Write  $K_V = \rho^*(K_X) + \Delta$  where  $\Delta$  is an effective Q-divisor and is  $\rho$ -exceptional. Thus  $\rho(\Delta)$  is the set of singularities on X. One gets  $-K_V = \rho^*(-K_X) - \Delta$ . For any given m > 0, write m = nr + t with  $n \geq 0, 0 \leq t < r = r_X$ . By Riemann-Roch formula on V ([5, P. 437]) and setting  $D := K_V + [(m+1)\rho^*(-K_X)]$ , one has:

$$P_{-m} = \frac{1}{12}D \cdot (D - K_V) \cdot (2D - K_V) + \frac{1}{12}D \cdot c_2(V) + \chi(\mathcal{O}_V) \\ = \frac{1}{12}m(m+1)(2m+1)(-K_X)^3 + A \cdot n + C(t)$$

where  $A := \frac{r}{12} \rho^*(-K_X) \cdot c_2(V)$  and C(t) is a constant only depending on t. Fix a number  $t < r, Q_t(n) := P_{-(nr+t)}$  can be viewed as a polynomial

in terms of n. Then we have

$$Q_t(n) = \hat{Q}_t(n) + A \cdot n + C(t)$$

where  $\hat{Q}_t(n) = \frac{1}{12}(nr+t)(nr+t+1)(2nr+2t+1)(-K_X)^3$ . For  $n \ge 1$  or n = 0 and  $t \ge 2$ , one has:

$$Q_t(n) \ge 0 \tag{4.2}$$

For  $n \leq -1$ , by [13, Corollary 5.25],  $\mathcal{O}_X((nr+t)K_X) := \omega_X^{[nr+t]}$  is Cohen-Macaulay and thus Serre duality gives:

$$\chi(X, \mathcal{O}_X((nr+t)K_X)) = -\chi(X, \mathcal{H}om(\mathcal{O}_X((nr+t)K_X), \omega_X)))$$
  
=  $-\chi(X, \mathcal{O}_X((1-nr-t)K_X))$ 

where the last equality is due to the fact that  $\mathcal{H}om(\mathcal{O}_X((nr+t)K_X), \omega_X)$  is reflexive (cf. [19, P. 150, Lemma 1.1.12]). Now the vanishing theorem [11, Theorem 1-2-5] gives:

$$Q_t(n) = \chi(\mathcal{O}_X((nr+t)K_X)) = -\chi(\mathcal{O}_X((1-nr-t)K_X)) = -P_{1-nr-t} \le 0$$
(4.3)

Let us estimate the lower bounds of both A and C(t). For any r, t, one has, by inequalities (4.2) and (4.3),

$$A + C(t) \ge -\hat{Q}_t(1) \ge -\frac{1}{12}(r+t)(r+t+1)(2r+2t+1)(-K_X)^3 \quad (4.4)$$

$$A - C(t) \ge \hat{Q}_t(-1) \ge \frac{1}{12}(-r+t)(-r+t+1)(-2r+2t+1)(-K_X)^3 \quad (4.5)$$

which imply:

$$A \ge -\frac{1}{12}(2r^2 + 6t^2 + 6t + 1)r(-K_X)^3.$$
(4.6)

Case 1. Assume  $t \ge 2$ . Clearly, one has  $r \ge 3$ . Then O(0) = P.

Clearly, one has  $r \ge 3$ . Then  $Q_t(0) = P_{-t} \ge 0$  gives:

$$C(t) \ge -\frac{1}{12}t(t+1)(2t+1)(-K_X)^3.$$

Thus (4.5) implies:

$$A \geq \frac{1}{12} \{ -2r^2 + (6t+3)r - (6t^2 + 6t + 1) \} r(-K_X)^3 \\ = \frac{1}{12} \{ -6t^2 + (6r-6)t + (-2r^2 + 3r - 1) \} r(-K_X)^3$$

for all  $2 \leq t < r$ . Noting that A is a constant independent of t, one should have:

$$A \ge \begin{cases} \frac{1}{12}(-\frac{1}{2}r^2 + \frac{1}{2})r(-K_X)^3, & \text{taking } t = \frac{r-1}{2} \text{ when } r \ge 5 \text{ is odd} \\ \frac{1}{12}(-\frac{1}{2}r^2 - 1)r(-K_X)^3, & \text{taking } t = \frac{r}{2} \text{ when } r \ge 4 \text{ is even.} \end{cases}$$

With the above inequalities, we are able to bound  $P_{-m}$  from below. In fact, we have

$$P_{-(r+2)} = Q_2(1) \ge \frac{1}{8}(r^2 + 10r + 24)r(-K_X)^3.$$

Clearly,  $P_{-(r+2)} > (r+2)r(-K_X)^3 + 1$  whenever  $r \ge 4$ . Thus Lemma 2.7 implies that  $|-(r+2)K_X|$  is not composed with a pencil when  $r \ge 4$  and this proves (3).

Case 2. Assume t = 0.

Inequalities (4.4) and (4.6) give:

$$P_{-nr} \ge \frac{1}{12}(n^2 - 1)(2nr + 3)r^2(-K_X)^3.$$

Clearly  $P_{-2r} > 2r \cdot r(-K_X)^3 + 1$  for r = 2, 3. Thus Lemma 2.7 implies that  $|-2rK_X|$  is not composed with a pencil when r = 2 or 3. This proves (1) and (2).

Besides, it is easy to get the lower bound of  $P_{-(nr+t)}$  for a given pair (n, t).

**Theorem 4.5.** Let X be a terminal weak Q-Fano 3-fold with  $r_X > 1$ . Assume dim  $\overline{\varphi_{-\hat{m}_0}(X)} = 1$  and  $|-m_1K_X|$  is not composed with a pencil for another integer  $m_1 > 0$ . Then  $\varphi_{-m}$  is birational onto its image for all  $m \ge \hat{m}_0 + m_1 + 2r_X$ .

*Proof.* By definition, we have  $\iota \geq 1$  and  $m_1 \geq \hat{m}_0$ .

Re-modify our original  $\pi$  in 2.2 such that the movable part  $M_{-m_1}$  of the linear system  $|\lfloor \pi^*(-m_1K_X) \rfloor|$  is base point free too. Pick a generic irreducible element S of  $|M_{-\hat{m}_0}|$  and take  $|G| := |M_{-m_1|S}|$ . Then we have  $\rho = \frac{1}{m_1}$ . Recall that we have seen  $\zeta \geq \frac{1}{r_X}$ .

Assume, for an integer

$$m \ge \begin{cases} \max\{m_1 + 6, \hat{m}_0 + m_1\} & \hat{m}_0 \ge 2\\ m_1 + 1, & \hat{m}_0 = 1, \end{cases}$$

one has already  $\varepsilon > 1$ . Proposition 3.6 and Proposition 3.9 imply that Assumption 3.4 is satisfied. In practice, take any  $m \ge \hat{m}_0 + m_1 + 2r$ , which naturally satisfies the above requests. Furthermore  $\varepsilon \ge (m + 1 - \hat{m}_0 - m_1)\zeta > 2$ . Consequently, Theorem 3.5 implies that  $\varphi_{-m}$  is birational for  $m \ge \hat{m}_0 + m_1 + 2r_X$ .

**Corollary 4.6.** Let X be a terminal weak  $\mathbb{Q}$ -Fano 3-fold with  $r_X > 1$ . Assume dim  $\overline{\varphi_{-\hat{m}_0}(X)} = 1$ . Then  $\varphi_{-m}$  is birational for

$$m \ge \begin{cases} \hat{m}_0 + 8, & r = 2;\\ \hat{m}_0 + 12, & r = 3;\\ \hat{m}_0 + 3r + 2, & r \ge 4. \end{cases}$$

Proof. Set

$$m_1 := \begin{cases} 2r, & \text{if } r = 2, 3; \\ r + 2, & \text{otherwise.} \end{cases}$$

The statement follows directly from Proposition 4.4 and Theorem 4.5.  $\hfill \Box$ 

If one does not request the "stable" birationality, then the slightly better result can be obtained as follows.

**Theorem 4.7.** Let X be a weak Q-Fano 3-fold with  $r_X > 1$ . Then  $\varphi_{-(\hat{m}_0+3r-1)}$  is birational onto its image.

Proof. Take  $m = \hat{m}_0 + 3r - 1$ . When  $r_X \ge 3$ , we have  $P_{-(m-\hat{m}_0)} > 0$  by Corollary 5.1. When  $r_X = 2$ , we have  $P_{-(m-\hat{m}_0)} = P_{-5} > 0$  by Proposition 4.4. Thus the proof of Proposition 3.6 implies that  $|K_Y + \lceil (m+1)\pi^*(-K_X)\rceil|$  can distinguish different irreducible elements of  $|M_{-\hat{m}_0}|$ . Thus we only need to prove the birationality of  $|K_S + \lceil \mathcal{L}_m\rceil|$  for a generic S. Now since

$$\mathcal{L}_m = 3r\pi^*(-K_X)|_S = 3L$$

and  $L := r\pi^*(-K_X)|_S$  is a nef and big Cartier divisor on S with  $L^2 \ge r \cdot r(\pi^*(-K_X)^2 \cdot S) \ge 2$  (see Lemma 2.4), Reider's theorem ([23]) says that  $|K_S + 3L|$  gives a birational map. We are done.

# 5. Appendix: Anti-plurigenus

Here we use all those formulae and inequalities and keep the same notation as in [3, Section 2]. The following result was used to prove our main results in this paper.

**Corollary 5.1.** Let X be a weak  $\mathbb{Q}$ -Fano 3-fold. Then  $P_{-m} > 0$  for all  $m \ge 6$ .

Proof. Denote by B := B(X), the virtual basket of singularities of Reid [22]. We shall study the formal basket  $\mathbf{B} := \{B, P_{-1}\}$  which was defined in [3, Section 2]. Then we know, from [3, 2.3], that  $-K^3(\mathbf{B}) = -K_X^3 > 0$  and  $P_{-m}(\mathbf{B}) = P_{-m}(X)$  for all  $m \ge 1$ .

**Case 1**. Assume  $P_{-2} > 0$ .

If  $P_{-5} > 0$ , then the statement is naturally true. So we only need to study the situation  $P_{-5} = 0$ . It then follows that  $P_{-1} = P_{-3} = 0$ .

Now  $\epsilon_5 \ge 0$  gives  $2 + P_{-2} \ge 2P_{-4} + \sigma_5$ . Since, by inequality [3, (2.3)],  $P_{-4} \ge 2P_{-2} - 1$ , we see  $P_{-2} = P_{-4} = 1$  and  $\sigma_5 \le 1$ .

First, consider the case  $\sigma_5 = 0$ . Then  $\epsilon_6 = 1 \neq 0$ , a contradiction. Thus we must have  $\sigma_5 = 1$ . Then we have  $\epsilon_5 = 0$  and

$$B^{(0)} = B^{(5)} = \{9 \times (1,2), (1,3), (1,r)\}, r \ge 5.$$

Clearly, since  $B^{(5)}$  admits no further prime packings, we have  $B = B^{(5)} = B^{(0)}$ . Now inequality [3, (2.1)] implies  $r \ge 7$ , while inequality [3, (2.2)] gives  $r \le 7$ . We see r = 7. Now our calculation shows:  $-K^3 = \frac{1}{42}$ ,  $P_{-6} = P_{-7} = 2$ ,  $P_{-8} = P_{-9} = 4$ ,  $P_{-10} = P_{-11} = 6$  and  $P_{-12} = 9$ . Clearly we can see  $P_{-m} > 0$  for all  $m \ge 6$ .

**Case 2**. Assume  $P_{-2} = 0$ .

According to our complete classification in [3, Theorem 3.5, Table A], there are exactly 23 cases as listed there. By the direct calculation,

we know  $P_{-m} > 0$  for all  $6 \le m \le 12$ . Thus it is easy to deduce  $P_{-m} > 0$  for all  $m \ge 6$ . We are done.

### References

- V. Alexeev, General elephants of Q-Fano 3-folds, Compositio Math. 91 (1994), 91-116.
- [2] T. Ando, Pluricanonical systems of algebraic varieties of general type of dimension ≤ 5. Algebraic geometry, Sendai, 1985, 1-10, Adv. Stud. Pure Math., 10, North-Holland, Amsterdam, 1987.
- [3] J. A. Chen, M. Chen, An optimal boundedness on weak Q-Fano 3-folds, Advances in Math. 219 (2008), 2086-2104.
- [4] M. Chen, Canonical stability in terms of singularity index for algebraic threefolds, Math. Proc. Camb. Phil. Soc. 13 (2001), 241-264.
- [5] R. Hartshorne, Algebraic Geometry, Graduate Texts in Mathematics 52, Springer-Verlag, New York, Berlin, Heidelberg, 1977.
- [6] V. A. Iskovskih, *Fano 3-folds, I*, Izv. Akad. Nauk SSSR Ser. Mat **41** (1977).
   English translation in Math. USSR izv. **11** (1977), 485-527.
- [7] V. A. Iskovskih, *Fano 3-folds, II*, Izv. Akad. Nauk SSSR Ser. Mat **42** (1978), English translation in Math. USSR izv. **12** (1978), 469-506.
- [8] V. A. Iskovskikh, Yu. G. Prokhorov, *Fano varieties*. Algebraic geometry, V, 1-247, Encyclopaedia Math. Sci., 47, Springer, Berlin, 1999.
- [9] Y. Kawamata, Boundedness of Q-Fano 3-folds. Proceedings of the International Conference on Algebra, Part 3 (Novosibirsk, 1989), 439–445, Contemp. Math., 131, Part 3, Amer. Math. Soc., Providence, RI, 1992.
- [10] Y. Kawamata, A generalization of Kodaira-Ramanujam's vanishing theorem, Math. Ann. 261(1982), 43-46.
- [11] Y. Kawamata, K. Matsuda, K. Matsuki, Introduction to the Minimal Model Problem, Advanced Studies in Pure Mathematics 10, 1987. Algebraic Geometry, Sendai, 1985, pp. 283-360.
- [12] J. Kollár, Y. Miyaoka, S. Mori, H. Takagi, Boundedness of canonical Q-Fano 3-folds, Proc. Japan Acad. 76, Ser. A (2000), 73-77.
- [13] J. Kollár, S. Mori, Birational geometry of algebraic varieties, 1998, Cambridge Univ. Press.
- [14] S. Mori, S. Mukai, Classification of Fano 3-folds with  $b_2 \ge 2$ , Algebraic and Topological Theories (Kinosaki, 1984), Kinokuniya, Tokyo, 1985, pp. 496-545.
- [15] S. Mukai, Curve and symmetric spaces, I, Amer. J. Math. 117 (1995), 1627-1644.
- [16] S. Mukai, New development of the theory of Fano threefolds: vector bundle method and moduli problem, Suguku 47 (1995), 125-144, English translation in Sugaku Expositions 15 (2002), 125-150.
- [17] N. Nakayama, Zariski-decomposition and abundance. MSJ Memoirs, 14. Mathematical Society of Japan, Tokyo, 2004. xiv+277 pp.
- [18] Y. Namikawa, Smoothing Fano 3-folds. J. Algebraic Geom. 6 (1997), no. 2, 307-324.
- [19] C. Okonek, M. Schneider, H. Spindler, Vector bundles on complex projective spaces, Progress in Math., 3, Birkhauser, Basel, 1978
- [20] Yu. G. Prokhorov, The degree of Q-Fano threefolds, Sbornik: Mathematics 198:11 (2007), 1683-1702.
- [21] M. Reid, Minimal models of canonical 3-folds, Adv. Stud. Pure Math. 1(1983), 131-180.
- [22] M. Reid, Young person's guide to canonical singularities, Proc. Symposia in pure Math. 46 (1987), 345-414.

- [23] I. Reider, Vector bundles of rank 2 and linear systems on algebraic surfaces, Ann. Math. 127(1988), 309-316.
- [24] V. V. Shokurov, The existence of a straight line on Fano 3-fold, Izv. Akad. Nauk SSSR Ser. Mat 43 (1979), 921-963, English translation in Math. USSR Izv. 15 (1980), 173-209.
- [25] V. V. Shokurov, 3-fold log flips, Izv. Russ. A. N. Ser. Mat. 56 (1992), 105-203.
- [26] V. V. Shokurov, Smoothness of a general anticanonical divisor on a Fano variety, Izv. Akad. Nauk SSSR, 14 (1980), 395-405.
- [27] K. Suzuki, On Fano indices of Q-Fano 3-folds. Manuscripta Math. 114 (2004), no. 2, 229–246
- [28] H. Takagi, Classification of primary Q-Fano threefolds with anti-canonical Du Val K3 surfaces, I, J. Alg. Geom. 15 (2006), 31-85.
- [29] S. G. Tankeev, On n-dimensional canonically polarized varieties and varieties of fundamental type, Izv. A. N. SSSR, Ser. Math. 35(1971), 31-44.
- [30] E. Viehweg, Vanishing theorems, J. reine angew. Math. 335(1982), 1-8.
- [31] Q. Zhang, Rational connectedness of log Q-Fano varieties. J. Reine Angew. Math. 590 (2006), 131-142.

Institute of Mathematics, School of Mathematical Sciences, Fudan University, Shanghai, 200433, China

*E-mail address*: mchen@fudan.edu.cn