Platonic polyhedra tune the 3-sphere: II. Harmonic analysis on cubic spherical 3-manifolds.

Peter Kramer, Institut fuer Theoretische Physik, University Tuebingen, Germany.

October 24, 2018

Abstract.

From the homotopy groups of two distinct cubic spherical 3-manifolds we construct the isomorphic groups of deck transformations acting on the 3-sphere. These groups become the cyclic group of order eight and the quaternion group respectively. By reduction of representations from the orthogonal group to the identity representation of these subgroups we provide two subgroup-periodic bases for the harmonic analysis on the 3-manifolds, which have applications to cosmic topology.

1 Introduction.

We view a spherical topological 3-manifold M , see [\[16\]](#page-22-0), as a prototile on its cover $\tilde{\mathcal{M}} = S^3$. We study in [\[11\]](#page-22-1) the isometric actions of $O(4, R)$ on the 3-sphere S^3 and give its basis as well-known homogeneous polynomials in [\[9\]](#page-22-2) eq.(37). From the homotopies of M we find as isomorphic images, see [\[14\]](#page-22-3), its deck transformations on S^3 . They form a subgroup H of $O(4, R)$. By group/subgroup representation theory with intermediate Coxeter groups we construct on $S³$ a H-periodic basis for the harmonic analysis on M.

Our approach yields in closed analytic form the onset, the selection rules, the multiplicity, projection operators, orthogonality rules and basis for each manifold. Among the Platonic polyhedra it was applied in [\[9\]](#page-22-2), [\[10\]](#page-22-4) to the Poincare dodecahedron with H the binary icosahedral group. An algorithm due to Everitt in $[7]$ describes homotopies for spherical 3-manifolds from five Platonic polyhedra. Following it we found and applied in [\[11\]](#page-22-1) for the tetrahedron as H the cyclic group C_5 .

One field of applications for harmonic analysis is cosmic topology, see [\[12\]](#page-22-5), [\[13\]](#page-22-6), [\[1\]](#page-21-1), [\[2\]](#page-21-2), [\[3\]](#page-21-3): The topology of a 3-manifold $\mathcal M$ is favoured if data from the Cosmic Microwave Background can be expanded in its harmonic basis.

Our work is well suited for comparing on $S³$ a family of topologies and their harmonic analysis. Here we turn to two distinct cubic spherical 3-manifolds, with homotopies described in [\[7\]](#page-21-0). We employ an intermediate Coxeter group, construct the groups of deck transformations, and derive and compare their harmonic analysis.

2 The Coxeter group G and the 8-cell on S^3 .

The cartesian coordinates $x = (x_0, x_1, x_2, x_3) \in E^4$ for S^3 we combine as in [\[9\]](#page-22-2), [\[11\]](#page-22-1) in the matrix form

$$
u = \begin{bmatrix} z_1 & z_2 \\ -\overline{z}_2 & \overline{z}_1 \end{bmatrix}, z_1 = x_0 - ix_3, z_2 = -x_2 - ix_1, z_1\overline{z}_1 + z_2\overline{z}_2 = 1.
$$
 (1)

For the group action we start from the Coxeter group $G < O(4, R)$ [\[8\]](#page-22-7), [\[7\]](#page-21-0) p. 254, with the diagram

$$
G = \circ \stackrel{4}{-} \circ \stackrel{3}{-} \circ \stackrel{3}{-} \circ. \tag{2}
$$

Fig. 1. The 8-cell projected to the plane according to [\[15\]](#page-22-8) pp. 177-8. Any face bounds two, any edge three, any vertex four out of the eight cubes.

For the Coxeter diagram eq. [2](#page-1-0) we give for the 4 Weyl reflections $W_s = W_{a_s}, s =$ 1, 2, 3, 4 the Weyl vectors in Table 2.1 and compute for each Weyl vector $a_s =$ $(a_{s0}, a_{s1}, a_{s2}, a_{s3})$ the matrix

$$
v_s := \begin{bmatrix} a_{s0} - ia_{s3} & -a_{s2} - ia_{s1} \ a_{s2} - ia_{s1} & a_{s0} + ia_{s3} \end{bmatrix} \in SU(2, R).
$$
 (3)

The matrices eq. [3](#page-1-1) will be used to relate, see [\[11\]](#page-22-1), the Weyl reflections to $SU^l(2, R) \times$ $SU^r(2, R)$ acting by left and right multiplication on the coordinates eq. [1.](#page-1-2) We include in Table 2.1 the inversion $\mathcal{J}_4 \in G$, and the additional Weyl operator W_0 .

G is isomorphic to the hyperoctahedral group, see $[4]$ p.90, which is a semidirect product of its normal subgroup $(C_2)^4$ of four reflections in the four coordinate axes with the subgroup $S(4)$ of all $4! = 24$ permutations of these axes,

$$
G = (C_2)^4 \times_s S(4). \tag{4}
$$

The group is of order $|G| = 2^4 4! = 384$. Its elements we denote as products $g =$ $\epsilon p, \epsilon \in (C_2)^4$, $p \in S(4)$ with the multiplication law

$$
\epsilon = (\epsilon_0, \epsilon_1, \epsilon_2, \epsilon_3), \ \epsilon_j = \pm 1,
$$

\n
$$
\epsilon p \ \epsilon' p' = \epsilon'' p'',
$$

\n
$$
\epsilon'' = \epsilon (p \epsilon' p^{-1}), \ p'' = p p',
$$

\n
$$
(p \epsilon p^{-1}) = (\epsilon_{p^{-1}(0)}, \epsilon_{p^{-1}(1)}, \epsilon_{p^{-1}(2)}, \epsilon_{p^{-1}(3)}).
$$
\n(5)

The hyperoctahedral form of G allows to study the irreducible representations. We write the permutation $p \in S(4)$ in cycle form but use the numbers $\{0, 1, 2, 3\}$ adapted to the enumeration of the coordinates in eq. [1.](#page-1-2) In Table 2.1 we give the action of the Weyl reflections on the coordinates x and the factors in $q = \epsilon p$.

Fig. 2. Enumeration of faces $F1, \ldots, F6$ and edges $1, \ldots, 12$ for the cubic prototile according to Everitt [\[7\]](#page-21-0) p. 260 Fig. 2.

The first three Weyl reflections from Table 2.1 generate, see [\[8\]](#page-22-7), the cubic Coxeter subgroup

$$
O = \circ \xrightarrow{4} \circ \xrightarrow{3} \circ,
$$
 (6)

isomorphic to the octahedral group $O \sim (C_2)^3 \times_s S(3)$ acting on $E^3 \in E^4$. This group from 48 simplices generates a spherical cube attached to a single vertex. We choose this cube as the prototile on S^3 .

We locate the center of the cubic prototile at $x = (1, 0, 0, 0) \in S^3$. When we include the action of W_4 along with the cubic Coxeter subgroup eq. [6,](#page-3-0) this cube is mapped into 7 companions which tile S^3 and generate the 8-cell tiling described in [\[15\]](#page-22-8) pp. 177-8 and shown in a projection in Fig. 1. The center positions of the 8 spherical cubes are located at the 8 points

$$
(\pm 1, 0, 0, 0), (0, \pm 1, 0, 0), (0, 0, \pm 1, 0), (0, 0, 0, \pm 1). \tag{7}
$$

Everitt [\[7\]](#page-21-0) has shown that $S³$ admits two cubic spherical 3-manifolds with inequivalent first homotopy. He enumerates the faces and edges and gives a graphical algorithm for their gluing, see Fig. 2. This gluing determines the generators for the first homotopy group.

Fig. 3. Axes $(x_1, x_2, x_3) \in E^3$ marked by arrows for the cubic prototile with face enumeration $(F1, F3, F2)$ according to Fig. 2. The three axes are cyclically rotated into one another by the rotation (W_2W_3) , $(W_2W_3)^3 = e$.

In Fig. 3, the three edges of the shaded triangle mark the intersections of the Weyl reflection (hyper-)planes for W_1, W_2, W_3 with the face F1 of the cube. Face F1 itself is part of the Weyl reflection (hyper-)plane for W_4 . In the homotopy group, there appears a gluing of opposite faces. Use of the inversion \mathcal{J}_3 in the center (black circle), followed by the Weyl reflection W_4 , converts this gluing of opposite faces into a standard deck operation $st(1 \Leftarrow 6)$ eq. [12.](#page-5-0) This operation maps the \mathcal{J}_3 -inverted and W_4 -reflected initial cube into a new position with its face F6 glued to face F1 of the initial cube.

Table 2.1 The Weyl vectors a_s , $s = 1, \dots, 4$ and a_0 for the Coxeter group G eq. [2,](#page-1-0) the 2×2 unitary matrices v_s eq. [3,](#page-1-1) the action on x, and the product form $g = \epsilon p$ with p in cycle form.

	Weyl vector a_s	matrix v_s	g x	ϵ	\mathcal{D}
	(0, 0, 0, 1)	$\begin{array}{c c} & \begin{array}{c} \cdot & 0 \\ 0 & i \end{array} \end{array}$	$(x_0, x_1, x_2, -x_3)$	$(+ + + -)$	
$2\,$	$ (0,0,-\sqrt{\frac{1}{2}},\sqrt{\frac{1}{2}})\ \sqrt{\frac{1}{2}}\ \begin{matrix} -i & 1 \\ -1 & i \end{matrix}$		(x_0, x_1, x_3, x_2)		(23)
	$3\left[\left(0,\sqrt{\frac{1}{2}},-\sqrt{\frac{1}{2}},0\right)\right]\sqrt{\frac{1}{2}}\left[\begin{array}{cc}0&1\\-1-i&0\end{array}\right]$		(x_0, x_2, x_1, x_3)		(12)
	$4\left[\left(-\sqrt{\frac{1}{2}}, \sqrt{\frac{1}{2}}, 0, 0\right)\right]$	$\sqrt{\frac{1}{2}}\begin{vmatrix} -1 & -i \\ -i & -1 \end{vmatrix}$	(x_1, x_0, x_2, x_3)		(01)
	(1,0,0,0)		$(-x_0, x_1, x_2, x_3)$		
	\mathcal{J}_4		$(-x_0, -x_1, -x_2, -x_3)$ $(- - -$		

3 The group isomorphism deck($C2 \sim \pi_1(C2) \sim C_8$ for the cubic spherical 3-manifold $(C2)$.

We start from the algorithm of [\[7\]](#page-21-0) p. 259 Table 3 on the first homotopy group $\pi_1(C2)$ for the cube and construct the explicit isomorphism to the group of deck transformations deck(C2). We denote the faces of the cube from Fig. 2 as $F1, \ldots, F6$. To the first three faces of $C2$, the glue partners are

$$
F3 \cup F1, \ F4 \cup F2, \ F5 \cup F3. \tag{9}
$$

We represent the four edges for a square face by the corresponding numbers from Fig. 2. Evaluating the glue algorithm for $C2$ in [\[7\]](#page-21-0), the first edge glue generator can be depicted from right to left as the map

$$
g_1 = g_1(1 \Leftarrow 3) : \begin{bmatrix} \overline{4} & \overline{1} & 2 \\ 7 & 7 & \end{bmatrix} \Leftarrow \begin{bmatrix} \overline{3} & 6 \\ 1 & 5 \end{bmatrix} . \tag{10}
$$

We use left arrows \Leftarrow in line with the usual sequence of operator products. Now we wish to map this glue generator $g_1(1 \Leftarrow 3)$ isomorphically into a generator from the

group deck $(C2)$ of deck transformation acting on the 3-sphere $S³$. We follow a similar method as employed in [\[9\]](#page-22-2) pp. 5322-4 for the Poincare dodecahedral 3-manifold. We refer to Fig. 3 for the position of the cube and for the three orthogonal directions 1, 2, 3. We factorize $g_1(1 \Leftarrow 3)$ into three actions: A positive rotation $R_3(\pi/2)$ around the 3-axis, followed by a standard glue operator $st(1 \leftarrow 6)$ eq. [12,](#page-5-0) followed by a positive rotation $R_1(\pi/2)$ around the 1-axis. In eq. [11](#page-5-1) we depict from right to left the images of the initial face $F3$. In the second line we write the corresponding triple product of operators.

$$
g_1 = g_1(1 \Leftarrow 3):
$$
\n
$$
\begin{bmatrix} \overline{1} \\ \overline{4} & 2 \\ 7 \end{bmatrix} \Leftarrow \begin{bmatrix} \overline{4} \\ 7 & 1 \\ 2 \end{bmatrix} \Leftarrow \begin{bmatrix} 10 \\ \overline{6} & 12 \\ \overline{11} & 2 \end{bmatrix} \Leftarrow \begin{bmatrix} \overline{3} \\ \overline{1} & 6 \\ 5 \end{bmatrix}
$$
\n
$$
R_1(\frac{\pi}{2}) \propto R_3(\frac{\pi}{2})
$$
\n(11)

The orientation of edges is counter-clockwise for the two right-hand diagrams but clockwise for the left-hand ones. The third operator maps $F6 \leftarrow F3$, the first one rotates F1 while preserving the center $x = (1, 0, 0, 0)$ of the initial cube. The second operator is crucial for the isomorphism $\pi_1(C2) \to deck(C2)$. It is constructed as the product

$$
st(1 \Leftarrow 6) = W_4 \mathcal{J}_3, \ \mathcal{J}_3 = W_0 \mathcal{J}_4,\tag{12}
$$

of the inversion operator $\mathcal{J}_3 \in H$ w.r.t. (x_1, x_2, x_3) , which can be factorized into the Weyl operator W_0 and the full coordinate inversion $\mathcal{J}_4 \in G$, followed by the Weyl reflection W_4 . The operator \mathcal{J}_3 in eq. [11](#page-5-1) inverts the cube in its center and so maps any face F_i of the cube into its opposite face, in the present case $F1 \leftarrow F6$. The final Weyl reflection W_4 following \mathcal{J}_3 reflects the inverted cube in its face $F1$ and produces an image of the initial cube with the new center $(0, 1, 0, 0)$, glued with its face $F3$ to the original face $F1$. The product eq. [12](#page-5-0) contains two Weyl reflections and therefore preserves the orientation.

For later use we list three standard glue operators of the cube in Table 3.1. We construct them from eq. [12](#page-5-0) by the conjugations

$$
st(2 \Leftarrow 4) = (W_3 W_2) st(1 \Leftarrow 6)(W_2 W_3), st(3 \Leftarrow 5) = (W_2 W_3) st(1 \Leftarrow 6)(W_3 W_2).
$$
\n(13)

listed in Table 3.2. Table 3.1: Standard glue operators in the Coxeter group G.

$$
\begin{array}{c|cc}\ng & g & \epsilon & p \\
\hline\nst(1 \Leftarrow 6) & (x_1, -x_0, -x_2, -x_3) & (+---) & (01) \\
st(2 \Leftarrow 4) & (x_2, -x_1, -x_0, -x_3) & (+---) & (02) \\
st(3 \Leftarrow 5) & (x_3, -x_1, -x_2, -x_0) & (+---) & (03)\n\end{array} \tag{14}
$$

The operations eq. [13](#page-5-2) are in G but not necessarily in $\text{deck}(C2)$. The rotations and the product W_4W_0 appearing in eq. [12](#page-5-0) are even in G, and so all operations preserve orientation.

Next we express the generator g_1 as a product of pairs of Weyl operators and find

$$
R_1(\pi/2) = (W_2W_1), \, st(1 \Leftarrow 6) = (W_4W_0)\mathcal{J}_4, R_3(\pi/2) = (W_2W_3)(W_2W_1)(W_3W_2), g_1 = (W_2W_1)(W_4W_0)\mathcal{J}_4(W_2W_3)(W_2W_1)(W_3W_2).
$$
 (15)

We use [\[11\]](#page-22-1) eqs. (60) for products of Weyl operators in eq. [15](#page-6-0) and the matrices v_s from Table 2.1 to rewrite the operator T_{g_1} in the form

$$
T_{g_1} = T_{(w_{l1}, w_{r1})},
$$
\n
$$
w_{l1} = (v_2 v_1^{-1})(v_4 v_0^{-1})(v_2 v_3^{-1})(v_2 v_1^{-1})(v_3 v_2^{-1}) = \sqrt{\frac{1}{2}} \begin{bmatrix} -\overline{a} & 0 \\ 0 & -a \end{bmatrix},
$$
\n
$$
w_{r1} = (-1)(v_2^{-1} v_1)(v_4^{-1} v_0)(v_2^{-1} v_3)(v_2^{-1} v_1)(v_3^{-1} v_2) = \sqrt{\frac{1}{2}} \begin{bmatrix} 0 & -a^3 \\ -a & 0 \end{bmatrix}, a = \exp(\pi i/4).
$$
\n(16)

By (w_l, w_r) we denote the elements of the $SU^l(2, R) \times SU^r(2, R)$ action $u \to w_l^{-1}uw_r$ on S^3 in the coordinates eq. [1.](#page-1-2) In this scheme, \mathcal{J}_4 in eq. [12](#page-5-0) yields the operator

$$
T_{\mathcal{J}_4} = T_{(e,-e)}.\tag{17}
$$

which commutes with all rotation operators. In eq. [16](#page-6-1) we have absorbed this operator into w_r . From eq. [16](#page-6-1) we easily see that $w_{l1}^8 = w_{r1}^8 = e$ so that $g_1, g_1^8 = e$ generates the cyclic group C_8 of order 8.

We can construct from the graphs given in [\[7\]](#page-21-0) the other glue generators of the homotopy group of C2 and map them isomorphically into generators of the group of deck transformations. It turns out that all of these glue generators become powers of the first glue generator $g_1 \in C_8$ eq. and its image eq. [16](#page-6-1) in $deck(C2)$. We list the actions of the eight powers g_1^t in **Table 3.2**.

1 Theorem: The homotopy group and the group of deck transformations for the spherical cubic 3-manifold C2 of Everitt [\[7\]](#page-21-0) p. 259 Table 3 are isomorphic to the cyclic group

$$
\text{deck}(C2) = C_8 = \langle g_1^t, t = 1, \dots, 8, g_1^8 = e \rangle \tag{18}
$$

with actions on $S³$ given in **Table 3.2**. The deck transformations generate fix-point free the 8-cell on S^3 . The group/subgroup scheme for the cubic 3-manifold $C2$ is

$$
O(4, R) > G > C_8.
$$
 (19)

3.1 The reduction $O(4, R) > C_8$ and harmonic analysis on $(C2).$

The irreducible representations of $O(4, R)$, their polynomial basis in terms of spherical harmonics, and the matrix elements of Weyl operators we adopt from [\[11\]](#page-22-1) section

4.2. Here we consider directly the reduction of representations for the group/subgroup pair

$$
O(4,R) > C_8. \tag{20}
$$

The operators for the generator of $g_1 \in C_8$ and its powers from [\[11\]](#page-22-1) eq. (60) have the form

$$
T_{g_1} = T_{(w_{l1}, w_{r1})}, \ T_{g_1^t} = T_{(w_{l1}^t, w_{r1}^t)}, \tag{21}
$$

given in **Table 3.2** in terms of the $SU^l(2, R) \times SU^r(2, R)$ action.

Table 3.2: The elements of the cyclic group C_8 of deck transformations of the manifold $C2$ and their actions on $S³$.

From eq. [21](#page-7-0) and from [\[11\]](#page-22-1) eq. (45) we find for the characters of the powers g_1^t in the irreducible representation $D^{(j,j)}$ in terms of characters of $SU(2, R)$ the result

$$
\chi^{(j,j)}(g_1^t) = \chi^j(w_{l1}^t)\chi^j(w_{r1}^t), \ t = 1, \dots, 8.
$$
 (23)

These expressions are easily evaluated with the help of [\[11\]](#page-22-1), Appendix. The multiplicity of the identity representation D^0 of C_8 is then given by

$$
m(C_8(j,j),0) = \frac{1}{8} \sum_{t=1}^{8} \chi^j(w_{l1}^t) \chi^j(w_{r1}^t).
$$
 (24)

In Table 3.3 we give the values of the characters of $SU^l(2, C), SU2(2, C)$ and the onset of multiplicities of C_8 -periodic states eq. [24](#page-7-1) as functions of the degree $2j$, $0 \leq$

 $j \leq 8$. The characters χ^j in this table can be divided into a set with the property $\chi^{j+4}(w_{l,r}^t) = \chi^j(w_{l,t}^t)$ and a set which increases with j. Treating these sets in eq. [24](#page-7-1) separately allows to derive the following recursion relation for the multiplicities:

$$
(2j) = \text{even} : m(C_8(j+4, j+4), 0) = m(C_8(j, j), 0) + 8j + 20 + 2(-1)^j. \tag{25}
$$

Multiplicities for odd degree $(2j)$ vanish: From **Table 3.2** one finds $g_1^4 = \mathcal{J}_4$, $\mathcal{J}_4 \in C_8$. The representation of any group element applied to a C_8 -periodic polynomial must give unity. For the inversion \mathcal{J}_4 in **Table 2.1** this allows only for even degree.

Table 3.3: The characters $\chi^j(w_l^t), \chi^j(w_r^t)$ $t = 1, ..., 8$ in $SU(2, R)$ for C_8 in the irreducible representations D^j of $SU(2, R)$. The characters $\chi^{(j,j)}(g_1^t)$ are given by eq[.23](#page-7-2) and the multiplicities $m(C_8(j, j)0)$ by eq. [24.](#page-7-1)

3.2 An orthogonal C_8 -periodic basis for the harmonic analysis on C2.

The spherical harmonics on S^3 are given from [\[11\]](#page-22-1) eq. (85) as the polynomials $D_{m_1,m_2}^j(z_1, z_2, \overline{z}_1, \overline{z}_2)$. Upon inserting the Euler angles (α, β, γ) by

$$
z_1 = \exp(i(\alpha + \gamma)/2)\cos(\beta/2)), z_2 = \exp(i(\alpha - \gamma)/2)\sin(\beta/2)),
$$
 (27)

into these spherical harmonics, we obtain the Wigner representation matrices of $u \in SU(2, C)$. Here we need them only for integer j. With the Euler angles we can then write the orthogonality relations from $[6]$ eq. $(4.6.1)$ as

$$
\frac{1}{8\pi^2} \int_0^{2\pi} \int_0^{\pi} \int_0^{2\pi} \overline{D}_{m'_1, m'_2}^{j'}(\alpha \beta \gamma) D_{m_1, m_2}^j(\alpha \beta \gamma) d\alpha \sin \beta d\beta d\gamma = \delta_{j', j} \ \delta_{m'_1, m_1} \ \delta_{m'_2, m_2} \frac{1}{2j+1}.
$$
\n(28)

With the representations of $SU^l(2, C) \times SU^r(2, C)$ given in [\[11\]](#page-22-1) eq. (85) we construct from Table 3.2 the representations of C_8 of the form

$$
(T_{(w_l^i, w_r^i)} D)^j_{m_1, m_2}(u) = D^j_{m_1, m_2}(w_l^{8-i} u w_r^i) = \sum_{m'_1, m'_2} D^j_{m'_1, m'_2}(u) D^j_{m_1, m'_1}(w_l^{8-i}) D^j_{m'_2, m_2}(w_r^i),
$$
\n(29)

whose coefficients are given in **Table 3.4**.

Table 3.4: Matrix elements in the representation $D^{(j,j)}$ eq. [29](#page-9-0) of the elements of C_8 for $i = 1, ..., 8$ with $l = 1, ..., 4$.

$$
\begin{array}{ll}\ni & D_{m_1,m'_1}^j(w_l^{8-i}) & D_{m'_2,m_2}^j(w_r^i) \\
\hline\n1 & i^{m_1} \delta_{m_1,m'_1} & (-1)^{2j}(-1)^{j+m_2} i^{m_2} \delta_{m'_2,-m_2} \\
2 & (-1)^{2j}(-1)^{m_1} \delta_{m_1,m'_1} & (-1)^{2j} \delta_{m'_2,m_2} \\
3 & (-1)^{m_1} i^{m_1} \delta_{m'_1,m_1} & (-1)^{j+m_2} i^{m_2} \delta_{m'_2,-m_2} \\
4 & (-1)^{2j} \delta_{m_1,m'_1} & \delta_{m'_2,m_2} \\
4 & (-1)^{2j} D_{m_1,m'_1}^j(w_l^{8-l}) & D_{m'_2,m_2}^j(w_r^l)\n\end{array} \tag{30}
$$

The projection operator on the identity representation D^0 of C_8 is

$$
P^{0} = \frac{1}{8} \sum_{i}^{8} T_{(w_{i}^{i}, w_{r}^{i})}.
$$
\n(31)

Its matrix elements from Table 3.4 for fixed j can be rewritten as

$$
(P^{0,j})_{m_1,m_2,m'_1,m'_2}
$$
\n
$$
= \frac{1}{2} \left[1 + (-1)^{2j} \right] \frac{1}{2} \left[1 + (-1)^{m_1} \right]
$$
\n
$$
\times \frac{1}{2} \left[\delta_{m_1,m'_1} \delta_{m'_2,m_2} + i^{m_1} (-1)^{j+m_2} i^{m_2} \delta_{m_1,m'_1} \delta_{m'_2,-m_2} \right]
$$
\n(32)

The two prefactors imply integer j and even m_1 . For this case we get by application of the projector

Table 3.5: The C_8 -periodic basis $\{\phi_{m_1,m_2}^j\}$ on S^3 for the harmonic analysis on the cubic spherical 3-manifold C2 in terms of spherical harmonics $D^{j}(u)$ on S^{3} .

$$
j = \text{integer}, m_1 = \text{even}, -j \le m_1 \le j, i^{m_1}(-1)^j = 1, m_2 = 0: \qquad (33)
$$

\n
$$
\phi_{m_1,0}^j = \frac{\sqrt{2j+1}}{\sqrt{8\pi}} D_{m_1,0}^j(u),
$$

\n
$$
j = \text{integer}, m_1 = \text{even}, -j \le m_1 \le j, 0 < m_2 \le j:
$$

\n
$$
\phi_{m_1,m_2}^j = \frac{\sqrt{2j+1}}{4\pi} \left[D_{m_1,m_2}^j(u) + i^{m_1}(-1)^{(j+m_2)} i^{m_2} D_{m_1,-m_2}^j(u) \right]
$$

3 Theorem: An orthonormal basis for the harmonic analysis on the cubic spherical manifold C2 is spanned by the C₈-periodic polynomials of degree $2j, j = 0, 1, 2, ...$ in Table 3.5.

4 The group isomorphism $\pi_1(C3) \sim$ deck $(C3) \sim Q$ for the cubic spherical 3-manifold $(C3)$.

The second possible homotopy group of the spherical 3-cube is given by Everitt [\[7\]](#page-21-0) in Table 3 p. 259 by a second graphical algorithm. We denote this cubic 3-manifold as C3. The order of the homotopy group and the group of deck transformations is again 8. For the homotopy of $C3$, opposite faces of the cube are glued,

$$
F6 \cup F1, F5 \cup F3, F4 \cup F2. \tag{34}
$$

From the edge gluing we find that each glue is followed by a left-handed rotation by $\pi/2$. We construct three glue generators q_1, q_2, q_3 from the prescription of [\[7\]](#page-21-0) p. 259 Table 3, depict the edge gluings from right to left as before and factorize them with the standard glue operators eq. [13:](#page-5-2)

$$
q_{1} = q_{1}(1 \Leftarrow 6):
$$
\n
$$
\begin{bmatrix} 7 \\ 2 & \overline{4} \\ \overline{1} & 1 \end{bmatrix} \Leftarrow \begin{bmatrix} 10 \\ \overline{6} & 12 \\ \overline{11} & 1 \end{bmatrix},
$$
\n
$$
\begin{bmatrix} 7 \\ 2 & \overline{4} \\ \overline{1} & 1 \end{bmatrix} \Leftarrow \begin{bmatrix} \overline{4} \\ 7 & \overline{1} \\ 2 & 1 \end{bmatrix} \Leftarrow \begin{bmatrix} \overline{4} \\ \overline{6} & 12 \\ \overline{11} & 12 \end{bmatrix}
$$
\n
$$
q_{2} = q_{2}(3 \Leftarrow 5):
$$
\n(35)

$$
\begin{bmatrix} 3 & \overline{5} \\ 3 & \overline{5} \\ 1 & \overline{1} \\ 3 & \overline{5} \\ 4 & \overline{6} \end{bmatrix} \leftarrow \begin{bmatrix} 8 & 7 \\ \overline{12} & \overline{5} \\ 1 & \overline{9} \\ 3 & \overline{5} \\ R_2(-\pi/2) \end{bmatrix},
$$
\n
$$
R_2(-\pi/2) \leftarrow \begin{bmatrix} 5 & 7 \\ 1 & 3 \\ 3 & \overline{6} \end{bmatrix},
$$
\n
$$
st(3 \Leftarrow 5) \left[\begin{bmatrix} 8 & 7 \\ \overline{12} & 5 \\ 9 & \overline{5} \\ 7 & \overline{3} \\ 8 & \overline{3} \\ 2 & 10 \\ 8 & \overline{3} \end{bmatrix} \right] \leftarrow \begin{bmatrix} 4 & 7 \\ 9 & 5 \\ 11 & \overline{5} \\ 11 & \overline{5} \\ 2 & \overline{3} \\ 8 & \overline{2} \end{bmatrix},
$$
\n
$$
R_3(-\pi/2) \leftarrow \begin{bmatrix} 4 & 7 \\ 3 & 5 \\ 7 & \overline{2} \\ 8 & \overline{2} \end{bmatrix} \leftarrow \begin{bmatrix} 4 & 7 \\ 10 & 5 \\ 2 & \overline{2} \\ 2 & \overline{2} \end{bmatrix} \right)
$$
\n
$$
st(2 \Leftarrow 4)
$$

The three generators in terms of Weyl reflections become

$$
q_1 = (W_1 W_2)(W_4 W_0) \mathcal{J}_4,
$$

\n
$$
q_2 = (W_3 W_2) q_1 (W_2 W_3),
$$

\n
$$
q_3 = (W_2 W_3) q_1 (W_3 W_2).
$$
\n(36)

Analysis of the products of Weyl operators in eq. [36](#page-11-0) in terms of actions of $SU^l(2, R)$ × $SU^{r}(2, R)$ given in eqs. [42,](#page-12-0) [43](#page-12-1) show that the generators q_1, q_2, q_3 act exclusively from the left on $u \in S^3$ eq. [1](#page-1-2) and in $SU^l(2, R)$ take a form equivalent according to

$$
(q_1, q_2, q_3) \sim (-\mathbf{k}, -\mathbf{j}, -\mathbf{i})
$$
\n(37)

with the standard 2×2 quaternion representation. As mentioned before, any apparent reversion of the edge orientation is compensated by a Weyl reflection. Expressed in G the generators from eq. [36](#page-11-0) are given in Table 4.1. Their multiplication yields the relations

$$
q_1^2 = q_2^2 = q_3^2 = q_1 q_2 q_3 = \mathcal{J}_4. \tag{38}
$$

These are exactly the relations characterizing the quaternion group Q , see [\[4\]](#page-21-4) p.8, with \mathcal{J}_4 playing the role of (-1) . The eight elements are

$$
\text{deck}(C3) = Q = \langle e, q_j, \mathcal{J}_4, \mathcal{J}_4 q_j, j = 1, 2, 3 \rangle. \tag{39}
$$

We therefore have shown

2 Theorem: The homotopy group and the group of deck transformations for the spherical cubic 3-manifold C3 of Everitt [\[7\]](#page-21-0) p. 259 Table 3 are isomorphic to the

quaternion group Q. The elements of the group $\text{deck}(C3)$ act on S^3 from the left as given in eqs. [42,](#page-12-0) [43](#page-12-1) and generate fix-point free the 8-cell on S^3 . By the construction of the generators of Q in Table 4.1 we also have shown the group/subgroup relation

$$
O(4, R) > G > \operatorname{deck}(C3) = Q.
$$
\n
$$
(40)
$$

4.1 The reduction $O(4, R) > Q$ and harmonic analysis on $(C3).$

The harmonic analysis for the 3-cube $(C3)$ has as its basis the Q-periodic spherical harmonics on S^3 of degree $(2j)$. Selection rules eliminate all contributions with $2j =$ odd. The periodic states are degenerate with respect to the subgroup $SU^r(2, R)$, but the degenerate states can be labelled by a representation index $m_2, -j \leq m_2 \leq$ j. The explicit construction of the Q -periodic polynomials can be found by the application of Young operators as described in [\[11\]](#page-22-1) section 4.7.

4.2 The multiplicity $m(Q(j, j), 0)$ of representations in $O(4, R) > Q$.

The reduction of representations we analyze in section 5 in the scheme eq. [40.](#page-12-2) Here we consider the reduction $O(4, R) > Q$.

To compute the multiplicity $m((j, j), 0)$ for the reduction of the irreducible representations $D^{(j,j)}$ of $O(4, R)$ to the identity representation of the subgroup Q, denoted as $D^{0}(Q)$, we need the characters $\chi^{(j,j)}$ for the 8 elements of Q eq. [39.](#page-11-1) The elements e, \mathcal{J}_4 have the characters

$$
\chi^{(j,j)}(e) = (2j+1)^2, \ \chi^{(j,j)}(\mathcal{J}_4) = (-1)^{2j}(2j+1)^2. \tag{41}
$$

The second equality arises because the basis of $D^{(j,j)}$ has the homogeneous degree $2j$. The elements q_1, q_2, q_3 by **Table 4.3** are conjugate not in Q but in $O(4, R)$ and so have the same character w.r.t. $D^{(j,j)}$. We choose $q_1 = W_2 W_1 W_0 W_4$ as representative. Application of [\[11\]](#page-22-1) eq.(60) for a product of four Weyl operators gives for q_1 in terms of the $SU^l(2, R) \times SU^{r}(2, R)$ action

$$
T_{q_1} = T_{W_2 W_1 W_0 W_4} = T_{(v_2 v_1^{-1} v_0 v_4^{-1}, v_2^{-1} v_1 v_0^{-1} v_4)}.
$$
\n
$$
(42)
$$

Evaluation of the two matrix products in eq. [42](#page-12-0) with $v_0 = e$ gives

$$
(v_2v_1^{-1})(v_0v_4^{-1}) = w_{l1} = \begin{bmatrix} 0 & -i \\ -i & 0 \end{bmatrix}, (v_2^{-1}v_1)(v_0^{-1}v_4) = w_{r1} = e.
$$
 (43)

and so from eq. [43](#page-12-1)

$$
T_{q_1} = T_{(w_{l1},e)}, \ \chi^{(j,j)}(q_1) = \chi^j(w_{l1})\chi^j(e) = \chi^j(w_{l1})(2j+1). \tag{44}
$$

The result eq. [43](#page-12-1) shows that q_1 operates on $u \in S^3$ exclusively by left action corresponding to the subgroup $SU^l(2, R) < (SU^l(2, R) \times SU^r(2, R))$. The same holds true for the operators T_{q_2}, T_{q_3} since the conjugations in eq. [36](#page-11-0) applied to T_{q_1} preserve the subgroup $SU^l(2, R)$.

For the characters of the element $w_l \in SU^l(2, R)$ one finds

$$
\chi^{1/2}(w_{l1}) = 2\cos(\phi/2) = 0, \chi^{0}(w_{l1}) = 1, \n\chi^{(j+2)}(w_{l1}) = \chi^{j}(w_{l1}) = (-1)^{j}
$$
\n(45)

where the period 2 in the last line arises from $\phi/2 = \pi/2$. With these expressions we find the multiplicity of the Q-periodic states for a given representation $D^{(j,j)}$ as

$$
m(Q(j, j), 0) = \frac{1}{8} \sum_{g \in Q} \chi^{(j, j)}(g)
$$

= $\frac{1}{8} (1 + (-1)^{(2j)})(2j + 1) [(2j + 1) + 3(-1)^j].$ (46)

The first prefactor eliminates all the states with $(2j) = \text{odd}$. Eq. [39](#page-11-1) states that $\mathcal{J}_4 \in Q$, and therefore any Q-periodic state must be even.

The second prefactor $(2j + 1)$ arises from the degeneracy with respect to the group $SU^{r}(2, R)$. For a Q-periodic state of polynomial degree $2j$, we can choose $(2j + 1)$ orthogonal basis states with respect to $SU^r(2, R)$, corresponding to the second label m_2 in the spherical harmonics $D^j_{m_1,m_2}(u)$ on S^3 .

In Table 4.2 we give the onset of the multiplicity eq. [46](#page-13-0) for the lowest values of $2j$.

Table 4.1: The three glue generators q_i of deck(C3) eq. [36](#page-11-0) as elements of the Coxeter group G and the corresponding pairs $(w_{li}, w_{ri}) \in SU^l(2, R) \times SU^r(2, R)$.

$i \mid q_i x$	w_{li}	w_{ri} ϵ		
				(47)
	1 $(x_1, -x_0, x_3, -x_2)$ $\begin{bmatrix} 0 & -i \\ -i & 0 \end{bmatrix}$ = -k e $(+ - + -)$ (01)(23) 2 $(x_2, -x_3, -x_0, x_1)$ $\begin{bmatrix} 0 & -1 \\ 0 & -1 \\ 1 & 0 \end{bmatrix}$ = -j e $(+ - +)$ (02)(13) 3 $(x_3, x_2, -x_1, -x_0)$ $\begin{bmatrix} -i & 0 \\ 0 & i \end{bmatrix}$ = -i e $(+ + - -)$			

Table 4.2 The character $\chi^j(w_{l1})$ and the multiplicity $m(Q(j, j), 0), j \leq 8, 2j$ = even, eq. [46](#page-13-0) of Q-periodic spherical harmonics on $S³$ as functions of j.

4.3 An orthogonal Q-periodic basis for the harmonic analysis on C3.

For the quaternionic group Q , the orthogonal basis for the harmonic analysis can be given in closed analytic form. We shall use the irreducible representation matrices $D_{m_1,m_2}^j(u)$ given in [\[11\]](#page-22-1) eq. (85). Since $Q < SU^l(2, C)$ acts from the left only, its representations are given by $D^j_{m_1,m_2}(g)$, $g \in Q$. These matrix elements are given in Table 4.3 for general j . In what follows we need them only for integer j .

Table 4.3: Representation matrices for $g \in Q$ and general j.

$$
\frac{g}{e} \frac{D_{m_1,m_2}^j(g), -j \le (m_1, m_2) \le j}{\delta_{m_1,m_2}} \n\frac{q_1}{\delta_{m_1,-m_2}(-1)^{j-m_1} \exp(-i\pi m_1)} \n\frac{q_2}{\delta_{m_1,-m_2}(-1)^{j+m_1}} \n\frac{q_3}{\delta_{m_1,m_2} \exp(-i\pi m_1)} \n\mathcal{J}_4 \begin{bmatrix} \delta_{m_1,m_2}(-1)^{2j} & \delta_{m_1,m_2}(-1)^{j} \end{bmatrix}
$$
\n(49)

We next construct a projection operator on the identity representation D^0 of Q :

$$
P^{0} := \frac{1}{8} \sum_{g \in Q} T_{g} = \frac{1}{4} \left[T_{e} + \sum_{i=1}^{3} T_{q_{i}} \right], (P^{0})^{2} = P^{0}.
$$
 (50)

The representation matrix of this operator for fixed integer j is obtained from Table 4.3 in the simple form

$$
(P^{0,j})_{m_1,m'_1} = \frac{1}{2} \left[1 + (-1)^{m_1} \right] \frac{1}{2} \left[\delta_{m_1,m'_1} + \delta_{m_1,-m'_1}(-1)^j \right]. \tag{51}
$$

The projection operator eq. [51](#page-14-0) applied to a spherical harmonic does not affect the column index $m₂$. For the projection it proves convenient to separate the even and odd j case. The linearly independent, orthogonal Q-periodic polynomials normalized according to eq. [28](#page-9-1) are then found by projection with eq. [51.](#page-14-0)

4 Theorem: An orthonormal basis for the harmonic analysis on the cubic spherical manifold C3 is spanned by the Q-periodic polynomials of degree $2j, j = 0, 1, 2, ...$ in Table 4.4.

Table 4.4: The Q-periodic orthonormal basis $\{\phi_{m_1,m_2}^{j \text{ odd}}, \phi_{m_1,m_2}^{j \text{ even}}\}$ for the harmonic analysis on the cubic spherical manifold C3 in terms of spherical harmonics $D^{j}(u)$ on S^3 .

$$
j = \text{odd}, j \ge 3, m_1 = \text{even}, 0 < m_1 \le j, -j \le m_2 \le j:
$$
\n
$$
\phi_{m_1, m_2}^{\text{jodd}} = \frac{\sqrt{2j+1}}{4\pi} \left[D_{m_1, m_2}^j(u) - D_{-m_1, m_2}^j(u) \right],
$$
\n
$$
(52)
$$

$$
m(Q(j, j), 0) = \frac{1}{2}(2j + 1)(j - 1),
$$

\n
$$
j = \text{even}, m_1 = 0, -j \le m_2 \le j:
$$

\n
$$
\phi_{0,m_2}^{j \text{even}} = \frac{\sqrt{2j + 1}}{\sqrt{8\pi}} D_{0,m_2}^j(u),
$$

\n
$$
j \ge 2, \text{even}, 0 < m_1 \le j, m_1 = \text{even}:
$$

\n
$$
\phi_{m_1,m_2}^{j \text{even}} = \frac{\sqrt{2j + 1}}{4\pi} \left[D_{m_1,m_2}^j(u) + D_{-m_1,m_2}^j(u) \right],
$$

\n
$$
m(Q(j,j), 0) = \frac{1}{2}(2j + 1)(j + 2).
$$

The orthogonality and normalization is obtained by use of eq. [28.](#page-9-1) The multiplicities $m(Q(j, j), 0)$ agree with the values derived before in **Table 4.2**.

5 Reduction from the Coxeter group G .

5.1 Representations of G .

For the construction of irreducible representations of G eq. [4](#page-2-0) we follow Coleman [\[5\]](#page-21-6) in the notion of the little group and little co-group. For the cyclic group C_2 we denote the two elements by $\epsilon = \pm 1$ and the two irreducible representations by

$$
\sigma^{\pm} : \sigma^{+}(\epsilon) = 1, \ \sigma^{-} : \sigma^{-}(\epsilon) = \epsilon. \tag{53}
$$

The irreducible representations of the direct product $(C_2)^4 < G$ are then denoted by

$$
\mu = {\mu_1, \mu_2, \mu_3, \mu_4}, \mu_j = \pm,
$$

\n
$$
D^{\mu}(\epsilon) = \sigma^{\mu_1}(\epsilon_0) \sigma^{\mu_2}(\epsilon_1) \sigma^{\mu_3}(\epsilon_2) \sigma^{\mu_4}(\epsilon_3).
$$
\n(54)

We choose for D^{μ} six representatives and give the little co-groups,

$$
D^{\mu},
$$

\n
$$
\mu = \{++++\}, \{---\}, K = S(4),
$$

\n
$$
\mu = \{+++-\}, \{---+\}, K = S(3) \times S(1),
$$

\n
$$
\mu = \{++--\}, \{---++\}, K = S(2) \times S(2).
$$
\n(55)

The little co-group K for the irreducible representation D^{μ} is defined [\[5\]](#page-21-6) p. 108 as the maximal subgroup $K < S(4)$ such that

$$
h \in H : D^{\mu}(h \epsilon h^{-1}) = D^{\mu}(\epsilon). \tag{56}
$$

The little group is the direct product $L = (C_2)^4 \times K < G$. Its irreducible representations are direct products of the chosen representation D^{μ} with the irreducible representations D^f of the little co-group H. Finally the irreducible representations of G are the representations induced from the little group $L < G$. This induction, denoted by the sign \uparrow , is obtained by choosing coset generators $c_i \in S(4)$, $j = 1, \dots, |S(4)|/|K|$ of the little co-group $K < S(4)$ and with their help constructing

$$
D_{is,jt}^{(\mu,f)\uparrow}(\epsilon p) = \delta(c_j^{-1}pc_i, k \in K)D^{\mu}(c_j^{-1}\epsilon c_j)D_{s,t}^f(k). \tag{57}
$$

The character of this representation is

$$
\chi^{(\mu,f)\uparrow}(\epsilon p) = \sum_{j} \delta(c_j^{-1} p c_j, k \in K) D^{\mu}(c_j^{-1} \epsilon c_j) \chi^f(k). \tag{58}
$$

To determine these characters we must specify for each little co-group its coset generators. The multiplicity of the identity representation of the subgroups $H = (C_8, Q)$ is then given by

$$
m((\mu, f) \uparrow, 0) = \frac{1}{8} \sum_{g \in H} \delta(g, \epsilon p) \chi^{(\mu, f) \uparrow}(\epsilon p).
$$
 (59)

We give the corresponding data and the multiplicities in the next subsection.

5.2 The reductions $G > C_8$, $G > Q$.

The factorizations $g = \epsilon p$ eq. [5](#page-2-1) of the subgroup elements are given in **Tables 3.2**, 4.1.

Representations of G with $\mu = \{++++\}, \{---\}, K = S(4)$: For all elements $g \in C_8, g \in Q : g = \epsilon p$ one finds in both representations $D^{\mu}(\epsilon) = 1$.

Table 5.1 Characters $\chi^{(\mu f)\uparrow}(g_1^t)$ for $g \in C_8$ and multiplicity $m(C_8) = (C_8 \mu f)$, 0) eq. [59](#page-16-0) of the identity representation of C_8 in the representations $\mu = \{+++\}, \{---\}$ for all partitions f of $S(4)$.

Table 5.2 Characters $\chi^{(\mu f)\uparrow}(g)$ for $g \in Q$ and multiplicity $m(Q) = m((\mu f), 0)$ eq[.59](#page-16-0) of the identity representation of Q in the representations $\mu = \{++++\}, \{---\}$. The elements (q_1, q_2, q_3) are conjugate in G and so their characters coincide.

Representations with $\mu = \{+++-\}, \{---+\}, K = S(3) \times S(1)$: In Table 5.3 we list the 4 coset generators c_i of $K = S(3) \times S(1) < S(4)$ in cycle form and give their action on $D^{\mu}(\epsilon)$.

Table 5.3 Coset generators c_j and their action on $D^{\mu}(\epsilon)$ for the representations of G with $\mu = \{+++-\}, \{---+\}, K = S(3) \times S(1).$

The representations of the little co-group $K = S(3) \times S(1)$ are $f = [3] \times [1]$, [111] \times $[1], [21] \times [1]$. The characters of the elements of $H = C_8$, Q are given by eq. [58.](#page-16-1) The conjugations $p \to c_j^{-1} p c_j$ cannot change the class of p encoded by its cycle expression. Therefore the condition $c_j^{-1}gc_j = k \in S(3) \times S(1)$ eliminates all contributions except for $q = (e, \mathcal{J}_4)$. The reduced set of characters and the multiplicities are given in the following Tables.

Table 5.4 Non-vanishing characters $\chi^{(\mu f)\dagger}(g_1^t)$ eq. [58](#page-16-1) for $g_1 \in C_8$ and multiplicity $m(C_8) = m((\mu f), 0)$ eq. [59](#page-16-0) of the identity representation of C_8 in the representations $\mu = \{+++-\}, \{---+\}$ for all partitions $f = f_1 \times f_2$ of $S(3) \times S(1)$.

Table 5.5 Non-vanishing characters $\chi^{(\mu f)\dagger}(g)$ for $g \in Q$ and multiplicity $m(Q)$ $m((\mu f), 0)$ eq[.59](#page-16-0) of the identity representation of Q in the representations

$$
\mu = \{++, + -\}, \{---+\}.
$$
\n
$$
\begin{array}{c|cc}\n\chi^{(\mu f)\dagger}(g), g \in Q & g & [3] \times [1] & [111] \times [1] & [21] \times [1] \\
\hline\ne & 4 & 4 & 4 \\
\hline\n\mathcal{J}_4 & -4 & -4 & -4 \\
\hline\nm(Q) & 0 & 0 & 0\n\end{array}
$$
\n(64)

Representations D^{μ} with $\mu = \{++--\}, \{--++\}, K = S(2) \times S(2)$: We list the 6 coset generator of the little co-group in the next Table.

Table 5.6 Coset generators c_j and their action on $D^{\mu}(\epsilon)$ for the representations of G with $\mu = \{++--\}, \{--++\}, K = S(2) \times S(2).$

For $g_1^l \text{ } \in C_8$ and the coset generators in **Table 5.6** we now check the condition $c_j^{-1}g_1^lc_j \in S(2) \times S(2)$. The cycle structure of $S(2) \times S(2)$ admits only the elements e , (01), (23), (01)(23), and this condition applies at most for the elements $g_1^2, g_1^4, g_1^6, g_1^8 = e$. For these elements one finds the non-vanishing contributions given in Table 5.7.

Table 5.7 Nonvanishing character contributions for C_8 for $\mu = \{++--\}, \{--++\}.$ The irreducible representations $f = f_1 \times f_2$ of $S(2) \times S(2)$ are $D^{f_1 \times f_2}((01)(23)) =$ $(-1)^{\rho_1+\rho_2}$, with $\rho_{1,2}=0,1$ for $f_{1,2}=[2],[11].$

Evaluation of the multiplicity eq. [59](#page-16-0) from this Table gives for both representations

$$
\mu = \{++--\}, \{--++\} : m(C_8) = m(C_8(\mu f) \uparrow, 0) = \frac{1}{8} \left[12 - 4(-1)^{\rho_1+\rho_2}\right].
$$
 (67)

Table 5.8 Nonvanishing character contributions for Q for $\mu = \{++--\}, \{--++\}$ in the notation of Table 5.7.

g				$c_j^1 c_i^{-1} \epsilon c_j$ $c_j^1 p c_j$ $D^{++--1}(c_i^{-1} \epsilon c_j)$ $D^{-+++1}(c_i^{-1} \epsilon c_j)$ $\chi^f(c_i^{-1} g c_j)$	
	e, \mathcal{J}_4 $\pm (++++)$ c_1c_6 e		\sim 1		
	$q_1, q_1, \mathcal{J}_4 \mid \pm (+ - + -) \quad c_1$	$(01)(23) -1$		-1	$(-1)^{\rho_1+\rho_2}$
	$\pm (+ - + -)$ c_6	$(01)(23) -1$		-1	$(-1)^{\rho_1+\rho_2}$
	$q_2, q_2 \mathcal{J}_4 \pm (+ - +) c_2$	$(01)(23) -1$		-1	$(-1)^{\rho_1+\rho_2}$
	$\pm(-++-)$ c_5	$(01)(23) -1$		-1	$(-1)^{\rho_1+\rho_2}$
	$q_3, q_3 \mathcal{J}_4 \mid \pm (+ - + -) \quad c_3$	$(01)(23) -1$		-1	$(-1)^{\rho_1+\rho_2}$
	$\pm (+ - + -)$ c_4	$(01)(23) -1$		-1	$(-1)^{\rho_1+\rho_2}$
					(68)

Evaluation of the multiplicity eq. [59](#page-16-0) from this Table gives for these representations

$$
\mu = \{++--\}, \{--++\} : m(Q) = m(Q(\mu f) \uparrow, 0) = \frac{1}{8} \left[12 - 12(-1)^{\rho_1+\rho_2}\right].
$$
 (69)

Table 5.9 Representations $D^{(\mu f) \uparrow}$ of G, their dimensions, and summary on multiplicities for C_{8} - and Q -periodic states.

$D^{(\mu,f)\uparrow}$	μ							
	$\{++++\}, \{-$	$\left 4\right $	[1111]	[31]	[211]	[22]		
$\dim((\mu f)\uparrow)$				3	3	$\overline{2}$	(70)	
$m(C_8)$			θ	θ	1			
m(Q)				θ	θ	$\overline{2}$		
$D^{(\mu,f)\uparrow}$	μ		$f = f_1 \times f_2$					
	$\{+++-\},\$	[3]	\times [1]	111	\times [1]	[21] \times	1	
$\dim((\mu f)\uparrow)$		4		4		8		
$m(C_8)$				0				
m(Q)		0		0		0		
$D^{(\mu,f)\uparrow}$	μ		$f = f_1 \times f_2$					
	$+ +$ $++$	[2]	\times [2]	$\left 2\right $	\times [11]	[2] 11 \times	$\left[11\right]$ 11 \times	
$\dim((\mu f)\uparrow)$		6		6		6	6	
$m(C_8)$				$\mathcal{D}_{\mathcal{A}}$		$\overline{2}$		
m(Q)				3		3	0	

The selection rules for the cubic spherical manifolds C_2, C_3 now eliminate full representations of the Coxeter group G. The multiplicities $m(C_8) = m(Q) = 0$ for the representations with $\mu = \{+++-\}, \{---+\}$ of dimensions $\{4, 8\}$ are easily understood. In these representations one finds for the element \mathcal{J}_4 of both subgroups from eq. [57](#page-16-2)

$$
D_{is,jt}^{(\mu f)\uparrow}(\mathcal{J}_4) = (-1)\delta_{ij}\delta_{st},\tag{71}
$$

which excludes the identity representation of C_8 and Q .

6 Alternative spherical harmonics on S^3 .

The authors from the Ulm group of scientists in [\[1\]](#page-21-1), [\[2\]](#page-21-2), [\[3\]](#page-21-3) have carried out a similar analysis of spherical 3-manifolds and combined it with a powerful astrophysical analysis. Here we briefly compare the present and their algebraic and topological concepts. The Laplacian Δ on S^3 used in [\[1\]](#page-21-1) equals a Casimir operator Λ^2 of $SO(4,R)$ analyzed in [\[9\]](#page-22-2) eq.(30). The index β is given by $\beta = 2j + 1$. The spherical harmonics on $S³$ of [\[1\]](#page-21-1) eq. (9) for fixed j are related to ours algebraically by

$$
\psi_{\beta lm}^{S^3}(u) = \delta_{\beta,2j+1} \sum_{m_1,m_2} D_{m_1,m_2}^j(u) \langle j - m_1 j m_2 | lm \rangle (-1)^{j-m_1},
$$
(72)

$$
l = 0, 1, 2, ..., 2j = \beta - 1,
$$

$$
D_{m_1,m_2}^j(u) = \delta_{\beta,2j+1} \sum_{l=0}^{2j}, \psi_{\beta lm}^{S^3}(u) \langle j - m_1 j m_2 | lm \rangle (-1)^{j-m_1}, m = -m_1 + m_2.
$$

The coefficients are the well-known Wigner coefficients of $SU(2, C)$, [\[6\]](#page-21-5) pp. 31-45. The spherical harmonics eq. [72](#page-20-0) are adapted to the conjugation action $T_{(g,g)}$: From eq. [72](#page-20-0) with [\[11\]](#page-22-1) eq. (44) follows the transformation law

$$
(T_{(g,g)}\psi_{\beta lm}^{S^3})(u) := \psi_{\beta lm}^{S^3}(g^{-1}ug) = \sum_{m'=-l}^{l} \psi_{\beta lm'}^{S^3}(u)D_{m',m}^l(g), \tag{73}
$$

corresponding to the law for spherical harmonics Y_{lm} on S^2 .

With respect to the topology, the authors of [\[1\]](#page-21-1), [\[2\]](#page-21-2) select homogeneous spaces which in our scheme arise from the left action of binary subgroups of $SU^l(2, C)$ on S^3 . For these cases the spherical harmonics $D_{m_1,m_2}^j(u)$ eq. [72](#page-20-0) provide the simplest transformation law and the degeneracy $(2j+1)$, and are used in the action of Q on the cubic 3-manifold C3 in section 4.

The groups considered in [\[9\]](#page-22-2), [\[11\]](#page-22-1) and here are derived from the homotopies of specific 3-manifolds and corresponding homogeneous spaces. The action of $C₅$ for the tetrahedral 3-manifold analyzed in [\[11\]](#page-22-1) and of C_8 on the cubic 3-manifold C_2 are outside the exclusive left-or-right-action schemes. They illustrate a greater variety of 3-manifolds, homotopies, and subgroup actions on $S³$. Our use of characters for multiplicities and of algebraic projection operators avoids a numerical analysis of multiplicities and basis functions.

7 Conclusion.

The harmonic analysis on spherical cubic 3-manifolds extends the comparative study of Platonic 3-manifolds beyond the dodecahedron in [\[9\]](#page-22-2),[\[10\]](#page-22-4) and the tetrahedron in [\[11\]](#page-22-1). Two distinct homotopy groups for cubic spherical 3-manifolds C2, C3, im-plicitly defined by Everitt [\[7\]](#page-21-0), are identified as the cyclic group $H = C_8$ and the quaternion group $H = Q$ respectively and mapped into their isomorphic groups of deck transformations. Using the general analysis of Weyl reflection operators and their representations given in [\[11\]](#page-22-1), we give on S^3 a C_8 -periodic orthonormal basis adapted to the harmonic analysis for the cubic spherical manifolds C2, and another Q-periodic orthonormal basis for C3. The distinct multiplicities, the onset, recursion and selection rules of H-periodic states are computed. When the Coxeter group G is placed in between the orthogonal group $O(4, R)$ and H, entire representations are ruled out and additional orthogonalities are provided.

Acknowledgment.

The author thanks T. Kramer, University of Regensburg, who did substantial algebraic computations for the Tables.

References

- [1] Aurich R, Lustig S, and Steiner F, *CMB Anisotropy of the Poincare' Dodecahedron.* Class Quant Grav 22 (2005) 2061
- [2] Aurich R, Lustig S, and Steiner F, *CMB Anisotropy of Spherical Spaces.* Class Quant Grav 22 (2005) 3443
- [3] Aurich R, Janzer H S, Lustig S, and Steiner F, *Do we live in a small Universe?* astro-phys/0708,1420v2
- [4] Coxeter H S M and Moser W O J, *Generators and relations for discrete groups*, Springer, Berlin 1965
- [5] Coleman A J, *Induced and subduced representations*, in: Group Theory and its applications, ed. E M Loebl, Academic Press, New York 1968, pp. 57-129
- [6] Edmonds A R, *Angular momentum in quantum mechanics*, Princeton University Press, Princeton 1957
- [7] Everitt B, *3-manifolds from Platonic solids*, Topology and its Applications 138 (2004), 253-63
- [8] Humphreys J E, *Reflection groups and Coxeter groups*, Cambridge University Press, Cambridge 1990
- [9] Kramer P, *An invariant operator due to F Klein quantizes H Poincare's dodecahedral manifold*, J Phys A: Math Gen 38 (2005) 3517-40
- [10] Kramer P, *Harmonic polynomials on the Poincare dodecahedral 3-manifold* J. of Geometry and Symmetry in Physics 6 (2006) 55-66
- [11] Kramer P, *Platonic polyhedra tune the 3-sphere: Harmonic analysis on simplices.* Physica Scripta (2009), to appear, arxiv: 0810.3403
- [12] Lachieze-Rey M and Luminet J-P, *Cosmic topology*, Phys Rep 254 (1995) 135- 214
- [13] Luminet J-P, Weeks J R, Riazuelo A, Lehoucq R and Uzan J-Ph, *Dodecahedral space topology as an explanation for weak wide-angle temperature correlations in the cosmic microwave background*, Nature 425 (2003) 593-5
- [14] Seifert H and Threlfall W, *Lehrbuch der Topologie*, Leipzig 1934, Chelsea Reprint 1980
- [15] Sommerville D M Y, *An introduction to the geometry of* N *dimensions*, Dover, New York 1958
- [16] Thurston W P, *Three-Dimensional Geometry and Topology*, Princeton University Press, Princeton 1997