

Spacetimes characterized by their scalar curvature invariants

Alan Coley¹, Sigbjørn Hervik², Nicos Pelavas¹

¹Department of Mathematics and Statistics,
Dalhousie University, Halifax, Nova Scotia,
Canada B3H 3J5

²Faculty of Science and Technology,
University of Stavanger,
N-4036 Stavanger, Norway

aac, pelavas@mathstat.dal.ca, sigbjorn.hervik@uis.no

August 17, 2009

Abstract

In this paper we determine the class of four-dimensional Lorentzian manifolds that can be completely characterized by the scalar polynomial curvature invariants constructed from the Riemann tensor and its covariant derivatives. We introduce the notion of an \mathcal{I} -non-degenerate spacetime metric, which implies that the spacetime metric is locally determined by its curvature invariants. By determining an appropriate set of projection operators from the Riemann tensor and its covariant derivatives, we are able to prove a number of results (both in the algebraically general and in algebraically special cases) of when a spacetime metric is \mathcal{I} -non-degenerate. This enables us to prove our main theorem that a spacetime metric is either \mathcal{I} -non-degenerate or a Kundt metric. Therefore, a metric that is not characterized by its curvature invariants must be of Kundt form. We then discuss the inverse question of what properties of the underlying spacetime can be determined from a given a set of scalar polynomial invariants, and some partial results are presented. We also discuss the notions of *strong* and *weak* non-degeneracy.

1 Introduction

In matters related to relativity and gravitational physics we are often interested in comparing various spacetime metrics. Often identical metrics (which, of course, would give identical physics) are given in different coordinates and will therefore be disguising their true equivalence. It is therefore of import to have an invariant way to distinguish spacetime metrics. The perhaps easiest way of distinguishing metrics is to calculate (some of) their scalar polynomial curvature invariants due to the fact that inequivalent invariants implies inequivalent metrics. However, if their scalar polynomial invariants are the same, what

conclusion can we draw about the (in)equivalence of the metrics? For example, if all such invariants are zero, can we say that the metric is flat? The answer to this question is known to be no, because all so-called VSI metrics have vanishing scalar invariants. Here, we will address the more general question: if two spacetimes have identical scalar polynomial curvature invariants, what can we say about these spacetimes? In particular, when do the invariants characterise the spacetime metric?

For a spacetime (\mathcal{M}, g) with a set of scalar polynomial curvature invariants, there are two conceivable ways in which the metric g can be altered such that the invariants remain the same. First, the metric can be continuously deformed in such a way that the invariants remain unchanged. This is what happens for the Kundt metrics for which we have free functions which do not affect the curvature invariants. Alternatively, a discrete transformation of the metric can leave the invariants the same. A simple example of when a discrete transformation can give another metric with the same set of invariants is the pair of metrics:

$$ds_1^2 = \frac{1}{z^2} (dx^2 + dy^2 + dz^2) - d\tau^2, \quad (1)$$

$$ds_2^2 = \frac{1}{z^2} (-dx^2 + dy^2 + dz^2) + d\tau^2. \quad (2)$$

One can straight-forwardly check that these metrics have identical invariants but are not diffeomorphic (over the reals). These discrete transformations are more difficult to deal with but the issue will be taken up in a later section.

Therefore, first we will consider the continuous metric deformations defined as follows.

Definition 1.1. For a spacetime (\mathcal{M}, g) , a (one-parameter) *metric deformation*, \hat{g}_τ , $\tau \in [0, \epsilon)$, is a family of smooth metrics on \mathcal{M} such that

1. \hat{g}_τ is continuous in τ ,
2. $\hat{g}_0 = g$; and
3. \hat{g}_τ for $\tau > 0$, is not diffeomorphic to g .

For any given spacetime (\mathcal{M}, g) we define the set of all scalar polynomial curvature invariants

$$\mathcal{I} \equiv \{R, R_{\mu\nu}R^{\mu\nu}, C_{\mu\nu\alpha\beta}C^{\mu\nu\alpha\beta}, R_{\mu\nu\alpha\beta;\gamma}R^{\mu\nu\alpha\beta;\gamma}, R_{\mu\nu\alpha\beta;\gamma\delta}R^{\mu\nu\alpha\beta;\gamma\delta}, \dots\}.$$

Therefore, we can consider the set of invariants as a function of the metric and its derivatives. However, we are interested in to what extent, or under what circumstances, this function has an inverse.

Definition 1.2. Consider a spacetime (\mathcal{M}, g) with a set of invariants \mathcal{I} . Then, if there does not exist a metric deformation of g having the same set of invariants as g , then we will call the set of invariants *non-degenerate*. Furthermore, the spacetime metric g , will be called *\mathcal{I} -non-degenerate*.

This implies that for a metric which is \mathcal{I} -non-degenerate the invariants characterize the spacetime uniquely, at least locally, in the space of (Lorentzian) metrics. This means that these metrics are characterized by their curvature invariants and therefore we can distinguish such metrics using their invariants. Since scalar curvature invariants are manifestly diffeomorphism-invariant we can thereby avoid the difficult issue whether a diffeomorphism exists connecting two spacetimes.

2 Main Theorems

Let us first state our main theorems which will be proven in the later sections. The theorems apply to four-dimensional (4D) Lorentzian manifolds. Such spacetimes are characterized algebraically by their Petrov [1, 2] and Segre [3, 2] types or, equivalently, in terms of their Ricci, Weyl (and Riemann) types [4, 5, 6]. The notation, which essentially follows that of the cited references, is briefly summarized in Appendix A. The proofs of these theorems, which are investigated on a case by case basis in terms of the algebraic type of the curvature tensors, are long and tedious and have therefore been placed in later sections. Once all of the various cases have been explored the theorems follow.

Furthermore, let us remark on the technical assumptions made in this paper. The following theorems hold on neighborhoods where the Riemann, Weyl and Segre types do not change. In the algebraically special cases we also need to assume that the algebraic type of the higher-derivative curvature tensors also do not change, up to the appropriate order. Most crucial is the definition of the curvature operators (see later) and in order for these to be well defined, the algebraic properties of the curvature tensors need to remain the same over a neighborhood.¹ Henceforth, we will therefore assume that we consider an open neighborhood where the algebraic properties of the curvature tensors do not change, up to the appropriate order (≤ 4).

The first theorem deals with the algebraic classification of the curvature tensors, and the relation to the \mathcal{I} -non-degenerate metrics.

Theorem 2.1 (Algebraically general). *If a spacetime metric is of Ricci type I , Weyl type I , or Riemann type I/G , the metric is \mathcal{I} -non-degenerate.*

This theorem indicates that the general metric is \mathcal{I} -non-degenerate and thus the metric is determined by its curvature invariants (at least locally, in the sense explained above). In the above, by Riemann type I/G , we are referring to the existence of a frame in which components of boost weight $+2$ vanish for Riemann type I , and in type G there does not exist a frame in which components with boost weight $+2$ or -2 vanish, in this case the Weyl and Ricci canonical frames are not aligned. For the algebraically special spacetimes, we need to consider covariant derivatives of the Riemann tensor.

Theorem 2.2 (Algebraically special). *If the spacetime metric is algebraically special, but ∇R , $\nabla^{(2)}R$, $\nabla^{(3)}R$, or $\nabla^{(4)}R$ is of type I or more general, the metric is \mathcal{I} -non-degenerate.*

In terms of the boost weight decomposition, an algebraically special metric has a Riemann tensor with zero positive boost weight components. In general, type I refers to the vanishing of boost weight components $+2$ and higher (but not boost weight $+1$ components). For example, we often use the notation $(\nabla R)_b = 0$, $b \geq 2$ to denote a ∇R of type I (but $(\nabla R)_1 \neq 0$). The above theorem indicates that if by taking covariant derivatives of the Riemann tensor you acquire positive boost weight components, then the metric is \mathcal{I} -non-degenerate. The remaining metrics which *do not* acquire a positive boost weight component when taking covariant derivatives, have a very special structure of their curvature tensors. Indeed, such metrics must be very special metrics:

¹Alternatively, we can assume that the spacetime is real analytic.

Theorem 2.3. *Consider a spacetime metric. Then either,*

1. *the metric is \mathcal{I} -non-degenerate; or,*
2. *the metric is contained in the Kundt class.*

This is a striking result because it tells us that metrics not determined by their curvature invariants must be of Kundt form. These Kundt metrics therefore correspond to degenerate metrics in the sense that many such spacetimes can have identical invariants. The Kundt class is defined by those metrics admitting a null vector ℓ that is geodesic, expansion-free, shear-free and twist-free (corresponding to the vanishing of the spin-coefficients κ , σ and ρ ; see also Appendix A)

$$\ell^\beta \ell_{\alpha;\beta} = 0 \quad \ell_\alpha{}^{;\alpha} = 0 \quad \ell^{(\alpha;\beta)} \ell_{\alpha;\beta} = 0 \quad \ell^{[\alpha;\beta]} \ell_{\alpha;\beta} = 0. \quad (3)$$

Any metric in the Kundt class can be written in the following canonical form [7, 4]:

$$ds^2 = 2du [dv + H(v, u, x^k)du + W_i(v, u, x^k)dx^i] + g_{ij}(u, x^k)dx^i dx^j. \quad (4)$$

For spacetimes with constant curvature invariants (CSI) Theorem 2.3 has an important consequence. For CSI metrics, \mathcal{I} -non-degenerate implies that the spacetime is curvature homogeneous to all orders; hence, an important corollary is a proof of the CSI-Kundt conjecture [7]:

Corollary 2.4 (CSI spacetimes). *Consider a 4-dimensional spacetime having all constant curvature invariants (CSI). Then either,*

1. *the spacetime is locally homogeneous; or,*
2. *a subclass of the Kundt spacetimes.*

These theorems imply that the Kundt spacetimes play a pivotal role in the question of which metrics are \mathcal{I} -non-degenerate. Indeed, the Kundt metrics are the *only metrics not determined by their curvature invariants* (in the sense explained above).

In fact, we can be somewhat more precise since only a subclass of the Kundt spacetimes have these exceptional properties. In the analysis (described below) it is found that a Kundt metric is \mathcal{I} -non-degenerate if the metric functions $W_i(v, u, x^k)$ in the canonical (*kinematic*) Kundt null frame are non-linear in v (i.e., $W_{i,vv} \neq 0$). Hence the exceptional spacetimes are the *aligned algebraically special type-II-Kundt* spacetimes or, in short (and consistent with the terminology of the above theorem) *degenerate Kundt* spacetimes, in which there exists a common null frame in which the geodesic, expansion-free, shear-free and twist-free null vector ℓ is also the null vector in which all positive boost weight terms of the Riemann tensor are zero (i.e., the kinematic Kundt frame and the Riemann type II aligned null frame are *aligned*). We note that the important Kundt-CSI and vanishing scalar invariant (VSI) spacetimes are degenerate Kundt spacetimes [8, 9, 10, 7].

3 Curvature operators and curvature projectors

In order to prove the main theorems we need to introduce some mathematical tools. These tools, although they are very simple, are extremely useful and powerful in proving these theorems.

A curvature operator, \mathbb{T} , is a tensor considered as a (pointwise) linear operator

$$\mathbb{T} : V \mapsto V,$$

for some vector space V , constructed from the Riemann tensor, its covariant derivatives, and the curvature invariants.

The archetypical example of a curvature operator is obtained by raising one index of the Ricci tensor. The Ricci operator is consequently a mapping of the tangent space $T_p\mathcal{M}$ into itself:

$$\mathbb{R} \equiv (R^\mu{}_\nu) : T_p\mathcal{M} \mapsto T_p\mathcal{M}.$$

Another example of a curvature operator is the Weyl tensor, considered as an operator, $\mathbb{C} \equiv (C^{\alpha\beta}{}_{\mu\nu})$, mapping bivectors onto bivectors.

For a curvature operator, \mathbb{T} , consider an eigenvector \mathbf{v} with eigenvalue λ ; i.e., $\mathbb{T}\mathbf{v} = \lambda\mathbf{v}$. If $d = \dim(V)$ and n is the dimension of the spacetime, then the eigenvalues of \mathbb{T} are $GL(d)$ invariant. Since the Lorentz transformations, $O(1, n-1)$, will act via a representation $\Gamma \subset GL(d)$ on \mathbb{T} , *the eigenvalues of a curvature operator is an $O(1, n-1)$ -invariant curvature scalar*. Therefore, curvature operators naturally provide us with a set of curvature invariants (not necessarily polynomial invariants) corresponding to the set of distinct eigenvalues: $\{\lambda_A\}$. Furthermore, the set of eigenvalues are uniquely determined by the polynomial invariants of \mathbb{T} via its characteristic equation. The characteristic equation, when solved, gives us the set of eigenvalues, and hence these are consequently determined by the invariants.²

We can now define a number of associated curvature operators. For example, for an eigenvector \mathbf{v}_A so that $\mathbb{T}\mathbf{v}_A = \lambda_A\mathbf{v}_A$, we can construct the annihilator operator:

$$\mathbb{P}_A \equiv (\mathbb{T} - \lambda_A\mathbb{1}).$$

Considering the Jordan block form of \mathbb{T} , the eigenvalue λ_A corresponds to a set of Jordan blocks. These blocks are of the form:

$$\mathbb{B}_A = \begin{bmatrix} \lambda_A & 0 & 0 & \cdots & 0 \\ 1 & \lambda_A & 0 & \ddots & \vdots \\ 0 & 1 & \lambda_A & \ddots & 0 \\ \vdots & \ddots & \ddots & \ddots & 0 \\ 0 & \cdots & 0 & 1 & \lambda_A \end{bmatrix}.$$

There might be several such blocks corresponding to an eigenvalue λ_A ; however, they are all such that $(\mathbb{B}_A - \lambda_A\mathbb{1})$ is nilpotent and hence there exists an $n_A \in \mathbb{N}$ such that $\mathbb{P}_A^{n_A}$ annihilates the whole vector space associated to the eigenvalue λ_A .

² Note that the 'corresponding eigenvalues' of the operators constructed from the covariant derivatives of the Riemann tensor are also related to scalar curvature invariants built from covariant derivatives.

This implies that we can define a set of operators $\tilde{\perp}_A$ with eigenvalues 0 or 1 by considering the products

$$\prod_{B \neq A} P_B^{n_B} = \Lambda_A \tilde{\perp}_A,$$

where $\Lambda_A = \prod_{B \neq A} (\lambda_A - \lambda_B)^{n_B} \neq 0$ (as long as $\lambda_B \neq \lambda_A$ for all B). Furthermore, we can now define

$$\perp_A \equiv 1 - \left(1 - \tilde{\perp}_A\right)^{n_A}$$

where \perp_A is a *curvature projector*. The set of all such curvature projectors obeys:

$$1 = \perp_1 + \perp_2 + \cdots + \perp_A + \cdots, \quad \perp_A \perp_B = \delta_{AB} \perp_A. \quad (5)$$

We can use these curvature projectors to decompose the operator \mathbb{T} :

$$\mathbb{T} = \mathbb{N} + \sum_A \lambda_A \perp_A. \quad (6)$$

The operator \mathbb{N} therefore contains all the information not encapsulated in the eigenvalues λ_A . From the Jordan form we can see that \mathbb{N} is nilpotent; i.e., there exists an $n \in \mathbb{N}$ such that $\mathbb{N}^n = 0$. In particular, if $\mathbb{N} \neq 0$, then \mathbb{N} is a negative/positive boost weight operator which can be used to lower/raise the boost weight of a tensor.

Considering the Ricci operator, or the Weyl operator, we can show that (where the type refers to either Ricci type or Weyl type):

- Type I: $\mathbb{N} = 0$, $\lambda_A \neq 0$.
- Type D: $\mathbb{N} = 0$, $\lambda_A \neq 0$.
- Type II: $\mathbb{N}^3 = 0$, $\lambda_A \neq 0$.
- Type III: $\mathbb{N}^3 = 0$, $\lambda_A = 0$.
- Type N: $\mathbb{N}^2 = 0$, $\lambda_A = 0$.
- Type O: $\mathbb{N} = 0$, $\lambda_A = 0$.

Consider a curvature projector $\perp : T_p \mathcal{M} \mapsto T_p \mathcal{M}$. Then, for a Lorentzian spacetime there are 4 categories:

1. Timelike: For all $v^\mu \in T_p \mathcal{M}$, $v_\nu (\perp)^\nu_\mu v^\mu \leq 0$.
2. Null: For all $v^\mu \in T_p \mathcal{M}$, $v_\nu (\perp)^\nu_\mu v^\mu = 0$.
3. Spacelike: For all $v^\mu \in T_p \mathcal{M}$, $v_\nu (\perp)^\nu_\mu v^\mu \geq 0$.
4. None of the above.

In the following, we shall consider a complete set of curvature projectors: $\perp_A : T_p\mathcal{M} \mapsto T_p\mathcal{M}$. These projectors can be of any of the aforementioned categories and we are going to use the Segre-like notation to characterize the set with a comma separating time and space. For example, $\{1, 111\}$ means we have 4 projectors: one timelike, and three spacelike. A bracket indicates that the image of the projectors are of dimension 2 or higher; e.g., $\{(1, 1)11\}$ means that we have two spacelike operators, and one with a 2 dimensional image. If there is a null projector, we automatically have a second null projector. Given an NP frame $\{\ell_\mu, n_\mu, m_\mu, \bar{m}_\mu\}$, then a null-projector can typically be:

$$(\perp_1)^\mu{}_\nu = -\ell^\mu n_\nu.$$

Note that $\perp_1^2 = \perp_1$, but it is not symmetric. Therefore, acting from the left and right gives two different operators. Indeed, defining

$$(\perp_2)^\mu{}_\nu \equiv g_{\nu\alpha} g^{\mu\beta} (\perp_1)^\alpha{}_\beta,$$

we get a second null-projector being orthogonal to \perp_1 . The existence of null-projectors enables us to pick out certain null directions; however, note that the null-operators, with respect to the aforementioned Newman-Penrose (NP) frame, are of boost weight 0 and so cannot be used to lower/raise the boost weights. In particular, considering the combination $\perp_1 + \perp_2$ we see that the existence of null-projectors implies the existence of projectors of type $\{(1, 1)(11)\}$.

The existence of curvature projectors is important due to the following result:

Theorem 3.1. *Consider a spacetime metric and assume that there exist curvature projectors of type $\{1, 111\}$, $\{1, 1(11)\}$ or $\{1, (111)\}$. Then the spacetime is \mathcal{I} -non-degenerate.*

Proof. Consider first the case $\{1, 111\}$. For any given curvature tensor, $R_{\alpha\beta\dots\delta}$, we can construct the curvature tensor

$$R[ij\dots k]_{\alpha\beta\dots\delta} \equiv R_{\mu\nu\dots\lambda} (\perp_i)^\mu{}_\alpha (\perp_j)^\nu{}_\beta \dots (\perp_k)^\lambda{}_\delta.$$

This enables us to consider the curvature invariant $R[ij\dots k]_{\alpha\beta\dots\delta} R[ij\dots k]^{\alpha\beta\dots\delta}$ which is, up to a constant factor, the square of the component $R_{ij\dots k}$. This implies that it is determined by the invariant (up to a sign) and we get that *the spacetime is \mathcal{I} -non-degenerate*.

Consider now the case $\{1, 1(11)\}$. We note that in this case we cannot isolate all components of the curvature tensors. However, we can uniquely define tensors $r_{IJ\dots K}^{(A)}$, $I, J, \dots = 3, 4$ by contractions with \perp_i . The curvature invariants will now be $SO(2)$ -invariant polynomials in the components of $r_{IJ\dots K}^{(A)}$. Hence, since $SO(2)$ is compact, the polynomials will separate the $SO(2)$ orbits. Hence, by a similar proof as in [11] we get that *the spacetime is \mathcal{I} -non-degenerate*.

Lastly, consider the case $\{1, (111)\}$. In this case we can define tensors $r_{IJ\dots K}^{(A)}$, $I, J, \dots = 2, 3, 4$ by contractions with \perp_i . The curvature invariants will be $SO(3)$ -invariant which is again compact. Hence, using a similar argument as in [11] we get that *the spacetime is \mathcal{I} -non-degenerate*. \square

4 Riemann type I/G

Let us consider first the case where the Riemann tensor is of type I or G . This corresponds to the three cases: Ricci type I , Weyl (Petrov) type I , and Ricci and Weyl canonical frames not aligned. We shall consider these in turn.

4.1 Ricci type I

This case consists of the following Segre types: $\{1, 111\}$, $\{1, 1(11)\}$, $\{1, (111)\}$, $\{z\bar{z}11\}$, $\{z\bar{z}(11)\}$.

4.1.1 Segre type $\{1, 111\}$:

Here the eigenvalues of the Ricci operator are all distinct and we can diagonalize the Ricci operator:

$$R = \text{diag}(\lambda_1, \lambda_2, \lambda_3, \lambda_4).$$

It now follows from Theorem 3.1 that *the spacetime is \mathcal{I} -non-degenerate*.

Indeed, to determine the spacetime it is sufficient to consider $R_{\mu\nu;\alpha}$. Choosing an orthonormal frame, \mathbf{e}_i , aligned with the eigendirections of R :

$$R_{ij;k} = \lambda_{i,k}g_{ij} + (\lambda_i - \lambda_j)\Gamma_{ijk},$$

where Γ_{ijk} are the connection coefficients, we find that all connection coefficients must be determined by the curvature invariants.

4.1.2 Segre type $\{1, 1(11)\}$:

This is the special case of above where we have $\lambda_3 = \lambda_4$. Using Theorem 3.1 *the spacetime is \mathcal{I} -non-degenerate*.

4.1.3 Segre type $\{1, (111)\}$:

Here we have $\lambda_2 = \lambda_3 = \lambda_4$ and from Theorem 3.1 we have that *the spacetime is \mathcal{I} -non-degenerate*.

4.1.4 Segre type $\{z\bar{z}11\}$ and $\{z\bar{z}(11)\}$:

In this case, the Ricci operator has two complex conjugate eigenvalues. We can always find an orthonormal frame $\{\mathbf{e}_i\}$, so that the Ricci operator takes the form

$$R = \begin{bmatrix} a & b & 0 & 0 \\ -b & a & 0 & 0 \\ 0 & 0 & \lambda_3 & 0 \\ 0 & 0 & 0 & \lambda_4 \end{bmatrix}.$$

We can now consider the complex transformation mapping the basis vectors \mathbf{e}_0 and \mathbf{e}_1 onto the eigenvectors \mathbf{v}_0 and \mathbf{v}_1 :

$$\mathbf{v}_0 = \frac{1}{\sqrt{2}}(\mathbf{e}_0 + i\mathbf{e}_1), \quad \mathbf{v}_1 = \frac{1}{\sqrt{2}}(i\mathbf{e}_0 + \mathbf{e}_1), \quad (7)$$

with inverse

$$\mathbf{e}_0 = \frac{1}{\sqrt{2}}(\mathbf{v}_0 - i\mathbf{v}_1), \quad \mathbf{e}_1 = \frac{1}{\sqrt{2}}(-i\mathbf{v}_0 + \mathbf{v}_1). \quad (8)$$

We note that $\mathbf{v}_0 \cdot \mathbf{v}_0 = -1$, $\mathbf{v}_1 \cdot \mathbf{v}_1 = 1$, $\mathbf{v}_0 \cdot \mathbf{v}_1 = 0$ and so the set $\{\mathbf{v}_0, \mathbf{v}_1, \mathbf{e}_2, \mathbf{e}_3\}$ can be considered as an orthonormal frame. In this frame the Ricci operator becomes diagonal:

$$R = \text{diag}(\lambda_1, \bar{\lambda}_1, \lambda_2, \lambda_3).$$

Therefore, we have a set of curvature projectors of the form $\{1, 111\}$ or $\{1, 1(11)\}$ and we can use Theorem 3.1. The only difference is that the invariants associated to the complex frame can now be complex; however, the result still stands. Using the inverse transformation, which induces a transformation between the invariants from the complex frame to the real frame, we obtain the curvature components of the real frame. Therefore we can conclude that *the spacetime is \mathcal{I} -non-degenerate*.

4.2 Weyl type I (Petrov type I)

For the Weyl tensor any non-trivial isotropy would make it algebraically special. The isotropy group of the Weyl tensor is the subgroup of the Lorentz group whose action on the Weyl tensor leaves it invariant; for example a Petrov type D Weyl tensor has a boost-spin isotropy group. So for the Weyl tensor to be of type I requires that the isotropy group is trivial. We therefore expect that we will be able to determine a unique frame using the curvature invariants.

We use the bivector formalism and write the Weyl tensor, $C_{\alpha\beta\mu\nu}$, as an operator in 6-dimensional bivector space, $C = (C^A_B)$. Using the following index convention:

$$[23] \leftrightarrow 1, [31] \leftrightarrow 2, [12] \leftrightarrow 3, [10] \leftrightarrow 4, [20] \leftrightarrow 5, [30] \leftrightarrow 6,$$

a type I Weyl tensor can always be put into the following canonical form [3]:

$$C = \begin{bmatrix} a_1 & 0 & 0 & b_1 & 0 & 0 \\ 0 & a_2 & 0 & 0 & b_2 & 0 \\ 0 & 0 & a_3 & 0 & 0 & b_3 \\ -b_1 & 0 & 0 & a_1 & 0 & 0 \\ 0 & -b_2 & 0 & 0 & a_2 & 0 \\ 0 & 0 & -b_3 & 0 & 0 & a_3 \end{bmatrix} \quad (9)$$

where $\sum_i a_i = \sum_i b_i = 0$ and not all of the a_i, b_i are zero.

First we note that the eigenvalues of C are $a_i \pm ib_i$. As explained above, a_i and b_i are uniquely determined by the zeroth order Weyl invariants. The eigenvectors are $F^A = \delta_i^A \pm i\delta_{3+i}^A$. We can therefore construct annihilator operators, $(C - \lambda 1)$, and projection operators as before (the only difference is that C is 6-dimensional). The eigenvectors correspond to (complex) antisymmetric tensors. For example, consider the eigenvector with eigenvalue $a_1 + ib_1$:

$$F = \frac{1}{2} F_{\mu\nu} \omega^\mu \wedge \omega^\nu = \omega^2 \wedge \omega^3 - i\omega^1 \wedge \omega^0.$$

Hence, from this we can construct an operator

$$P_1 = (F^\mu_\alpha \bar{F}^\alpha_\nu) = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & -1 \end{bmatrix}. \quad (10)$$

For the other eigenvectors we then get (analogously):

$$P_2 = \text{diag}(1, -1, 1, -1), \quad P_3 = \text{diag}(1, -1, -1, 1).$$

Thus the linear set $\{1, P_i\}$ span all diagonal matrices; in particular, we can construct the projection operators:

$$\begin{aligned} \perp_1 &= \frac{1}{4}(1 + P_1 + P_2 + P_3), \\ \perp_2 &= \frac{1}{4}(1 + P_1 - P_2 - P_3), \\ \perp_3 &= \frac{1}{4}(1 - P_1 + P_2 - P_3), \\ \perp_4 &= \frac{1}{4}(1 - P_1 - P_2 + P_3). \end{aligned}$$

It is clear that we will get 3 operators, P_i , as long as the 3 sets of complex eigenvalues, $\lambda_i = a_i + ib_i$, are all different. Since $\sum_i \lambda_i = 0$, this can only fail when:

$$\begin{aligned} \lambda_1 = \lambda_2, \quad \lambda_3 = -2\lambda_1, \\ \lambda_1 = \lambda_2 = \lambda_3 = 0. \end{aligned}$$

The first of these is actually Weyl type D , while the latter is Weyl type O ; hence, these are excluded by assumption.

Therefore, we can conclude that as long as the Weyl type is I (and not simpler), we can define 4 projection operators of type $\{1, 111\}$. Therefore, from Theorem 3.1, *the spacetime is \mathcal{I} -non-degenerate*.

At this stage we wish to remark on a certain subtlety in the choice of eigenvectors. From the Weyl tensor we can actually only determine the product $F_{\mu\nu}F_{\alpha\beta}$. Therefore, we can only construct the ‘‘square’’ $P_1 \otimes P_1$. So in order to get the operator P_1 there is an ambiguity in the choice of sign. Regarding the question of \mathcal{I} -non-degeneracy as defined above this has no consequence; however, it may have an effect on discrete changes to the metric. This sign ambiguity results in a permutation of the axes; essentially, we don’t know which axis corresponds to time. We will get back to this issue later but note that this phenomenon will recur in several cases below.

4.3 Ricci and Weyl canonical frames not aligned

Consider now the case where both the Ricci tensor and the Weyl tensor are algebraically special but where there does not exist a null-frame such that both the Ricci tensor and the Weyl tensor has only non-positive boost weights.

First, assume the Weyl type is D and choose the Weyl canonical frame. For Weyl type D the Weyl operator is of the form of eq. (9) with $\lambda_1 = \lambda_2$, $\lambda_3 = -2\lambda_1$. This immediately implies we have projection operators of type $\{(1, 1)(11)\}$.

In the Weyl canonical frame, the Ricci tensor must have both positive *and* negative boost weight components (or else there would exist a frame where they are aligned). Now, by symmetry, we can consider three cases for the Ricci tensor (see Appendix A for $(R)_b$ notation):

1. $(R)_{+2} \neq 0$, $(R)_{-2} \neq 0$: Here, we use the $(1, 1)$ -projection operator and we get a reduced Ricci operator of the form (in the $\{\ell, n\}$ frame):

$$\tilde{R} = \begin{bmatrix} a & b \\ c & a \end{bmatrix}, \quad bc \neq 0. \quad (11)$$

This gives two distinct eigenvalues $\lambda = a \pm \sqrt{bc}$, and hence, two additional projection operators. This case therefore reduces to the case $\{1, 1(11)\}$ or $\{z\bar{z}(11)\}$ presented earlier. This spacetime is therefore \mathcal{I} -non-degenerate.

2. $(R)_{+2} = 0, (R)_{+1} \neq 0, (R)_{-2} \neq 0$: For this case we note that the square $R^\mu_\alpha R^\alpha_\nu$ necessarily has boost weight $+2$ components. Therefore, using either $R^\mu_\alpha R^\alpha_\nu$ or $R^\mu_\alpha R^\alpha_\nu + qR^\mu_\nu$, where q is a parameter, we can use the results of the previous paragraph. This case is consequently \mathcal{I} -non-degenerate.
3. $(R)_{\pm 2} = 0, (R)_{+1} \neq 0, (R)_{-1} \neq 0$: Here we consider $R^\mu_\alpha R^\alpha_\nu$ which necessarily has non-zero boost weight -2 and $+2$ components. This case is therefore also \mathcal{I} -non-degenerate.

Assume now Weyl type N and choose the Weyl canonical frame such that $C = (C)_{-2}$. In this frame, either $(R)_{+1}$ or $(R)_{+2}$ is non-zero. If $(R)_{+2}$ is zero, we replace the Ricci tensor $R_{\mu\nu}$ with the square $R^\mu_\alpha R^\alpha_\nu$ in what follows. Therefore, assume that $(R)_{+2} \neq 0$. Consider now the operator

$$(T^\mu_\nu) \equiv C^\mu_{\alpha\nu\beta} R^{\alpha\beta}$$

Under the above assumptions, this operator can be used to get projectors of type $\{(1, 1)11\}$. Indeed, these projectors are aligned with the Weyl canonical frame. We can now use one of the spacelike projectors, \perp_3 (say), to construct the symmetric operator:

$$(\hat{T}^\mu_\nu) \equiv C^{\mu\alpha}_{\nu\beta} (\perp_3)^\beta_\alpha + qR^\mu_\nu,$$

where q is a parameter. We can use this operator to construct the remaining projection operators so that we have a set $\{1, 111\}$. This case is therefore \mathcal{I} -non-degenerate.

For Weyl type II we can decompose the Weyl tensor:

$$C = N + \sum_A \lambda_A \perp_A,$$

where the operator N is a ‘‘Weyl’’ operator of type N while the piece $\sum_A \lambda_A \perp_A$ is a ‘‘Weyl’’ operator of type D . By assumption, the Ricci tensor is not aligned with the Weyl canonical frame; therefore, using the above results, this case is also \mathcal{I} -non-degenerate.

Lastly, for Weyl type III , we can consider the square C^2 which is a Weyl operator of type N . The above results imply that this case is \mathcal{I} -non-degenerate.

4.3.1 Summary

Therefore, we have shown that: *If a 4-dimensional spacetime (\mathcal{M}, g) is either Ricci type I , Weyl type I or Riemann type I/G , then it is \mathcal{I} -non-degenerate.*

5 Algebraically special cases

For the algebraically special cases the Riemann tensor itself does not give enough information to provide us with all the required projection operators. Indeed,

in the algebraically special cases it is also necessary to calculate the covariant derivatives. The strategy is as follows: we will consider the two cases of Weyl type D and N in detail. The second Bianchi identity will not be imposed at this time because we aim to use these results on more general tensors with the same symmetries, not necessarily the Weyl tensor itself. Weyl type II and III will now follow from these computations and Weyl type O will be treated last.

We should also point out that for any symmetric tensor $S_{\mu\nu}$ we can always construct a Weyl-like tensor with the same symmetries as the Weyl tensor. If $S_{\mu\nu}$ is the trace-free Ricci tensor, the corresponding Weyl-like tensor is the so-called Plebański tensor. Explicitly, given the trace-free part of $S_{\mu\nu}$, denoted $\hat{S}_{\mu\nu}$, the Plebański tensor is given by

$$W^{\alpha\beta}{}_{\mu\nu} \equiv \hat{S}^{\alpha}{}_{[\mu} \hat{S}^{\beta]}{}_{\nu]} + \delta^{\alpha}{}_{[\mu} \hat{S}_{\nu]\gamma} \hat{S}^{\beta]\gamma} - \frac{1}{6} \delta^{\alpha}{}_{[\mu} \delta^{\beta]}{}_{\nu]} \hat{S}_{\gamma\epsilon} \hat{S}^{\gamma\epsilon}. \quad (12)$$

Therefore, to any symmetric tensor there is an associated ‘‘Plebański’’ tensor.

Henceforth we are going to use the NP-formalism where we introduce a null frame $\{\ell, n, m, \bar{m}\}$. (We will use the notation of [2]; also see Appendix A). In order to get the desired results we introduce the canonical frames for the various algebraic types. For the Weyl tensor, C , this means that we express its components in terms of the Weyl scalars Ψ_i . Then using the NP-connection coefficients, we can express the covariant derivative ∇C in terms of Ψ_i and the connection coefficients. At this stage it is useful not to assume anything about the connection ∇ (i.e., the tensor C need not be the Weyl tensor of the connection). The advantage of this is that we can utilise the full formalism of projection operators without worrying about the compatibility of the Weyl tensor and the connection. Furthermore, the results obtained here will therefore be more general than what is indicated. These expressions are then utilised to obtain the required results for the curvature tensor. Another important thing to note is that when taking covariant derivatives, some of the components have terms which are partial derivatives of Ψ_i , while other terms are algebraic in Ψ_i and Γ_{jk}^i . These algebraic terms are most useful simply because they give algebraic relations rather than differential ones.

5.1 Weyl (Petrov) type D

We choose the canonical frame for which $\Psi_2 \neq 0$. From the Weyl operator C we can construct projectors of type $\{(1,1)(11)\}$ where the $(1,1)$ -projector projects onto the $\ell - n$ -plane, while the (11) -projector projects onto the $m - \bar{m}$ -plane. In the following let us use the indices $a, b, ..$ for projections onto the $\ell - n$ plane and the indices $i, j, ...$ for projections onto the $m - \bar{m}$ plane.

Calculating ∇C we get the boost weight decomposition

$$\nabla C = (\nabla C)_{+2} + (\nabla C)_{+1} + (\nabla C)_0 + (\nabla C)_{-1} + (\nabla C)_{-2}.$$

The key observation is that the positive boost weight components vanish if and only if $\ell^\mu \nabla_\mu \Psi_2 = 0$ and $\kappa = \sigma = \rho = 0$. Therefore, the idea is to define the appropriate operators so that we can isolate the necessary components.

Consider the (projected) tensor:

$$T_{iab} \equiv C^j{}_{ij(a;b)}$$

This tensor has the following structure,

$$T_{iab} = v_i n_a n_b + t_i (\ell_a n_b + n_a \ell_b) + w_i \ell_a \ell_b,$$

where

$$v_i \equiv 3(\bar{\Psi}_2 \bar{\kappa} m_i + \Psi_2 \kappa \bar{m}_i), \quad (13)$$

$$t_i \equiv -\frac{3}{2}(\pi \Psi_2 - \bar{\tau} \bar{\Psi}_2) m_i - \frac{3}{2}(\bar{\pi} \bar{\Psi}_2 - \tau \Psi_2) \bar{m}_i, \quad (14)$$

$$w_i \equiv -3(\Psi_2 \nu m_i + \bar{\Psi}_2 \bar{\nu} \bar{m}_i). \quad (15)$$

$$(16)$$

Furthermore, define the trace-free tensor $\hat{T}_{iab} \equiv T_{iab} + (1/2)(\ell_a n_b + n_a \ell_b) T_{i \ c}^c$, and then the tensor

$$S_{abcd} = \hat{T}_{ab}^i \hat{T}_{icd}.$$

This tensor can be considered as an operator $\mathbf{S} = (S_B^A)$ mapping symmetric trace-free tensors onto symmetric trace-free tensors. For simplicity, let us also consider the trace-free part of S_{abcd} so that

$$\hat{S}_{abcd} = v^i v_i n_a n_b n_c n_d + w^i w_i \ell_a \ell_b \ell_c \ell_d.$$

Consider the trace-free tensor $M^{ab} = x n^a n^b + y \ell^a \ell^b$. The operator $\hat{\mathbf{S}}$ has eigenvalues $\lambda = \pm |v| |w|$, 0. Therefore, as long as both v^i and w^j are non-zero, there are three distinct eigenvalues. Assuming $\lambda \neq 0$, M^{ab} is an eigentensor if $x = |v|$ and $y = |w|$. In this case we can consider the curvature projectors (up to scaling), $M^{ab} M_{cd}$. The eigentensor M^{ab} can again be considered as an operator $\mathbf{M} = (M_b^a)$ mapping vectors onto vectors. The eigenvalues of \mathbf{M} are $\lambda = \pm i |v| |w|$; hence, this reduces to the case of two complex eigenvalues.

We note that $v^i v_i = 0$ if and only if $\kappa = 0$. Furthermore, if either of $|v|$ or $|w|$ is non-zero we can assume, by using the discrete symmetry defined later by eq. (26), that $w^i w_i \neq 0$. Therefore, $\kappa \neq 0$ (so that $|v| \neq 0$ also) implies that the spacetime is *\mathcal{I} -non-degenerate*.

Therefore, assume $\kappa = 0$ and consider the symmetric tensor

$$Q_b^a = C^{ijka;l} C_{ijkb;l}.$$

The trace-free part of this tensor is

$$\hat{Q}_{ab} \propto |\Psi_2|^2 (|\sigma|^2 + |\rho|^2) n_a n_b + |\Psi_2|^2 (|\lambda|^2 + |\mu|^2) \ell_a \ell_b.$$

If $(|\sigma|^2 + |\rho|^2) (|\lambda|^2 + |\mu|^2) \neq 0$, then this tensor is of type *I*. So from the Ricci type *I* analysis, this would imply that the spacetime is *\mathcal{I} -non-degenerate*.

Let us next consider the non-aligned case where $\kappa = 0$, $\lambda = \mu = 0$ and $|\sigma|^2 + |\rho|^2 \neq 0$. We can now consider the mixed tensor:

$$\hat{Q}_{ab} + \hat{S}_{abcd} \hat{Q}^{cd}.$$

This tensor is of type *I* if $(w^i w_i) (|\sigma|^2 + |\rho|^2) \neq 0$ and consequently, the spacetime is *\mathcal{I} -non-degenerate*.

Assume now that $w^i = 0$, $\kappa = 0$, for which we still have an unused discrete symmetry (eq.(26)). If $(|\sigma|^2 + |\rho|^2) (|\lambda|^2 + |\mu|^2) = 0$ we can therefore assume that $\rho = \sigma = 0$. This spacetime is thus *Kundt*.

Lastly, consider the case when $w^i w_i \neq 0$, $\kappa = \rho = \sigma = 0$. This automatically implies that the spacetime is *Kundt*.

Let us also consider the differential $\ell^\mu \nabla_\mu \Psi_2$ which in general (not assuming the Bianchi identities are satisfied) also contributes to $(\nabla C)_{+1}$. We note that the Weyl invariant I , for a Weyl type D tensor, is given by $I = 3\Psi_2^2$. Therefore, we can consider the curvature tensor defined by the gradient $\nabla_\mu I = 6\Psi_2 \Psi_{2,\mu}$. We can now use the $(1,1)$ -projector and project this gradient onto the $\ell - n$ -plane: $x_a \equiv \nabla_a I$.

1. If x_a is either time-like or space-like (and consequently $\ell^\mu \nabla_\mu \Psi_2 \neq 0$), we can construct another curvature projector (by considering the operator $x^a x_b$) so that we have a set $\{1, 1(11)\}$. Therefore, this case is *\mathcal{I} -non-degenerate*.
2. If x_a is null or zero, then either $n^a x_a = 0$ or $\ell^a x_a = 0$. If $\ell^a x_a = 0$ then $\ell^\mu \nabla_\mu \Psi_2 = 0$ and does not contribute to positive boost weight components. Assume therefore that $\ell^a x_a \neq 0$, which implies that $x_a \propto n_a$. If $\nu = \lambda = \mu = 0$, we can use the discrete symmetry eq.(26) so that $\ell^\mu \nabla_\mu \Psi_2 = 0$.

If any of ν , λ or μ is non-zero, then $(\nabla C)_{b<0}$ is non-zero. Hence, by contracting with x_a , we can straight-forwardly construct another projection operator so that we get a set $\{1, 1(11)\}$. Therefore, this case is *\mathcal{I} -non-degenerate*.

5.1.1 Summary Weyl type D :

A Weyl type D spacetime is either *\mathcal{I} -non-degenerate* or *Kundt*. Moreover, for a Weyl type D spacetime, if ∇C is of type I or more general, then it is *\mathcal{I} -non-degenerate*.

5.2 Weyl (Petrov) type II

The Weyl type II tensor can be decomposed as

$$C = N + \sum_A \lambda_A \perp_A.$$

By using the annihilator operators and the projection operators we can, up to scaling, isolate each term in this decomposition. Each term can thus be considered a curvature operator in its own right.

In particular, by considering only the curvature tensor $\sum_A \lambda_A \perp_A$, this tensor is identical to a Weyl type D tensor. We can therefore use these results. In addition to these results we do have an additional boost weight -2 operator N . This breaks the discrete symmetry present in the Weyl type D tensor and therefore restricts the choice even more. However, with minor modifications we obtain: *a Weyl type II spacetime is either \mathcal{I} -non-degenerate or Kundt*.

We also note that for a Weyl type II spacetime, the Weyl invariant $I = 3\Psi_2^2$ as for type D . Therefore, using a similar argument, *a Weyl type II spacetime, if ∇C is of type I or more general, then it is \mathcal{I} -non-degenerate*.

5.3 Weyl (Petrov) type III

For the Weyl type III case we get no non-trivial curvature operators from the Weyl tensor itself. The first non-trivial projection operators appears at first covariant derivative; however, in order to delineate this case completely we need to consider second covariant derivatives.

We note that for the type III Weyl operator, $C^2 \neq 0$ and is of type N . The proof for this case is therefore contained in the Weyl type N case considered below.

5.4 Weyl (Petrov) type N

Consider first the tensor

$$T_{\nu}^{\mu} = \nabla_{\gamma} C^{\alpha\beta\gamma\mu} \nabla^{\delta} C_{\alpha\beta\delta\nu},$$

whose boost weight 0 components are of the form (it has no positive boost weight components)

$$(T_{\mu\nu})_0 \propto \bar{\kappa}^2 \bar{\Psi}_4^2 \bar{m}_{\mu} \bar{m}_{\nu} + \kappa^2 \Psi_4^2 m_{\mu} m_{\nu}.$$

Therefore, if $\kappa \neq 0$, we can construct curvature operators of type $\{(1, 1)11\}$. The curvature operator $T_{\mu\nu}$ gives rise to a ‘‘Plebański’’ tensor of type D . Therefore, by considering second covariant derivatives, it follows from the Weyl type D analysis that if $\kappa \neq 0$, the spacetime is \mathcal{I} -non-degenerate.

Henceforth, assume that $\kappa = 0$ (and therefore we have no projectors from first derivatives). Consider $\square C_{\mu\nu\alpha\beta}$, which has the same symmetries as the Weyl tensor itself. This tensor has no positive boost weight components. Considering the boost weight 0 components, we note that $\square C_{\mu\nu\alpha\beta}$ is of type II if and only if $\rho\sigma \neq 0$. Therefore, if $\rho\sigma \neq 0$ we can use the Weyl type II analysis, and calculate $\nabla\square C$; hence, this spacetime is \mathcal{I} -non-degenerate.

Therefore, consider the case where either σ or ρ are zero. Define

$$W_{\epsilon\eta}^{\mu\nu} \equiv C^{\alpha\beta\gamma\delta;(\mu\nu)} C_{\alpha\beta\gamma\delta;(\epsilon\eta)}.$$

To get a projection operator we note that the boost weight 0 components of $W^{\mu\alpha\beta\gamma} W_{\nu\alpha\beta\gamma}$ (it has no positive boost weight components) is of the form

$$(W^{\mu\alpha\beta\gamma} W_{\nu\alpha\beta\gamma})_0 \propto |\Psi_4|^4 |\rho \text{ or } \sigma|^8 (\bar{m}^{\mu} m_{\nu} + m^{\mu} \bar{m}_{\nu})$$

Therefore, if either ρ or σ are non-zero, we can use this operator and we get (at least) two curvature projectors \perp_1 and \perp_2 of type $\{(1, 1)(11)\}$. This means that we can construct a Weyl-like tensor of type D . Hence, we can use the type D results. Therefore, by considering third derivatives of the curvature tensors, if σ or ρ is non-zero, then the spacetime is \mathcal{I} -non-degenerate.

The remaining case, for which $\kappa = \rho = \sigma = 0$, is a *Kundt* spacetime.

5.4.1 Summary Weyl type III or N:

Therefore, we can conclude that a Weyl type III or N spacetime is either \mathcal{I} -non-degenerate or *Kundt*.

5.5 Algebraically special Ricci type

Using the trace-free Ricci tensor, we can construct the Plebański tensor, which is a Weyl-like tensor. The corresponding algebraic classification of the Plebański tensor is called the Plebański-Petrov (PP) classification. For the various algebraically special PP types we have the following Segre types:

- PP type *II*: $\{211\}$,
- PP type *D*: $\{(1,1)(11)\}$, $\{(1,1)11\}$, $\{2(11)\}$, $\{z\bar{z}(11)\}$, $\{1,1(11)\}$
- PP type *III*: $\{31\}$
- PP type *N*: $\{(31)\}$, $\{(21)1\}$
- PP type *O*: $\{(1,11)1\}$, $\{1,(111)\}$, $\{(211)\}$, $\{(1,111)\}$.

Now, since the Plebański tensor is a tensor with the same symmetries as the Weyl tensor, we can use the previous results for the PP types *II*, *D*, *III* and *N*. There is consequently only the Weyl (Petrov) type *O* and PP type *O* case left to consider.

5.6 Weyl type *O*, PP-type *O*

Let us consider the remaining Segre types assuming Weyl (Petrov) type *O* and PP type *O*.

5.6.1 Segre type $\{(1,11)1\}$

Using the Bianchi identities we get several differential constraints on the spin coefficients. For this Segre type we have that $\Phi_{11} \neq 0$, so the Bianchi identities immediately imply $\kappa + \bar{\kappa} = \nu + \bar{\nu} = 0$. In addition, we get the following restrictions:

$$\rho + \bar{\sigma} = s \quad (\text{real}), \quad \mu + \bar{\lambda} = m \quad (\text{real}).$$

Furthermore, after some manipulation of the remaining Bianchi identities, we get

$$\epsilon - \bar{\epsilon} = \gamma - \bar{\gamma} = \tau + \bar{\tau} + \pi + \bar{\pi} = 0,$$

and

$$DR = 24s\Phi_{11}, \quad \Delta R = -24m\Phi_{11}.$$

We now split the analysis into 3 cases, according to whether $\nabla_\mu R$ is timelike, spacelike or null.

If $\nabla_\mu R$ is timelike, we immediately have that this spacetime is \mathcal{I} -non-degenerate since we can use $(\perp_1)^\mu{}_\nu = (\nabla^\mu R)(\nabla_\nu R)$ as a timelike operator, and hence we obtain operators of type $\{1,1(11)\}$.

If $\nabla_\mu R$ is spacelike, we can always use the remaining freedom to choose $DR = \Delta R = 0$. This implies that $m = s = 0$. Furthermore, $\delta R - \bar{\delta} R \neq 0$ which means we have an additional spacelike projection operator. Therefore, we have a set $\{(1,1)11\}$, which can be used to give a ‘‘PP-type’’ D tensor. Hence, using second covariant derivatives, we find that this is either Kundt or \mathcal{I} -non-degenerate.

Lastly, $\nabla_\mu R$ is null. If $\nabla_\mu R$ is zero, from the Bianchi identities we find that this is a symmetric space, and hence, is actually locally homogeneous (and

Kundt). If $\nabla_\mu R$ is null, we consider $\delta\Phi_{11} - \bar{\delta}\Phi_{11}$. If this is non-zero we get an additional projection operator and thus a set $\{(1, 1)11\}$. This would therefore give a ‘‘PP-type’’ D and hence, by considering second derivatives, this is either Kundt or \mathcal{I} -non-degenerate. Therefore, let us assume $\delta\Phi_{11} - \bar{\delta}\Phi_{11} = 0$. The Bianchi identities now imply that $\alpha - \bar{\beta} = 0$. Using the symmetric operator $R^{\alpha\beta;\nu}R_{\alpha\beta;\mu}$ we get that this is either of types $\{2(11)\}$, $\{(1, 1)(11)\}$ or $\{(211)\}$. The first two of these give a type D ‘‘Plebański’’ tensor which means, by considering second derivatives, they are either \mathcal{I} -non-degenerate or Kundt. For the last case, $\{(211)\}$, we can combine with the Ricci operator to break the symmetry down to type $\{(21)1\}$ which gives a ‘‘Plebański’’ of type N. Therefore, by considering third and fourth derivatives, we get that this is \mathcal{I} -non-degenerate or Kundt.³

5.6.2 Segre type $\{1, (111)\}$

This is actually Ricci type I and is therefore \mathcal{I} -non-degenerate.

5.6.3 Segre type $\{(211)\}$

Choosing a frame where Φ_{22} is a constant we get, after using the Bianchi identities (and some manipulation), $\kappa = \sigma = 0$. We then calculate the second derivatives and compute the operator $R^{\alpha\beta;(\gamma\mu)}R_{\alpha\beta;(\gamma\nu)}$, which gives operators of type $\{(1, 1)(11)\}$, assuming $\rho \neq 0$. This again gives ‘‘PP-type’’ D tensor and hence, by calculating third derivatives, this is \mathcal{I} -non-degenerate or Kundt.

5.6.4 Segre type $\{(1, 111)\}$

This is the maximally symmetric case and is therefore Kundt.

In addition to Weyl-type I , Ricci-type I or Riemann-type I/G , we have shown that there are \mathcal{I} -non-degenerate metrics with algebraically special curvature types and further conditions on the spin-coefficients. These are summarized in Tables (1) and (2).

6 Curvature invariants

We have addressed the question of what is the class of Lorentzian manifolds that can be completely characterized by the scalar polynomial invariants constructed from the Riemann tensor and its covariant derivatives. Let us now consider the ‘inverse’ question: given a set of scalar polynomial invariants, what can we say about the underlying spacetime? In practice, it is somewhat tedious and a lengthy ordeal to determine the spacetime from the set of invariants. However, in most circumstances we only need some partial results or we are dealing with special cases. Let us discuss how to determine, from the invariants, whether the spacetime is \mathcal{I} -non-degenerate.

We remind the reader that the zeroth order Weyl invariants are I and J , and if all Weyl invariants up to order k vanish, we will denote this by VSI_k^W .

³Note that this is the only case in which we have needed to utilize fourth derivatives; it is possible, by explicitly calculating the components of $R_{\mu\nu;\alpha\beta\gamma}$, that we need only consider third derivatives, but we have not done this here.

P-type	Conditions
I	—
D or II	$\kappa \neq 0$ $\kappa = 0, (\sigma ^2 + \rho ^2)(\lambda ^2 + \mu ^2) \neq 0$ $\kappa = \lambda = \mu = 0, \sigma ^2 + \rho ^2 \neq 0$
N or III	$\kappa \neq 0$ $\kappa = 0; \sigma = 0 \text{ or } \rho = 0 \text{ (not both)}$

Table 1: In each Petrov-type we list conditions on the spin-coefficients yielding distinct subcases for which the metric is \mathcal{I} -non-degenerate.

Proposition 6.1. *If $27J^2 \neq I^3$, then the spacetime is \mathcal{I} -non-degenerate.*

This follows easily from the fact that if $27J^2 \neq I^3$ then the spacetime is of Weyl (Petrov) type I.

If $27J^2 = I^3 \neq 0$ (Weyl type II or D), we need to go to higher order invariants in order to check whether it is \mathcal{I} -non-degenerate or not. Ideally, we would like to have a set of syzygies which gives the appropriate condition for this to be the case. Such a complete set is not known. However, we have found two such syzygies which gives a *sufficient* condition for \mathcal{I} -non-degeneracy. A number of invariants of ∇C were constructed with degrees ranging from 2 to 4 (see Appendix C for details). Imposing the minimal number of conditions required for the normal form of a ∇C -type II (boost weight $+3, +2, +1$ vanish) or D (only boost weight 0 is nonzero) results in a degree 8 syzygy, $S_1 = 0$, and a degree 16 syzygy, $S_2 = 0$, amongst our invariants. Therefore if $S_1 \neq 0$ or $S_2 \neq 0$ then ∇C is not of type II or D . Next, we showed that using the normal form of a ∇C -type G (all components nonzero) or H (boost weight $+3$ vanish) or I (boost weight $+3, +2$ vanish) then $S_1 \neq 0$ and $S_2 \neq 0$. It is important to note that this implication refers only to the general types of G, H and I and there is no consideration of a secondary alignment type or any further algebraic specialization within these types. Indeed, it is possible that there is an algebraically special subcase, for example of a ∇C -type I , that results in $S_1 = S_2 = 0$. A stronger statement relating invariants of ∇C to its algebraic type may be achieved by considering a different basis of invariants and a finer algebraic classification of ∇C . Initially, one would attempt to construct a set of pure ∇C invariants that was complete within each ∇C algebraic type G, H, I and II , including special subcases. We have excluded type D since such a set of invariants is equivalent to type II , and also types III, N or O since these invariants vanish. By completeness of the set, an algebraic specialization would result in a dependence amongst invariants and hence syzygies arise characterizing the algebraically special type. We now have the following invariant characterizations of \mathcal{I} -non-degeneracy.

Proposition 6.2. *If $27J^2 = I^3$, but $S_1 \neq 0$ or $S_2 \neq 0$, then the spacetime is*

Segre type of $R_{\mu\nu}$	Conditions
$\{1, (111)\}$	—
$\{(211)\}$	Φ_{22} const., $\kappa = \sigma = 0$, $\rho \neq 0$ (3rd deriv.)
$\{(1, 11)1\}$	$\nabla_\mu R$ timelike
	$\nabla_\mu R$ spacelike : $DR = \Delta R = \kappa + \bar{\kappa} = \rho + \bar{\sigma} = \mu + \bar{\lambda} = 0$ (2nd deriv. ; at least one of κ , σ or ρ is nonzero)
	$\nabla_\mu R = 0$ (symmetric space)
	$\nabla_\mu R$ null : $\delta\Phi_{11} - \bar{\delta}\Phi_{11} \neq 0$ (2nd deriv.) $\nabla_\mu R \neq 0$: $\delta\Phi_{11} - \bar{\delta}\Phi_{11} = \alpha - \bar{\beta} = 0$ (2nd or 3rd and 4th deriv.)

Table 2: Within P-type O and PP-type O we list the Segre types that contain \mathcal{I} -non-degenerate metrics. The n th derivative conditions indicate that higher order constraints exist on the spin-coefficients arising from n th order curvature operators. These higher-order constraints provide sufficient conditions for the metric to be \mathcal{I} -non-degenerate. In all cases, at least one of κ , σ or ρ is nonzero.

\mathcal{I} -non-degenerate.

The remaining cases are when both I and J are zero, and hence, the spacetime is VSI_0^W :

Proposition 6.3. *Assume a spacetime is VSI_0^W . Then:*

1. *If it is not VSI_1^W , it is \mathcal{I} -non-degenerate.*
2. *If it is VSI_1^W , but not VSI_2^W , it is \mathcal{I} -non-degenerate.*

To prove the final result below, we shall assume for simplicity that the spacetime is Einstein, so that $R_{\mu\nu} = \lambda g_{\mu\nu}$. We therefore only have to consider the Ricci scalar ($= 4\lambda$) and the Weyl invariants. If this is not the case, then we would need to include the Ricci and mixed invariants. This can be done in a straight-forward manner. A summary of these results is given in Figure 1.

Proposition 6.4. *Assume a spacetime is Einstein. Then:*

3. *If it is VSI_2^W , then it is Kundt.*

From the above results we have conditions on the scalar invariants (in terms of the Weyl tensor and its covariant derivatives) to determine whether the spacetime is \mathcal{I} -non-degenerate. Consequently, we have a number of conditions in




Figure 1: Using invariants in terms of the Weyl tensor and its covariant derivatives to determine whether the spacetime is \mathcal{I} -non-degenerate.

terms of scalar invariants that can be used to determine when a spacetime is not \mathcal{I} -non-degenerate and hence an aligned algebraically special type-*II* (or *degenerate*) Kundt spacetime.

Let us further consider to what extent the class of degenerate Kundt spacetimes can be characterized by their scalar curvature invariants. Clearly such spacetimes are algebraically special and of type *II* (or more special) and hence $27J^2 = I^3$. If $I = J = 0$, then if the spacetime is of Weyl type *N*, then $\mathcal{I}_1 = \mathcal{I}_2 = 0$ if and only if $\kappa = \rho = \sigma = 0$ from the results in [8] (the definitions of the invariants \mathcal{I}_1 and \mathcal{I}_2 are given therein). Similar results follow for Weyl type *III* spacetimes (in terms of the invariants $\tilde{\mathcal{I}}_1$ and $\tilde{\mathcal{I}}_2$) and in the conformally flat (but non-vacuum) case (in terms of similar invariants \mathcal{I}_1 and \mathcal{I}_2 constructed from the Ricci tensor) [8]. If $27J^2 = I^3 \neq 0$ (Weyl types *II* and *D*): essentially if $\kappa = \rho = \sigma \neq 0$, we can construct positive boost weight terms in the derivatives of the curvature and determine an appropriate set of scalar

curvature invariants. For example, consider the positive boost weight terms of the first covariant derivative of the Riemann tensor, $\nabla(Riem)$. If the spacetime is \mathcal{I} -non-degenerate, then each component of $\nabla(Riem)$ is related to a scalar curvature invariant. In this case, in principle we can solve (for the positive boost weight components of $\nabla(Riem)$) to uniquely determine κ, ρ, σ in terms of scalar invariants, and we can therefore find necessary conditions for the spacetime to be degenerate Kundt (there are two cases to consider, corresponding to whether $\Psi_2 + \frac{2}{3}\Phi_{11}$ is zero or non-zero). We note that even if the invariants exist in principle, it may not be possible to construct them in practice.

7 Weakly and Strongly \mathcal{I} -non-degenerate

Until now we have only considered \mathcal{I} -non-degeneracy in terms of a local deformation of the metric. It is also of interest to know whether a \mathcal{I} -non-degenerate metric is unique under a discrete transformation. We shall call a spacetime such that the set of invariants *uniquely* specifies the metric *strongly \mathcal{I} -non-degenerate*. Similarly, we shall call a spacetime such that the set of invariants only defines a unique metric up to discrete transformations *weakly \mathcal{I} -non-degenerate*.

Let us revisit the examples given by eqs. (1) and (2) in the Introduction. These two examples are both of Weyl type O , but they are of Segre type $\{1, (111)\}$ and $\{(1, 11)1\}$. Hence, the eigenvalues of the Ricci operator is the same but we cannot, from the invariants alone, determine which eigenvalue is associated with the timelike direction and which is associated with the spacelike direction. This is linked to the fact that the map where we swap time with a space direction is not a Lorentz transformation. Note that permuting any two axes in the Riemannian-signature case is an $O(n)$ transformation, while permuting time and space in the Lorentzian case is *not* an $O(1, n-1)$ transformation. Therefore, there is no distinction between weakly and strongly \mathcal{I} -non-degenerate in the Riemannian case.

In most cases we do actually have a frame in which we know which direction is time. However, if we are only handed a set of invariants we would not have such a frame and, a priori, we would not know which eigenvalue is associated with time. We also note that the ambiguity in choosing a projection operator in certain cases is linked to the same problem; we do not necessarily know which eigenvalue is associated with time.

Therefore, the question of which \mathcal{I} -non-degenerate metrics are strongly \mathcal{I} -non-degenerate is linked to the question of when the time direction can be uniquely specified from the set of invariants.

Consider an invariant I . Then we can consider the gradient, $v_\mu \equiv \nabla_\mu I$, which is a curvature “vector”. Assume that the metric is \mathcal{I} -non-degenerate, in which case we always have a timelike projection operator, \perp_1 . Therefore, we can consider $\perp_1 \mathbf{v}$. Now, if $\perp_1 \mathbf{v} \neq 0$ then clearly it is timelike and the invariant $(\perp_1 \mathbf{v})^\mu (\perp_1 \mathbf{v})_\mu < 0$. Therefore, we could uniquely specify time, because $\perp_1 \mathbf{v}$ would give us the time direction. So if there exists an invariant I for which $(\perp_1)^\mu \nabla_\mu I$ is timelike (and non-zero), this spacetime is strongly \mathcal{I} -non-degenerate.

A similar conclusion is reached if we have three spacelike projection operators and all of these have similar non-zero gradients. To be more precise:

Proposition 7.1. *Consider a (weakly) \mathcal{I} -non-degenerate spacetime. Then, if either:*

1. *there exists an invariant $I = v_\mu v^\mu$, where v_μ is a curvature 1-tensor, such that $I < 0$; or,*
2. *there exist curvature 1-tensors v_μ , u_μ and w_μ such that the invariants $I_1 = v_\mu v^\mu > 0$, $I_2 = u_\mu u^\mu > 0$, $I_3 = w_\mu w^\mu > 0$, and*

$$I_4 = v_\alpha u_\beta w_\gamma g_{\mu\nu}^{\alpha\beta} g_{\lambda}^{\gamma] v^\mu u^\nu w^\lambda \neq 0;$$

then the spacetime is strongly \mathcal{I} -non-degenerate.

Proof. In case (1) we can construct a timelike projection operator, and the result follows. In case (2) there exist three spacelike projection operators, and the condition that $I_4 \neq 0$ ensures that these are linearly independent. Hence, the timelike vector is orthogonal to these three and the result follows. \square

Therefore, the only spacetimes that are weakly \mathcal{I} -non-degenerate but not strongly \mathcal{I} -non-degenerate must have a timelike and a spacelike derivative which annihilate all invariants. If the spacetime is weakly \mathcal{I} -non-degenerate, but not strongly \mathcal{I} -non-degenerate, there must consequently exist a timelike vector, ξ_1 , and a spacelike vector, ξ_2 , for which

$$\xi_1(I) = \xi_2(I) = 0,$$

for all scalar invariants I . If $[\xi_1, \xi_2] = \xi_3$, it also follows that $\xi_3(I) = 0$. Therefore, there will be a set of vectors, $\{\xi_i\}$, closed under commutation (consequently, the Jacobi identity will also be satisfied), which annihilates all curvature invariants. This has several consequences. First, this set will span a timelike (sub)manifold of dimension 2, 3 or 4. We can therefore locally introduce normal coordinates, so that the invariants only depend on the normal coordinates; i.e., $I = I(x, y)$ (dim 2), $I = I(x)$ (dim 3) or $I = \text{constant}$ (dim 4, and the spacetime is a CSI spacetime). Second, by the assumption that this spacetime is weakly \mathcal{I} -non-degenerate, and the fact that these invariants only depend on the coordinates (x, y) , there exists an orthonormal frame such that *all components of the curvature tensors only depend on the normal coordinates (x, y)* [12, 2].

This indicates that these vectors that annihilate all invariants have a special geometric meaning. First, let us consider an arbitrary curvature tensor of rank (n, n) , $\mathcal{R}^{\alpha_1 \dots \alpha_n}_{\beta_1 \dots \beta_n}$, being a sum, tensor products and contractions of the Riemann tensor and its covariant derivatives. Since this tensor has as many covariant as contravariant indices, we can interpret this as a curvature operator, $\mathbf{R} \equiv \mathcal{R}^{\alpha_1 \dots \alpha_n}_{\beta_1 \dots \beta_n} \mathbf{e}_{\alpha_1} \otimes \dots \otimes \mathbf{e}_{\alpha_n} \otimes \boldsymbol{\omega}^{\beta_1} \otimes \dots \otimes \boldsymbol{\omega}^{\beta_n}$, mapping rank n contravariant tensors into rank n contravariant tensors. Let us denote \mathbf{T}^{End} as the tensor algebra of *all* such curvature operators. It is clear that all polynomial curvature invariants can be considered as complete contractions of operators in \mathbf{T}^{End} .

Theorem 7.2. *Consider a spacetime which is (weakly) \mathcal{I} -non-degenerate, and a vector field ξ . Then the following conditions are locally equivalent:*

1. $\xi(I) = 0$ for all curvature invariants I .

2. The Lie derivative of any curvature operator $R \in \mathbf{T}^{End}$ with respect to ξ , vanishes; i.e.,

$$\mathcal{L}_\xi R = 0.$$

Proof. (1) \Rightarrow (2): Assume that $\xi(I) = 0$ for all curvature invariants I . Consider the 1-parameter group of diffeomorphisms, ϕ_t , associated with the vector field ξ . Then

$$\xi(I) = \mathcal{L}_\xi(I) = \lim_{t \rightarrow 0} \frac{1}{t} [I - \phi_t^*(I)] = 0.$$

Assuming the conditions hold over a neighborhood U , this can be integrated and we get, at a point $p \in U$, $I(p) = I(\phi_{-t}(p))$. Hence, along the integral curves the value of the invariants do not change. Consider now the Lie derivative of an arbitrary curvature operator $R = \mathcal{R}^{\alpha_1 \dots \alpha_n}_{\beta_1 \dots \beta_n} \mathbf{e}_{\alpha_1} \otimes \dots \otimes \mathbf{e}_{\alpha_n} \otimes \boldsymbol{\omega}^{\beta_1} \otimes \dots \otimes \boldsymbol{\omega}^{\beta_n}$ (e.g., see [12]):

$$\mathcal{L}_\xi R = \lim_{t \rightarrow 0} \frac{1}{t} [R - \hat{R}_t],$$

where \hat{R}_t is the ϕ_t -transformed tensor defined by:

$$\hat{R}_t = \phi_t(R) = \phi_t^*(\mathcal{R}^{\alpha_1 \dots \alpha_n}_{\beta_1 \dots \beta_n})(\phi_{t*} \mathbf{e}_{\alpha_1}) \otimes \dots \otimes (\phi_{t*} \mathbf{e}_{\alpha_n}) \otimes (\phi_t^* \boldsymbol{\omega}^{\beta_1}) \otimes \dots \otimes (\phi_t^* \boldsymbol{\omega}^{\beta_n}).$$

The action of ϕ_t preserves the form and symmetries of a tensor. Thus the transformed tensor \hat{R}_t will be a curvature tensor of the same kind as R . The curvature invariants at p will be $I(p)$ for R and $I(\phi_{-t}(p))$ for \hat{R}_t . From the above, these invariants are the same and, from the assumption of \mathcal{I} -non-degeneracy, the invariants characterise the spacetime, which means that there exists a frame such that the components of the curvature tensors do not change along $\phi_{-t}(p)$. This frame essentially is the eigenvalue frame of the curvature tensors. In particular, the projection operators define this frame.

If v is an eigenvector of R , then

$$Rv - \lambda v = 0, \quad \Rightarrow \quad \hat{R}_t \hat{v}_t - \lambda \hat{v}_t = 0,$$

where hatted quantities are transformed under ϕ_t . Eigenvectors are therefore transformed onto eigenvectors of \hat{R}_t . Using the fact that there exists a frame so that $\phi_t^*(\mathcal{R}^{\alpha_1 \dots \alpha_n}_{\beta_1 \dots \beta_n}) = \mathcal{R}^{\alpha_1 \dots \alpha_n}_{\beta_1 \dots \beta_n}$, means that the components remain the same in this frame. For a symmetric operator the eigenvectors are orthogonal and we can introduce a basis of orthonormal eigenvectors $\{\mathbf{e}_I\}$ with duals $\{\boldsymbol{\omega}^I\}$. Consider now a symmetric projection operator, \perp , written in the eigenvector basis:

$$\perp = \delta^A_B \mathbf{e}_A \otimes \boldsymbol{\omega}^B, \quad \hat{\perp}_t = \delta^A_B \hat{\mathbf{e}}_A \otimes \hat{\boldsymbol{\omega}}^B,$$

where the indices run over a subset of eigenvectors with the same eigenvalue, and the hatted basis is the transformed basis. From the above discussion we see that the eigenspaces are ϕ_t -invariant, and hence there is a transformation matrix M^A_B such that $\hat{\mathbf{e}}_A = M^{\hat{A}}_A \mathbf{e}_{\hat{A}}$, and $\hat{\boldsymbol{\omega}}^B = (M^{-1})^B_{\hat{B}} \boldsymbol{\omega}^{\hat{B}}$. Consequently,

$$\perp = \hat{\perp}_t,$$

and the curvature projection operators are ϕ_t -invariant. Therefore, since all $R \in \mathbf{T}^{End}$ can be expanded in terms of these projection operators and the

curvature invariants (since it is \mathcal{I} -non-degenerate), we have that $\mathbf{R} = \hat{\mathbf{R}}_t$ and (2) follows.

(2) \Rightarrow (1): This follows trivially from the observation that $\boldsymbol{\xi}(I) = \mathcal{L}_{\boldsymbol{\xi}}(I)$ and the properties of the Lie derivative. \square

Corollary 7.3. *If there exists a non-zero vector field, $\boldsymbol{\xi}$, fulfilling*

$$\mathcal{L}_{\boldsymbol{\xi}}\mathbf{R} = 0,$$

for all curvature operators $\mathbf{R} \in \mathbf{T}^{End}$, then the spacetime possesses a Killing vector field, $\hat{\boldsymbol{\xi}}$.

Proof. This follows from the equivalence principle [2]. The Cartan scalars are related to the components of the Riemann tensor and its derivatives, and along the integral curves of $\boldsymbol{\xi}$ we can use ϕ_t at any given point p . We want to compare the tensors at p and $q \equiv \phi_t(p)$. Consider an arbitrary even-ranked curvature tensor \mathbf{R} . By raising or lowering indices appropriately, we get an operator \mathbf{R} . Since the Lie derivative of \mathbf{R} along $\boldsymbol{\xi}$ vanishes, there is a frame such that \mathbf{R}_q and \mathbf{R}_p has identical components. Therefore, by raising and lowering the indices appropriately, the components of \mathbf{R}_q and \mathbf{R}_p are also the same. The Cartan invariants of \mathbf{R}_q and \mathbf{R}_p are therefore the same. For a curvature tensor, \mathbf{R} , of odd rank we consider $\mathbf{R} \otimes \mathbf{R}$, which is of even rank, and use the fact that ϕ_t is continuous in t . Therefore, there exists a frame such that all the components of any curvature tensor are identical at p and q . The equivalence principle now implies that ϕ_t , for any given t , is an isometry; hence, there must exist a Killing vector field $\hat{\boldsymbol{\xi}}$ which generates an isometry $\hat{\phi}_t$ such that $\hat{\phi}_t(p) = \phi_t(p)$. \square

Note that in most cases $\hat{\boldsymbol{\xi}}$ and $\boldsymbol{\xi}$ are the same. However, in some very special cases with additional symmetries they need not be (although locally they are of the same causality; e.g., they are both timelike or both spacelike). For example, for flat space the curvature vanishes identically; hence, $\mathcal{L}_{\boldsymbol{\xi}}\mathbf{R} = 0$ for all $\boldsymbol{\xi}$ and any curvature tensor \mathbf{R} , although not all $\boldsymbol{\xi}$ are Killing vectors. However, in these special cases there will always exist at least two Killing vectors.

Therefore, to conclude:

Corollary 7.4. *If a spacetime is weakly \mathcal{I} -non-degenerate but not strongly \mathcal{I} -non-degenerate, then it possesses locally (at least) one timelike Killing vector and one spacelike Killing vector.*

8 Conclusions

In this paper we have addressed the question of what is the class of Lorentzian manifolds that can be completely characterized by the scalar polynomial invariants constructed from the Riemann tensor and its covariant derivatives. In the Riemannian case the manifold is always locally characterized by the scalar polynomial invariants and, therefore, all of the Cartan invariants are related to the scalar curvature invariants [2]. We have generalized these results to the Lorentzian case.

We have introduced the important notion of \mathcal{I} -non-degenerate spacetime metrics. In order to prove the main theorems, which is done on a case-by-case

(depending on the algebraic type) using a boost weight decomposition, we have introduced an appropriate set of curvature operators and curvature projectors. In the (algebraically) general case we have shown that if a 4D spacetime is either Ricci type I , Weyl type I or Riemann type I/G , then it is \mathcal{I} -non-degenerate, which implies that the spacetime metric is determined by its curvature invariants (at least locally, in the sense explained above).

For the algebraically special cases the Riemann tensor itself does not give enough information to provide us with all the required projection operators, and it is also necessary to consider the covariant derivatives. In terms of the boost weight decomposition, for an algebraically special metric (which has a Riemann tensor with zero positive boost weight components) which is not Kundt, by taking covariant derivatives of the Riemann tensor positive boost weight components are acquired and a set of higher derivative projection operators are obtained. Consequently, we found that if the spacetime metric is algebraically special, but ∇R , $\nabla^{(2)}R$, $\nabla^{(3)}R$ or $\nabla^{(4)}R$ is of type I or more general, the metric is \mathcal{I} -non-degenerate.

The remaining metrics which *do not* acquire a positive boost weight component when taking covariant derivatives have a very special curvature structure. Indeed, in our main theorem we proved that a spacetime metric is either \mathcal{I} -non-degenerate or the metric is a Kundt metric. This is very striking result because it implies that a metric that is not determined by its scalar curvature invariants must be of Kundt form. The Kundt metrics which are not \mathcal{I} -non-degenerate therefore correspond to degenerate metrics in the sense that many such metrics can have identical scalar invariants. This exceptional property of the the degenerate Kundt metrics essentially follows from the fact that they do not define a unique timelike curvature operator.

The results in the case of Petrov type I spacetimes in 4D follow from the above theorems. Although these results were not previously known, some partial results for 4D Weyl (Petrov) type I spacetimes, which are consistent with the above analysis, can be deduced from previous work. This is discussed in the next section (also see Appendix D).

Therefore, if a spacetime is \mathcal{I} -non-degenerate and the algebraic type is explicitly known (using, for example, the Plebański notion for the Segre type in which commas are used to distinguish between timelike and spacelike eigenvectors and their associated eigenvalues, as is common in general relativity), the spacetime can be completely classified in terms of its scalar curvature invariants.

There are a number of important consequences of the results obtained. A corollary of the main theorem applied to spacetimes with constant curvature invariants (CSI) is a proof of the CSI-Kundt conjecture in 4D [13]. In future work we will study CSI spacetimes in more detail [14].

We then considered the inverse question: given a set of scalar polynomial invariants, what can we say about the underlying spacetime? In 4D we can partially characterize the Petrov type in terms of scalar curvature invariants. In most circumstances we only need some partial results or necessary conditions. For example, we found that if $27J^2 \neq I^3$, or if $27J^2 = I^3$ but the invariants $S_1 \neq 0$ or $S_2 \neq 0$, then the spacetime is \mathcal{I} -non-degenerate. Some results were then presented in the remaining cases when both I and J are zero, and hence the spacetime is VSI_0^W .

We also discussed whether a \mathcal{I} -non-degenerate metric is unique under a discrete transformation. We introduced the notion *strong* and *weak* non-degeneracy.

We provided a necessary criterion to determine spacetimes that are weakly \mathcal{I} -non-degenerate but not strongly \mathcal{I} -non-degenerate.

Having determined when a spacetime is completely characterized by its scalar curvature invariants, it is also of interest to determine the minimal set of such invariants needed for this classification. For example, in 4D there are results on determining the Riemann tensor in terms of zeroth order scalar curvature invariants (and determining a minimal set of such invariants) [15]. It is also of interest to study when a spacetime can be explicitly *constructed* from scalar curvature invariants.

This work is also of importance to the equivalence problem of characterizing Lorentzian spacetimes (in terms of their Cartan scalars) [2]. Clearly, by knowing which spacetimes can be characterized by their scalar curvature invariants alone, the computations of the invariants (i.e., simple polynomial scalar invariants) is much more straightforward and can be done algorithmically (i.e., the full complexity of the equivalence method is not necessary). On the other hand, the Cartan equivalence method also contains, at least in principle, the conditions under which the classification is complete (although in practice carrying out the classification for the more general spacetimes is difficult, if not impossible). Therefore, in a sense, the full machinery of the Cartan equivalence method is only necessary for the classification of the degenerate Kundt spacetimes (which we shall address in future work).

Let us briefly discuss this further in the context of two simple examples, which also serve to illustrate the results of the main theorem:

1. The Schwarzschild vacuum type D spacetime is an example of an \mathcal{I} -non-degenerate spacetime. In the canonical coordinate form of the metric as given in [16], the two scalar polynomial invariants $C^2 \equiv C_{abcd}C^{abcd} = 48m^2r^{-6}$ and $(\nabla C)^2 \equiv C_{abcd;e}C^{abcd;e} = 720(r - 2m)m^2r^{-9}$ are functionally independent and can be used to solve for r and m , and all of the algebraically independent Cartan scalars Ψ_2 , $\nabla^2\Psi_{20'}$, $\nabla^2\Psi_{31'}$, and $\nabla^2\Psi_{42'}$ are consequently related to the polynomial curvature invariants C^2 and $(\nabla C)^2$ [16]. In particular, $\Psi_2 = -mr^{-3}$, $\nabla^2\Psi_{20'} = 12m^2r^{-6} - 6mr^{-5}$, so that $48(\Psi_2)^2 = C^2$ and $120(\Psi_2)(\nabla^2\Psi_{20'}) = -(\nabla C)^2$. We note that the second derivative Cartan scalars have the following boost weights: $\nabla^2\Psi_{20'}$ is $+2$, $\nabla^2\Psi_{42'}$ is -2 and $\nabla^2\Psi_{31'}$ is 0 .
2. A spatially homogeneous vacuum plane wave, which is a special subcase of a Petrov type N vacuum spacetime admitting a covariantly constant null vector, belongs to the class of vanishing scalar invariant (VSI) spacetimes [8] and is consequently an example of a degenerate Kundt spacetime. Since it is a VSI spacetime, all scalar polynomial invariants are zero. However, distinct VSI spacetimes give rise to a distinct set of Cartan scalars [2] (e.g., in flat space all of the Cartan scalars are zero). A spatially homogeneous vacuum plane wave has two non-trivial Cartan scalars, $\nabla\Psi_{00'}$ and $\nabla^2\Psi_{00'}$.

9 Discussion

We have addressed the question of what is the class of Lorentzian manifolds that can be completely characterized by the scalar polynomial invariants constructed from the Riemann tensor and its covariant derivatives. In particular, we proved

the result that this is true in the case of Petrov type I spacetimes in 4D. This result was not previously known. However, some partial results for 4D Weyl (Petrov) type I spacetimes are known, which are consistent with the above analysis. Let us review these results.

Essentially, in the case of Petrov (Weyl) type I, there exists a unique frame so that all components of the Riemann tensor are related to curvature invariants. Indeed, in general there are four different curvature invariants (e.g., corresponding to the complex invariants I and J), so that all invariants (which depend on 4 coordinates) are functionally dependent on these four invariants. Problems arise in degenerate cases and cases with symmetries. It is also known that all Petrov type I spacetimes are completely backsolvable [15].

Let us consider the Petrov type I case in more detail. From [17, 3] (also see Appendix D) it follows that if a 4D spacetime is of Petrov type I it can be classified according to its rank and it is either:

1. curvature class A (and the holonomy group is general and of type R_{15}),
2. curvature class C (and of holonomy type R_{10} or R_{13} , with restricted Segre type).

Now, suppose the components of the Riemann tensor $R^a{}_{bcd}$ are given in a coordinate domain U with metric g . In case (1), where the curvature class is of type A, for any other metric g' with the same components $R^a{}_{bcd}$ it follows that $g'_{ab} = \alpha g_{ab}$ (where α is a constant); i.e., the metric is determined up to a constant conformal factor and the connection is uniquely determined. This implies that all higher order covariant derivatives of the Riemann tensor are completely determined; i.e., given $R^a{}_{bcd}$, all of the components of the covariant derivatives are determined and we only need classify the Riemann tensor itself. (Note that all of the scalar polynomial curvature invariants are then determined, at least up to an overall constant factor).

We can then pass to the frame formalism and determine the frame components of the Riemann tensor (to do this we need the metric to determine the orthogonality of the frame vectors and hence construct the frame; since g is specified up to an overall constant conformal factor, orthogonality is unique). The Petrov type I case is completely backsolvable [15] and hence the frame components are completely determined by the zeroth order scalar invariants. Therefore, it follows that the spacetime is completely characterized by its scalar curvature invariants in this case.

Let us now consider case (2), where the curvature class is C . Again, let us suppose that the $R^a{}_{bcd}$ are given in U with metric g . If g' is any other metric with the same $R^a{}_{bcd}$, it follows that

$$g'_{ab} = \alpha g_{ab} + \beta k_a k_b$$

(where α and β are constants). The equation

$$R^a{}_{bcd} k^d = 0, \tag{17}$$

has a unique non-trivial solution for $k \in T_m M$. Note that $R^a{}_{bcd} k_a = 0$ implies that $I_1 k_e = 0$ and hence $I_1 = 0$, where I_1 is the Euler density:

$$I_1 \equiv [R^{abcd} R_{abcd} - 4R^{ab} R_{ab} + R^2].$$

If $R^a{}_{bcd;e}k_a \neq 0$, then $\beta = 0$ and the metric is determined up to a constant conformal factor (and the holonomy type is R_{15}). This is similar to the first case discussed above, but now some information on the covariant derivative of the Riemann tensor is necessary (to ensure $R^a{}_{bcd;e}k_a \neq 0$). Hence, first order curvature invariants are needed for the classification of the spacetime. Since $R^a{}_{bcd;e}k_a = 0$ implies that $I_2 k_e = 0$, where

$$I_2 \equiv [R^{abcd;e}R_{abcd;e} - 4R^{ab;c}R_{ab;c} + R^a{}R_{,a}],$$

it follows that the invariant $I_2 \neq 0$ implies that $R^a{}_{bcd;e}k_a \neq 0$ in this case.

If $R^a{}_{bcd;e}k_a = 0$, then $R^a{}_{bcd}k_{a;e} = 0$, and since eqn. (17) has a unique solution, k_a is recurrent. If k_a is null, the spacetime is algebraically special, and since we assume that the Petrov type is I, this is not possible. Hence, k_a is (a) timelike (TL) or (b) spacelike (SL) and is, in fact, covariant constant (CC).

In case (2a), the spacetime admits a TL CC vector field k_a . The holonomy is R_{13} , with a TL holonomy invariant subspace which is non-degenerately reducible, and M is consequently locally (1+3) decomposable (and static). There exist local coordinates (with $k = \frac{\partial}{\partial t}$) such that the metric is given by

$$- dt^2 + g_{\alpha\beta}(x^\gamma)dx^\alpha dx^\beta \quad (\alpha = 1, 2, 3) \quad (18)$$

where $g_{\alpha\beta}$ is independent of t . The metric is unique up to an overall constant scaling and a time translation $t \rightarrow \lambda t$, where $\lambda^2 = 1 + \beta/\alpha$ (reflecting the non-uniqueness of the TL CC vector up to a constant scaling λ). All of the non-trivial components of the Riemann tensor and its covariant derivatives are constructed from the 3D positive definite metric $g_{\alpha\beta}$, and can be classified by the corresponding 3D Riemann curvature invariants. In this case (and case (2b)) there is an ignorable coordinate and all invariants are functions of 3 independent functions; $R^a{}_{bcd;e}$ must be used to uniquely fix the frame, and hence we need information from the first order scalar invariants.

In case (2b), the spacetime admits a SL CC vector field k_a . The holonomy is R_{10} , there exists a holonomy invariant *SL* vector k_a which is non-degenerately reducible, and M is this locally (3+1) decomposable. Choosing local coordinates in which the SL CC vector $k = \frac{\partial}{\partial x}$, the metric is given by

$$dx^2 + g_{\bar{\alpha}\bar{\beta}}dx^{\bar{\alpha}}dx^{\bar{\beta}} \quad (\bar{\alpha} = 0, 2, 3) \quad (19)$$

and $g_{\alpha\beta}$ is independent of x . The metric is unique up to an overall constant conformal factor and a space translation $x \rightarrow \lambda x$ ($\lambda^2 = 1 + \beta/\alpha$). Classification now reduces to the classification of the class of 3D Lorentzian spacetimes with Lorentzian metric $g_{\bar{\alpha}\bar{\beta}}$ (the subclass such that (2) is of Petrov type *I*). We can now iterate the procedure for 3D Lorentzian spacetimes (such that (2) is Petrov type *I*). In the degenerate cases in which additional KV are admitted, we will be led to the locally homogeneous case, and hence the 4D Petrov type I locally homogeneous spacetimes (which are characterized by their constant scalar invariants). Indeed, in 3D the Riemann tensor is completely determined by the Ricci tensor. There always exists a frame in which the components of the Ricci tensor are constants [7] and so in this case the 4D spacetime is Petrov type *I* and CH_0 (curvature homogeneous [18]), and hence generically locally homogeneous.

Acknowledgments

We would like to thank Robert Milson for useful comments and questions on our manuscript, and to Lode Wylleman for pointing out a mistake. This work was supported by the Natural Sciences and Engineering Research Council of Canada.

A Notation

Throughout we have used a Newman-Penrose (NP) tetrad given by $e_a = \{\ell, n, m, \bar{m}\}$ with inner product

$$\eta_{ab} = \begin{bmatrix} 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & -1 \\ 0 & 0 & -1 & 0 \end{bmatrix} \quad (20)$$

and directional derivatives defined by

$$D = \ell^\mu \nabla_\mu, \quad \Delta = n^\mu \nabla_\mu, \quad \delta = m^\mu \nabla_\mu. \quad (21)$$

Associated with an NP tetrad are the following definitions for the connection coefficients that appear frequently above

$$\kappa = m^\mu D\ell_\mu, \quad \sigma = m^\mu \delta\ell_\mu, \quad \rho = m^\mu \bar{\delta}\ell_\mu \quad (22)$$

with the remaining ones being similarly defined. Given the frame components $R_{abcd} = R_{\alpha\beta\gamma\delta} e_a^\alpha e_b^\beta e_c^\gamma e_d^\delta$, we have the definitions for the Weyl scalars

$$\Psi_0 = -C_{1313}, \quad \Psi_1 = -C_{1213}, \quad \Psi_2 = -C_{1342}, \quad \Psi_3 = -C_{1242}, \quad \Psi_4 = -C_{2424}$$

and the Ricci scalars

$$\begin{aligned} \Phi_{00} &= \frac{1}{2}R_{11}, & \Phi_{01} &= \frac{1}{2}R_{13}, & \Phi_{02} &= \frac{1}{2}R_{33} \\ \Phi_{11} &= \frac{1}{4}(R_{12} + R_{34}), & \Phi_{12} &= \frac{1}{2}R_{23}, & \Phi_{22} &= \frac{1}{2}R_{22}. \end{aligned} \quad (23)$$

Given a covariant tensor T with respect to an NP tetrad (or null frame), the effect of a boost $\ell \mapsto e^\lambda \ell$, $n \mapsto e^{-\lambda} n$ allows T to be decomposed according to its boost weight

$$T = \sum_b (T)_b \quad (24)$$

where $(T)_b$ denotes the boost weight b components of T . An algebraic classification of tensors T has been developed [6, 5] which is based on the existence of certain normal forms of (24) through successive application of null rotations and spin-boost. In the special case where T is the Weyl tensor in four dimensions, this classification reduces to the well-known Petrov classification. However, the boost weight decomposition can be used in the classification of any tensor T in arbitrary dimensions. As an application, a Riemann tensor of type G has the following decomposition

$$R = (R)_{+2} + (R)_{+1} + (R)_0 + (R)_{-1} + (R)_{-2} \quad (25)$$

in every null frame. A Riemann tensor is algebraically special if there exists a frame in which certain boost weight components can be transformed to zero, these are summarized in Table 3.

A useful discrete symmetry is the following (orientation-preserving) Lorentz transformation:

$$\ell \leftrightarrow n, \quad m \leftrightarrow \bar{m}, \quad (26)$$

which interchanges the boost weights, $(T)_b \leftrightarrow (T)_{-b}$, and makes the replacements

$$(\kappa, \sigma, \rho, \tau, \epsilon, \beta) \leftrightarrow -(\nu, \lambda, \mu, \pi, \gamma, \alpha). \quad (27)$$

Riemann type	Conditions
G	—
I	$(R)_{+2} = 0$
II	$(R)_{+2} = (R)_{+1} = 0$
III	$(R)_{+2} = (R)_{+1} = (R)_0 = 0$
N	$(R)_{+2} = (R)_{+1} = (R)_0 = (R)_{-1} = 0$
D	$(R)_{+2} = (R)_{+1} = (R)_{-1} = (R)_{-2} = 0$
O	all vanish (Minkowski space)

Table 3: The relation between Riemann types and the vanishing of boost weight components. For example, $(R)_{+2}$ corresponds to the frame components $R_{1313}, R_{1414}, R_{1314}$.

B Some special operators

Consider the case where we have a tensor $S_{\mu\nu\alpha\beta}$, where

$$S_{\mu\nu\alpha\beta} = S_{(\mu\nu)(\alpha\beta)} = S_{\alpha\beta\mu\nu},$$

This tensor can be considered as an operator:

$$\mathbf{S} = (S^{\mu\nu}_{\alpha\beta}) : V \mapsto V,$$

where V is the vector space of symmetric 2-tensors $M^{\mu\nu}$.

Therefore, we can consider the eigentensors of this map in the standard manner. We can construct a set of projectors \perp_A projecting onto each corresponding eigenspace. Assume that \perp_1 is of rank 1 (as an operator). If $M^{\mu\nu}$ is the corresponding (normalized) eigenvector, this means that

$$(\perp_1)_{\mu\nu\alpha\beta} = M_{\mu\nu} M_{\alpha\beta}.$$

We can now consider the eigenvectors of $\mathbf{M} \equiv M^{\mu}_{\nu}$. We are actually not considering the operator \mathbf{M} itself, but rather \perp_1 . However, \perp_1 can also be considered as an operator:

$$\mathbf{P} : N^{\mu\nu} \mapsto M^{\mu}_{\alpha} M^{\nu}_{\beta} N^{\alpha\beta}$$

Assume that v^μ and w^ν are eigenvectors of M with eigenvalues λ_v and λ_w , respectively. Then, if $N^{\mu\nu} = v^\mu w^\nu$,

$$M^\mu_\alpha M^\nu_\beta N^{\alpha\beta} = \lambda_v \lambda_w N^{\mu\nu},$$

and is therefore an eigenvector of P with eigenvalue $\lambda = \lambda_v \lambda_w$. Clearly, $v^\mu w^\nu$ has the same eigenvalue as $w^\mu v^\nu$, so we will not be able to distinguish these using projection operators. Furthermore, if $\lambda_v = \pm \lambda_w$, then $v^\mu v^\nu$ has the same eigenvalue as $w^\mu w^\nu$.

The above construction is useful in several cases. An example that recurs is the case where M has two one-dimensional eigenspaces spanned by v^μ and w^μ , say. Assume also that $\lambda_v = -\lambda_w$. Then, P has two projection operators:

$$(P_1)_{\mu\nu\alpha\beta} \propto v_\mu w_\nu v_\alpha w_\beta + w_\mu v_\nu w_\alpha v_\beta, \quad (28)$$

$$(P_2)_{\mu\nu\alpha\beta} \propto v_\mu v_\nu v_\alpha v_\beta + w_\mu w_\nu w_\alpha w_\beta, \quad (29)$$

We see that this is somewhat unfortunate because in spite of the fact that M sees the difference between the vectors v^μ and w^ν , P does not. This is related to the fact that for some spacetimes there exists a discrete symmetry which interchanges two spacetimes with identical curvature invariants. Here this manifests itself in that we cannot actually determine which eigenvector correspond to which eigenvalue.

C Algebraically special ∇C

The relationship between the invariants of the Weyl tensor and the Petrov type is well known; however, this is not the case for the covariant derivative of the Weyl tensor. A similar analysis for ∇C would require an algebraic classification based on its boost weight decomposition, and a complete set of its first order invariants. We do not attempt to solve this general problem but rather provide some relations relevant to our paper. Restricting attention to four dimensions we define the following tensors

$$\overset{2}{T}_{abe}{}^{fgh} = C_{abcd;e} C^{cdfg;h} \quad (30)$$

$$\overset{3}{T}_{abe}{}^{ijk}{}^h = \overset{2}{T}_{abe}{}^{fgh} C_{fgij;k} \quad (31)$$

$$\overset{3,0}{T}_{abe}{}^{ij}{}^h = \overset{2}{T}_{abe}{}^{fgh} C_{fgij} \quad (32)$$

$$\overset{4}{T}_{abe}{}^{hlmn}{}^k = \overset{3}{T}_{abe}{}^{ijk}{}^h C^{ijlm;n} \quad (33)$$

where the number above the tensor refers to the degree in ∇C or C . All of these tensors are constructed purely from ∇C with the exception of $\overset{3,0}{T}$ (which

involves C). Next, we consider the following first order invariants

$$w_{2,1} = T_{abe}^2 \begin{matrix} abe \\ \end{matrix} \quad w_{2,2} = T_{abe}^2 \begin{matrix} eab \\ \end{matrix} \quad w_{2,3} = T_{a \begin{matrix} e \ a \ h \\ e \ h \end{matrix}}^2 \quad (34)$$

$$w_{3,1} = T_{abe}^{3,0} \begin{matrix} eab \\ \end{matrix} \quad w_{3,2} = T_{abe}^{3,0} \begin{matrix} bea \\ \end{matrix} \quad w_{3,3} = T_{a \begin{matrix} e \ b \ a \\ e \ b \end{matrix}}^{3,0} \quad (35)$$

$$w_{4,1} = T_{abe}^4 \begin{matrix} e \ abk \\ \ k \end{matrix} \quad w_{4,2} = T_{abe}^4 \begin{matrix} heab \\ \ h \end{matrix} \quad w_{4,3} = T_{abe}^4 \begin{matrix} h \ abe \\ \ h \end{matrix} \quad (36)$$

$$w_{4,4} = T_{abe}^4 \begin{matrix} h \ eba \\ \ h \end{matrix} \quad w_{4,5} = T_{abe}^4 \begin{matrix} abem \\ \ m \end{matrix} \quad w_{4,6} = T_{abe}^4 \begin{matrix} a \ ebm \\ \ m \end{matrix} \quad (37)$$

$$w_{4,7} = T_{abe}^4 \begin{matrix} maeb \\ \ m \end{matrix} \quad (38)$$

in which $w_{n,i}$ denotes the i th invariant of degree n in ∇C or C . Since the aligned frames of ∇C and C need not be the same, the $w_{3,i}$ are mixed invariants and the remaining invariants are pure ∇C invariants. For ∇C and C algebraically general (type G) we obtain the following syzygies

$$w_{2,1} + 2w_{2,2} - 2w_{2,3} = 0, \quad w_{3,1} + 2w_{3,2} + 2w_{3,3} = 0, \quad w_{4,1} - w_{4,3} = 0$$

which are the result of identities, symmetries and dimensionally dependent relations⁴ [19]. In subsequent calculations we always impose these syzygies so that our set reduces to ten invariants. Now consider ∇C of algebraically special type, which is obtained by setting the minimal number of appropriate boost weight components to vanish. We obtain the following results:

1. If ∇C is type II or D (i.e., boost weight $+3, +2, +1$ components vanish) then the syzygies $S_1 = 0$ and $S_2 = 0$ hold.
2. If ∇C is type G or type H (i.e., boost weight $+3$ vanish), or type I (i.e., boost weight $+3, +2$ vanish) then, in general, $S_1 \neq 0$ and $S_2 \neq 0$.

The second statement refers to the most general types of G , H or I where no further algebraically special subcases are taken into account. Below are the expressions for S_1 and S_2 . Note that S_1 is linear in $w_{4,5}$, and when $S_1 = 0$, we use this syzygy in the derivation⁵ of S_2 ; hence these two invariant expressions are generally independent. In type II or D we can regard $S_1 = 0$ as expressing the dependency of $w_{4,5}$ in terms of the other invariants of S_1 . In S_2 each of the $w_{2,i}$ appear quadratically whereas each of $w_{4,i}$ appear quartically therefore one of these invariants is dependent with respect to the other invariants in S_2 . Since these syzygies are of degree 8 and 16, and the invariants considered here are of maximum degree 4, one would expect S_1 and S_2 to attain a simpler form if expressed in terms of higher degree invariants. These calculations were

⁴Thanks to Jose M. Martin-Garcia for pointing this out to us.

⁵ $w_{4,5}$ does not appear in S_2 .

performed with the aid of *GRTensorII* [20].

$$\begin{aligned}
S_1 = & -14464w_{2,3}w_{2,2}w_{4,4} - 992w_{2,3}w_{2,2}w_{4,1} + 15872w_{2,3}w_{2,2}w_{4,6} \\
& + 7424w_{2,3}w_{2,2}w_{4,7} - 1600w_{2,3}w_{2,2}w_{4,2} + 1216w_{2,2}^2w_{4,2} \\
& + 640w_{2,3}^2w_{4,2} + (21504w_{4,4} + 2304w_{2,2}^2 - 24576w_{4,6} - 4608w_{2,3}w_{2,2} \\
& - 6144w_{4,7} + 2304w_{2,3}^2 - 768w_{4,1} + 768w_{4,2})w_{4,5} - 6656w_{2,2}^2w_{4,7} \\
& - 154112w_{4,6}w_{4,4} - 8960w_{2,3}^2w_{4,6} - 464w_{4,7}w_{4,1} - 2272w_{4,7}w_{4,2} \\
& - 15104w_{2,2}^2w_{4,6} + 224w_{2,2}^2w_{4,1} + 7760w_{4,4}w_{4,2} - 2432w_{4,6}w_{4,1} \\
& - 10240w_{4,6}w_{4,2} + 15232w_{2,2}^2w_{4,4} + 2680w_{4,4}w_{4,1} + 56320w_{4,7}w_{4,6} \\
& + 512w_{2,3}^2w_{4,1} + 6400w_{2,3}^2w_{4,4} - 41408w_{4,7}w_{4,4} - 36w_{4,2}w_{4,1} \\
& - 2816w_{2,3}^2w_{4,7} + 180w_{4,2}^2 + 6464w_{4,7}^2 + 58832w_{4,4}^2 + 94208w_{4,6}^2 - 171w_{4,1}^2
\end{aligned}$$

$$\begin{aligned}
S_2 = & -5364449280w_{4,6}^2w_{2,3}w_{2,2}w_{4,2} + 2223360w_{2,3}^2w_{4,2}^3 \\
& -3893760w_{4,2}^3w_{2,2}^2 - 603625881600w_{4,6}^3w_{4,4} - 20101201920w_{4,6}^3w_{2,3}^2 + 55490641920w_{4,6}^3w_{2,2}^2 \\
& -22171852800w_{4,7}^3w_{4,4} - 4885920w_{4,2}^3w_{4,1} + 148414464000w_{4,6}^2w_{4,7}^2 + 568104468480w_{4,6}^2w_{4,4}^2 \\
& -44177817600w_{4,6}^3w_{4,2} - 28282060800w_{4,6}^3w_{4,1} + 3288600w_{4,2}w_{4,1}^3 - 81312860160w_{4,7}w_{4,4}^3 \\
& +68296366080w_{4,7}^2w_{4,4}^2 + 2340126720w_{2,2}^2w_{4,7}^3 + 3975480w_{4,2}^2w_{4,1}^2 + 305528832000w_{4,6}^3w_{4,7} \\
& +187499520w_{4,6}^2w_{4,1}^2 + 3116666880w_{4,6}^2w_{4,2}^2 - 227302871040w_{4,6}w_{4,4}^3 + 12399045120w_{4,4}^3w_{4,2} \\
& -80592261120w_{4,4}^3w_{2,2}^2 - 194522400w_{4,4}^2w_{4,1}^2 + 2817964800w_{4,4}^3w_{4,1} + 2603059200w_{4,4}^3w_{2,3}^2 \\
& -16178400w_{4,1}^3w_{4,7} + 22063680w_{4,1}^3w_{2,2}^2 + 62340480w_{4,1}^2w_{4,7}^2 + 41428800w_{4,1}^3w_{4,6} \\
& -10962000w_{4,1}^3w_{4,4} - 1406880w_{4,1}^3w_{2,3}^2 + 1134028800w_{4,7}^3w_{4,1} - 1420185600w_{4,7}^3w_{4,2} \\
& +31371264000w_{4,7}^3w_{4,6} - 612679680w_{4,7}^3w_{2,3}^2 - 30101760w_{4,2}^3w_{4,7} + 71660160w_{4,2}^3w_{4,4} \\
& +310187520w_{4,2}^2w_{4,7}^2 + 1551254400w_{4,2}^2w_{4,4}^2 - 96145920w_{4,2}^3w_{4,6} + 1095120w_{4,2}^4 \\
& +231211008000w_{4,6}^4 + 32785562880w_{4,4}^4 + 496125w_{4,1}^4 + 2437632000w_{4,7}^4 \\
& -101231493120w_{4,6}^2w_{2,3}w_{2,2}w_{4,4} + 41724149760w_{4,6}^2w_{2,3}w_{2,2}w_{4,7} - 10441359360w_{4,6}^2w_{2,3}w_{2,2}w_{4,1} \\
& +8616960w_{2,3}w_{2,2}w_{4,2}^2w_{4,7} + 109117440w_{2,3}w_{2,2}w_{4,2}^2w_{4,6} + 61908480w_{2,3}w_{2,2}w_{4,2}^2w_{4,4} \\
& -266860800w_{2,3}w_{2,2}w_{4,1}^2w_{4,7} + 1940244480w_{2,3}w_{2,2}w_{4,1}^2w_{4,7} + 2091409920w_{2,3}w_{2,2}w_{4,1}^2w_{4,4} \\
& +414351360w_{2,3}w_{2,2}w_{4,1}^2w_{4,6} + 52652160w_{2,3}w_{2,2}w_{4,1}^2w_{4,2} + 106024320w_{2,3}w_{2,2}w_{4,1}^2w_{4,4} \\
& +50016960w_{2,3}w_{2,2}w_{4,1}^2w_{4,2} + 643184640w_{4,2}w_{4,1}w_{2,2}^2w_{4,7} + 201784320w_{4,2}w_{4,1}w_{4,6}w_{4,4} \\
& -150036480w_{4,2}w_{4,1}w_{2,3}^2w_{4,6} + 1132830720w_{4,2}w_{4,1}w_{2,2}^2w_{4,6} - 1692518400w_{4,2}w_{4,1}w_{4,7}w_{4,6} \\
& -1386869760w_{4,2}w_{4,1}w_{2,2}^2w_{4,4} - 74626560w_{4,2}w_{4,1}w_{2,3}^2w_{4,4} + 1610933760w_{4,2}w_{4,1}w_{4,7}w_{4,4} \\
& +195978240w_{4,2}w_{4,1}w_{2,3}^2w_{4,7} - 4386816000w_{2,3}^2w_{4,2}w_{4,6}w_{4,4} + 2208890880w_{2,3}^2w_{4,2}w_{4,7}w_{4,6} \\
& -1048596480w_{2,3}^2w_{4,2}w_{4,7}w_{4,4} + 26542080000w_{4,6}w_{4,4}^2w_{2,3}w_{2,2} - 141363118080w_{4,6}w_{4,4}^2w_{2,2}w_{4,7} \\
& +3734138880w_{4,6}w_{4,4}w_{4,7}w_{4,1} + 60439633920w_{4,6}w_{4,4}w_{4,7}w_{4,2} + 19318947840w_{4,6}w_{4,4}w_{2,2}^2w_{4,1} \\
& +18244730880w_{4,6}w_{4,4}w_{2,2}^2w_{4,2} - 2918891520w_{4,6}w_{4,4}w_{2,3}^2w_{4,1} + 28718530560w_{4,6}w_{4,4}w_{2,3}^2w_{4,7} \\
& -14858588160w_{4,7}w_{4,4}^2w_{2,3}w_{2,2} + 3924910080w_{4,7}w_{4,4}w_{2,3}w_{2,2} + 9899274240w_{4,7}w_{4,4}w_{2,2}^2w_{4,1} \\
& +6060810240w_{4,7}w_{4,4}w_{2,2}^2w_{4,2} + 304588800w_{4,7}w_{4,4}w_{2,3}^2w_{4,1} + 1175224320w_{2,3}^2w_{4,6}w_{4,7}w_{4,1} \\
& -7813693440w_{2,2}^2w_{4,7}w_{4,6}w_{4,1} - 5205196800w_{2,2}^2w_{4,7}w_{4,6}w_{4,2} + 2413071360w_{4,4}^2w_{2,3}w_{2,2}w_{4,2} \\
& +13086720w_{2,3}w_{2,2}w_{4,7}^2w_{4,2} + 5733089280w_{2,3}w_{2,2}w_{4,7}^2w_{4,6} - 1048320w_{2,3}w_{2,2}w_{4,2}^3 \\
& -312531840w_{4,2}w_{4,1}w_{4,4}^2 - 25586565120w_{4,6}^2w_{2,3}^2w_{4,7} - 21538897920w_{4,6}^2w_{4,4}^2w_{2,3}^2 \\
& -204650496000w_{4,6}w_{4,4}w_{4,7}^2 - 17712000w_{4,2}^2w_{4,1}w_{2,3}^2 + 78222827520w_{4,7}w_{4,4}^2w_{2,2}^2 \\
& -84564000w_{4,2}w_{4,1}^2w_{4,4} + 824785920w_{2,3}^2w_{4,2}w_{4,4}^2 + 426528000w_{4,6}w_{4,4}w_{4,1}^2 \\
& -7596933120w_{4,6}^2w_{2,2}^2w_{4,2} - 202144481280w_{4,6}^2w_{2,2}^2w_{4,4} + 152928000w_{4,2}^2w_{4,1}w_{4,6} \\
& +90594754560w_{4,6}^2w_{4,4}w_{4,2} - 158883840w_{2,3}^2w_{4,2}^2w_{4,6} - 4459392000w_{4,6}w_{4,4}w_{4,2}^2 \\
& +2974187520w_{4,6}^2w_{2,3}^2w_{4,1} - 33359040w_{4,2}w_{4,1}^2w_{4,7} - 46275840w_{4,2}^2w_{4,1}w_{2,2}^2 \\
& +91186560w_{4,2}^2w_{4,1}w_{4,7} + 277770240w_{2,3}^2w_{4,2}w_{4,7}^2 + 59454259200w_{4,6}^2w_{2,2}^2w_{4,7} \\
& +202020480w_{4,2}w_{4,1}^2w_{4,6} + 38804520960w_{4,6}^2w_{2,3}^2w_{4,4} - 560862720w_{4,2}w_{4,1}w_{4,7}^2 \\
& +78681600w_{2,3}^2w_{4,2}^2w_{4,4} - 4388981760w_{4,7}^2w_{4,4}w_{4,1} - 20584074240w_{4,7}w_{4,4}^2w_{4,2}
\end{aligned}$$

$$\begin{aligned}
& -23960862720w_{4,7}^2w_{4,4}w_{2,2}^2 + 396922429440w_{4,6}w_{4,4}^2w_{4,7} - 43977600w_{4,2}w_{4,1}^2w_{2,2}^2 \\
& + 456929280w_{4,6}^2w_{4,2}w_{4,1} + 664450560w_{4,7}w_{4,4}^2w_{4,1} + 3913482240w_{4,6}^2w_{2,3}^2w_{4,2} \\
& + 495043200w_{4,7}w_{4,4}w_{4,1}^2 - 20418462720w_{4,6}w_{4,4}^2w_{4,1} - 614758809600w_{4,6}^2w_{4,7}w_{4,4} \\
& - 143570880w_{4,2}^2w_{4,1}w_{4,4} - 1454814720w_{4,7}w_{4,4}w_{4,2}^2 - 5780275200w_{4,6}^2w_{2,2}^2w_{4,1} \\
& - 12847680w_{4,2}w_{4,1}^2w_{2,3}^2 + 44401582080w_{4,6}^2w_{4,4}w_{4,1} + 9839646720w_{4,7}^2w_{4,4}w_{4,2} \\
& - 59617105920w_{4,6}w_{4,4}^2w_{4,2} - 7641907200w_{4,6}^2w_{4,7}w_{4,1} + 205920w_{2,3}w_{2,2}w_{4,1}^3 \\
& - 42624000w_{2,3}^2w_{4,2}^2w_{4,7} - 43020288000w_{4,6}^2w_{4,7}w_{4,2} + 226496839680w_{4,6}w_{4,4}^2w_{2,2}^2 \\
& - 5489233920w_{4,7}w_{4,4}^2w_{2,3}^2 + 3500236800w_{4,7}^2w_{4,4}w_{2,3}^2 - 7591034880w_{2,3}^2w_{4,6}w_{4,7}^2 \\
& - 91699200w_{2,3}^2w_{4,6}w_{4,1}^2 + 1986324480w_{4,2}^2w_{4,7}w_{4,6} - 377994240w_{4,2}^2w_{2,2}^2w_{4,4} \\
& + 10468362240w_{4,4}^3w_{2,3}w_{2,2} - 10165155840w_{4,4}^2w_{2,2}^2w_{4,2} - 856350720w_{2,2}^2w_{4,7}^2w_{4,2} \\
& + 20755906560w_{2,2}^2w_{4,7}^2w_{4,6} + 101836800w_{2,2}^2w_{4,7}w_{4,2}^2 + 237242880w_{2,2}^2w_{4,7}w_{4,1}^2 \\
& + 321085440w_{2,2}^2w_{4,6}w_{4,2}^2 - 167454720w_{2,2}^2w_{4,6}w_{4,1}^2 - 2234142720w_{2,2}^2w_{4,7}^2w_{4,1} \\
& - 152616960w_{2,3}w_{2,2}w_{4,7}^3 - 228591360w_{2,2}^2w_{4,4}w_{4,1}^2 - 12494730240w_{2,2}^2w_{4,4}^2w_{4,1} \\
& - 13230720w_{2,3}^2w_{4,4}w_{4,1}^2 + 260743680w_{2,3}^2w_{4,4}^2w_{4,1} + 68417280w_{2,3}^2w_{4,7}w_{4,1}^2 \\
& + 65399685120w_{4,6}^3w_{2,3}w_{2,2} + 6782976000w_{2,3}w_{2,2}w_{4,1}w_{4,6}w_{4,4} - 534067200w_{2,3}^2w_{4,7}^2w_{4,1} \\
& - 1123038720w_{4,1}^2w_{4,7}w_{4,6} + 4393267200w_{4,7}^2w_{4,6}w_{4,1} - 13674700800w_{4,7}^2w_{4,6}w_{4,2} \\
& - 644705280w_{2,3}w_{2,2}w_{4,1}w_{4,7}w_{4,2} + 780894720w_{2,3}w_{2,2}w_{4,1}w_{4,4}w_{4,2} \\
& - 204963840w_{2,3}w_{2,2}w_{4,1}w_{4,6}w_{4,2} + 14745600w_{2,3}w_{2,2}w_{4,1}w_{4,7}w_{4,6} \\
& - 4408104960w_{2,3}w_{2,2}w_{4,1}w_{4,7}w_{4,4} + 1975910400w_{4,6}w_{4,4}w_{2,3}w_{2,2}w_{4,2} \\
& - 19641139200w_{4,6}w_{4,4}w_{2,3}w_{2,2}w_{4,7} - 1053757440w_{4,7}w_{4,4}w_{2,3}w_{2,2}w_{4,2} \\
& - 1527644160w_{2,3}w_{2,2}w_{4,2}w_{4,7}w_{4,6}
\end{aligned}$$

D Curvature

Let M be a 4-dimensional smooth connected Hausdorff manifold admitting a global smooth Lorentz metric h with associated curvature tensor R . It will be convenient to describe a simple algebraic classification of R according to its rank (relative to h). This classification is easily described geometrically and is a pointwise classification [3].

A skew-symmetric tensor F of type $(0, 2)$ or $(2, 0)$ at $m \in T_m M$ is called a *bivector*. If $F (\neq 0)$ is such a bivector, the rank of any of its (component) matrices is either two or four. In the former case, one may write (e.g. in the $(2, 0)$ case) $F^{ab} = 2r^{[a} s^{b]}$ for $r, s \in T_m M$ (or alternatively, $F = r \wedge s$) and F is called *simple*, with the 2-dimensional subspace (2-space) of $T_m M$ spanned by r, s referred to as the *blade* of F . In the latter case, F is called *non-simple*.

The metric $h(m)$ converts $T_m M$ into a Lorentz inner product space and thus it makes sense to refer to vectors in $T_m M$ and covectors in the cotangent space $T_m^* M$ to M at m (using $h(m)$ to give a unique isomorphism $T_m M \leftrightarrow T_m^* M$, that is, to raise and lower tensor indices) as being *timelike*, *spacelike*, *null* or *orthogonal*, using the signature $(-, +, +, +)$. The same applies to 1-dimensional subspaces (*directions*) and 2- and 3-dimensional subspaces of $T_m M$ or $T_m^* M$. A *simple* bivector at m is then called *timelike* (respectively, *spacelike* or *null*) if

its blade at m is a timelike (respectively a spacelike or null) 2-space at m . A *non-simple* bivector F at m may be shown to uniquely determine an orthogonal pair of 2-spaces at m , one spacelike and one timelike, and which are referred to as the *canonical pair of blades* of F . A tetrad (l, n, x, y) of members of $T_m M$ is called a null tetrad at m if the only non-vanishing inner products between its members at m are $h(l, n) = h(x, x) = h(y, y) = 1$. Thus l and n are null.

D.1 Classification

Define a linear map f from the 6-dimensional vector space of type $(2, 0)$ bivectors at m into the vector space of type $(1, 1)$ tensors at m by $f : F^{ab} \rightarrow R^a{}_{bcd} F^{cd}$. The condition (2) shows that if a tensor T is in the range of f then

$$h_{ae} T^e{}_b + h_{be} T^e{}_a = 0 \quad (\Rightarrow T_{ab} = -T_{ba}, \quad T_{ab} = h_{ae} T^e{}_b) \quad (39)$$

and so T can be regarded as a member of the matrix representation of the Lie algebra of the pseudo-orthogonal (Lorentz) group of $h(m)$. Using f one can divide the curvature tensor $R(m)$ into five classes.

Class A This is the most general curvature class and the curvature will be said to be of (curvature) class A at $m \in M$ if it is not in any of the classes B , C , D or O below.

Class B The curvature tensor is said to be of (curvature) class B at $m \in M$ if the range of f is 2-dimensional and consists of all linear combinations of type $(1, 1)$ tensors F and G where $F^a{}_b = x^a y_b - y^a x_b$ and $G^a{}_b = l^a n_b - n^a l_b$ with l, n, x, y a null tetrad at m . The curvature tensor at m can then be written as

$$R_{abcd} \equiv h_{ae} R^e{}_{bcd} = \frac{\alpha}{2} F_{ab} F_{cd} - \frac{\beta}{2} G_{ab} G_{cd} \quad (40)$$

where $\alpha, \beta \in \mathbb{R}$, $\alpha \neq 0 \neq \beta$.

Class C The curvature tensor is said to be of (curvature) class C at $m \in M$ if the range of f is 2- or 3-dimensional and if there exists $0 \neq k \in T_m M$ such that each of the type $(1, 1)$ tensors in the range of f contains k in its kernel (i.e. each of their matrix representations F satisfies $F^a{}_b k^b = 0$).

Class D The curvature tensor is said to be of (curvature) class D at $m \in M$ if the range of f is 1-dimensional. It follows that the curvature components satisfy $R_{abcd} = \lambda F_{ab} F_{cd}$ at m ($0 \neq \lambda \in \mathbb{R}$) for some bivector F at m which then satisfies $F_{a[b} F_{cd]} = 0$ and is thus simple.

Class O The curvature tensor is said to be of (curvature) class O at $m \in M$ if it vanishes at m .

The following results are useful [3]:

1. For the classes A and B there does **not** exist $0 \neq k \in T_m M$ such that $F^a{}_b k^b = 0$ for *every* F in the range of f .
2. For class A , the range of f has dimension at least two and if this dimension is four or more the class is necessarily A .

3. The vector k in the definition of class C is unique up to a scaling.
4. For the classes A and B there does **not** exist $0 \neq k \in T_m M$ such that $R^a{}_{bcd}k^d = 0$, whereas this equation has exactly one independent solution for class C and two for class D .
5. The five classes A, B, C, D and O are mutually exclusive and exhaustive for the curvature tensor at m . If the curvature class is the same at each $m \in M$ then M will be said to be of that class.

D.2 Properties

Suppose that the components of the Riemann tensor $R^a{}_{bcd}$ are given in a coordinate domain U with metric h . Suppose that h' is another metric with the same components $R^a{}_{bcd}$. It follows from [17] that:

$$\text{Class A} \quad h'_{ab} = \alpha h_{ab} \tag{42a}$$

$$\text{Class B} \quad h'_{ab} = \alpha h_{ab} + 2\beta l_{(a}n_{b)} = (\alpha + \beta)h_{ab} - \beta(x_a x_b + y_a y_b) \tag{42b}$$

$$\text{Class C} \quad h'_{ab} = \alpha h_{ab} + \beta k_a k_b \tag{42c}$$

$$\text{Class D} \quad h'_{ab} = \alpha h_{ab} + \beta r_a r_b + \gamma s_a s_b + 2\delta r_{(a} s_{b)} \tag{42d}$$

where $\alpha, \beta, \gamma, \delta \in \mathbb{R}$.

Note that $h_{ab;c} = h_{ab}w_c$ for some smooth 1-form w on the open subset A . Using condition (2) above and the Ricci identity we get $h_{ab;[cd]} = 0$, which implies $h_{ab}w_{[c;d]} = 0$; thus $w_{[c;d]} = 0$ and so w_a is locally a gradient. Hence, for each $m \in A$, there is an open neighborhood W of m on which $w_a = w_{,c}$ for some smooth function w . Then on W , $g_{ab} = e^{-w}h_{ab}$ satisfies $g_{ab;c} = 0$. Further, if g' is any other local metric defined on some neighborhood W' of m and compatible with Γ then g' satisfies condition (2) on W' and hence, on $W \cap W'$, $g' = \phi g$ for some positive smooth function ϕ . From this and the result $g'_{ab;c} = 0$ it follows that g' is a constant multiple of g on $W \cap W'$.

References

- [1] A.Z. Petrov, *Einstein spaces* (Pergamon, 1969)
- [2] H. Stephani, D. Kramer, M. A. H. MacCallum, C. A. Hoenselaers, E. Herlt 2003 *Exact solutions of Einstein's field equations, second edition* (Cambridge University Press; Cambridge).
- [3] G S Hall, 2004, *Symmetries and curvature structure in general Relativity* (World Science, Singapore).
- [4] A. Coley, 2008, *Class. Quant. Grav.* **25**, 033001.
- [5] R. Milson, A. Coley, V. Pravda and A. Pravdova, 2005, *Int. J. Geom. Meth. Mod. Phys.* **2**, 41.
- [6] A. Coley, R. Milson, V. Pravda and A. Pravdova, 2004, *Class. Quant. Grav.* **21**, L35.
- [7] A. Coley, S. Hervik and N. Pelavas, 2006, *Class. Quant. Grav.* **23**, 3053.

- [8] V. Pravda, A. Pravdová, A. Coley and R. Milson, 2002, *Class. Quant. Grav.* **19**, 6213.
- [9] A. Coley, R. Milson, V. Pravda and A. Pravdova, 2004, *Class. Quant. Grav.* **21**, 5519.
- [10] A. Coley, A. Fuster, S. Hervik, N. Pelavas, 2006, *Class. Quant. Grav.* **23**, 7431
- [11] F. Prüfer, F. Tricerri and L. Vanhecke, 1996, *Trans. American Math. Soc.*, **348**, 4643.
- [12] S Kobayashi and K. Nomizu, 1963, *Foundations of Differential Geometry, Volume 1* (Interscience Publishers).
- [13] A. Coley, S. Hervik and N. Pelavas, 2008, *Class. Quantum Grav.* **25**, 025008.
- [14] A. Coley, S. Hervik and N. Pelavas, 2009, *Class.Quant.Grav.* **29** 125011.
- [15] E. Zakhary and J. Carminati, 2001, *J. Math. Phys.* **42**, 1474; J. Carminati, E. Zakhary, and R. G. McLenaghan, 2002, *J. Math. Phys.* **43**, 492; J. Carminati and E. Zakhary, 2002, *J. Math. Phys.* **43**, 4020
- [16] F. M. Paiva, M. J. Reboucas and M. A. H. MacCallum, 1993, *Class. Quant. Grav.* **10**, 1165.
- [17] G S Hall and W Kay, 1988, *J. Math. Phys.* **29**, 428
- [18] R. Milson and N. Pelavas, 2008, *Class. Quantum Grav.* **25** 012001; *ibid.* arXiv:0711.3851 ; *ibid.* arXiv:gr-qc/0702152.
- [19] J. M. Martin-Garca, D. Yllanes and R. Portugal, 2008, *Comp. Phys. Commun.* **179** 586-590, arXiv:0802.1274 [cs.SC]
- [20] This is a package which runs within Maple. It is entirely distinct from packages distributed with Maple and must be obtained independently. The GRTensorII software and documentation is distributed freely on the World-Wide-Web from the address <http://grtensor.org>