

Global well-posedness for a modified critical dissipative quasi-geostrophic equation

Changxing Miao* and Liutang Xue†

Abstract

In this paper we consider the following modified quasi-geostrophic equation

$$\partial_t \theta + u \cdot \nabla \theta + \nu |D|^\alpha \theta = 0, \quad u = |D|^{\alpha-1} \mathcal{R}^\perp \theta, \quad x \in \mathbb{R}^2$$

with $\nu > 0$ and $\alpha \in]0, 1[\cup]1, 2[$. When $\alpha \in]0, 1[$, the equation was firstly introduced by Constantin, Iyer and Wu in [10]. Here, by using the modulus of continuity method, we prove the global well-posedness of the system with the smooth initial data. As a byproduct, we also show that for every $\alpha \in]0, 2[$, the Lipschitz norm of the solution has a uniform exponential bound.

MSC(2000): 76U05, 76B03, 35Q35

Keywords: Modified quasi-geostrophic equation, Modulus of continuity, Blow-up criterion, Global well-posedness.

1 Introduction

In this paper we focus on the following modified 2D dissipative quasi-geostrophic equation

$$\begin{cases} \partial_t \theta + u \cdot \nabla \theta + \nu |D|^\alpha \theta = 0 \\ u = |D|^{\alpha-1} \mathcal{R}^\perp \theta, \quad \theta|_{t=0} = \theta_0(x) \end{cases} \quad (1.1)$$

with $\nu > 0$, $\alpha \in]0, 1[\cup]1, 2[$, $|D|^\beta = (-\Delta)^{\frac{\beta}{2}}$ is defined via the Fourier transform

$$\widehat{|D|^\beta f}(\zeta) = |\zeta|^\beta \hat{f}(\zeta)$$

and

$$\mathcal{R}^\perp \theta = (-\mathcal{R}_2 \theta, \mathcal{R}_1 \theta) = |D|^{-1} (\partial_2 \theta, -\partial_1 \theta)$$

where $\mathcal{R}_i (i = 1, 2)$ are the usual Riesz transforms (cf. [15]).

When $\alpha = 0$, this model describes the evolution of the vorticity of a two dimensional damped inviscid incompressible fluid. The case of $\alpha = 1$ just is the critical dissipative

*Institute of Applied Physics and Computational Mathematics, P.O. Box 8009, Beijing 100088, P.R. China. Email: miao.changxing@iapcm.ac.cn.

†The Graduate School of China Academy of Engineering Physics, P.O. Box 2101, Beijing 100088, P.R. China. Email: xue_lt@163.com.

quasi-geostrophic equation which arises in the geostrophic study of rotating fluids (cf. [7]). Although when $\alpha = 2$ the flow term in (1.1) vanishes, we can still view the model introduced in [16] as a meaningful generalization of this endpoint case, where the model is derived from the study of the full magnetohydrodynamic equations and the divergence-free three-dimensional velocity u satisfies $u = M[\theta]$ with M a nonlocal differential operator of order 1.

For convenience, we here recall the well-known 2D quasi-geostrophic equation

$$(QG)_\alpha \quad \begin{cases} \partial_t \theta + u \cdot \nabla \theta + \nu |D|^\alpha \theta = 0 \\ u = \mathcal{R}^\perp \theta, \quad \theta(0, x) = \theta_0(x) \end{cases}$$

where $\nu \geq 0$ and $0 \leq \alpha \leq 2$. When $\nu > 0$, $\alpha \in]0, 1[\cup]1, 2[$, we observe that the system (1.1) is almost the same with the quasi-geostrophic equation, and its only difference lies on introducing an extra $|D|^{\alpha-1}$ in the definition of u . When $\alpha \in]0, 1[$, $|D|^{\alpha-1}$ is a negative derivative operator and always plays a good role; while when $\alpha \in]1, 2[$, $|D|^{\alpha-1}$ is a positive derivative operator and always takes a bad part. Moreover, corresponding to the dissipation operator $|D|^\alpha$, this additional operator makes the equation $(QG)_\alpha$ be a new balanced state: the flow term $u \cdot \nabla \theta$ scale the same way as the dissipative term $|D|^\alpha \theta$, i.e., the equation (1.1) is scaling invariant under the transformation

$$\theta(t, x) \rightarrow \theta_\lambda(t, x) := \theta(\lambda^\alpha t, \lambda x), \quad \text{with } \lambda > 0.$$

We note that in the sense of scaling invariance, the system (1.1) is similar to the critical dissipative quasi-geostrophic equation.

Recently, when $\alpha \in]0, 1[$, Constantin, Iyer and Wu in [10] introduced this modified quasi-geostrophic equation and proved the global regularity of Leray-Hopf weak solutions to the system with L^2 initial data. Basically, they use the method from Caffarelli-Vasseur [3] which deals with the same issue of 2D critical dissipative quasi-geostrophic equation $(QG)_1$. We also remark that partially because of its simple form and its internal analogy with the 3D Euler/Navier-Stokes equations, the quasi-geostrophic equation $(QG)_\alpha$, especially the critical one $(QG)_1$, has been extensively considered (see e.g. [1, 3, 6, 7, 8, 9, 11, 13, 17, 23] and references therein). While global regularity of Navier-Stokes equations remains an outstanding challenge in mathematical physics, the global issue of the 2D critical dissipative quasi-geostrophic equation has been in a satisfactory state. In [9] Constantin, Cordoba and Wu showed the global well-posedness of the classical solution under the condition that the zero-dimensional L^∞ norm of the data is small. This smallness assumption was firstly removed by Kiselev, Nazarov and Volberg in [17], where they obtained the global well-posedness for the arbitrary periodic smooth initial data by using a modulus of continuity method. Almost at the same time, Caffarelli and Vasseur in [3] resolved the problem to establish the global regularity of weak solutions associated with L^2 initial data by exploiting the De Giorgi method. We also cite the work of Abidi-Hmidi [1] and Dong-Du [13], as extended work of [17], in which the authors proved the global well-posedness with the initial data belonging to the (critical) space $\dot{B}_{\infty,1}^0$ and H^1 respectively without the additional periodic assumption.

The main goal in this paper is to prove the global well-posedness of the smooth solutions for the system (1.1) with $\alpha \in]0, 1[\cup]1, 2[$. In contrast with the work of [10], we here basically follow the pathway of [17] to obtain the global results by constructing suitable moduli of continuity. Precisely, we have

Theorem 1.1. Let $\nu > 0$, $\alpha \in]0, 2[$ and $\theta_0 \in H^m$, $m > 2$, then there exists a unique global solution

$$\theta \in \mathcal{C}([0, \infty[; H^m) \cap L_{\text{loc}}^2([0, \infty[; H^{m+\frac{\alpha}{2}}) \cap \mathcal{C}^\infty(]0, \infty[\times \mathbb{R}^2)$$

to the modified quasi-geostrophic equation (1.1). Moreover, we get the uniform bound of the Lipschitz norm

$$\sup_{t \geq 0} \|\nabla \theta(t)\|_{L^\infty} \leq C \|\nabla \theta_0\|_{L^\infty} \exp\{C \|\theta_0\|_{L^\infty}\}, \quad (1.2)$$

where C is an absolute constant depending only on α, ν .

The proof is divided into two parts. First through applying the classical method, we obtain the local existence results (Proposition 4.1) and further build the blowup criterion (Proposition 4.2). Then we adopt the nonlocal maximum principle method of Kiselev-Nazarov-Volberg and finally manage to remove all the possible breakdown scenarios by constructing suitable moduli of continuity.

Remark 1.1. The main new ingredient in the global existence consists of modulus of continuity with the explicit formula (5.11) which is suitable for every $\alpha \in]0, 2[$. This MOC has a logarithmic growth near infinity, and further yields the uniform exponential bound of the Lipschitz norm of the solution. In particular, when $\alpha = 1$, (1.2) is a slight improvement of the corresponding bound in [20], where it is a double exponential type.

Remark 1.2. When $\alpha \in]1, 2[$, from the viewpoint of weak solutions, the authors in [22] find that the regularity criterion of (1.1) in terms of Hölder continuous solutions is worse than that of $(QG)_1$, that is, we a priori need $\theta \in L^\infty([t_0, t_1]; C^\sigma(\mathbb{R}^2))$ with $\sigma > \frac{\alpha-1}{2}$ to ensure that θ is a smooth solution in $]t_0, t_1]$ (in contrast with $\sigma > 0$ when $\alpha = 1$). Thus if we rely on this criterion, it is not sufficient to obtain the global regularity of (1.1) with $\alpha \in]1, 2[$ by merely applying the method of [3].

The paper is organized as follows. In Section 2, we present some preparatory results. In Section 3, some facts about modulus of continuity are discussed. In Section 4, we obtain the local results and establish blowup criterion. Finally, we prove the global existence in Section 5.

2 Preliminaries

In this preparatory section, we present the definitions and some related results of the Sobolev spaces and the Besov spaces, also we provide some important estimates which will be used later.

We begin by introducing some notations.

◇ Throughout this paper C stands for a constant which may be different from line to line. We sometimes use $A \lesssim B$ instead of $A \leq CB$, and use $A \lesssim_{\beta, \gamma, \dots} B$ instead of $A \leq C(\beta, \gamma, \dots)B$ with $C(\beta, \gamma, \dots)$ a constant depending on β, γ, \dots . For $A \approx B$ we mean $A \lesssim B \lesssim A$.

◇ Denote by $\mathcal{S}(\mathbb{R}^n)$ the Schwartz space of rapidly decreasing smooth functions, $\mathcal{S}'(\mathbb{R}^n)$ the space of tempered distributions, $\mathcal{S}'(\mathbb{R}^n)/\mathcal{P}(\mathbb{R}^n)$ the quotient space of tempered distributions which modulo polynomials.

◇ $\mathcal{F}f$ or \hat{f} denotes the Fourier transform, that is $\mathcal{F}f(\zeta) = \hat{f}(\zeta) = \int_{\mathbb{R}^n} e^{-ix \cdot \zeta} f(x) dx$, while $\mathcal{F}^{-1}f$ the inverse Fourier transform, namely, $\mathcal{F}^{-1}f(x) = (2\pi)^{-n} \int_{\mathbb{R}^n} e^{ix \cdot \zeta} f(\zeta) d\zeta$.

Now we give the definition of L^2 based Sobolev space. For $s \in \mathbb{R}$, the inhomogeneous Sobolev space

$$H^s := \left\{ f \in \mathcal{S}'(\mathbb{R}^n); \|f\|_{H^s}^2 := \int_{\mathbb{R}^n} (1 + |\zeta|^2)^s |\hat{f}(\zeta)|^2 d\zeta < \infty \right\}$$

Also one can define the corresponding homogeneous space:

$$\dot{H}^s := \left\{ f \in \mathcal{S}'(\mathbb{R}^n)/\mathcal{P}(\mathbb{R}^n); \|f\|_{\dot{H}^s}^2 := \int_{\mathbb{R}^n} |\zeta|^{2s} |\hat{f}(\zeta)|^2 d\zeta < \infty \right\}$$

The following calculus inequality is well-known(see [2])

Lemma 2.1. $\forall m \in \mathbb{R}^+$, there exists a constant $c_m > 0$ such that

$$\|fg\|_{H^m} \leq c_m (\|f\|_{L^\infty} \|g\|_{H^m} + \|f\|_{H^m} \|g\|_{L^\infty}). \quad (2.1)$$

To define Besov space we need the following dyadic partition of unity (see e.g. [5]). Choose two nonnegative radial functions $\chi, \varphi \in \mathcal{D}(\mathbb{R}^n)$ be supported respectively in the ball $\{\zeta \in \mathbb{R}^n : |\zeta| \leq \frac{4}{3}\}$ and the shell $\{\zeta \in \mathbb{R}^n : \frac{3}{4} \leq |\zeta| \leq \frac{8}{3}\}$ such that

$$\chi(\zeta) + \sum_{j \geq 0} \varphi(2^{-j}\zeta) = 1, \quad \forall \zeta \in \mathbb{R}^n; \quad \sum_{j \in \mathbb{Z}} \varphi(2^{-j}\zeta) = 1, \quad \forall \zeta \neq 0.$$

For all $f \in \mathcal{S}'(\mathbb{R}^n)$ we define the nonhomogeneous Littlewood-Paley operators

$$\Delta_{-1}f := \chi(D)f; \quad \Delta_j f := \varphi(2^{-j}D)f, \quad S_j f := \sum_{-1 \leq k \leq j-1} \Delta_k f, \quad \forall j \in \mathbb{N},$$

And the homogeneous Littlewood-Paley operators can be defined as follows

$$\dot{\Delta}_j f := \varphi(2^{-j}D)f; \quad \dot{S}_j f := \sum_{k \in \mathbb{Z}, k \leq j-1} \dot{\Delta}_k f \quad \forall j \in \mathbb{Z}.$$

Now we introduce the definition of Besov spaces . Let $(p, r) \in [1, \infty]^2$, $s \in \mathbb{R}$, the nonhomogeneous Besov space

$$B_{p,r}^s := \left\{ f \in \mathcal{S}'(\mathbb{R}^n); \|f\|_{B_{p,r}^s} := \left\| \{2^{js} \|\Delta_j f\|_{L^p}\}_{j \geq -1} \right\|_{\ell^r} < \infty \right\}$$

and the homogeneous space

$$\dot{B}_{p,r}^s := \left\{ f \in \mathcal{S}'(\mathbb{R}^n)/\mathcal{P}(\mathbb{R}^n); \|f\|_{\dot{B}_{p,r}^s} := \left\| \{2^{js} \|\dot{\Delta}_j f\|_{L^p}\}_{j \in \mathbb{Z}} \right\|_{\ell^r(\mathbb{Z})} < \infty \right\}.$$

We point out that for all $s \in \mathbb{R}$, $B_{2,2}^s = H^s$ and $\dot{B}_{2,2}^s = \dot{H}^s$.

The classical space-time Besov space $L^\rho([0, T], B_{p,r}^s)$, abbreviated by $L_T^\rho B_{p,r}^s$, is the set of tempered distribution f such that

$$\|f\|_{L_T^\rho B_{p,r}^s} := \left\| \left\| \{2^{js} \|\Delta_j f\|_{L^p}\}_{j \geq -1} \right\|_{\ell^r} \right\|_{L^\rho([0, T])} < \infty.$$

We can similarly extend to the homogeneous one $L_T^\rho \dot{B}_{p,r}^s$.

Bernstein's inequality is fundamental in the analysis involving Besov spaces (see [5])

Lemma 2.2. *Let $f \in L^a$, $1 \leq a \leq b \leq \infty$. Then for every $(k, q) \in \mathbb{N}^2$ there exists a constant $C > 0$ such that*

$$\sup_{|\alpha|=k} \|\partial^\alpha S_q f\|_{L^b} \leq C 2^{q(k+n(\frac{1}{a}-\frac{1}{b}))} \|f\|_{L^a},$$

$$C^{-1} 2^{qk} \|f\|_{L^a} \leq \sup_{|\alpha|=k} \|\partial^\alpha \Delta_q f\|_{L^a} \leq C 2^{qk} \|f\|_{L^a}$$

Finally we state an important maximum principle for the transport-diffusion equation (cf. [11])

Proposition 2.3. *Let u be a smooth divergence-free vector field and f be a smooth function. Assume that θ is the smooth solution of the equation*

$$\partial_t \theta + u \cdot \nabla \theta + \nu |D|^\alpha \theta = f, \quad \operatorname{div} u = 0,$$

with initial datum θ_0 and $\nu \geq 0$, $0 \leq \alpha \leq 2$, then for every $p \in [1, \infty]$ we have

$$\|\theta(t)\|_{L^p} \leq \|\theta_0\|_{L^p} + \int_0^t \|f(\tau)\|_{L^p} d\tau. \quad (2.2)$$

3 Moduli of Continuity

In this section, we discuss the moduli of continuity which play a key role in our global existence part.

We suppose that ω is a modulus of continuity, that is, a continuous, increasing, concave function on $[0, \infty)$ such that $\omega(0) = 0$. We say that a function $f : \mathbb{R}^n \rightarrow \mathbb{R}^m$ has modulus of continuity if $|f(x) - f(y)| \leq \omega(|x - y|)$ for all $x, y \in \mathbb{R}^n$ and that f has strict modulus of continuity if the inequality is strict for $x \neq y$.

Next we introduce the pseudo-differential operators $\mathcal{R}_{\alpha, j}$ which may be termed as the modified Riesz transforms

Proposition 3.1. *Let $\alpha \in]0, 2[$, $1 \leq j \leq n$, $n \geq 2$, then for every $f \in \mathcal{S}(\mathbb{R}^n)$*

$$\mathcal{R}_{\alpha, j} f(x) = |D|^{\alpha-1} \mathcal{R}_j f(x) = c_{\alpha, n} \text{p.v.} \int_{\mathbb{R}^n} \frac{y_j}{|y|^{n+\alpha}} f(x-y) dy, \quad (3.1)$$

where $c_{\alpha, n}$ is the normalization constant such that

$$\widehat{\mathcal{R}_{\alpha, j} f}(\zeta) = -i \frac{\zeta_j}{|\zeta|^{2-\alpha}} \hat{f}(\zeta).$$

The proof is placed in the appendix. Also note that when $\alpha \in]0, 1[$, we do not need to introduce the principle value of integral expression in the formula (3.1).

The pseudo-differential operators like the modified Riesz transforms do not preserve the moduli of continuity generally, but they also do not destroy them too much either. Precisely, similarly as the Lemma in [17], we have

Lemma 3.2. *If the function θ has the modulus of continuity ω , then $u = (-\mathcal{R}_{\alpha,2}\theta, \mathcal{R}_{\alpha,1}\theta)$ ($\alpha \in]0, 2[$) has the modulus of continuity*

$$\Omega(\xi) = A_\alpha \left(\int_0^\xi \frac{\omega(\eta)}{\eta^\alpha} d\eta + \xi \int_\xi^\infty \frac{\omega(\eta)}{\eta^{1+\alpha}} d\eta \right) \quad (3.2)$$

with some absolute constant $A_\alpha > 0$ that may depend on α .

Proof. The modified Riesz transforms are pseudo-differential operators with kernels $K(x) = \frac{S(x')}{|x|^{n-1+\alpha}}$ (in our special case, $n = 2$ and $S(x') = \frac{x_j}{|x|}$, $j = 1, 2$), where $x' = \frac{x}{|x|} \in \mathbb{S}^{n-1}$. The function $S \in C^1(\mathbb{S}^{n-1})$ and $\int_{\mathbb{S}^{n-1}} S(x') d\sigma(x') = 0$. Assume that the function $f : \mathbb{R}^n \rightarrow \mathbb{R}^m$ has some modulus of continuity ω , that is $|f(x) - f(y)| \leq \omega(|x - y|)$ for all $x, y \in \mathbb{R}^n$. Then take any x, y with $|x - y| = \xi$, and consider the difference

$$\int K(x - t)f(t)dt - \int K(y - t)f(t)dt. \quad (3.3)$$

First due to the cancelation property of S we have

$$\left| \int_{|x-t| \leq 2\xi} K(x-t)f(t)dt \right| = \left| \int_{|x-t| \leq 2\xi} K(x-t)(f(t) - f(x))dt \right| \leq C \int_0^{2\xi} \frac{\omega(r)}{r^\alpha} dr$$

since ω is concave, we obtain

$$\int_0^{2\xi} \frac{\omega(r)}{r^\alpha} dr \leq 2^{2-\alpha} \int_0^\xi \frac{\omega(r)}{r^\alpha} dr \quad (3.4)$$

A similar estimate holds for the second integral in (3.3). Next, set $z = \frac{x+y}{2}$, then

$$\begin{aligned} & \left| \int_{|x-t| \geq 2\xi} K(x-t)f(t)dt - \int_{|y-t| \geq 2\xi} K(y-t)f(t)dt \right| \\ &= \left| \int_{|x-t| \geq 2\xi} K(x-t)(f(t) - f(z))dt - \int_{|y-t| \geq 2\xi} K(y-t)(f(t) - f(z))dt \right| \\ &\leq \int_{|z-t| \geq 3\xi} |K(x-t) - K(y-t)| |f(t) - f(z)| dt \\ &\quad + \int_{\frac{3\xi}{2} \leq |z-t| \leq 3\xi} (|K(x-t)| + |K(y-t)|) |f(t) - f(z)| dt \\ &= I_1 + I_2 \end{aligned}$$

To estimate the first integral, we use the smoothness condition of S to get

$$|K(x-t) - K(y-t)| \leq C \frac{|x-y|}{|z-t|^{n+\alpha}} \quad \text{when } |z-t| \geq 3\xi$$

thus

$$I_1 \leq C\xi \int_{3\xi}^\infty \frac{\omega(r)}{r^{1+\alpha}} dr \leq C3^{-\alpha}\xi \int_\xi^\infty \frac{\omega(3r)}{r^{1+\alpha}} dr \leq C\xi \int_\xi^\infty \frac{\omega(r)}{r^{1+\alpha}} dr$$

For the second integral, using the concavity of ω and (3.4), we have

$$\begin{aligned} I_2 &\leq 2C\omega(3\xi)\xi^{1-\alpha} \int_{\xi \leq |x-t| \leq \frac{7}{2}\xi} \frac{1}{|x-t|^n} dt \\ &\leq C\omega(\xi)\xi^{1-\alpha} \leq C2^\alpha \int_\xi^{2\xi} \frac{\omega(r)}{r^\alpha} dr \leq C \int_0^\xi \frac{\omega(r)}{r^\alpha} dr \end{aligned}$$

□

Now we consider the action of the fractional differential operators $|D|^\alpha$ ($\alpha \in]0, 2[$) on the function having modulus of continuity. Precisely,

Lemma 3.3. *If the function $\theta : \mathbb{R}^2 \rightarrow \mathbb{R}$ has modulus of continuity ω , and especially satisfies $\theta(x) - \theta(y) = \omega(\xi)$ at some $x, y \in \mathbb{R}^2$ with $|x - y| = \xi > 0$, then we have*

$$\begin{aligned} [(-|D|^\alpha)\theta](x) - [(-|D|^\alpha)\theta](y) &\leq B_\alpha \int_0^{\frac{\xi}{2}} \frac{\omega(\xi + 2\eta) + \omega(\xi - 2\eta) - 2\omega(\xi)}{\eta^{1+\alpha}} d\eta \\ &+ B_\alpha \int_{\frac{\xi}{2}}^\infty \frac{\omega(2\eta + \xi) - \omega(2\eta - \xi) - 2\omega(\xi)}{\eta^{1+\alpha}} d\eta \end{aligned} \quad (3.5)$$

where $B_\alpha > 0$ is an absolute constant.

Remark 3.1. In fact this result has occurred in [24], as a generalization of the one in [17]. For convenience, we prove it again for the general n -dimensional case and place the proof in the appendix. Also note that due to concavity of ω , both terms on the righthand side of (3.5) are strictly negative.

4 Local existence and Blowup criterion

Our purpose in this section is to prove the following local result

Proposition 4.1. *Let $\nu > 0$, $0 < \alpha < 2$ and the initial data $\theta_0 \in H^m$, $m > 2$. Then there exists a positive T depending only on α , ν and $\|\theta_0\|_{H^m}$ such that the modified quasi-geostrophic equation (1.1) generates a unique solution $\theta \in \mathcal{C}([0, T], H^m) \cap L^2([0, T], H^{m+\frac{\alpha}{2}})$. Moreover we have $t^\gamma \theta \in L^\infty([0, T], H^{m+\gamma\alpha})$ for all $\gamma \geq 0$, which implies $\theta \in \mathcal{C}^\infty([0, T] \times \mathbb{R}^2)$.*

We further obtain the following criterion for the breakdown of smooth solutions

Proposition 4.2. *Let T^* be the maximal existence time of θ in $\mathcal{C}([0, T^*), H^m) \cap L^2([0, T^*), H^{m+\frac{\alpha}{2}})$. If $T^* < \infty$ then we necessarily have*

$$\int_0^{T^*} \|\nabla\theta(t, \cdot)\|_{L^\infty}^\alpha dt = \infty. \quad (4.1)$$

The method of proof for the Proposition 4.1 is to regularize the equation (1.1) by the standard Friedrich method, and then pass to the limit for the regularization parameter.

Denote the frequency cutoff operator $\mathcal{J}_\epsilon : L^2(\mathbb{R}^2) \rightarrow H^m(\mathbb{R}^2)$, $\epsilon > 0$, $m \geq 0$ by

$$(\mathcal{J}_\epsilon f)(x) = \mathcal{F}^{-1}(\hat{f}(\cdot)1_{B_{1/\epsilon}}(\cdot))(x) = (2\pi)^{-2} \int_{\mathbb{R}^2} e^{ix \cdot \zeta} \hat{f}(\zeta) 1_{\{|\cdot| \leq \frac{1}{\epsilon}\}}(\zeta) d\zeta.$$

The following properties of \mathcal{J}_ϵ are obvious.

Lemma 4.3. *Let \mathcal{J}_ϵ be the projection operator defined as above, $m \in \mathbb{R}^+$, $k \in \mathbb{R}^+$, $\delta \in [0, m[$. Then*

$$(i) \text{ for all } f \in H^m, \lim_{\epsilon \rightarrow 0} \|\mathcal{J}_\epsilon f - f\|_{H^m} = 0.$$

- (ii) for all $f \in H^m$, $|D|^m(\mathcal{J}_\epsilon f) = \mathcal{J}_\epsilon(|D|^m f)$ and $\Delta_j(\mathcal{J}_\epsilon f) = \mathcal{J}_\epsilon(\Delta_j f)$.
- (iii) for all $f \in H^m$, $\|\mathcal{J}_\epsilon f - f\|_{H^{m-\delta}} \lesssim \epsilon^\delta \|f\|_{H^m}$ and $\|\mathcal{J}_\epsilon f\|_{H^{m+k}} \lesssim \frac{1}{\epsilon^k} \|f\|_{H^m}$.

Then we regularize the modified quasi-geostrophic equation (1.1) as follows

$$\begin{cases} \theta_t^\epsilon + \mathcal{J}_\epsilon((\mathcal{J}_\epsilon u^\epsilon) \cdot \nabla(\mathcal{J}_\epsilon \theta^\epsilon)) + \nu \mathcal{J}_\epsilon |D|^\alpha \theta^\epsilon = 0 \\ u^\epsilon = |D|^{\alpha-1} \mathcal{R}^\perp \theta^\epsilon, \quad \theta^\epsilon|_{t=0} = \mathcal{J}_\epsilon \theta_0. \end{cases} \quad (4.2)$$

For this approximate system, we have

Proposition 4.4. *Let the initial data $\theta_0 \in L^2$. Then for any $\epsilon > 0$ there exists a unique global solution $\theta^\epsilon \in \mathcal{C}^1([0, \infty[, H^\infty)$ to the regularized equation (4.2).*

Proof. We can write (4.2) as follows

$$\frac{d}{dt} \theta^\epsilon = F_\epsilon(\theta^\epsilon), \quad \theta^\epsilon|_{t=0} = \mathcal{J}_\epsilon \theta_0, \quad (4.3)$$

with

$$F_\epsilon(\theta^\epsilon) = -\mathcal{J}_\epsilon((\mathcal{J}_\epsilon u^\epsilon) \cdot \nabla(\mathcal{J}_\epsilon \theta^\epsilon)) - \nu \mathcal{J}_\epsilon |D|^\alpha \theta^\epsilon.$$

For every $\epsilon > 0$, we can show that

$$\|F_\epsilon(f)\|_{L^2} \lesssim_{\epsilon, \nu} \|f\|_{L^2} + \|f\|_{L^2}^2,$$

and

$$\|F_\epsilon(f_1, f_2)\|_{L^2} \lesssim_{\epsilon, \nu, \|f_i\|_{L^2}} \|f_1 - f_2\|_{L^2},$$

where f, f_1, f_2 are all in L^2 . This means that F_ϵ maps L^2 into L^2 and F_ϵ is locally Lipschitz continuous on L^2 . Hence the Cauchy-Lipschitz theorem ensures that for every $\theta_0 \in L^2$, there exists a unique solution $\theta^\epsilon \in \mathcal{C}^1([0, T_\epsilon[, L^2)$ with $T_\epsilon > 0$ is the maximum existence time.

Moreover, using the L^2 energy method, from $\operatorname{div} u^\epsilon = 0$ and $\mathcal{J}_\epsilon \theta^\epsilon \in \mathcal{C}^1([0, T_\epsilon[, H^\infty)$, we get

$$\frac{1}{2} \frac{d}{dt} \|\theta^\epsilon\|_{L^2}^2 + \| |D|^{\alpha/2} \mathcal{J}_\epsilon \theta^\epsilon \|_{L^2}^2 = 0$$

Thus

$$\sup_{t \in [0, T_\epsilon[} \|\theta^\epsilon(t)\|_{L^2} \leq \|\mathcal{J}_\epsilon \theta_0\|_{L^2} \leq \|\theta_0\|_{L^2}.$$

Then the classical continuation criterion guarantees $T_\epsilon = \infty$.

Moreover, since $\mathcal{J}_\epsilon \theta^\epsilon$ is also a solution of (4.2), from the uniqueness of θ^ϵ we find $\theta^\epsilon = \mathcal{J}_\epsilon \theta^\epsilon$. \square

Remark 4.1. From the proof we know $\theta^\epsilon = \mathcal{J}_\epsilon \theta^\epsilon$, thus (4.2) will be written as follows

$$\begin{cases} \theta_t^\epsilon + \mathcal{J}_\epsilon(u^\epsilon \cdot \nabla \theta^\epsilon) + \nu |D|^\alpha \theta^\epsilon = 0 \\ u^\epsilon = |D|^{\alpha-1} \mathcal{R}^\perp \theta^\epsilon, \quad \theta^\epsilon|_{t=0} = \mathcal{J}_\epsilon \theta_0. \end{cases} \quad (4.4)$$

In the sequel we shall instead consider this form.

Next, we prove the main result in this section.

Proof of Proposition 4.1. Step 1: Uniform Bounds.

We claim that: the regularized solution $\theta^\epsilon \in \mathcal{C}^1([0, \infty[, H^\infty)$ to equation (4.2) satisfies

$$\frac{d}{2dt} \|\theta^\epsilon\|_{B_{2,2}^m}^2 + \frac{\nu}{2} \left\| |D|^{\frac{\alpha}{2}} \theta^\epsilon \right\|_{B_{2,2}^m}^2 \lesssim_{\nu, \alpha} \frac{1}{\nu} \|\nabla \theta^\epsilon\|_{L^\infty}^\alpha \|\theta^\epsilon\|_{L^\infty}^{2-\alpha} \|\theta^\epsilon\|_{B_{2,2}^m}^2 + \|\theta^\epsilon\|_{L^2}^2 \|\theta^\epsilon\|_{B_{2,2}^m}. \quad (4.5)$$

Indeed, for every $q \in \mathbb{N}$, applying dyadic operator Δ_q to both sides of regularized equation (4.4) yields

$$\partial_t \Delta_q \theta^\epsilon + \mathcal{J}_\epsilon((S_{q+1} u^\epsilon) \cdot \nabla \Delta_q \theta^\epsilon) + \nu |D|^\alpha \Delta_q \theta^\epsilon = \mathcal{J}_\epsilon(F_q(u^\epsilon, \theta^\epsilon)),$$

where

$$F_q(u^\epsilon, \theta^\epsilon) = (S_{q+1} u^\epsilon) \cdot \nabla \Delta_q \theta^\epsilon - \Delta_q(u^\epsilon \cdot \nabla \theta^\epsilon).$$

Taking the L^2 inner product in the above equality with $\Delta_q \theta^\epsilon$ and using the divergence free property, we have

$$\begin{aligned} \frac{1}{2} \frac{d}{dt} \|\Delta_q \theta^\epsilon\|_{L^2}^2 + \nu \left\| |D|^{\frac{\alpha}{2}} \Delta_q \theta^\epsilon \right\|_{L^2}^2 &\leq \left| \int_{\mathbb{R}^2} (F_q(u^\epsilon, \theta^\epsilon))(x) \mathcal{J}_\epsilon \Delta_q \theta^\epsilon(x) dx \right| \\ &\leq 2^{-q \frac{\alpha}{2}} \|F_q(u^\epsilon, \theta^\epsilon)\|_{L^2} 2^{q \frac{\alpha}{2}} \|\mathcal{J}_\epsilon \Delta_q \theta^\epsilon\|_{L^2} \\ &\lesssim 2^{-q \frac{\alpha}{2}} \|F_q(u^\epsilon, \theta^\epsilon)\|_{L^2} \left\| |D|^{\frac{\alpha}{2}} \Delta_q \theta^\epsilon \right\|_{L^2}. \end{aligned}$$

Then by virtue of Young inequality, we deduce

$$\frac{1}{2} \frac{d}{dt} \|\Delta_q \theta^\epsilon\|_{L^2}^2 + \frac{\nu}{2} \left\| |D|^{\frac{\alpha}{2}} \Delta_q \theta^\epsilon \right\|_{L^2}^2 \leq \frac{C_0}{\nu} \left(2^{-q \frac{\alpha}{2}} \|F_q(u^\epsilon, \theta^\epsilon)\|_{L^2} \right)^2. \quad (4.6)$$

From the inequality (6.2) in the appendix, we know that

$$\begin{aligned} &2^{-q \frac{\alpha}{2}} \|F_q(u^\epsilon, \theta^\epsilon)\|_{L^2} \\ &\lesssim \left\| |D|^{1-\frac{\alpha}{2}} u^\epsilon \right\|_{L^\infty} \sum_{q' \geq q-4} 2^{(q-q')(1-\frac{\alpha}{2})} \|\Delta_{q'} \theta^\epsilon\|_{L^2} + \left\| |D|^{\frac{\alpha}{2}} \theta^\epsilon \right\|_{L^\infty} \sum_{|q'-q| \leq 4} \|\Delta_{q'} \theta^\epsilon\|_{L^2} \end{aligned} \quad (4.7)$$

Also notice that for some number $K \in \mathbb{N}$

$$\begin{aligned} \left\| |D|^{1-\frac{\alpha}{2}} u^\epsilon \right\|_{L^\infty} + \left\| |D|^{\frac{\alpha}{2}} \theta^\epsilon \right\|_{L^\infty} &\lesssim \left\| |D|^{1-\frac{\alpha}{2}} |D|^{\alpha-1} \mathcal{R}^\perp \theta^\epsilon \right\|_{\dot{B}_{\infty,1}^0} + \left\| |D|^{\frac{\alpha}{2}} \theta^\epsilon \right\|_{\dot{B}_{\infty,1}^0} \\ &\lesssim \sum_{k=-\infty}^{K-1} 2^{k\alpha/2} \left\| \dot{\Delta}_k \theta^\epsilon \right\|_{L^\infty} + \sum_{k=K}^{\infty} 2^{-k(1-\frac{\alpha}{2})} \left\| \dot{\Delta}_k \nabla \theta^\epsilon \right\|_{L^\infty} \\ &\lesssim 2^{K\alpha/2} \|\theta^\epsilon\|_{L^\infty} + 2^{K(\frac{\alpha}{2}-1)} \|\nabla \theta^\epsilon\|_{L^\infty}, \end{aligned}$$

thus choosing K satisfying $\|\theta^\epsilon\|_{L^\infty} 2^K \approx \|\nabla \theta^\epsilon\|_{L^\infty}$, we deduce

$$\left\| |D|^{1-\frac{\alpha}{2}} u^\epsilon \right\|_{L^\infty} + \left\| |D|^{\frac{\alpha}{2}} \theta^\epsilon \right\|_{L^\infty} \lesssim \|\nabla \theta^\epsilon\|_{L^\infty}^{\frac{\alpha}{2}} \|\theta^\epsilon\|_{L^\infty}^{1-\frac{\alpha}{2}}. \quad (4.8)$$

Plunging the above two estimates (4.8) and (4.7) into inequality (4.6), then multiplying both sides by 2^{2qm} and summing up over $q \in \mathbb{N}$, we obtain

$$\frac{1}{2} \frac{d}{dt} \sum_{q \in \mathbb{N}} 2^{2qm} \|\Delta_q \theta^\epsilon\|_{L^2}^2 + \frac{\nu}{2} \sum_{q \in \mathbb{N}} 2^{2qm} \left\| |D|^{\frac{\alpha}{2}} \Delta_q \theta^\epsilon \right\|_{L^2}^2 \lesssim \frac{1}{\nu} \|\nabla \theta^\epsilon\|_{L^\infty}^\alpha \|\theta^\epsilon\|_{L^\infty}^{2-\alpha} \|\theta^\epsilon\|_{B_{2,2}^m}^2. \quad (4.9)$$

On the other hand, we apply the low frequency operator Δ_{-1} to the regularized system (4.2) to get

$$\partial_t \Delta_{-1} \theta^\epsilon + \nu |D|^\alpha \Delta_{-1} \theta^\epsilon = -\mathcal{J}_\epsilon \Delta_{-1} (u^\epsilon \cdot \nabla \theta^\epsilon).$$

Multiplying both sides by $\Delta_{-1} \theta^\epsilon$ and integrating in the spatial variable, we obtain

$$\begin{aligned} \frac{1}{2} \frac{d}{dt} \|\Delta_{-1} \theta^\epsilon\|_{L^2}^2 + \nu \left\| |D|^{\frac{\alpha}{2}} \Delta_{-1} \theta^\epsilon \right\|_{L^2}^2 &\leq \left| \int_{\mathbb{R}^2} \operatorname{div} \Delta_{-1} (u^\epsilon \theta^\epsilon)(x) \Delta_{-1} \mathcal{J}_\epsilon \theta^\epsilon(x) dx \right| \\ &\lesssim \|u^\epsilon\|_{L^\infty} \|\theta^\epsilon\|_{L^2}^2. \end{aligned}$$

We see that

$$\begin{aligned} \|u^\epsilon\|_{L^\infty} &\leq \left(\sum_{j \leq -1} + \sum_{j \geq 0} \right) \left\| \dot{\Delta}_j |D|^{\alpha-1} \mathcal{R}^\perp \theta^\epsilon \right\|_{L^\infty} \\ &\lesssim \sum_{j \leq -1} 2^{j\alpha} \left\| \dot{\Delta}_j \theta^\epsilon \right\|_{L^2} + \sum_{j \geq 0} 2^{j(\alpha-2)} \left\| \dot{\Delta}_j \nabla \theta^\epsilon \right\|_{L^\infty} \\ &\lesssim \|\theta^\epsilon\|_{L^2} + \|\nabla \theta^\epsilon\|_{L^\infty}, \end{aligned} \tag{4.10}$$

thus we have

$$\frac{1}{2} \frac{d}{dt} \|\Delta_{-1} \theta^\epsilon\|_{L^2}^2 + \frac{\nu}{2} \left\| |D|^{\frac{\alpha}{2}} \Delta_{-1} \theta^\epsilon \right\|_{L^2}^2 \lesssim \|\theta^\epsilon\|_{B_{2,2}^m} \|\theta^\epsilon\|_{L^2}^2. \tag{4.11}$$

Multiplying (4.11) by 2^{-2m} and combining it with (4.9) leads to (4.5).

Next, we prove that the solution family (θ^ϵ) is uniformly bounded in H^m . Indeed, from estimate (4.5), Besov embedding and the fact that $\|\cdot\|_{B_{2,2}^m}^2 / C_0 \leq \|\cdot\|_{H^m}^2 \leq C_0 \|\cdot\|_{B_{2,2}^m}^2$ with C_0 a universal number, we have

$$\begin{aligned} \frac{d}{dt} \left(\|\theta^\epsilon(t)\|_{H^m}^2 + \int_0^t \|\theta^\epsilon(\tau)\|_{H^{m+\frac{\alpha}{2}}}^2 d\tau \right) &\leq C \left(\|\nabla \theta^\epsilon\|_{L^\infty}^\alpha \|\theta^\epsilon\|_{L^\infty}^{2-\alpha} \|\theta^\epsilon\|_{H^m} + \|\theta^\epsilon\|_{L^2} \right) \|\theta^\epsilon\|_{H^m}^2 \\ &\leq C_1 (1 + \|\theta^\epsilon(t)\|_{H^m}^2) \|\theta^\epsilon(t)\|_{H^m}^2, \end{aligned} \tag{4.12}$$

where C_1 depends only on m, α, ν . Gronwall inequality yields that

$$\sup_{0 \leq t \leq T} \|\theta^\epsilon\|_{H^m}^2 + \|\theta^\epsilon\|_{L_T^2 H^{m+\frac{\alpha}{2}}}^2 \leq \frac{\|\theta_0\|_{H^m}^2}{(\|\theta_0\|_{H^m}^2 + 1)e^{-CT} - \|\theta_0\|_{H^m}^2}. \tag{4.13}$$

Thus for some

$$T < \frac{1}{C} \log(1 + 1/\|\theta_0\|_{H^m}^2),$$

the family (θ^ϵ) is uniformly bounded in $\mathcal{C}([0, T], H^m) \cap L^2([0, T]; H^{m+\frac{\alpha}{2}})$, $m > 2$.

Step 2: Strong Convergence

We firstly claim that the solutions (θ^ϵ) to the approximate equation (4.4) converge in $\mathcal{C}([0, T], L^2(\mathbb{R}^2))$. Indeed for all $0 < \tilde{\epsilon} < \epsilon$, we assume that θ^ϵ and $\theta^{\tilde{\epsilon}}$ are two approximate solutions, then from a direct calculation

$$(\theta_t^\epsilon - \theta_t^{\tilde{\epsilon}}, \theta^\epsilon - \theta^{\tilde{\epsilon}}) = -\nu(|D|^\alpha \theta^\epsilon - |D|^\alpha \theta^{\tilde{\epsilon}}, \theta^\epsilon - \theta^{\tilde{\epsilon}}) - \left((\mathcal{J}_\epsilon(u^\epsilon \cdot \nabla \theta^\epsilon) - \mathcal{J}_{\tilde{\epsilon}}(u^{\tilde{\epsilon}} \cdot \nabla \theta^{\tilde{\epsilon}})), \theta^\epsilon - \theta^{\tilde{\epsilon}} \right),$$

we have

$$\begin{aligned}
& \frac{1}{2} \frac{d}{dt} \|\theta^\epsilon(t) - \theta^{\tilde{\epsilon}}(t)\|_{L^2}^2 + \nu \left\| |D|^{\frac{\alpha}{2}} (\theta^\epsilon - \theta^{\tilde{\epsilon}}) \right\|_{L^2}^2 \\
&= \left((\mathcal{J}_\epsilon - \mathcal{J}_{\tilde{\epsilon}})(u^\epsilon \cdot \nabla \theta^\epsilon), \theta^\epsilon - \theta^{\tilde{\epsilon}} \right) + \left(\mathcal{J}_{\tilde{\epsilon}}((u^\epsilon - u^{\tilde{\epsilon}}) \cdot \nabla \theta^\epsilon), \theta^\epsilon - \theta^{\tilde{\epsilon}} \right) \\
&\quad + \left(\mathcal{J}_{\tilde{\epsilon}}(u^{\tilde{\epsilon}} \cdot \nabla (\theta^\epsilon - \theta^{\tilde{\epsilon}})), \theta^\epsilon - \theta^{\tilde{\epsilon}} \right) \\
&:= II_1 + II_2 + II_3.
\end{aligned}$$

We set $\delta_0 := \min\{m - \alpha, 1\}$, then for II_1 , by means of the calculus inequality (2.1), divergence free condition and the following simple inequality

$$\|u^\epsilon\|_{H^{m-\alpha+1}} = \left\| |D|^{\alpha-1} R^\perp \theta^\epsilon \right\|_{H^{m-\alpha+1}} \lesssim \|\theta^\epsilon\|_{H^m} \lesssim M,$$

we have

$$\begin{aligned}
|II_1| &\lesssim \epsilon^{\delta_0} \|u^\epsilon \theta^\epsilon\|_{H^{1+\delta_0}} \|\theta^\epsilon - \theta^{\tilde{\epsilon}}\|_{L^2} \\
&\lesssim \epsilon^{\delta_0} (\|u^\epsilon\|_{H^{1+\delta_0}} + \|\theta^\epsilon\|_{H^{1+\delta_0}}) \|\theta^\epsilon - \theta^{\tilde{\epsilon}}\|_{L^2} \\
&\lesssim_M \epsilon^{\delta_0} \|\theta^\epsilon - \theta^{\tilde{\epsilon}}\|_{L^2}.
\end{aligned}$$

For II_2 , we directly obtain

$$\begin{aligned}
|II_2| &\leq \|(u^\epsilon - u^{\tilde{\epsilon}}) \cdot \nabla \theta^\epsilon\|_{\dot{H}^{-\frac{\alpha}{2}}} \left\| |D|^{\frac{\alpha}{2}} (\theta^\epsilon - \theta^{\tilde{\epsilon}}) \right\|_{L^2} \\
&\leq C_\alpha \left\| |D|^{\alpha-1} \mathcal{R}^\perp (\theta^\epsilon - \theta^{\tilde{\epsilon}}) \right\|_{\dot{H}^{1-\alpha}}^2 \|\nabla \theta^\epsilon\|_{\dot{H}^{\frac{\alpha}{2}}}^2 + \frac{\nu}{2} \left\| |D|^{\frac{\alpha}{2}} (\theta^\epsilon - \theta^{\tilde{\epsilon}}) \right\|_{L^2}^2 \\
&\leq C_{M,\alpha} \|\theta^\epsilon - \theta^{\tilde{\epsilon}}\|_{L^2}^2 + \frac{\nu}{2} \left\| |D|^{\frac{\alpha}{2}} (\theta^\epsilon - \theta^{\tilde{\epsilon}}) \right\|_{L^2}^2,
\end{aligned}$$

where in the second line we have used the classical product estimate (cf. [14]) that for every $s, t < 1$ and $s + t > 0$,

$$\|fg\|_{\dot{H}^{s+t-1}} \lesssim_{s,t} \|f\|_{\dot{H}^s} \|g\|_{\dot{H}^t}.$$

For the last term, II_3 , from the divergence free fact of $u^{\tilde{\epsilon}}$ and $\mathcal{J}_{\tilde{\epsilon}} \theta^\epsilon = \theta^\epsilon$ we get

$$II_3 = \left((u^{\tilde{\epsilon}} \cdot \nabla (\theta^\epsilon - \theta^{\tilde{\epsilon}})), \mathcal{J}_{\tilde{\epsilon}} (\theta^\epsilon - \theta^{\tilde{\epsilon}}) \right) = \frac{1}{2} \left(u^{\tilde{\epsilon}}, \nabla (\theta^\epsilon - \theta^{\tilde{\epsilon}})^2 \right) = 0$$

Putting all these estimates together yields that for $\delta_0 = \min\{m - \alpha, 1\}$

$$\frac{1}{2} \frac{d}{dt} \|\theta^\epsilon - \theta^{\tilde{\epsilon}}\|_{L^2}^2 \lesssim_M (\epsilon^{\delta_0} + \|\theta^\epsilon - \theta^{\tilde{\epsilon}}\|_{L^2}) \|\theta^\epsilon - \theta^{\tilde{\epsilon}}\|_{L^2}.$$

Furthermore

$$\frac{d}{dt} \|\theta^\epsilon - \theta^{\tilde{\epsilon}}\|_{L^2} \leq C(M) (\epsilon^{\delta_0} + \|\theta^\epsilon - \theta^{\tilde{\epsilon}}\|_{L^2}).$$

Thus the Grönwall inequality leads to the desired result:

$$\begin{aligned}
\sup_{0 \leq t \leq T} \|\theta^\epsilon - \theta^{\tilde{\epsilon}}\|_{L^2} &\leq e^{C(M)T} (\epsilon^{\delta_0} + \|\theta_0^\epsilon - \theta_0^{\tilde{\epsilon}}\|_{L^2}) \\
&\lesssim_{T, \|\theta_0\|_{H^m}} a(\epsilon),
\end{aligned} \tag{4.14}$$

where $a(\epsilon) := \epsilon^{\delta_0} + \|(Id - \mathcal{J}_\epsilon) \theta_0\|_{L^2}$ satisfies that $a(\epsilon) \rightarrow 0$ as $\epsilon \rightarrow 0$.

From (4.14), we deduce that the solution family (θ^ϵ) is Cauchy sequence in $\mathcal{C}([0, T], L^2(\mathbb{R}^2))$, so that it converges strongly to a function $\theta \in \mathcal{C}([0, T], L^2(\mathbb{R}^2))$. This result combined with uniform bounds (4.13) and the interpolation inequality in Sobolev spaces gives that for all $0 \leq s < m$

$$\begin{aligned} \sup_{0 \leq t \leq T} \|\theta^\epsilon - \theta\|_{H^s} &\leq C_s \sup_{0 \leq t \leq T} (\|\theta^\epsilon - \theta\|_{L^2}^{1-s/m} \|\theta^\epsilon - \theta\|_{H^m}^{s/m}) \\ &\lesssim_{s, T, \|\theta_0\|_{H^m}} a(\epsilon)^{1-s/m}. \end{aligned}$$

Hence we obtain the strong convergence in $\mathcal{C}([0, T], H^s(\mathbb{R}^2))$ for all $s < m$. With $2 < s < m$, this specially implies strong convergence in $\mathcal{C}([0, T], \mathcal{C}^1(\mathbb{R}^2))$. Also from the equation

$$\theta_t^\epsilon = -\nu|D|^\alpha \theta^\epsilon - \mathcal{J}_\epsilon(u^\epsilon \cdot \nabla \theta^\epsilon),$$

we find that θ_t^ϵ strongly converges to $-\nu|D|^\alpha \theta - u \cdot \nabla \theta$ in $\mathcal{C}([0, T], L^2(\mathbb{R}^2))$. Since $\theta^\epsilon \rightarrow \theta$, the distribution limit of θ_t^ϵ has to be θ_t . Thus $\theta \in \mathcal{C}^1([0, T], L^2(\mathbb{R}^2)) \cap \mathcal{C}([0, T], \mathcal{C}^1(\mathbb{R}^2))$ is a solution to the original equation (1.1). Using Fatou's Lemma, from (4.13), we also have $\theta \in L^\infty([0, T], H^m(\mathbb{R}^2)) \cap L^2([0, T], H^{m+\frac{\alpha}{2}}(\mathbb{R}^2))$.

Next, we show that $\theta \in \mathcal{C}([0, T], H^m(\mathbb{R}^2))$ indeed. The proof is classical (cf. [18]). We first prove that $\theta(t) \rightarrow \theta_0$ weakly in H^m as $t \rightarrow 0$. Let $\psi(x) \in \mathcal{C}_0^\infty(\mathbb{R}^2)$, denote

$$F^\epsilon(t, \psi) := (\theta^\epsilon, \psi) = \int_{\mathbb{R}^2} \theta^\epsilon(t, x) \psi(x) dx.$$

Clearly $F^\epsilon(\cdot, \psi) \in \mathcal{C}([0, T])$. And by taking the inner product of (4.4) with ψ , we get

$$\frac{d}{dt} F^\epsilon(t, \psi) = -(u^\epsilon \theta^\epsilon, \mathcal{J}_\epsilon \nabla \psi) - \nu(\theta^\epsilon, |D|^\alpha \psi),$$

thus for every $p \in [1, 2]$

$$\int_0^T |F_t^\epsilon|^p dt \leq T^{\frac{1}{p}-\frac{1}{2}} \|u^\epsilon\|_{L_T^2 L^2} \|\theta^\epsilon\|_{L_T^\infty L^2} \|\psi\|_{H^3} + \nu \|\theta^\epsilon\|_{L_T^p L^2} \|\psi\|_{H^\alpha}.$$

From the L^2 energy estimate $\|\theta^\epsilon\|_{L_T^\infty L^2}^2 + \|\theta^\epsilon\|_{L_T^2 \dot{H}^{\alpha/2}}^2 \leq \|\theta_0\|_{L^2}^2$, we know $\|F_t^\epsilon(\cdot, \psi)\|_{L^p([0, T])} \lesssim_{T, \|\psi\|_{H^3}} 1$. Hence by Arzela-Ascoli theorem, $\{F^\epsilon(t, \psi)\}_{\epsilon > 0}$ is compact in $\mathcal{C}([0, T])$, and we can choose a subsequence $F^{\epsilon_j}(t, \psi)$ converging to a function $F(t, \psi) \in \mathcal{C}([0, T])$ uniformly in t . In particular, from $\theta^\epsilon \rightarrow \theta$ in $\mathcal{C}([0, T]; L^2)$, we can further find a subsequence (still denote F^{ϵ_j}) such that $F(t, \psi) = (\theta(t), \psi)$ for all $t \in [0, T]$. Next, since $\mathcal{C}_0^\infty(\mathbb{R}^2)$ is dense in $H^{-m}(\mathbb{R}^2)$ and $\|\theta^\epsilon(t)\|_{H^m}$ is uniformly bounded in $[0, T]$, $F^{\epsilon_j}(t, \psi)$ converges to $F(t, \psi)$ for every $\psi \in H^{-m}$. Then for every $t > 0$ and $\psi \in H^{-m}$

$$|(\theta(t) - \theta_0, \psi)| \leq |(\theta(t) - \theta^{\epsilon_j}(t), \psi)| + |(\theta^{\epsilon_j}(t) - \theta_0^{\epsilon_j}, \psi)| + |(\theta_0^{\epsilon_j} - \theta_0, \psi)|.$$

All the three terms in the RHS can be made small for sufficiently small ϵ_j and t , thus $\theta(t)$ converges to θ_0 weakly in H^m as $t \rightarrow 0$. So we have

$$\|\theta_0\|_{H^m} \leq \liminf_{t \rightarrow 0} \|\theta(t)\|_{H^m}. \quad (4.15)$$

Furthermore, from (4.12) we infer that for every $\epsilon > 0$ the function $\|\theta^\epsilon(t)\|_{H^m}^2$ is below the graph of the solution of the equation

$$\frac{d}{dt} y(t) = C y(t) + C y^2(t), \quad y(0) = \|\theta_0\|_{H^m}^2.$$

By construction, the same holds for $\|\theta(t)\|_{H^m}^2$. Thus from the continuity of $y(t)$, we find $\|\theta_0\|_{H^m} \geq \limsup_{t \rightarrow 0} \|\theta(t)\|_{H^m}$. Therefore $\|\theta_0\|_{H^m} = \lim_{t \rightarrow 0} \|\theta(t)\|_{H^m}$, and the conclusion follows from this fact combined with the weak convergence.

Step 3: Uniqueness

Let $\theta^1, \theta^2 \in L^\infty([0, T], H^m(\mathbb{R}^2))$ be two smooth solutions to the modified quasi-geostrophic equation (1.1) with the same initial data. Denote $u^i = |D|^{\alpha-1} R^\perp \theta^i$, $i = 1, 2$, $\delta\theta = \theta^1 - \theta^2$, $\delta u = u^1 - u^2$, then we write the difference equation as

$$\partial_t \delta\theta + u^1 \cdot \nabla \delta\theta + \nu |D|^\alpha \delta\theta = -\delta u \cdot \nabla \theta^2, \quad \delta\theta|_{t=0} = 0$$

We also use the L^2 energy method, and in a similar way as treating the term II_3 , we obtain

$$\frac{d}{dt} \|\delta\theta\|_{L^2} \leq C_\alpha \|\nabla \theta^2\|_{\dot{H}^{\frac{\alpha}{2}}}^2 \|\delta\theta\|_{L^2} \leq C_\alpha \|\theta^2\|_{H^m}^2 \|\delta\theta\|_{L^2}.$$

Thus the Grönwall inequality ensures $\delta\theta \equiv 0$, that is, $\theta^1 \equiv \theta^2$.

Step 4: Smoothing Effect

Precisely, we have that for all $\gamma \in \mathbb{R}^+$ and $t \in [0, T]$

$$\|t^\gamma \theta(t)\|_{L_T^\infty H^{m+\gamma\alpha}}^2 + \|t^\gamma \theta(t)\|_{L_T^2 H^{m+\alpha/2+\gamma\alpha}}^2 \leq C e^{C(\gamma+1)(T\|\theta\|_{L_T^\infty H^{m+T}}^2)} \|\theta_0\|_{H^m}^2, \quad (4.16)$$

where C is an absolute constant depending only on α, ν, m . Notice that $t^\gamma \theta$ ($\gamma > 0$) satisfies

$$\partial_t(t^\gamma \theta) + u \cdot \nabla(t^\gamma \theta) + \nu |D|^\alpha(t^\gamma \theta) = \gamma t^{\gamma-1} \theta, \quad (t^\gamma \theta)|_{t=0} = 0. \quad (4.17)$$

which is a linear transport-diffusion equation with the velocity $u = |D|^{\alpha-1} R^\perp \theta$, $\alpha \in]0, 2[$. We first treat the case $\gamma \in \mathbb{Z}^+$. For $\gamma = 1$, in a similar way as obtaining (4.5), and using the Sobolev embedding we infer

$$\begin{aligned} \frac{d}{dt} \|t\theta(t)\|_{B_{2,2}^{m+\alpha}}^2 + \|t\theta(t)\|_{B_{2,2}^{m+\frac{3}{2}\alpha}}^2 &\lesssim (\|\nabla \theta(t)\|_{L^\infty}^\alpha \|\theta(t)\|_{L^\infty}^{2-\alpha} + \|\theta(t)\|_{L^2}) \|t\theta(t)\|_{B_{2,2}^{m+\alpha}}^2 + \|\theta(t)\|_{B_{2,2}^{m+\frac{\alpha}{2}}}^2 \\ &\lesssim (\|\theta(t)\|_{H^m}^2 + 1) \|t\theta(t)\|_{B_{2,2}^{m+\alpha}}^2 + \|\theta(t)\|_{B_{2,2}^{m+\frac{\alpha}{2}}}^2. \end{aligned}$$

Gronwall inequality yields that

$$\|t\theta(t)\|_{B_{2,2}^{m+\alpha}}^2 + \|t\theta(t)\|_{L_T^2 B_{2,2}^{m+\frac{3}{2}\alpha}}^2 \lesssim e^{CT+CT\|\theta\|_{L_T^\infty H^m}^2} \int_0^T \|\theta(\tau)\|_{B_{2,2}^{m+\frac{\alpha}{2}}}^2 d\tau. \quad (4.18)$$

Meanwhile, similarly as obtaining (4.12), we get

$$\|\theta(t)\|_{H^m}^2 + \|\theta\|_{L_T^2 H^{m+\frac{\alpha}{2}}}^2 \leq \|\theta_0\|_{H^m}^2 e^{CT+CT\|\theta\|_{L_T^\infty H^m}^2}. \quad (4.19)$$

Thus (4.16) with $\gamma = 1$ follows from (4.18) and (4.19) and the fact that the space $B_{2,2}^s$ is equivalent with H^s , $s \in \mathbb{R}$. Now suppose estimate (4.16) holds for $\gamma = N$, we shall consider the case $N + 1$. We use the equation (4.17) with $\gamma = N + 1$. Similarly as above, and observing that the constant C in (4.18) is independent of N if $\theta(t)$ is replaced by $t^N \theta(t)$ and m by $m + N\alpha$, we have

$$\begin{aligned} \|t^{N+1} \theta(t)\|_{H^{m+(N+1)\alpha}}^2 + \|t^{N+1} \theta(t)\|_{L_T^2 H^{m+(N+1)\alpha+\frac{\alpha}{2}}}^2 &\lesssim e^{CT+CT\|\theta\|_{L_T^\infty H^m}^2} \|t^N \theta(t)\|_{L_T^2 H^{m+(N+\frac{1}{2})\alpha}}^2 \\ &\lesssim e^{C(N+2)(T+T\|\theta\|_{L_T^\infty H^m}^2)} \|\theta_0\|_{H^m}^2. \end{aligned}$$

Thus the induction method ensures the estimate (4.16) for all $\gamma \in \mathbb{Z}^+$. Also notice that for $\gamma = 0$ the inequality (4.16) is also satisfied. Hence we obtain estimate (4.16) for all $\gamma \in \mathbb{N}$. For the general $\gamma \geq 0$, we set $[\gamma] \leq \gamma < [\gamma] + 1$, where $[\gamma]$ denotes the integer part of γ , and use the interpolation inequality in Sobolev spaces to get

$$\begin{aligned} \|t^\gamma \theta\|_{L_T^\infty H^{m+\gamma\alpha}}^2 &\leq \|t^{[\gamma]} \theta\|_{L_T^\infty H^{m+[\gamma]\alpha}}^{2([\gamma]+1-\gamma)} \|t^{[\gamma]+1} \theta\|_{L_T^\infty H^{m+([\gamma]+1)\alpha}}^{2(\gamma-[\gamma])} \\ &\lesssim e^{C(\gamma+1)(T+T\|\theta\|_{L_T^\infty H^m}^2)} \|\theta_0\|_{H^m}^2. \end{aligned}$$

Similar estimate holds for $\|t^\gamma \theta\|_{L_T^2 H^{m+(\gamma+\frac{1}{2})\alpha}}^2$.

Therefore, we conclude the Proposition 4.1. \square

Now, we are devoted to building the blowup criterion.

Proof of Proposition 4.2. We first note that the equation has a natural blowup criterion: if $T^* < \infty$ then necessarily

$$\|\theta\|_{L^\infty([0, T^*], H^m)} + \|\theta\|_{L^2([0, T^*], H^{m+\frac{\alpha}{2}})} = \infty.$$

Otherwise from the local result, the solution will continue over T^* .

In the same way as obtaining the estimate (4.5), we get the similar result for the original equation

$$\frac{1}{2} \frac{d}{dt} \|\theta(t)\|_{B_{2,2}^m}^2 + \frac{\nu}{2} \|\theta(t)\|_{B_{2,2}^{m+\frac{\alpha}{2}}}^2 \leq C_{m,\alpha} \left(\frac{1}{\nu} \|\nabla \theta\|_{L^\infty}^\alpha \|\theta\|_{L^\infty}^{2-\alpha} \|\theta\|_{B_{2,2}^m}^2 + \|\theta\|_{L^2}^2 \|\theta\|_{B_{2,2}^m} \right).$$

Also due to the maximum principle Proposition 2.3, we have

$$\frac{d}{dt} \left(\|\theta(t)\|_{B_{2,2}^m}^2 + \nu \int_0^t \|\theta(\tau)\|_{B_{2,2}^{m+\frac{\alpha}{2}}}^2 d\tau \right) \lesssim_{\alpha,\nu,m} (\|\nabla \theta(t)\|_{L^\infty}^\alpha + 1) \|\theta(t)\|_{B_{2,2}^m}^2.$$

This together with the Grönwall inequality leads to

$$\begin{aligned} \sup_{0 \leq t \leq T} \|\theta(t)\|_{H^m}^2 + \|\theta\|_{L^2([0, T], H^{m+\frac{\alpha}{2}})}^2 &\leq C_0 \sup_{0 \leq t \leq T} \|\theta(t)\|_{B_{2,2}^m}^2 + C_0 \|\theta\|_{L^2([0, T], B_{2,2}^{m+\frac{\alpha}{2}})}^2 \\ &\leq C \exp \left\{ CT + C \int_0^T \|\nabla \theta(t)\|_{L^\infty}^\alpha dt \right\}. \end{aligned}$$

Further, if $T^* < \infty$ and the integral $\int_0^{T^*} \|\nabla \theta(t)\|_{L^\infty}^\alpha dt < \infty$, then from the above estimate we directly have

$$\sup_{0 \leq t < T^*} \|\theta(t)\|_{H^m} + \|\theta\|_{L^2([0, T^*], H^{m+\frac{\alpha}{2}})} < \infty.$$

Clearly this contradicts the upper natural blowup criterion. Thus, if $T^* < \infty$, we necessarily have the equality $\int_0^{T^*} \|\nabla \theta(t)\|_{L^\infty}^\alpha dt = \infty$. \square

5 Global Existence

In this section, we use the modulus of continuity argument developed by Kiselev, Nazarov and Volberg [17] to prove the global result. Throughout this section, we assume T^* be the maximal existence time of the solution in $\mathcal{C}([0, T^*), H^m) \cap L^2([0, T^*), H^{m+\frac{\alpha}{2}})$.

Let $\lambda > 0$ be a real number which will be chosen later, then we define the set

$$\mathcal{I} := \{T \in [0, T^*) \mid \forall t \in [0, T], \forall x, y \in \mathbb{R}^2, x \neq y, |\theta(t, x) - \theta(t, y)| < \omega_\lambda(|x - y|)\},$$

where ω is a strict modulus of continuity also satisfying that $\omega'(0) < \infty$, $\lim_{\eta \searrow 0} \omega''(\eta) = -\infty$ and

$$\omega_\lambda(|x - y|) = \omega(\lambda|x - y|).$$

The explicit expression of ω will be shown later (i.e. (5.10)).

We first show that the set \mathcal{I} is nonempty, that is, at least $0 \in \mathcal{I}$. The proof is almost the same with the one in [1] only by setting T_1 there to be 0. We omit it here and only note that to fit our purpose λ can be taken

$$\lambda = \frac{\omega^{-1}(3\|\theta_0\|_{L^\infty})}{2\|\theta_0\|_{L^\infty}} \|\nabla\theta_0\|_{L^\infty}. \quad (5.1)$$

Thus \mathcal{I} is an interval of the form $[0, T_*)$, where $T_* := \sup_{T \in \mathcal{I}} T$. We have three possibilities:

- (a) $T_* = T^*$
- (b) $T_* < T^*$ and $T_* \in \mathcal{I}$
- (c) $T_* < T^*$ and $T_* \notin \mathcal{I}$

For case (a), we necessarily have $T^* = \infty$, since the Lipschitz norm of θ does not blow up from the definition of \mathcal{I} which contradicts with (4.1). This is our goal.

For case (b), we observe that this is just the case treated in [1] or [13] showing that it is impossible. The proof only needs very small modification, so we omit it either. We just point out in this case the smoothing effects will be used, since we need the fact that $\|\nabla^2\theta(T_*)\|_{L^\infty}$ is finite.

Then our task is reduced to get rid of the case (c). We prove by contradiction. If the case (c) is satisfied, then by the time continuity of θ , we necessarily get

$$\sup_{x, y \in \mathbb{R}^2, x \neq y} \frac{|\theta(T_*, x) - \theta(T_*, y)|}{\omega_\lambda(|x - y|)} = 1.$$

We further have the following assertion (with its proof in the end of this section).

Lemma 5.1. *If $T_* < T^*$ is the first time that the strict modulus of continuity ω_λ is lost (i.e. case (c)), then there exists $x, y \in \mathbb{R}^2$, $x \neq y$ such that*

$$\theta(T_*, x) - \theta(T_*, y) = \omega_\lambda(\xi), \quad \text{with } \xi := |x - y|. \quad (5.2)$$

Moreover, let $\ell = \frac{x-y}{|x-y|}$ and $v \in \mathbb{S}^1$ be the unit vector perpendicular to ℓ , we have

$$\partial_\ell \theta(T_*, x) = \partial_\ell \theta(T_*, y) = \omega'_\lambda(\xi), \quad \partial_v \theta(T_*, x) = \partial_v \theta(T_*, y) = 0, \quad (5.3)$$

where $\partial_\ell = \ell \cdot \nabla$ and $\partial_v = v \cdot \nabla$ are the directional derivatives along ℓ and v respectively.

We shall show that this scenario (5.2) can not happen, more precisely, we shall prove

$$f'(T_*) < 0, \quad \text{with} \quad f(t) := \theta(t, x) - \theta(t, y).$$

This is impossible because we necessarily have $f(t) \leq f(T_*)$, for all $0 \leq t \leq T_*$ from the definition of \mathcal{I} .

We see that the modified quasi-geostrophic equation (1.1) can be defined in the classical sense (from the smoothing effect), and thus

$$\begin{aligned} f'(T_*) &= - \left[(u \cdot \nabla \theta)(T_*, x) - (u \cdot \nabla \theta)(T_*, y) \right] + \nu \left[(-|D|^\alpha \theta)(T_*, x) - (-|D|^\alpha \theta)(T_*, y) \right] \\ &:= \mathcal{A}_1 + \mathcal{A}_2 \end{aligned}$$

with

$$u = |D|^{\alpha-1} \mathcal{R}^\perp \theta = \mathcal{R}_\alpha^\perp \theta := (-\mathcal{R}_{\alpha,2} \theta, \mathcal{R}_{\alpha,1} \theta)$$

where $\mathcal{R}_{\alpha,j}$ are the modified Riesz transforms introduced in the section 3.

For the first term, \mathcal{A}_1 , from (5.2), we find that

$$\begin{aligned} \mathcal{A}_1 &= [(u(T_*, x) - u(T_*, y)) \cdot \ell] \omega'_\lambda(\xi) \\ &= [(u(T_*, x) - u(T_*, y)) \cdot \ell] \lambda \omega'(\lambda \xi). \end{aligned}$$

Lemma 3.2 gives us a rough estimate as follows

$$|\mathcal{A}_1| \leq \Omega_\lambda(\xi) \lambda \omega'(\lambda \xi) = \lambda^\alpha (\Omega \omega')(\lambda \xi),$$

where $\Omega_\lambda(\xi)$ is defined from (3.2), i.e.

$$\Omega_\lambda(\xi) = A \left(\int_0^\xi \frac{\omega_\lambda(\eta)}{\eta^\alpha} d\eta + \xi \int_\xi^\infty \frac{\omega_\lambda(\eta)}{\eta^{1+\alpha}} d\eta \right) = \lambda^{\alpha-1} \Omega(\lambda \xi). \quad (5.4)$$

For the second term, \mathcal{A}_2 , from Lemma 3.3 we get

$$\mathcal{A}_2 \leq \nu \lambda^\alpha \Upsilon(\lambda \xi),$$

where

$$\begin{aligned} \Upsilon(\xi) &:= B \int_0^{\frac{\xi}{2}} \frac{\omega(\xi + 2\eta) + \omega(\xi - 2\eta) - 2\omega(\xi)}{\eta^{1+\alpha}} d\eta \\ &\quad + B \int_{\frac{\xi}{2}}^\infty \frac{\omega(2\eta + \xi) - \omega(2\eta - \xi) - 2\omega(\xi)}{\eta^{1+\alpha}} d\eta. \end{aligned}$$

Thus we obtain

$$f'(T_*) \leq \lambda^\alpha (\Omega \omega' + \nu \Upsilon)(\lambda \xi). \quad (5.5)$$

Observe that when $\alpha \in]1, 2[$ and ξ is a large number, the integral from 0 to ξ in the expression of Ω always produce much difficulty, roughly speaking, to ensure the RHS of (5.5) is negative, we need that the contribution from this part $\omega(\xi)\omega'(\xi) \ll \frac{\omega(\xi)}{\xi^\alpha}$, thus it seems impossible to construct an appropriate unbounded MOC. However, basically following a idea from [20], we can further develop the contribution of the dissipative term and use the additional dissipation to control the "bad" part of the nonlinearity so that improved estimates of \mathcal{A}_1 and \mathcal{A}_2 can be obtained. Meanwhile, when $\alpha = 1$ this method can also slightly improve the MOC constructed in [17]. Precisely,

Lemma 5.2. *Under the condition of Lemma 5.1 and for $\alpha \in [1, 2[$, we have*

$$\mathcal{A}_2 \leq \nu\lambda^\alpha\Upsilon(\lambda\xi) + \nu\lambda^\alpha\Upsilon^\perp(\lambda\xi), \quad (5.6)$$

where $\Upsilon^\perp \leq 0$ is a meaningful integral defined from θ and ω (with its explicit formula cf. Lemma 5.5 of [22]). Correspondingly, we can treat the drift term as follows

$$|(u(T_*, x) - u(T_*, y)) \cdot \ell| \leq \lambda^{\alpha-1}\tilde{\Omega}(\lambda\xi) \quad (5.7)$$

with

$$\tilde{\Omega}(\xi) = A\left(-\xi\Upsilon^\perp(\xi) + \xi \int_\xi^\infty \frac{\omega(\eta)}{\eta^{1+\alpha}} d\eta + \xi^{-\alpha+1}\omega(\xi)\right), \quad (5.8)$$

where A is an absolute constant that may depend on α .

Proof. This is a direct consequence of Lemma 5.5 and Lemma 5.6 in [22] when $\alpha \in]1, 2[$, and they can simply extend to the case $\alpha = 1$. \square

Hence when $\alpha \in [1, 2[$, based on Lemma 5.2, we also get

$$f'(T_*) \leq \lambda^\alpha(\tilde{\Omega}\omega' + \nu\Upsilon + \nu\Upsilon^\perp)(\lambda\xi). \quad (5.9)$$

Next we shall construct our special modulus of continuity in the spirit of [17]. Let $0 < \gamma < \delta < 1$ be two small positive numbers chosen later, and define the continuous functions ω as follows

$$\text{MOC} \begin{cases} \omega(\xi) = \xi - \xi^{1+\frac{\alpha}{2}} & \text{if } 0 \leq \xi \leq \delta, \\ \omega'(\xi) = \frac{\gamma}{4\xi} & \text{if } \xi > \delta, \end{cases} \quad (5.10)$$

equivalently,

$$\omega(\xi) = \begin{cases} \xi - \xi^{1+\frac{\alpha}{2}} & \text{if } 0 \leq \xi \leq \delta, \\ \delta - \delta^{1+\frac{\alpha}{2}} + \frac{\gamma}{4} \log \frac{\xi}{\delta} & \text{if } \xi > \delta. \end{cases} \quad (5.11)$$

Note that, for small δ , the left derivative of ω at δ is about 1, while the right derivative equals $\frac{\gamma}{4\delta} < \frac{1}{4}$. So ω is concave if δ is small enough. Clearly, $\omega(0) = 0$, $\omega'(0) = 1$ and $\lim_{\eta \rightarrow 0^+} \omega''(\eta) = -\infty$, and ω is unbounded (it has the logarithmic growth at infinity).

Then our target is to show that, for this MOC ω , when $\alpha \in]0, 1[$

$$\Omega(\xi)\omega'(\xi) + \nu\Upsilon(\xi) < 0 \quad \text{for all } \xi > 0, \quad (5.12)$$

and when $\alpha \in [1, 2[$

$$\tilde{\Omega}(\xi)\omega'(\xi) + \nu\Upsilon(\xi) + \nu\Upsilon^\perp(\xi) < 0 \quad \text{for all } \xi > 0. \quad (5.13)$$

In the following we shall carefully check these two formulae.

Case I: when $\alpha \in]0, 1[$

Precisely, we shall check the following inequality

$$A \left[\int_0^\xi \frac{\omega(\eta)}{\eta^\alpha} d\eta + \xi \int_\xi^\infty \frac{\omega(\eta)}{\eta^{1+\alpha}} d\eta \right] \omega'(\xi) + \nu B \int_0^{\frac{\xi}{2}} \frac{\omega(\xi + 2\eta) + \omega(\xi - 2\eta) - 2\omega(\xi)}{\eta^{1+\alpha}} d\eta \\ + \nu B \int_{\frac{\xi}{2}}^\infty \frac{\omega(2\eta + \xi) - \omega(2\eta - \xi) - 2\omega(\xi)}{\eta^{1+\alpha}} d\eta < 0 \quad \text{for all } \xi > 0.$$

We further divide it into two cases.

Case I.1: $\alpha \in]0, 1[$ and $0 < \xi \leq \delta$

Since $\frac{\omega(\eta)}{\eta} \leq \omega'(0) = 1$ for all $\eta > 0$ and $\eta \leq \eta^\alpha$ for $\eta \leq \delta < 1$, we have

$$\int_0^\xi \frac{\omega(\eta)}{\eta^\alpha} d\eta \leq \int_0^\xi \frac{\omega(\eta)}{\eta} d\eta \leq \xi,$$

and

$$\int_\xi^\delta \frac{\omega(\eta)}{\eta^{1+\alpha}} d\eta \leq \int_\xi^\delta \frac{1}{\eta^\alpha} d\eta = \frac{1}{1-\alpha} (\delta^{1-\alpha} - \xi^{1-\alpha}) \leq \frac{1}{1-\alpha}.$$

Furthermore,

$$\int_\delta^\infty \frac{\omega(\eta)}{\eta^{1+\alpha}} d\eta = \frac{1}{\alpha} \frac{\omega(\delta)}{\delta^\alpha} + \frac{1}{\alpha} \int_\delta^\infty \frac{\gamma}{4\eta^{1+\alpha}} d\eta \leq \frac{1}{\alpha} + \frac{1}{\alpha^2} \frac{\gamma}{\delta^\alpha} \leq \frac{2}{\alpha},$$

if $\gamma < \alpha\delta$. Clearly $\omega'(\xi) \leq \omega'(0) = 1$, so we get that the positive part is bounded by $A\xi \frac{2}{\alpha(1-\alpha)}$.

For the negative part, we have

$$\nu B \int_0^{\frac{\xi}{2}} \frac{\omega(\xi + 2\eta) + \omega(\xi - 2\eta) - 2\omega(\xi)}{\eta^{1+\alpha}} d\eta \leq \nu B \int_0^{\frac{\xi}{2}} \frac{\omega''(\xi) 2\eta^2}{\eta^{1+\alpha}} d\eta \\ = -\nu B \frac{\alpha(2+\alpha)}{2^{1-\alpha}(2-\alpha)} \xi^{1-\frac{\alpha}{2}} \leq -\frac{\alpha}{2} \nu B \xi^{1-\frac{\alpha}{2}}.$$

But, clearly $\xi \left(A \frac{2}{\alpha(1-\alpha)} - \frac{\alpha}{2} \nu B \xi^{-\frac{\alpha}{2}} \right) < 0$ on $]0, \delta]$ when δ is small enough.

Case I.2: $\alpha \in]0, 1[$ and $\xi \geq \delta$

For $\eta \leq \delta < 1$ we still use $\eta^\alpha \geq \eta$ and for $\delta \leq \eta \leq \xi$ we use $\omega(\eta) \leq \omega(\xi)$, then

$$\int_0^\xi \frac{\omega(\eta)}{\eta^\alpha} d\eta \leq \delta + \frac{\omega(\xi)}{1-\alpha} (\xi^{1-\alpha} - \delta^{1-\alpha}) \leq \omega(\xi) \left(\frac{2}{\alpha} + \frac{\xi^{1-\alpha}}{1-\alpha} \right),$$

where the last inequality is due to $\frac{\alpha}{2}\delta < \omega(\delta) \leq \omega(\xi)$ if δ is small enough (i.e. $\delta < (1 - \frac{\alpha}{2})^{2/\alpha}$).

Also

$$\int_\xi^\infty \frac{\omega(\eta)}{\eta^{1+\alpha}} d\eta = \frac{1}{\alpha} \frac{\omega(\xi)}{\xi^\alpha} + \frac{1}{\alpha} \int_\xi^\infty \frac{\gamma}{4\eta^{1+\alpha}} d\eta \leq \frac{1}{\alpha} \frac{\omega(\xi)}{\xi^\alpha} + \frac{1}{\alpha^2} \frac{\gamma}{2\xi^\alpha} \leq \frac{2}{\alpha} \frac{\omega(\xi)}{\xi^\alpha}$$

if $\gamma < \alpha^2 \delta$ (thus $\gamma/2 \leq \alpha \omega(\xi)$) and δ is small enough. Thus the positive term is bounded from above by

$$A\omega(\xi) \left(\frac{2}{\alpha} + \left(\frac{1}{1-\alpha} + \frac{2}{\alpha} \right) \xi^{1-\alpha} \right) \omega'(\xi) \leq A \frac{\omega(\xi)}{\xi^\alpha} \frac{2}{\alpha(1-\alpha)} (\xi + \xi^\alpha) \frac{\gamma}{4\xi} \leq \frac{A\delta^{\alpha-1}\gamma}{\alpha(1-\alpha)} \frac{\omega(\xi)}{\xi^\alpha}.$$

For the negative part, we first observe that for $\xi \geq \delta$,

$$\omega(2\xi) = \omega(\xi) + \int_{\xi}^{2\xi} \omega'(\eta) d\eta = \omega(\xi) + \frac{\log 2}{2} \gamma \leq \frac{3}{2} \omega(\xi),$$

under the same assumptions on δ and γ as above. Also, taking advantage of the concavity we obtain $\omega(2\eta + \xi) - \omega(2\eta - \xi) \leq \omega(2\xi)$ for all $\eta \geq \frac{\xi}{2}$. Therefore

$$\nu B \int_{\frac{\xi}{2}}^{\infty} \frac{\omega(2\eta + \xi) - \omega(2\eta - \xi) - 2\omega(\xi)}{\eta^{1+\alpha}} d\eta \leq -\nu B \frac{\omega(\xi)}{2} \int_{\frac{\xi}{2}}^{\infty} \frac{1}{\eta^{1+\alpha}} d\eta = -\nu B \frac{2^\alpha}{2\alpha} \frac{\omega(\xi)}{\xi^\alpha}.$$

But $\frac{\omega(\xi)}{\xi^\alpha} \left(\frac{A\delta^{\alpha-1}\gamma}{\alpha(1-\alpha)} - \nu B \frac{2^\alpha}{2\alpha} \right) < 0$ if γ is small enough (i.e. $\gamma < \min\{\alpha^2 \delta, \frac{\nu(1-\alpha)B2^\alpha}{2A} \delta^{1-\alpha}\}$).

Case II: when $\alpha \in [1, 2[$

Precisely, we shall check the following inequality

$$A \left[-\xi \Upsilon^\perp(\xi) + \xi^{-\alpha+1} \omega(\xi) + \xi \int_{\xi}^{\infty} \frac{\omega(\eta)}{\eta^{1+\alpha}} d\eta \right] \omega'(\xi) + \nu B \int_0^{\frac{\xi}{2}} \frac{\omega(\xi + 2\eta) + \omega(\xi - 2\eta) - 2\omega(\xi)}{\eta^{1+\alpha}} d\eta \\ + \nu B \int_{\frac{\xi}{2}}^{\infty} \frac{\omega(2\eta + \xi) - \omega(2\eta - \xi) - 2\omega(\xi)}{\eta^{1+\alpha}} d\eta + \nu \Upsilon^\perp(\xi) < 0 \quad \text{for all } \xi > 0.$$

We also further divide it into two cases.

Case II.1: $\alpha \in [1, 2[$ and $0 < \xi \leq \delta$

Since $\frac{\omega(\eta)}{\eta} \leq \omega'(0) = 1$ for all $\eta > 0$ and $-\Upsilon^\perp(\xi) \geq 0$, we have $-\xi \Upsilon^\perp(\xi) \leq -\delta \Upsilon^\perp(\xi)$ and $\xi^{-\alpha+1} \omega(\xi) \leq \xi^{2-\alpha}$ and

$$\int_{\xi}^{\delta} \frac{\omega(\eta)}{\eta^{1+\alpha}} d\eta \leq \int_{\xi}^{\delta} \frac{1}{\eta^\alpha} d\eta \leq \begin{cases} \frac{1}{\alpha-1} \xi^{1-\alpha}, & \alpha \in]1, 2[, \\ \log(\delta/\xi), & \alpha = 1. \end{cases}$$

Further, integration by parts leads to

$$\int_{\delta}^{\infty} \frac{\omega(\eta)}{\eta^{1+\alpha}} d\eta = \frac{1}{\alpha} \frac{\omega(\delta)}{\delta^\alpha} + \frac{1}{\alpha} \int_{\delta}^{\infty} \frac{\gamma}{4\eta^{\alpha+1}} d\eta \\ \leq \frac{1}{\alpha} \frac{1}{\delta^{\alpha-1}} + \frac{\gamma}{4\alpha^2} \frac{1}{\delta^\alpha} \leq 2 \frac{1}{\delta^{\alpha-1}} \leq 2\xi^{1-\alpha}.$$

Clearly $\omega'(\xi) \leq \omega'(0) = 1$, so we get that the positive part is bounded by

$$\begin{cases} A \left(-\delta \Upsilon^\perp(\xi) + \xi^{2-\alpha} \frac{4}{\alpha-1} \right), & \alpha \in]1, 2[, \\ A \left(-\delta \Upsilon^\perp(\xi) + \xi \left(3 + \log \frac{\delta}{\xi} \right) \right), & \alpha = 1. \end{cases}$$

For the negative part, we have

$$\begin{aligned} \nu B \int_0^{\frac{\xi}{2}} \frac{\omega(\xi + 2\eta) + \omega(\xi - 2\eta) - 2\omega(\xi)}{\eta^{1+\alpha}} d\eta &\leq \nu B \int_0^{\frac{\xi}{2}} \frac{\omega''(\xi) 2\eta^2}{\eta^{1+\alpha}} d\eta \\ &= -\nu B \frac{\alpha(2+\alpha)}{2^{1-\alpha}(2-\alpha)} \xi^{1-\frac{\alpha}{2}} \leq -3\nu B \xi^{1-\frac{\alpha}{2}}. \end{aligned}$$

But, clearly if δ is chosen small enough, we find that for every $\xi \in]0, \delta]$

$$\begin{cases} (-A\delta + \nu)\Upsilon^\perp(\xi) + \xi^{2-\alpha} \left(A \frac{4}{\alpha-1} - 3\nu B \xi^{\frac{\alpha}{2}-1} \right) < 0, & \alpha \in]1, 2[, \\ (-A\delta + \nu)\Upsilon^\perp(\xi) + \xi \left(3A + A \log \frac{\delta}{\xi} - 3\nu B \xi^{-\frac{1}{2}} \right) < 0, & \alpha = 1. \end{cases}$$

Case II.2: $\alpha \in [1, 2[$ and $\xi \geq \delta$

For the positive part we have

$$\begin{aligned} \int_\xi^\infty \frac{\omega(\eta)}{\eta^{1+\alpha}} d\eta &= \frac{1}{\alpha} \frac{\omega(\xi)}{\xi^\alpha} + \frac{1}{\alpha} \int_\xi^\infty \frac{\gamma}{4\eta^{\alpha+1}} d\eta \\ &\leq \frac{1}{\alpha} \frac{\omega(\xi)}{\xi^\alpha} + \frac{\gamma}{4\alpha^2} \frac{1}{\xi^\alpha} \leq 2 \frac{\omega(\xi)}{\xi^\alpha}, \end{aligned}$$

where we have used the simple fact that $\gamma \leq \frac{\delta}{2} \leq \omega(\delta) \leq \omega(\xi)$. Thus the positive term is bounded from above by

$$A \left(-\xi \Upsilon^\perp(\xi) + 3\omega(\xi) \xi^{1-\alpha} \right) \omega'(\xi) = A \left(-\xi \Upsilon^\perp(\xi) + 3\omega(\xi) \xi^{1-\alpha} \right) \frac{\gamma}{4\xi} \leq -A\gamma \Upsilon^\perp(\xi) + A\gamma \frac{\omega(\xi)}{\xi^\alpha}.$$

For the negative part, in a similar way as treating the corresponding part in case I.2, we have

$$\nu B \int_{\frac{\xi}{2}}^\infty \frac{\omega(2\eta + \xi) - \omega(2\eta - \xi) - 2\omega(\xi)}{\eta^{1+\alpha}} d\eta \leq -\nu B \frac{\omega(\xi)}{2} \int_{\frac{\xi}{2}}^\infty \frac{1}{\eta^{1+\alpha}} d\eta \leq -\frac{\nu B \omega(\xi)}{2} \frac{1}{\xi^\alpha}.$$

But, clearly $(-A\gamma + \nu)\Upsilon^\perp(\xi) + \frac{\omega(\xi)}{\xi^\alpha} (A\gamma - \frac{\nu B}{2}) < 0$ if γ is small enough.

Therefore both case I and case II yield $f'(T_*) < 0$.

Finally, only case (a) occurs and we obtain $T^* = \infty$. Moreover

$$\|\nabla\theta(t)\|_{L^\infty} < \lambda, \quad \forall t \in [0, \infty[,$$

where the value of $\lambda \sim C \|\nabla\theta_0\|_{L^\infty} e^{C\|\theta_0\|_{L^\infty}}$ is given by (5.1).

Proof of Lemma 5.1. Set $C' := \omega^{-1}(3\|\theta_0\|_{L^\infty})$, then from the maximum principle (2.2), we get

$$\lambda|x - y| \geq C' \Rightarrow |\theta(T_*, x) - \theta(T_*, y)| < \frac{2}{3} \omega_\lambda(|x - y|). \quad (5.14)$$

Since $\nabla\theta(t) \in \mathcal{C}([0, T^*), H^{m-1}(\mathbb{R}^2))$, then for every $\epsilon > 0$, there exists $R > 0$ such that

$$\|\nabla\theta(T_*)\|_{L^\infty(\mathbb{R}^2 \setminus B_R)} \leq C_0 \|\nabla\theta(T_*)\|_{H^{m-1}(\mathbb{R}^2 \setminus B_R)} \leq \epsilon,$$

where B_R is a ball centered at the origin with the radius R and $\mathbb{R}^2 \setminus B_R$ is its complement. Thus for every $x, y (x \neq y)$ satisfying that $\lambda|x - y| \leq C'$ and x or y belongs to $\mathbb{R}^2 \setminus B_{R+C'/\lambda}$, we get

$$|\theta(T_*, x) - \theta(T_*, y)| \leq \|\nabla\theta(T_*)\|_{L^\infty(\mathbb{R}^2 \setminus B_R)} |x - y| \leq \epsilon|x - y|.$$

Taking advantage of the following inequality from the concavity of ω

$$\frac{\omega(C')}{C'}\lambda|x - y| \leq \omega_\lambda(|x - y|),$$

we can take ϵ small enough such that $\epsilon < \frac{1}{2}\frac{\omega(C')}{C'}\lambda$ to obtain

$$\lambda|x - y| \leq C', x \text{ or } y \in \mathbb{R}^2 \setminus B_{R+\frac{C'}{\lambda}}; \Rightarrow |\theta(T_*, x) - \theta(T_*, y)| < \frac{1}{2}\omega_\lambda(|x - y|). \quad (5.15)$$

Now it remains to consider the case when $x, y \in B_{R+\frac{C'}{\lambda}}$. From the smoothing effect we know $\|\nabla^2\theta(T_*)\|_{L^\infty} < \infty$, thus we have (cf. [17])

$$\|\nabla\theta(T_*)\|_{L^\infty(B_{R+\frac{C'}{\lambda}})} < \lambda\omega'(0).$$

Let $\delta' \ll 1$ small enough, then we see

$$\|\theta(T_*)\|_{L^\infty(B_{R+\frac{C'}{\lambda}})} < \lambda(1 - \delta')\frac{\omega(\delta')}{\delta'}.$$

Thus for every $x, y (x \neq y)$ satisfying that $\lambda|x - y| \leq \delta'$ and both x, y belongs to $B_{R+C'/\lambda}$, we have

$$\begin{aligned} |\theta(T_*, x) - \theta(T_*, y)| &\leq \|\nabla\theta(T_*)\|_{L^\infty(B_{R+\frac{C'}{\lambda}})} |x - y| \\ &< (1 - \delta')\frac{\omega(\delta')}{\delta'}\lambda|x - y| \leq (1 - \delta')\omega_\lambda(|x - y|). \end{aligned} \quad (5.16)$$

We set

$$\Omega := \{(x, y) \in \mathbb{R}^2 \times \mathbb{R}^2 : \max\{|x|, |y|\} \leq R + \frac{C'}{\lambda}, |x - y| \geq \frac{\delta'}{\lambda}\},$$

then from the above results we necessarily have

$$1 = \sup_{x \neq y} \frac{|\theta(T_*, x) - \theta(T_*, y)|}{\omega_\lambda(|x - y|)} = \sup_{(x, y) \in \Omega} \frac{|\theta(T_*, x) - \theta(T_*, y)|}{\omega_\lambda(|x - y|)}.$$

Thus the conclusion follows from the compactness of Ω .

For (5.3), it is from a direct computation under the scenario (5.2) (cf. Proposition 2.4 in [20]). \square

6 Appendix

6.1 The formula for $\mathcal{R}_{\alpha, j} f$

Proof of Proposition 3.1. The pseudo-differential operator $\mathcal{R}_{\alpha, j}$ ($\alpha \in]0, 2[$) is the composition of two operators $|D|^{\alpha-1}$ and \mathcal{R}_j , which both are (constant coefficient) pseudo-differential operators, thus the symbol of $\mathcal{R}_{\alpha, j}$ is $-i\zeta_j/|\zeta|^{2-\alpha}$. Now we want to know the explicit formula of $\mathcal{F}^{-1}(-i\zeta_j/|\zeta|^{2-\alpha})$.

From the equality in the distributional sense

$$\frac{\partial}{\partial x_j} |x|^{-(n+\alpha-2)} = -(n+\alpha-2) \text{p.v.} \frac{x_j}{|x|^{n+\alpha}},$$

and the known formula that for every $0 < a < n$ (c.f. [15])

$$(|x|^{-a})^\wedge(\zeta) = \frac{2^{n-a} \pi^{n/2} \Gamma(\frac{n-a}{2})}{\Gamma(\frac{a}{2})} |\zeta|^{-n+a},$$

we directly have

$$\begin{aligned} (\text{p.v.} \frac{x_j}{|x|^{n+\alpha}})^\wedge(\zeta) &= -\frac{1}{n+\alpha-2} (\partial_{x_j} |x|^{-n-\alpha+2})^\wedge(\zeta) \\ &= -\frac{i\zeta_j}{n+\alpha-2} (|x|^{-n-\alpha+2})^\wedge(\zeta) \\ &= -\frac{i\zeta_j}{n+\alpha-2} \frac{2^{2-\alpha} \pi^{n/2} \Gamma(\frac{2-\alpha}{2})}{\Gamma(\frac{n+\alpha-2}{2})} |\zeta|^{\alpha-2} \\ &= -i \frac{2^{1-\alpha} \pi^{n/2} \Gamma(\frac{2-\alpha}{2})}{\Gamma(\frac{n+\alpha}{2})} \cdot \frac{\zeta_j}{|\zeta|^{2-\alpha}}. \end{aligned}$$

□

6.2 A commutator estimate

The key to the proof of the uniform estimate is the following commutator estimate

Lemma 6.1. *Let v be a divergence free vector field over \mathbb{R}^n . For every $q \in \mathbb{N}$, denote*

$$F_q(v, f) := S_{q+1} v \cdot \nabla \Delta_q f - \Delta_q(v \cdot \nabla f).$$

Then for every $\beta \in]0, 1[$, there exists a positive constant C such that

$$\begin{aligned} &2^{-q\beta} \|F_q(v, f)\|_{L^2} \\ &\leq C \left\| |D|^{1-\beta} v \right\|_{L^\infty} \left(\sum_{q' \leq q+4} 2^{q'-q} \|\Delta_{q'} f\|_{L^2} + \sum_{q' \geq q-4} 2^{(q-q')(1-\beta)} \|\Delta_{q'} f\|_{L^2} \right), \end{aligned} \quad (6.1)$$

Epecially, in the case $n = 2$ and $v = |D|^{\alpha-1} \mathcal{R}^\perp f$ ($\alpha \in]0, 2[$), we further have for every $\beta \in]\max\{0, \alpha - 1\}, 1[$ and every $q \in \mathbb{N}$

$$\begin{aligned} &2^{-q\beta} \|F_q(v, f)\|_{L^2} \\ &\leq C \left(\left\| |D|^{1-\beta} v \right\|_{L^\infty} \sum_{q' \geq q-4} 2^{(q-q')(1-\beta)} \|\Delta_{q'} f\|_{L^2} + \left\| |D|^{\alpha-\beta} f \right\|_{L^\infty} \sum_{|q'-q| \leq 4} \|\Delta_{q'} f\|_{L^2} \right). \end{aligned} \quad (6.2)$$

Moreover, when $\beta = 0$, $\alpha \in]0, 1[$, (6.1) and (6.2) hold if we replace $\left\| |D|^{1-\beta} v \right\|_{L^\infty}$ by $\|\nabla v\|_{L^\infty}$; and when $\beta = 1$, $\alpha = 2$, then (6.1) and (6.2) hold if we make such a modification

$$\left\| |D|^{1-\beta} v \right\|_{L^\infty} \rightarrow \|v\|_{B_{\infty,1}^0}, \quad \left\| |D|^{\alpha-\beta} f \right\|_{L^2} \rightarrow \|\nabla f\|_{L^\infty}.$$

Proof. Using Bony decomposition, we decompose $F_q(v, f)$ into $\sum_{i=1}^6 F_q^i(v, f)$, where

$$\begin{aligned} F_q^1(v, f) &= (S_{q+1}v - v) \cdot \nabla \Delta_q f, & F_q^2(v, f) &= [\Delta_{-1}v, \Delta_q] \cdot \nabla f, \\ F_q^3(v, f) &= \sum_{q' \in \mathbb{N}} [S_{q'-1}\tilde{v}, \Delta_q] \cdot \nabla \Delta_{q'} f, & F_q^4(v, f) &= \sum_{q' \geq -1} \Delta_{q'} \tilde{v} \cdot \nabla \Delta_q S_{q'+2} f, \\ F_q^5(v, f) &= - \sum_{q' \in \mathbb{N}} \Delta_q \left(\Delta_{q'} \tilde{v} \cdot \nabla S_{q'-1} f \right), & F_q^6(v, f) &= - \sum_{q' \geq -1} \operatorname{div} \Delta_q \left(\Delta_{q'} \tilde{v} \sum_{i \in \{\pm 1, 0\}} \Delta_{q'+i} f \right), \end{aligned}$$

where $[A, B] := AB - BA$ denotes the commutator operator and $\tilde{v} := v - \Delta_{-1}v$ denotes the high frequency part of v .

For F_q^1 , from the divergence-free property of v we directly obtain that when $1 - \beta > 0$

$$\begin{aligned} 2^{-q\beta} \|F_q^1(v, f)\|_{L^2} &\lesssim \sum_{q' \geq q+1} 2^{(1-\beta)(q-q')} 2^{q'(1-\beta)} \|\Delta_{q'} v\|_{L^\infty} \|\Delta_q f\|_{L^2} \\ &\lesssim \| |D|^{1-\beta} v \|_{L^\infty} \|\Delta_q f\|_{L^2}. \end{aligned}$$

For F_q^2 , since $F_q^2(v, f) = \sum_{|q'-q| \leq 1} [\Delta_{-1}v, \Delta_q] \cdot \nabla \Delta_{q'} f$, then from the expression formula of Δ_q and mean value theorem, we get that when $\beta > 0$

$$\begin{aligned} 2^{-q\beta} \|F_q^2(v, f)\|_{L^2} &\lesssim 2^{-q\beta} 2^{-q} \|\nabla \Delta_{-1}v\|_{L^\infty} \sum_{|q'-q| \leq 1} 2^{q'} \|\Delta_{q'} f\|_{L^2} \\ &\lesssim \sum_{-\infty \leq j \leq -1} 2^{j\beta} \| |D|^{1-\beta} \Delta_j v \|_{L^\infty} \sum_{|q'-q| \leq 1} \|\Delta_{q'} f\|_{L^2} \\ &\lesssim \| |D|^{1-\beta} v \|_{L^\infty} \sum_{|q'-q| \leq 1} \|\Delta_{q'} f\|_{L^2}. \end{aligned}$$

For F_q^3 , similarly as estimating F_q^2 , we infer

$$\begin{aligned} 2^{-q\beta} \|F_q^3(v, f)\|_{L^2} &\lesssim 2^{-q\beta} \sum_{|q'-q| \leq 4} 2^{-q} \|\nabla S_{q'-1}\tilde{v}\|_{L^\infty} 2^{q'} \|\Delta_{q'} f\|_{L^2} \\ &\lesssim \sum_{|q'-q| \leq 4} \sum_{q'' \leq q'-2} 2^{(q''-q')\beta} \| |D|^{1-\beta} \Delta_{q''}\tilde{v} \|_{L^\infty} \|\Delta_{q'} f\|_{L^2} \\ &\lesssim \| |D|^{1-\beta} v \|_{L^\infty} \sum_{|q'-q| \leq 4} \|\Delta_{q'} f\|_{L^2}. \end{aligned}$$

For F_q^4 and F_q^5 , from the spectral property and the fact $2^{q'(1-\beta)} \|\Delta_{q'} \tilde{v}\|_{L^\infty} \approx \| \Delta_{q'} |D|^{1-\beta} \tilde{v} \|_{L^\infty}$, we have

$$2^{-q\beta} \|F_q^4(v, f)\|_{L^2} \lesssim \sum_{q' \geq q-2} 2^{(q-q')(1-\beta)} 2^{q'(1-\beta)} \|\Delta_{q'} \tilde{v}\|_{L^\infty} \|\Delta_q f\|_{L^2} \lesssim \| |D|^{1-\beta} v \|_{L^\infty} \|\Delta_q f\|_{L^2}.$$

$$\begin{aligned} 2^{-q\beta} \|F_q^5(v, f)\|_{L^2} &\lesssim 2^{-q\beta} \sum_{|q'-q| \leq 4} 2^{q'} \|\Delta_{q'} \tilde{v}\|_{L^\infty} \sum_{q'' \leq q'-2} 2^{q''-q'} \|\Delta_{q''} f\|_{L^2} \\ &\lesssim \| |D|^{1-\beta} v \|_{L^\infty} \sum_{q'' \leq q+2} 2^{q''-q} \|\Delta_{q''} f\|_{L^2}. \end{aligned}$$

Besides, for F_q^5 when $v = |D|^{\alpha-1} \mathcal{R}^\perp f$, we alternatively have the following improvement that when $\beta > \alpha - 1$

$$\begin{aligned} 2^{-q\beta} \|F_q^5(v, f)\|_{L^2} &\leq 2^{-q\beta} \sum_{|q'-q|\leq 4} \left\| \Delta_{q'} (Id - \Delta_{-1}) |D|^{\alpha-1} \mathcal{R}^\perp f \right\|_{L^2} \|\nabla S_{q'-1} f\|_{L^\infty} \\ &\lesssim \sum_{|q'-q|\leq 4} \|\Delta_{q'} f\|_{L^2} \sum_{-\infty \leq q'' \leq q'-2} 2^{(\alpha-1-\beta)(q'-q'')} \left\| |D|^{\alpha-\beta} \dot{\Delta}_{q''} f \right\|_{L^\infty} \\ &\lesssim \left\| |D|^{\alpha-\beta} f \right\|_{L^\infty} \sum_{|q'-q|\leq 4} \|\Delta_{q'} f\|_{L^2}. \end{aligned}$$

Finally, for F_q^6 we easily have

$$\begin{aligned} 2^{-q\beta} \|F_q^6(v, f)\|_{L^2} &\lesssim \sum_{q' \geq q-3} 2^{(q-q')(1-\beta)} 2^{q'(1-\beta)} \|\Delta_{q'} \tilde{v}\|_{L^\infty} \sum_{i \in \{\pm 1, 0\}} \|\Delta_{q'+i} f\|_{L^2} \\ &\lesssim \left\| |D|^{1-\beta} v \right\|_{L^\infty} \sum_{q' \geq q-4} 2^{(q-q')(1-\beta)} \|\Delta_{q'} f\|_{L^2}. \end{aligned}$$

Combining the above estimates appropriately yields the inequalities (6.1) and (6.2). \square

6.3 Proof of Lemma 3.3

Proof. We treat the general n -dimensional case. Let $x = (x_1, \tilde{x}) = (x_1, x_2, \dots, x_n)$ and the Fourier variable $\zeta = (\zeta_1, \tilde{\zeta}) = (\zeta_1, \zeta_2, \dots, \zeta_n)$. First we observe that for every $\alpha \in]0, 2[$ (cf. [4])

$$(-|D|^\alpha) \theta = \frac{d}{dh} e^{-h|D|^\alpha} \theta \Big|_{h=0} = \frac{d}{dh} \mathcal{P}_{h,n}^\alpha * \theta \Big|_{h=0}$$

where

$$\mathcal{P}_{h,n}^\alpha(x) := c'_{n,\alpha} \frac{h}{(|x|^2 + \alpha^2 h^{\frac{2}{\alpha}})^{\frac{n+\alpha}{2}}}$$

and $c'_{n,\alpha}$ is the normalization constant such that $\int \mathcal{P}_{h,n}^\alpha dx = 1 (= e^{-h|\zeta|^\alpha}|_{\zeta=0})$. In the following we take $\mathcal{P}_{h,n}$ instead of $\mathcal{P}_{h,n}^\alpha$ for brevity. Thus our task reduces to estimate

$$(\mathcal{P}_{h,n} * \theta)(x) - (\mathcal{P}_{h,n} * \theta)(y).$$

Due to the translation and rotation invariant properties, we may assume that $x = (\frac{\xi}{2}, 0, \dots, 0)$ and $y = (-\frac{\xi}{2}, 0, \dots, 0)$. Then from the symmetry and monotonicity of the kernel $\mathcal{P}_{h,n}$ and the fact

$$\int_{\mathbb{R}^{n-1}} \mathcal{P}_{h,n}(x_1, \tilde{x}) d\tilde{x} = \mathcal{F}^{-1}(\widehat{\mathcal{P}_{h,n}}|_{\tilde{\zeta}=0})(x_1) = \mathcal{F}^{-1}(e^{-h|\zeta_1|^\alpha})(x_1) = \mathcal{P}_{h,1}(x_1)$$

we have

$$\begin{aligned}
& (\mathcal{P}_{h,n} * \theta)(x) - (\mathcal{P}_{h,n} * \theta)(y) \\
&= \iint_{\mathbb{R}^n} [\mathcal{P}_{h,n}(\frac{\xi}{2} - \eta, -\tilde{\eta}) - \mathcal{P}_{h,n}(-\frac{\xi}{2} - \eta, -\tilde{\eta})] \theta(\eta, \tilde{\eta}) d\eta d\tilde{\eta} \\
&= \int_{\mathbb{R}^{n-1}} d\tilde{\eta} \int_0^\infty [\mathcal{P}_{h,n}(\frac{\xi}{2} - \eta, \tilde{\eta}) - \mathcal{P}_{h,n}(-\frac{\xi}{2} - \eta, \tilde{\eta})] [\theta(\eta, \tilde{\eta}) - \theta(-\eta, \tilde{\eta})] d\eta \\
&\leq \int_{\mathbb{R}^{n-1}} d\tilde{\eta} \int_0^\infty [\mathcal{P}_{h,n}(\frac{\xi}{2} - \eta, \tilde{\eta}) - \mathcal{P}_{h,n}(-\frac{\xi}{2} - \eta, \tilde{\eta})] \omega(2\eta) d\eta \\
&= \int_0^\infty [\mathcal{P}_{h,1}(\frac{\xi}{2} - \eta) - \mathcal{P}_{h,1}(-\frac{\xi}{2} - \eta)] \omega(2\eta) d\eta \\
&= \int_0^{\frac{\xi}{2}} \mathcal{P}_{h,1}(\eta) [\omega(2\eta + \xi) + \omega(\xi - 2\eta)] d\eta + \int_{\frac{\xi}{2}}^\infty \mathcal{P}_{h,1}(\eta) [\omega(2\eta + \xi) - \omega(2\eta - \xi)] d\eta
\end{aligned}$$

Because of $\int_0^\infty \mathcal{P}_{h,1}(\eta) d\eta = \frac{1}{2}$, we have the estimate of the difference

$$\begin{aligned}
& (\mathcal{P}_{h,n} * \theta)(x) - (\mathcal{P}_{h,n} * \theta)(y) - \omega(\xi) \\
&\leq \int_0^{\frac{\xi}{2}} \mathcal{P}_{h,1}(\eta) [\omega(2\eta + \xi) + \omega(\xi - 2\eta) - 2\omega(\xi)] d\eta \\
&\quad + \int_{\frac{\xi}{2}}^\infty \mathcal{P}_{h,1}(\eta) [\omega(2\eta + \xi) - \omega(2\eta - \xi) - 2\omega(\xi)] d\eta
\end{aligned}$$

Hence from the above estimates and the explicit formula of kernel $\mathcal{P}_{h,1}$, we can conclude that

$$\begin{aligned}
& [(-|D|^\alpha)\theta](x) - [(-|D|^\alpha)\theta](y) \\
&= \lim_{h \rightarrow 0} \frac{[(\mathcal{P}_{h,n} * \theta)(x) - \theta(x)] - [(\mathcal{P}_{h,n} * \theta)(y) - \theta(y)]}{h} \\
&= \lim_{h \rightarrow 0} \frac{(\mathcal{P}_{h,n} * \theta)(x) - (\mathcal{P}_{h,n} * \theta)(y) - \omega(\xi)}{h} \\
&\lesssim_{\alpha,n} \int_0^{\frac{\xi}{2}} \frac{\omega(\xi + 2\eta) + \omega(\xi - 2\eta) - 2\omega(\xi)}{\eta^{1+\alpha}} d\eta + \int_{\frac{\xi}{2}}^\infty \frac{\omega(2\eta + \xi) - \omega(2\eta - \xi) - 2\omega(\xi)}{\eta^{1+\alpha}} d\eta
\end{aligned}$$

□

Acknowledgments: The authors would like to thank Prof. P.Constantin for helpful advice and discussion. They would also like to express their deep gratitude to the anonymous referees for their kind suggestions. The authors were partly supported by the NSF of China (No.10725102).

References

- [1] H. Abidi, T. Hmidi, On the global wellposedness of the critical quasi-geostrophic equation, SIAM J. Math. Anal. **40**(2008), 167-185.
- [2] A.L. Bertozzi and A.J.Majda, *Vorticity and Incompressible Flow*, Cambridge University Press, Cambridge, (2002).

- [3] L. Caffarelli and V. Vasseur, Drift diffusion equations with fractional diffusion and the quasi-geostrophic equations. Arxiv, math.AP/0608447, To appear in Annals of Math.
- [4] L. Caffarelli and L. Silvestre. An extension problem related to the fractional Laplacian. Communications in PDE, **32**, Issue 8(2007), 1245-1260
- [5] J-Y. Chemin, Perfect incompressible fluids, Clarendon press, Oxford, (1998).
- [6] Q. Chen, C. Miao and Z. Zhang, A new Bernstein's inequality and the 2D dissipative quasi-geostrophic equation. Comm. Math. Phys. **271**(2007), 821-838
- [7] P. Constantin, A.J. Majda and E. Tabak, Formation of strong fronts in the 2-D quasigeostrophic thermal active scalar. Nonlinearity **7**(1994), 1495-1533.
- [8] P. Constantin and J. Wu, Behavior of solutions of 2D quasi-geostrophic equations. SIAM J. Math. Anal. **30**(1999), 937-948.
- [9] P. Constantin, D. Cordoba and J. Wu, On the critical dissipative quasi-geostrophic equation. Indiana Univ. Math. J. **50**(2001), 97-107.
- [10] P. Constantin, G. Iyer and J. Wu, Global regularity for a modified critical dissipative quasi-geostrophic equation. Indiana Univ. Math. Journal, **57**(2008), 2681-2692.
- [11] A. Córdoba and D. Córdoba, A maximum principle applied to the quasi-geostrophic equations. Comm. Math. Phys. **249**(2004) 511-528
- [12] R. Danchin and M. Paicu. Global existence results for the anisotropic Boussinesq system in dimension two. Arxiv, math.AP/08094964.
- [13] H. Dong and D. Du, Global well-posedness and a decay estimate for the critical dissipative quasi-geostrophic equation in the whole space. Discrete Contin. Dyn. Syst. **21** (2008) no. 4, 1095-1101.
- [14] H. Dong, Dissipative quasi-geostrophic equations in critical Sobolev spaces: smoothing effect and global well-posedness. Discrete Contin. Dyn. Syst. **26** Issue 4(2010), 1197-1211.
- [15] J. Duoandikoetxea. *Fourier analysis* (Translated and revised by D. Cruz-Uribe). GSM 29, AMS, Providence, RI, 2001.
- [16] S. Friedlander and V. Vicol. Global well-posedness for an advection-diffusion equation arising in magneto-geostrophic dynamics. Arxiv, math.AP/1007.1211v1.
- [17] A. Kiselev, F. Nazarov and A. Volberg, Global well-posedness for the critical 2D dissipative quasi-geostrophic equation, Invent. Math. **167**(2007), 445-453.
- [18] A. Kiselev, F. Nazarov and R. Shterenberg, On blow up and regularity in dissipative Burgers equation. Dynamics of PDE, **5**(2008), 211-240.
- [19] A. Kiselev, Regularity and blow up for active scalars, Math. Model. Math. Phenom. **5** (2010), 225-255.
- [20] A. Kiselev, Nonlocal maximum principle for active scalars. Arxiv, math.AP/1009.0542.
- [21] R. May. Global well-posedness for a modified 2D dissipative quasi-geostrophic equation with initial data in the critical Sobolev space H^1 , Arxiv, math.AP/0910.0998v1.
- [22] C. Miao and L. Xue, On the regularity of a class of generalized quasi-geostrophic equations, Arxiv, math.AP/1011.6214.

- [23] J. Wu, Global solutions of the 2D dissipative quasi-geostrophic equation in Besov spaces. *SIAM J. Math. Anal.* **36**(2004), 1014-1030.
- [24] X. Yu, Remarks on the global regularity for the super-critical 2D dissipative quasi-geostrophic equation, *J. Math. Anal. Appl.* **339**(2008), 359-371.