

Unitaries in a Simple C^* -algebra of Tracial Rank One

Huaxin Lin

Abstract

Let A be a unital separable simple infinite dimensional C^* -algebra with tracial rank no more than one and with the tracial state space $T(A)$ and let $U(A)$ be the unitary group of A . Suppose that $u \in U_0(A)$, the connected component of $U(A)$ containing the identity. We show that, for any $\epsilon > 0$, there exists a selfadjoint element $h \in A_{s,a}$ such that

$$\|u - \exp(ih)\| < \epsilon.$$

We also study the problem when u can be approximated by unitaries in A with finite spectrum.

Denote by $CU(A)$ the closure of the subgroup of unitary group of $U(A)$ generated by its commutators. It is known that $CU(A) \subset U_0(A)$. Denote by \hat{a} the affine function on $T(A)$ defined by $\hat{a}(\tau) = \tau(a)$. We show that u can be approximated by unitaries in A with finite spectrum if and only if $u \in CU(A)$ and $u^n + \widehat{(u^n)^*}, i(u^n - \widehat{(u^n)^*}) \in \overline{\rho_A(K_0(A))}$ for all $n \geq 1$. Examples are given that there are unitaries in $CU(A)$ which can not be approximated by unitaries with finite spectrum. Significantly these results are obtained in the absence of amenability.

1 Introduction

Let M_n be the C^* -algebra of $n \times n$ matrices and let $u \in M_n$ be a unitary. Then u can be diagonalized, i.e., $u = \sum_{k=1}^n e^{i\theta_k} p_k$, where $\theta_k \in \mathbb{R}$ and $\{p_1, p_2, \dots, p_n\}$ are mutually orthogonal projections. As a consequence, $u = \exp(ih)$, where $h = \sum_{k=1}^n \theta_k p_k$ is a selfadjoint matrix. Now let A be a unital C^* -algebra and let $U(A)$ be the unitary group of A . Denote by $U_0(A)$ the connected component of $U(A)$ containing the identity. Suppose that $u \in U_0(A)$. Even in the case that A has real rank zero, $sp(u)$ can have infinitely many points and it is impossible to write u as an exponential, in general. However, it was shown ([3]) that u can be approximated by unitaries in A with finite spectrum if and only if A has real rank zero. This is an important and useful feature for C^* -algebras of real rank zero. In this case, u is a norm limit of exponentials.

Tracial rank for C^* -algebras was introduced (see [4]) in the connection with the program of classification of separable amenable C^* -algebras, or otherwise known as the Elliott program. Unital separable simple amenable C^* -algebras with tracial rank no more than one which satisfy the universal coefficient theorem have been classified by the Elliott invariant ([1] and [5]). A unital separable simple C^* -algebra A with $TR(A) = 1$ has real rank one. Therefore a unitary $u \in U_0(A)$ may not be approximated by unitaries with finite spectrum. We will show that, in a unital infinite dimensional simple C^* -algebra A with tracial rank no more than one, if u can be approximated by unitaries in A with finite spectrum then u must be in $CU(A)$, the closure of the subgroup generated by commutators of the unitary group. A related problem is whether every unitary $u \in U_0(A)$ can be approximated by unitaries which are exponentials. Our first result is to show that, there are selfadjoint elements $h_n \in A_{s,a}$ such that

$$u = \lim_{n \rightarrow \infty} \exp(ih_n)$$

(converge in norm). It should be mentioned that exponential rank has been studied quite extendedly (see [14], [11], [12], [13], etc.). In fact, it was shown by N. C. Phillips that a unital simple C^* -algebra A which is an inductive limit of finite direct sums of C^* -algebras with the form $C(X_{i,n}) \otimes M_{i,n}$ with the dimension of $X_{i,n}$ is bounded has exponential rank $1 + \epsilon$, i.e., every unitary $u \in U_0(A)$ can be approximated by unitaries which are exponentials (see [11]). These simple C^* -algebras have tracial rank one or zero. Theorem 3.3 was proved without assuming A is an AH-algebra, in fact, it was proved in the absence of amenability.

Let $T(A)$ be the tracial state space of A . Denote by $\text{Aff}(T(A))$ the space of all real affine continuous functions on $T(A)$. Denote by $\rho_A : K_0(A) \rightarrow \text{Aff}(T(A))$ the positive homomorphism induced by $\rho_A([p])(\tau) = \tau(p)$ for all projections in $M_k(A)$ (with $k = 1, 2, \dots$) and for all $\tau \in T(A)$. It was introduced by de la Harpe and Scandalis ([2]) a determinant like map Δ which maps $U_0(A)$ into $\text{Aff}(T(A))/\overline{\rho_A(K_0(A))}$. By a result of K. Thomsen ([15]) the de la Harpe and Scandalis determinant induces an isomorphism between $\text{Aff}(T(A))/\overline{\rho_A(K_0(A))}$ and $U_0(A)/CU(A)$. We found out that if u can be approximated by unitaries in A with finite spectrum then u must be in $CU(A)$. But can every unitary in $CU(A)$ be approximated by unitaries with finite spectrum? To answer this question, we consider even simpler question: when can a self-adjoint element in a unital separable simple C^* -algebra with $TR(A) = 1$ be approximated by self-adjoint elements with finite spectrum? Immediately, a necessary condition for a self-adjoint element $a \in A$ to be approximated by self-adjoint elements with finite spectrum is that $\widehat{h^n} \in \overline{\rho_A(K_0(A))}$ (for all $n \in \mathbb{N}$). Given a unitary $u \in U_0(A)$, there is an affine continuous map from $\text{Aff}(T(C(\mathbb{T})))$ into $\text{Aff}(T(A))$ induced by u . Let $\Gamma(u) : \text{Aff}(T(C(\mathbb{T}))) \rightarrow \text{Aff}(T(A))/\overline{\rho_A(K_0(A))}$ be the map given by u . Then it is clear that $\Gamma(u) = 0$ is a necessary condition for u being approximated by unitaries with finite spectrum. Note that $\Gamma(u) = 0$ if and only if $\widehat{u^n + (u^n)^*}, i\widehat{(u^n - (u^n)^*)} \in \overline{\rho_A(K_0(A))}$ for all positive integers n . By applying a uniqueness theorem together with classification results in simple C^* -algebras, we show that the condition is also sufficient. From this, we show that a unitary $u \in CU(A)$ can be approximated by unitaries with finite spectrum if and only if $\Gamma(u) = 0$. We also show that $\Delta(u) = 0$ is not sufficient for $\Gamma(u) = 0$. Therefore, there are unitaries in $CU(A)$ which can not be approximated by unitaries with finite spectrum (see 4.7). Perhaps more interesting fact is that $\Gamma(u) = 0$ does not imply that $\Delta(u) = 0$ for $u \in U_0(A)$ (see 3.7 and 4.9).

2 Preliminaries

2.1. Denote by \mathcal{I} the class of C^* -algebras which are finite direct sums of C^* -subalgebras with the form $M_k(C([0, 1]))$ or M_k , $k = 1, 2, \dots$

Definition 2.2. Recall that a unital simple C^* -algebra A is said to have tracial rank no more than one (or $TR(A) \leq 1$), if for any $\epsilon > 0$, any $a \in A_+ \setminus \{0\}$ and any finite subset $\mathcal{F} \subset A$, there exists a projection $p \in A$ and a C^* -subalgebra B with $1_B = p$ such that

- (1) $\|px - xp\| < \epsilon$ for all $x \in \mathcal{F}$;
- (2) $\text{dist}(pxp, B) < \epsilon$ for all $x \in \mathcal{F}$ and
- (3) $1 - p$ is Murry-von Neumann equivalent to a projection in \overline{aAa} .

Recall that, in the above definition, if B can always be chosen to have finite dimension, then A has tracial rank zero ($TR(A) = 0$). If $TR(A) \leq 1$ but $TR(A) \neq 0$, we write $TR(A) = 1$.

Every unital simple AH-algebra with very slow dimension growth has tracial rank no more than one (see Theorem 2.5 of [5]). There are C^* -algebras with tracial rank no more than one which are not amenable.

Definition 2.3. Suppose that $u \in U(A)$. We will use \bar{u} for the image of u in $U(A)/CU(A)$. If

$x, y \in U(A)/CU(A)$, define

$$\text{dist}(x, y) = \inf\{\|u - v\| : \bar{u} = x \text{ and } \bar{v} = y\}.$$

Let C be another unital C^* -algebra and let $\varphi : C \rightarrow A$ be a unital homomorphism. Denote by $\varphi^\ddagger : U(C)/CU(C) \rightarrow U(A)/CU(A)$ the homomorphism induced by φ .

2.4. Let A be a unital separable simple C^* -algebra with $TR(A) \leq 1$, then A is quasi-diagonal, stable rank one, weakly unperforated $K_0(A)$ and, if $p, q \in A$ are two projections, then p is equivalent to a projection $p' \leq q$ whenever $\tau(p) < \tau(q)$ for all tracial states τ in $T(A)$ (see [4]).

For unitary group of A , we have the following:

- (i) $CU(A) \subset U_0(A)$ (Lemma 6.9 of [5]);
- (ii) $U_0(A)/CU(A)$ is torsion free and divisible (Theorem 6.11 and Lemma 6.6 of [5]);

Theorem 2.5. (Theorem 3.4 of [9]) *Let A be a unital separable simple C^* -algebra with $TR(A) \leq 1$ and let $e \in A$ be a non-zero projection. Then the map $u \mapsto u + (1 - e)$ induces an isomorphism j from $U(eAe)/CU(eAe)$ onto $U(A)/CU(A)$.*

Corollary 2.6. *Let A be a unital separable simple C^* -algebra with $TR(A) \leq 1$. Then the map $j : a \rightarrow \text{diag}(a, \overbrace{1, 1, \dots, 1}^m)$ from A to $M_n(A)$ induces an isomorphism from $U(A)/CU(A)$ onto $U(M_n(A))/CU(M_n(A))$ for any integer $n \geq 1$.*

Definition 2.7. Let $u \in U_0(A)$. There is a piece-wise smooth and continuous path $\{u(t) : t \in [0, 1]\} \subset A$ such that $u(0) = u$ and $u(1) = 1$. Define

$$R(\{u(t)\})(\tau) = \frac{1}{2\pi i} \int_0^1 \tau\left(\frac{du(t)}{dt} u(t)^*\right) dt.$$

$R(\{u(t)\})(\tau)$ is real for every τ .

Definition 2.8. Let A be a unital C^* -algebra with $T(A) \neq \emptyset$. As in [2] and [15], define a homomorphism $\Delta : U_0(A) \rightarrow \text{Aff}(T(A))/\overline{\rho_A(K_0(A))}$ by

$$\Delta(u) = \Pi\left(\frac{1}{2\pi} \int_0^1 \tau\left(\frac{du(t)}{dt} u(t)^*\right) dt\right),$$

where $\Pi : \text{Aff}(T(A)) \rightarrow \text{Aff}(T(A))/\overline{\rho_A(K_0(A))}$ is the quotient map and where $\{u(t) : t \in [0, 1]\}$ is a piece-wise smooth and continuous path of unitaries in A with $u(0) = u$ and $u(1) = 1_A$. This is well-defined and is independent of the choices of the paths.

The following is a combination of a result of K. Thomsen ([15]) and the work of [2]. We state here for the convenience.

Theorem 2.9. *Let A be a unital separable simple C^* -algebra with $TR(A) \leq 1$. Suppose that $u \in U_0(A)$. Then the following are equivalent:*

- (1) $u \in CU(A)$;
- (2) $\Delta(u) = 0$;
- (3) for some piecewise continuous path of unitaries $\{u(t) : t \in [0, 1]\} \subset A$ with $u(0) = u$ and $u(1) = 1_A$,

$$R(\{u(t)\}) \in \overline{\rho_A(K_0(A))},$$

- (4) for any piecewise continuous path of unitaries $\{u(t) : t \in [0, 1]\} \subset A$ with $u(0) = u$ and $u(1) = 1_A$,

$$R(\{u(t)\}) \in \overline{\rho_A(K_0(A))}.$$

(5) there are $h_1, h_2, \dots, h_m \in A_{s.a.}$ such that

$$u = \prod_{j=1}^m \exp(ih_j) \quad \text{and} \quad \sum_{j=1}^m \widehat{h}_j \in \overline{\rho_A(K_0(A))}.$$

(6) $\sum_{j=1}^m \widehat{h}_j \in \overline{\rho_A(K_0(A))}$ for any $h_1, h_2, \dots, h_m \in A_{s.a.}$ for which

$$u = \prod_{j=1}^m \exp(ih_j).$$

Proof. Equivalence of (2), (3), (4), (5) and (6) follows from the definition of the determinant and follows from the Bott periodicity (see [2]). The equivalence of (1) and (2) follows from 3.1 of [15]. \square

The following is a consequence of 2.9.

Theorem 2.10. *Let A be a unital simple separable C^* -algebra with $TR(A) \leq 1$. Then $\ker \Delta = CU(A)$. The de la Harpe and Skandalis determinant gives an isomorphism:*

$$\overline{\Delta} : U_0(A)/CU(A) \rightarrow \text{Aff}(T(A))/\overline{\rho_A(K_0(A))}.$$

Moreover, one has the following short exact (splitting) sequence

$$0 \rightarrow \text{Aff}(T(A))/\overline{\rho_A(K_0(A))} \xrightarrow{\overline{\Delta}^{-1}} U(A)/CU(A) \rightarrow K_1(A) \rightarrow 0.$$

(Note that $U_0(A)/CU(A)$ is divisible in this case, by 6.6 of [5].)

3 Exponentials and approximate unitary equivalence orbit of unitaries

Theorem 3.1. *Let A be a unital simple C^* -algebra with $TR(A) \leq 1$ and let $\gamma : C(\mathbb{T})_{s.a} \rightarrow \text{Aff}(T(A))$ be a (positive) affine continuous map.*

For any $\epsilon > 0$, there exists $\delta > 0$ and there exists a finite subset $\mathcal{F} \subset C(\mathbb{T})_{s,a}$ satisfying the following: If $v \in U_0(A)$ with

$$|\tau(f(u)) - \gamma(f)(\tau)| < \delta, \quad \text{for all } f \in \mathcal{F} \text{ and } \tau \in T(A), \quad \text{and} \quad (\text{e3.1})$$

$$\text{dist}(\bar{u}, \bar{v}) < \delta \text{ in } U_0(A)/CU(A). \quad (\text{e3.2})$$

Then there exists a unitary $W \in U(A)$ such that

$$\|u - W^*vW\| < \epsilon. \quad (\text{e3.3})$$

Proof. The lemma follows immediately from 3.11 of [6]. See also 11.5 of [7] and 3.15 of [6]. Note that, in 3.15 of [6], we can replace the given map h_1 (in this case a given unitary) by a given map γ . \square

Corollary 3.2. *Let A be a unital simple C^* -algebra with $TR(A) \leq 1$ and let $u \in U_0(A)$ be a unitary. For any $\epsilon > 0$, there exists $\delta > 0$ and there exists an integer $N \geq 1$ satisfying the following: If $v \in U_0(A)$ with*

$$|\tau(u^k) - \tau(v^k)| < \delta, \quad k = 1, 2, \dots, N \text{ for all } \tau \in T(A) \text{ and} \quad (\text{e3.4})$$

$$\text{dist}(\bar{u}, \bar{v}) < \delta \text{ in } U_0(A)/CU(A). \quad (\text{e3.5})$$

Then there exists a unitary $W \in U(A)$ such that

$$\|u - W^*vW\| < \epsilon. \quad (\text{e3.6})$$

Proof. Note that (e3.4),

$$|\tau(u^k) - \tau(v^k)| < \delta \quad k = \pm 1, \pm 2, \dots, \pm N. \quad (\text{e3.7})$$

For any subset $\mathcal{G} \subset C(S^1)$ and any $\eta > 0$, there exists $N \geq 1$ and $\delta > 0$ such that

$$|\tau(g(u)) - \tau(g(v))| < \eta \quad \text{for all } \tau \in T(A)$$

if (e3.7) holds.

Then the lemma follows from 3.1 (or 3.16 of [6]). □

Theorem 3.3. *Let A be a unital simple C^* -algebra with $TR(A) \leq 1$. Suppose that $u \in U_0(A)$, then, for any $\epsilon > 0$, there exists a selfadjoint element $a \in A_{s,a}$ such that*

$$\|u - \exp(ia)\| < \epsilon. \quad (\text{e3.8})$$

Proof. Since $u \in U_0(A)$, we may write

$$u = \prod_{j=1}^k \exp(ih_j). \quad (\text{e3.9})$$

Let $M = \max\{\|h_j\| : j = 1, 2, \dots, k\} + 1$. Let $\delta > 0$ and N be given in 3.2 for u . We may assume that $\delta < 1$ and $N \geq 3$. We may also assume that $\delta < \epsilon$. Since $TR(A) \leq 1$, there exists a projection $p \in A$ and a C^* -subalgebra $B \in A$ with $1_B = p$ such that $B \cong \oplus_{i=1}^m C(X_i, M_{r(i)})$, where $X_i = [0, 1]$ or a point, and

$$\|pu - up\| < \frac{\delta}{16NMk}, \quad (\text{e3.10})$$

$$\|(1-p)u(1-p) - (1-p) \prod_{j=1}^k \exp(i((1-p)h_j(1-p)))\| < \frac{\delta}{16NMk}, \quad (\text{e3.11})$$

$$pup \in \frac{\delta}{16NMk} B \quad \text{and} \quad \tau(1-p) < \frac{\delta}{2NMk} \quad \text{for all } \tau \in T(A). \quad (\text{e3.12})$$

There exist unitary $u_1 \in B$ such that

$$\|pup - u_1\| < \frac{\delta}{8NMk} \quad (\text{e3.13})$$

Put $u_2 = (1-p) \prod_{j=1}^k \exp(i(1-p)h_j(1-p))$. Since $u_1 \in B$, it is well known that there exists a selfadjoint element $b \in B_{s,a}$ such that

$$\|u_1 - p \exp(ib)\| < \frac{\delta}{16NMk}. \quad (\text{e3.14})$$

Let $v_0 = (1-p) + p \exp(ib)$ and $u_0 = p \exp(ib) + u_2$. Then, by (e3.10), (e3.11), (e3.13) and (e3.14),

$$\|u_0 - u\| < \|u - pup - (1-p)u(1-p)\| \quad (\text{e3.15})$$

$$+ \|(pup - p \exp(ib)) + ((1-p)u(1-p) - u_2)\| \quad (\text{e3.16})$$

$$< \frac{\delta}{8NMk} + \frac{\delta}{8NMk} + \frac{\delta}{16NMk} = \frac{3\delta}{8NMk}. \quad (\text{e3.17})$$

and

$$u_0 v_0^* = \prod_{j=1}^k \exp(i(1-p)h_j(1-p)). \quad (\text{e 3.18})$$

Note that

$$|\tau(\sum_{j=1}^k (1-p)h_j(1-p))| \leq \sum_{j=1}^k |\tau((1-p)h_j(1-p))| \quad (\text{e 3.19})$$

$$= k\tau(1-p) \max\{\|h_j\| : j = 1, 2, \dots, k\} < \delta/16N \quad (\text{e 3.20})$$

for all $\tau \in T(A)$. It follows that

$$\text{dist}(\bar{u}_0, \bar{v}_0) < \delta/16N \text{ in } U_0(A)/CU(A). \quad (\text{e 3.21})$$

It follows from that

$$\text{dist}(\bar{u}, \bar{v}_0) < \delta/8N. \quad (\text{e 3.22})$$

On the other hand, for each $s = 1, 2, \dots, N$, by (e 3.18), (e 3.17) and (e 3.12)

$$|\tau(u^s) - \tau(v_0^s)| \leq |\tau(u^s) - \tau(u_0^s)| + |\tau(u_0^s) - \tau(v_0^s)| \quad (\text{e 3.23})$$

$$\leq \|u^s - u_0^s\| + |\tau((1-p) - (1-p) \prod_{j=1}^k \exp(i(1-p)sh_j(1-p)))| \quad (\text{e 3.24})$$

$$\leq N\|u - u_0\| + 2\tau(1-p) \quad (\text{e 3.25})$$

$$< \frac{3\delta}{8Mk} + \frac{\delta}{MNk} < \delta \quad (\text{e 3.26})$$

for all $\tau \in T(A)$. From the above inequality and (e 3.22) and applying 3.2, one obtains a unitary $W \in U(A)$ such that

$$\|u - W^* v_0 W\| < \epsilon. \quad (\text{e 3.27})$$

Put $a = W^*((1-p) + b)W$. Then

$$\|u - \exp(ia)\| < \epsilon. \quad (\text{e 3.28})$$

□

Note that Theorem 3.3 does not assume that A is amenable, in particular, it may not be a simple AH-algebra. The proof used a kind of uniqueness theorem for unitaries in a unital simple C^* -algebra A with $TR(A) \leq 1$. This bring us to the following theorem which is an immediate consequence of 3.2.

Theorem 3.4. *Let A be a unital simple C^* -algebra with $TR(A) \leq 1$. Let u and v be two unitaries in $U_0(A)$. Then they are approximately unitarily equivalent if and only if*

$$\Delta(u) = \Delta(v) \text{ and} \quad (\text{e 3.29})$$

$$\tau(u^k) = \tau(v^k) \text{ for all } \tau \in T(A), \quad (\text{e 3.30})$$

$k = 1, 2, \dots$

Since $\Delta : U_0(A)/CU(A) \rightarrow \text{Aff}(T(A))/\overline{\rho_A(K_0(A))}$ is an isomorphism, one may ask if (e 3.30) implies that $\Delta(u) = \Delta(v)$? In other words, would $\tau(f(u)) = \tau(f(v))$ for all $f \in C(S^1)$ imply that $\Delta(u) = \Delta(v)$? This becomes a question only in the case that $\overline{\rho_A(K_0(A))} \neq \text{Aff}(T(A))$. Thus we would like to recall the following:

Theorem 3.5. (cf. Theorem [4])

Let A be a unital simple C^* -algebra with $TR(A) \leq 1$. Then the following are equivalent:

- (1) $TR(A) = 0$,
- (2) $\overline{\rho_A(K_0(A))} = \text{Aff}(T(A))$ and
- (3) $CU(A) = U_0(A)$.

However, when $TR(A) = 1$, at least, one has the following:

Proposition 3.6. Let A be a unital simple infinite dimensional C^* -algebra with $TR(A) \leq 1$. If $a \in \overline{\rho_A(K_0(A))}$, then

$$ra \in \overline{\rho_A(K_0(A))} \quad (\text{e 3.31})$$

for all $r \in \mathbb{R}$. In fact, $\overline{\rho_A(K_0(A))}$ is a closed \mathbb{R} -linear subspace of $\text{Aff}(T(A))$.

Proof. Note that $\overline{\rho_A(K_0(A))}$ is an additive subgroup of $\text{Aff}(T(A))$. It suffices to prove the following: Given any projection $p \in A$, any real number $0 < r_1 < 1$ and $\epsilon > 0$, there exists a projection $q \in A$ such that

$$|r_1\tau(p) - \tau(q)| < \epsilon \text{ for all } \tau \in T(A). \quad (\text{e 3.32})$$

Choose $n \geq 1$ such that

$$|m/n - r_1| < \epsilon/2 \text{ and } 1/n < \epsilon/2 \quad (\text{e 3.33})$$

for some $1 \leq m < n$.

Note that $TR(pAp) \leq 1$. By Theorem 5.4 or Lemma 5.5 of [5], there are mutually orthogonal projections $q_0, p_1, p_2, \dots, p_n$ with $[q_0] \leq [p_1]$ and $[p_1] = [p_i]$, $i = 1, 2, \dots, n$ and $\sum_{i=1}^n p_i + q_0 = p$. Put $q = \sum_{i=1}^m p_i$. We then compute that

$$|r_1\tau(p) - \tau(q)| < \epsilon \text{ for all } \tau \in T(A). \quad (\text{e 3.34})$$

□

Theorem 3.7. Let A be a unital simple infinite dimensional C^* -algebra with $TR(A) = 1$. Then there exist unitaries $u, v \in U_0(A)$ with

$$\tau(u^k) = \tau(v^k) \text{ for all } \tau \in T(A), \quad k = 0, \pm 1, \pm 2, \dots, \pm n, \dots$$

such that $\Delta(u) \neq \Delta(v)$. In particular, u and v are not approximately unitarily equivalent.

Proof. Since we assume that $TR(A) = 1$, then, by 3.5, $\text{Aff}(T(A)) \neq \overline{\rho_A(K_0(A))}$ and $U_0(A)/CU(A)$ are not trivial.

Let $\kappa_1, \kappa_2 : K_1(C(\mathbb{T})) \rightarrow U_0(A)/CU(A)$ be two different homomorphisms. Fix an affine continuous map $s : T(A) \rightarrow T_f(C(\mathbb{T}))$, where $T_f(C(\mathbb{T}))$ is the space of strictly positive normalized Borel measures on \mathbb{T} . Denote by $\gamma_0 : \text{Aff}(T(C(\mathbb{T}))) \rightarrow \text{Aff}(T(A))$ the positive affine continuous map induced by $\gamma_0(f)(\tau) = f(s(\tau))$ for all $f \in \text{Aff}(T(C(\mathbb{T})))$ and $\tau \in T(A)$. Let

$$\overline{\gamma_0} : U_0(C(\mathbb{T}))/CU(C(\mathbb{T})) = \text{Aff}(T(C(\mathbb{T})))/\mathbb{Z} \rightarrow \text{Aff}(T(A))/\overline{\rho_A(K_0(A))} = U_0(A)/CU(A)$$

be the map induced by γ_0 . Write

$$U(C(\mathbb{T}))/CU(C(\mathbb{T})) = U_0(C(\mathbb{T}))/CU(C(\mathbb{T})) \oplus K_1(C(\mathbb{T})).$$

Define $\lambda_i : U(C(\mathbb{T}))/CU(C(\mathbb{T})) \rightarrow U_0(A)/CU(A)$ by

$$\lambda_i(x \oplus z) = \overline{\gamma_0}(x) + \kappa_i(z)$$

for $x \in U_0(C(\mathbb{T}))/CU(C(\mathbb{T}))$ and $z \in K_1(C(\mathbb{T}))$, $i = 1, 2$. It follows from 8.4 of [10] that there are two unital monomorphisms $\varphi_1, \varphi_2 : C(\mathbb{T}) \rightarrow A$ such that

$$(\varphi_1)_{*i} = 0, \varphi_i^\dagger = \lambda_i \text{ and } \varphi_i^\natural = s, \quad (\text{e 3.35})$$

$i = 1, 2$. Let z be the standard unitary generator of $C(S^1)$. Define $u = \varphi_1(z)$ and $v = \varphi_2(z)$. Then $u, v \in U_0(A)$. The condition that $\varphi_i^\natural = s$ implies that

$$\tau(u^k) = \tau(v^k) \text{ for all } \tau \in T(A), \quad k = 0, \pm 1, \pm 2, \dots, \pm n, \dots$$

But since $\lambda_1 \neq \lambda_2$,

$$\Delta(u) \neq \Delta(v).$$

Therefore u and v are not approximately unitarily equivalent. □

Remark 3.8. Given any continuous affine map $s : T(A) \rightarrow T_f(C(\mathbb{T}))$, let $\gamma_0 : \text{Aff}(T(C(\mathbb{T}))) \rightarrow \text{Aff}(T(A))$ by defined by $\gamma_0(f)(\tau) = f(s(\tau))$ for all $f \in \text{Aff}(T(C(\mathbb{T})))$ and $\tau \in T(A)$. This further induces a homomorphism $\lambda : U_0(C(\mathbb{T}))/CU(C(\mathbb{T})) \rightarrow U_0(A)/CU(A)$.

Given any element $x \in \text{Aff}(T(A))/\overline{\rho_A(K_0(A))}$, the proof of the above theorem actually says that there is a unitary $u \in U_0(A)$ such that $\Delta(u) = x$ and

$$\tau(f(u)) = f(s(\tau))$$

for all $f \in C(\mathbb{T})_{s.a}$ and $\tau \in T(A)$. Moreover, u induces λ .

4 Approximated by unitaries with finite spectrum

Now we consider the problem when a unitary $u \in U_0(A)$ in a unital simple infinite dimensional C^* -algebra A with $TR(A) \leq 1$ can be approximated by unitaries with finite spectrum. When $TR(A) = 0$, A has real rank zero, it was proved ([3]) that every unitary in $U_0(A)$ can be approximated by unitaries with finite spectrum. When, $TR(A) = 1$, even a selfadjoint element in A may not be approximated by those selfadjoint with finite spectrum. As stated in 3.5, in this case, $\rho_A(K_0(A))$ is not dense in $\text{Aff}(T(A))$. It turns out that that is the only issue.

Lemma 4.1. *Let A be a unital separable simple infinite dimensional C^* -algebra with $TR(A) \leq 1$ and let $h \in A$ be a self-adjoint element. Then h can be approximated by self-adjoint elements with finite spectrum if and only if $\widehat{h^n} \in \overline{\rho_A(K_0(A))}$, $n = 1, 2, \dots$*

Proof. If h can be approximated by self-adjoint elements so can h^n . By 3.6, $\overline{\rho_A(K_0(A))}$ is a closed linear subspace. Therefore $\widehat{h^n} \in \overline{\rho_A(K_0(A))}$ for all n .

Now we assume that $\widehat{h^n} \in \overline{\rho_A(K_0(A))}$, $n = 1, 2, \dots$. The Stone-Weierstrass theorem implies that $\widehat{f(h)} \in \overline{\rho_A(K_0(A))}$ for all real-value functions $f \in C(sp(h))$. For any $\epsilon > 0$, by Lemma 2.4 of [5], there is $f \in C(sp(x))_{s.a.}$ such that

$$\|f(h) - h\| < \epsilon$$

and $sp(f(h))$ consists of a union of finitely many closed intervals and finitely many points.

Thus, to simplify notation, we may assume that $X = sp(h)$ is a union of finitely many intervals and finitely many points. Let $\psi : C(X) \rightarrow A$ be the homomorphism defined by $\psi(f) = f(h)$. Let $s : T(A) \rightarrow T_f(C(X))$ be the affine map defined by $f(s(\tau)) = \psi(f)(\tau)$ for all $f \in \text{Aff}(C(X))$ and $\tau \in T(A)$.

Let B be a unital simple AH-algebra with real rank zero, stable rank one and

$$(K_0(B), K_0(B)_+, [1_B], K_1(B)) \cong (K_0(A), K_0(A)_+, [1_A], K_1(A)).$$

In particular, $K_0(B)$ is weakly unperforated. The proof of Theorem 10.4 of [5] provides a unital homomorphism $\iota : B \rightarrow A$ which carries the above identification. This can be done by applying Proposition 9.10 of [5] and the uniqueness theorem Theorem 8.6 of [5], or better by corollary 11.7 of [7] because $TR(B) = 0$, the maps φ^\ddagger is not needed since $U(B) = CU(B)$ and the map on traces is determined by the map on $K_0(B)$. This also follows immediately from Lemma 8.5 of [10].

Note that $\text{Aff}(T(B)) = \overline{\rho_B(K_0(B))}$. By identifying B with a unital C^* -subalgebra of A , we may write $\overline{\rho_B(K_0(B))} = \overline{\rho_A(K_0(A))}$.

Let $\psi^\natural : \text{Aff}(T(C(X))) \rightarrow \overline{\rho_A(K_0(A))}$ be the map induced by ψ . This gives a an affine map $\gamma : \text{Aff}(T(C(X))) \rightarrow \overline{\rho_B(K_0(B))}$. It follows from Lemma 5.1 of [8] that there exists a unital monomorphism $\varphi : C(X) \rightarrow B$ such that

$$\iota \circ \varphi_{*0} = \psi_{*0} \quad \text{and} \quad (\iota \circ \varphi)^\natural = \psi^\natural,$$

where $(\iota \circ \varphi)^\natural : \text{Aff}(T(C(X))) \rightarrow \text{Aff}(T(A))$ defined by $(\iota \circ \varphi)^\natural(a)(\tau) = \tau(\iota \circ \varphi)(a)$ for all $a \in A_{s.a.}$. It follows from Corollary 11.7 of [7] that ψ and $\iota \circ \varphi$ are approximately unitarily equivalent. On the hand, since B has real rank zero, φ can be approximated by homomorphisms with finite dimensional range. It follows that h can be approximated by self-adjoint elements with finite spectrum. □

Theorem 4.2. *Let A be a unital separable simple infinite dimensional C^* -algebra with $TR(A) \leq 1$ and let $u \in U_0(A)$. Then u can be approximated by unitaries with finite spectrum if and only if $u \in CU(A)$ and*

$$\widehat{u^n + (u^n)^*}, i(\widehat{u^n - (u^n)^*}) \in \overline{\rho_A(K_0(A))}, \quad n = 1, 2, \dots$$

Proof. Suppose that there exists a sequence of unitaries $\{u_n\} \subset A$ with finite spectrum such that

$$\lim_{n \rightarrow \infty} u_n = u.$$

There are mutually orthogonal projections $p_{1,n}, p_{2,n}, \dots, p_{m(n),n} \in A$ and complex numbers $\lambda_{1,n}, \lambda_{2,n}, \dots, \lambda_{m(n),n} \in \mathbb{C}$ with $|\lambda_{i,n}| = 1$, $i = 1, 2, \dots, m(n)$ and $n = 1, 2, \dots$, such that

$$\lim_{n \rightarrow \infty} \left\| u - \sum_{i=1}^{m(n)} \lambda_{i,n} p_{i,n} \right\| = 0.$$

It follows that

$$\lim_{n \rightarrow \infty} \left\| ((u^*)^n + u^n) - \sum_{i=1}^{m(n)} 2\text{Re}(\lambda_{i,n}) p_{i,n} \right\| = 0.$$

By 3.6,

$$\sum_{i=1}^{m(n)} 2\operatorname{Re}(\lambda_{i,n})\widehat{p_{i,n}} \in \overline{\rho_A(K_0(A))}.$$

Thus $\widehat{\operatorname{Re}(u^n)} \in \overline{\rho_A(K_0(A))}$. Similarly, $\widehat{\operatorname{Im}(u^n)} \in \overline{\rho_A(K_0(A))}$.

To show that $u \in CU(A)$, consider a unitary $v = \sum_{i=1}^m \lambda_i p_i$, where $\{p_1, p_2, \dots, p_m\}$ is a set of mutually orthogonal projections such that $\sum_{i=1}^m p_i = 1$, and where $|\lambda_i| = 1$, $i = 1, 2, \dots, m$. Write $\lambda_j = e^{i\theta_j}$ for some real number θ_j , $j = 1, 2, \dots$. Define

$$h = \sum_{j=1}^m \theta_j p_j.$$

Then

$$v = \exp(ih).$$

By 3.6, $\widehat{h} \in \overline{\rho_A(K_0(A))}$. It follows from 2.9 that $v \in CU(A)$. Since u is a limit of those unitaries with finite spectrum, $u \in CU(A)$.

Now assume $u \in CU(A)$ and $u^n + (u^n)^*$, $i(u^n - (u^n)^*) \in \overline{\rho_A(K_0(A))}$ for $n = 1, 2, \dots$. If $sp(u) \neq \mathbb{T}$, then the problem is reduced to the case in 4.1. So we now assume that $sp(u) = \mathbb{T}$. Define a unital monomorphism $\varphi : C(\mathbb{T}) \rightarrow A$ by $\varphi(f) = f(u)$. By the Stone-Weierstrass theorem and 3.6, every real valued function $f \in C(\mathbb{T})$, $\widehat{\varphi(f)} \in \overline{\rho_A(K_0(A))}$.

As in the proof of 4.1, one obtains a unital C^* -subalgebra $B \subset A$ which is a unital simple AH-algebra with tracial rank zero such that the embedding $\iota : B \rightarrow A$ gives an identification:

$$(K_0(B), K_0(B)_+, [1_B], K_1(B)) = (K_0(A), K_0(A)_+, [1_A], K_1(A)).$$

Moreover, by Lemma 5.1 of [8] that there is a unital monomorphism $\psi : C(\mathbb{T}) \rightarrow B$ such that

$$\psi_{*1} = 0 \text{ and } (\iota \circ \psi)^\natural = \varphi^\natural.$$

Note also

$$(\iota \circ \psi)^\sharp = \varphi^\sharp$$

(both are trivial, since $u \in CU(A)$).

It follows from 3.4 (see also Theorem 11.7 of [7]) that $\iota \circ \psi$ and φ are approximately unitarily equivalent. However, since $\psi_{*1} = 0$, in B , by [3], ψ can be approximated by homomorphisms with finite dimensional range. It follows that u can be approximated by unitaries with finite spectrum. □

If A is a finite dimensional simple C^* -algebra, then $TR(A) = 0$. Of course, every unitary in A has finite spectrum. But $CU(A) \neq U_0(A)$. To unify the two cases, we note that $K_0(A) = \mathbb{Z}$. Instead of using $\overline{\rho_A(K_0(A))}$, one may consider the following definition:

Definition 4.3. Let A be a unital C^* -algebra. Denote by $V(\rho_A(K_0(A)))$, the closed \mathbb{R} -linear subspace of $\operatorname{Aff}(T(A))$ generated by $\rho_A(K_0(A))$. Let $\Pi : \operatorname{Aff}(T(A)) \rightarrow \operatorname{Aff}(T(A))/V(\rho_A(K_0(A)))$ be the quotient map. Define the new determinant

$$\tilde{\Delta} : U_0(A) \rightarrow \operatorname{Aff}(T(A))/V(\rho_A(K_0(A)))$$

by

$$\tilde{\Delta}(u) = \Pi \circ \Delta(u) \text{ for all } u \in U_0(A).$$

Note that if A is a finite dimensional C^* -algebra $\text{Aff}(T(A)) = V(\rho_A(K_0(A)))$. Thus $\tilde{\Delta} = 0$. If A is a unital simple infinite dimensional C^* -algebra with $TR(A) \leq 1$, by 3.6,

$$V(\rho_A(K_0(A))) = \overline{\rho_A(K_0(A))}.$$

Definition 4.4. Suppose that $u \in A$ is a unitary with $X = \text{sp}(u)$. Then it induces a positive affine continuous map from $\gamma_0 : C(X)_{s.a.} \rightarrow \text{Aff}(T(A))$ defined by

$$\gamma_0(f(u))(\tau) = \tau(f(u))$$

for all $f \in C(X)_{s.a.}$ and all $\tau \in T(A)$. Let $\Pi : \text{Aff}(T(A)) \rightarrow \text{Aff}(T(A))/V(\rho_A(K_0(A)))$. Put $\Gamma(u) = \Pi \circ \gamma_0$. Then $\Gamma(u)$ is a map from $C(X)_{s.a.}$ into $\text{Aff}(T(A))/V(\rho_A(K_0(A)))$.

It is clear that, $\Gamma(u) = 0$ if and only if $u^n + \widehat{(u^n)^*}, i(u^n + \widehat{(u^n)^*}) \in V(\rho_A(K_0(A)))$ for all $n \geq 1$.

Thus, we may state the following:

Corollary 4.5. *Let A be a unital simple C^* -algebra with $TR(A) \leq 1$ and let $u \in U_0(A)$. Then u can be approximated by unitaries with finite spectrum if and only if*

$$\tilde{\Delta}(u) = 0 \text{ and } \Gamma(u) = 0.$$

4.6. Suppose that $u = \exp(ih)$ for some self-adjoint element $h \in A$. If $u \in CU(A)$, then, by 2.9, $\tilde{\Delta}(u) = 0$, i.e., $\widehat{h} \in V(\rho_A(K_0(A)))$. So one may ask if there are unitaries with $\tilde{\Delta}(u) = 0$ but $\Gamma(u) \neq 0$. Proposition 4.7 below says that this could happen.

Proposition 4.7. *For any unital separable simple C^* -algebra A with $TR(A) = 1$, there is a unitary u with $\tilde{\Delta}(u) = 0$ (or $u \in CU(A)$) such that $\Gamma(u) \neq 0$ and which is not a limit of unitaries with finite spectrum.*

Proof. Let $e \in A$ be a non-zero projection such that there is a projection $e_1 \in (1 - e)A(1 - e)$ such that $[e] = [e_1]$. Then $TR(eAe) \leq 1$ by 5.3 of [4]. Since A does not have real rank zero, one has $TR(eAe) = 1$.

It follows from 3.5 that

$$\text{Aff}(T(eAe)) \neq \overline{\rho_A(K_0(eAe))} = \overline{\rho_A(K_0(A))}.$$

Choose $h \in (eAe)_{s.a.}$ with $\|h\| \leq 1$ such that h is not a norm limit of self-adjoint elements with finite spectrum.

If $\widehat{h} \in \overline{\rho_A(K_0(eAe))}$, then define

$$u = \exp(ih).$$

Then, $\Delta(u) = 0$ and by Theorem 2.9, $u \in CU(A)$. Since h can not be approximated by self-adjoint elements with finite spectrum, nor u can be approximated by unitaries with finite spectrum since $h = (1/i) \log(u)$ for a continuous branch of the logarithm (note that $\text{sp}(u) \neq \mathbb{T}$).

Now suppose that $\widehat{h} \notin \overline{\rho_A(K_0(eAe))}$.

We also have, by 3.6, $2\pi\widehat{h} \notin \overline{\rho_A(K_0(A))}$. We claim that there is a rational number $0 < r \leq 1$ such that $r\widehat{h}^2 - 2\pi\widehat{h} \notin \overline{\rho_A(K_0(eAe))}$.

In fact, if $\widehat{h}^2 \in \overline{\rho_A(K_0(eAe))}$, then the claim follows easily. So we assume that $\widehat{h}^2 \notin \overline{\rho_A(K_0(eAe))}$. Suppose that, for some $0 < r_1 < 1$, $r_1\widehat{h}^2 - 2\pi\widehat{h} \in \overline{\rho_A(K_0(eAe))}$. Then $(1 - r_1)\widehat{h}^2 \notin \overline{\rho_A(K_0(eAe))}$. Hence

$$\widehat{h}^2 - 2\pi\widehat{h} = (1 - r_1)\widehat{h}^2 + (r_1\widehat{h}^2 - 2\pi\widehat{h}) \notin \overline{\rho_A(K_0(eAe))}.$$

This proves the claim.

Now define $h_1 = rh + 2\pi e_1 - w^*rhw$, where $w \in A$ is a unitary such that $w^*ew = e_1$. Put

$$u = \exp(ih_1)$$

It follows from 3.6 that

$$2\pi\widehat{e}_1 \in \overline{\rho_A(K_0(eAe))}.$$

Thus $\tau(h_1) = 2\pi\tau(e_1) \in \overline{\rho_A(K_0(eAe))}$. Therefore, by 2.9, $u \in CU(A)$. Since

$$\widehat{h_1^2} = \widehat{r^2h^2} + 4\pi^2\widehat{e}_1 - 4\pi\widehat{rh} + \widehat{r^2h^2} \quad (\text{e 4.36})$$

$$= 2r(\widehat{rh^2} - 2\pi\widehat{h}) - 4\pi^2\widehat{e}_1 \notin \overline{\rho_A(K(A))}. \quad (\text{e 4.37})$$

Therefore, by 4.1, h_1 can not be approximated by self-adjoint elements with finite spectrum. It follows that u can not be approxiamted by unitaries with finite spectrum. \square

Another question is whether $\Gamma(u) = 0$ is sufficient for $\Delta(u) = 0$. For the case that $\text{sp}(u) \neq \mathbb{T}$, one has the following. But in general, 4.9 gives a negative answer.

Proposition 4.8. *Let A be a unital separable simple C^* -algebra with $TR(A) \leq 1$. Suppose that $u \in U_0(A)$ with $\text{sp}(u) \neq \mathbb{T}$. If $\Gamma(u) = 0$, then $\tilde{\Delta}(u) = 0$, $u \in CU(A)$ and u can be approximated by unitaries with finite spectrum.*

Proof. Since $\text{sp}(u) \neq \mathbb{T}$, there is a real valued continuous function $f \in C(\text{sp}(u))$ such that $u = \exp(if(u))$. Thus the condition that $\Gamma(u) = 0$ implies that $\widehat{f(u)} \in \overline{\rho_A(K_0(A))}$. By 2.9, $u \in CU(A)$. \square

Proposition 4.9. *Let A be a unital infinite dimensional separable simple C^* -algebra with $TR(A) = 1$. Then there are unitaries $u \in U_0(A)$ with $\Gamma(u) = 0$ such that $u \notin CU(A)$. In particular, $\tilde{\Delta}(u) \neq 0$ and u can not be approximated by unitaries with finite spectrum.*

Proof. There exists a unital C^* -subalgebra $B \subset A$ with tracial rank zero such that the embedding gives the following identification:

$$(K_0(B), K_0(B)_+, [1_B], K_1(B)) = (K_0(A), K_0(A)_+, [1_A], K_1(A)).$$

Note that $\text{Aff}(T(B)) = \overline{\rho_B(K_0(B))} = \overline{\rho_A(K_0(A))}$.

Let $w \in U_0(B)$ be a unitary with $\text{sp}(w) = \mathbb{T}$. Thus $\Gamma(w) = 0$. Let $\gamma : \text{Aff}(T(C(\mathbb{T}))) \rightarrow \text{Aff}(T(A))$ be given by $\gamma(f)(\tau) = \tau(f(u))$ for $f \in C(T)_{s.a.}$ and $\tau \in T(A)$. Since $TR(A) = 1$, by 2.9, there are unitaries $u_0 \in U_0(A) \setminus CU(A)$. By the proof of 3.7 (see also 3.8), there is a unitary $u \in U_0(A)$ such that

$$\overline{u} = \overline{u_0} \quad \text{and}$$

$$\tau(f(u)) = \tau(f(w)) \quad \text{for all } \tau \in T(A)$$

and for all $f \in C(\mathbb{T})_{s.a.}$. Thus $\tilde{\Delta}(u) \neq 0$ and $\Gamma(u) = \Gamma(w) = 0$. By 4.2, u can not be approximated by unitaries with finite spectrum. \square

References

- [1] G. A. Elliott, G. Gong and L. Li, *On the classification of simple inductive limit C^* -algebras. II, The isomorphism theorem*, Invent. Math. **168** (2007), 249–320.
- [2] P. de la Harpe, P. and G. Skandalis, G. *Déterminant associé à une trace sur une algèbre de Banach*, Ann. Inst. Fourier (Grenoble) **34** (1984), 1, 241–260.
- [3] H. Lin, *Exponential rank of C^* -algebras with real rank zero and the Brown-Pedersen conjectures*, J. Funct. Anal. **114** (1993), 1–11.
- [4] H. Lin, *The tracial topological rank of C^* -algebras* Proc. London Math. Soc. **83** (2001), 199–234.
- [5] H. Lin, *Simple nuclear C^* -algebras of tracial topological rank one*, J. Funct. Anal. **251** (2007), 601–679.
- [6] H. Lin and H. Matui, *Minimal dynamical systems on the product of the Cantor set and the circle. II*, Selecta Math. (N.S.) **12** (2006), 199–239.
- [7] H. Lin, *Approximate Unitary Equivalence in Simple C^* -algebras of Tracial Rank One*, preprint (arXiv:0801.2929).
- [8] H. Lin, *The Range of Approximate Unitary Equivalence Classes of Homomorphisms from AH -algebras*, to appear (arXiv:0801.3858).
- [9] H. Lin, *Homotopy of unitaries in simple C^* -algebras with tracial rank one*, preprint (arXiv:0805.0583).
- [10] H. Lin, *Asymptotically Unitary Equivalence and Classification of Simple Amenable C^* -algebras*, preprint (arXiv:0806.0636).
- [11] N. C. Phillips, *Reduction of exponential rank in direct limits of C^* -algebras*, Canad. J. Math. **46** (1994), 818–853.
- [12] N. C. Phillips, *A survey of exponential rank. C^* -algebras: 1943–1993* (San Antonio, TX, 1993), 352–399, Contemp. Math., **167**, Amer. Math. Soc., Providence, RI, 1994.
- [13] N. C. Phillips, *How many exponentials?* Amer. J. Math. **116** (1994), 1513–1543.
- [14] N. C. Phillips and J. R. Ringrose, *Exponential rank in operator algebras. Current topics in operator algebras* (Nara, 1990), 395–413, World Sci. Publ., River Edge, NJ, 1991.
- [15] K. Thomsen, *Traces, unitary characters and crossed products by Z* Publ. Res. Inst. Math. Sci. **31** (1995), 1011–1029.