

On Brownian motion on the plane with membranes on rays with a common endpoint

Olga V. Aryasova, Andrey Yu. Pilipenko¹

Abstract

We consider a Brownian motion on the plane with semipermeable membranes on n rays that have a common endpoint in the origin. We obtain the necessary and sufficient conditions for the process to reach the origin and we show that the probability of hitting the origin is equal to zero or one.

Introduction

Let c_1, \dots, c_n be given rays in \mathbb{R}^2 such that they share a common endpoint. Suppose polar coordinates in \mathbb{R}^2 are denoted by (r, φ) , $r \geq 0$, $\varphi \in [0, 2\pi)$ and

$$c_k = \{(r, \varphi) : r \geq 0, \varphi = \varphi_k\},$$

where $0 \leq \varphi_k < 2\pi$, $\varphi_j \neq \varphi_k$, $j \neq k$, $j, k \in \{1, \dots, n\}$.

Let $n_k, k = 1, \dots, n$, denote the unit vector normal to c_k that points anticlockwise and let v_k be a vector in \mathbb{R}^2 such that $(v_k, n_k) = 1$. The angle between n_k and v_k denoted by $\theta_k \in (-\frac{\pi}{2}, \frac{\pi}{2})$ is referred to as a positive if and only if v_k points towards the origin. The case of $n = 5, \theta_1 > 0, \theta_2 > 0$ is shown in Figure 1.

Let $\gamma_k, k = 1, \dots, n$, be a real constant such that $|\gamma_k| \leq 1$. Consider the stochastic differential equation in \mathbb{R}^2

$$dx(t) = dw(t) + \sum_{k=1}^n \gamma_k v_k dL_x^{c_k}(t) \quad (1)$$

with initial condition $x(0) = x^0$, $x^0 \in \mathbb{R}^2$.

In this equation $(w(t))_{t \geq 0}$ is a Wiener process in \mathbb{R}^2 , $(L_x^{c_k}(t))_{t \geq 0}$ is a local time of the process $(x(t))_{t \geq 0}$ at the ray c_k , and it is defined by the formula

$$L_x^{c_k}(t) = \lim_{\varepsilon \downarrow 0} \frac{1}{2\varepsilon} \int_0^t \mathbb{I}_{A_\varepsilon^k}(x(s)) ds, \quad (2)$$

¹Research partially supported by Ministry of Education and Science of Ukraine, Grant F26/433-2008, and National Academy of Sciences of Ukraine, Grant 104-2008.




Figure 1.

where $A_\varepsilon^k = \{x \in \mathbb{R}^2 : \exists y \in c_k, s \in [-1, 1] \text{ such that } x = y + \varepsilon sn_k\}$.

The solution of this equation can be regarded as a Wiener process in \mathbb{R}^2 skewing on the rays c_1, \dots, c_n until the time it reaches the origin (see Section 1). The skew on c_k is defined by γ_k being a coefficient of permeability. When γ_k is equal to 1 we have a Wiener process reflecting instantaneously at c_k in the direction of v_k . If γ_k is equal to -1 the process reflects at c_k in the direction of $-v_k$. If $\gamma_k \in (-1, 1)$ a semipermeable membrane is on c_k .

The aim of the investigation is to show that the process hits the origin, when starting away from it, with probability zero or one. We obtain an explicit expression in variables $\theta_1, \dots, \theta_n, \gamma_1, \dots, \gamma_n, \xi_1, \dots, \xi_n$ whose sign this probability depends on.

The case of the wedge (two rays) with reflection on its sides was studied by Varadhan and Williams [1]. They proved (in our notation) that the process does not reach the origin if $\theta_1 - \theta_2 \leq 0$

and does reach the origin if $\theta_1 - \theta_2 > 0$. In this case our result coincides with that of Varadhan and Williams (see Example 2).

1. Construction of the process

In this Section we build the process which is a solution of equation (1) up to the time it hits the origin.

Conventionally, we put $\varphi_{nl+k} = \varphi_k, c_{nl+k} = c_k, \gamma_{nl+k} = \gamma_k, k = 1, \dots, n, l \in \mathbb{N}$. Define $\xi_k = \varphi_{k+1} - \varphi_k$. Let $S_k = \{(r, \varphi) : r > 0, \varphi_k \leq \varphi \leq \varphi_{k+1}\}, k = 1, \dots, n$, be the wedge $c_k 0 c_{k+1}$. Let $\varphi_1 = 0$.

Remark 1. Without loss of generality we can assume that $\xi_k \leq \pi, k = 1, \dots, n$. Indeed, otherwise we can introduce additional ray

$$c_{ad} = \{(r, \varphi) : r \geq 0, \varphi = \pi\},$$

and put $\gamma_{ad} = 0$.

So, throughout this Section we assume $\xi_k \leq \pi/2, k = 1, \dots, n$.

Lemma 1. *Given $x^0 \in \mathbb{R}^2$, there exists a unique strong solution of the equation (1) up to the first time of hitting the origin.*

Proof. (i) The process $(x(t))_{t \geq 0}$ is an ordinary Wiener process up to the first time of hitting one of the rays c_1, \dots, c_n . Define

$$\tau_1 = \inf \{t \geq 0 : x(t) \in c_{j_1} \text{ for some } j_1 \in \{1, \dots, n\}\}.$$

Without loss of generality, let $x(\tau_1) \in c_1$. Until the process reaches another ray only the local time at the ray c_1 can increase. Put

$$\tau_2 = \inf \{t > \tau_1 : x(t) \in c_{j_2} \text{ for some } j_2 \in \{1, \dots, n\}, j_2 \neq j_1\}.$$

Then for all $t < \tau_2$ equation (1) has the form

$$dx_1(t) = dw_1(t) - \gamma_1 \tan \theta_1 dL_x^{c_1}(t), \quad (3)$$

$$dx_2(t) = dw_2(t) + \gamma_1 dL_x^{c_1}(t), \quad (4)$$

where $(x_1(t), x_2(t))$ are the cartesian coordinates of the process $(x(t))_{t \geq 0}$, $(w_1(t))_{t \geq 0}$ and $(w_2(t))_{t \geq 0}$ are independent one-dimensional Wiener processes. Note that by Remark 1

$$L_x^{c_1}(t) = L_{x_2}^0(t) := \lim_{\varepsilon \downarrow 0} \frac{1}{2\varepsilon} \int_0^t \mathbb{1}_{[-\varepsilon, \varepsilon]}(x_2(s)) ds, \quad t \leq \tau_2,$$

where $(L_{x_2}^0(t))_{t \geq 0}$ is a usual local time of the process $(x_2(t))_{t \geq 0}$ at zero. There exists a unique strong solution of equation (4) (cf.[4]). Substitution of solution to (4) into (3) allows us to get the solution of the latter. Hence we have proved existence and uniqueness of solution to equation (1) up to the time τ_2 .

(ii) Put for $m \geq 2$

$$\tau_m = \inf \{t > \tau_{m-1} : x(t) \in c_{j_m} \text{ for some } j_m \in \{1, \dots, n\}, j_m \neq j_{m-1}\}.$$

Assume $x(\tau_2) \in c_2$. Then for $\tau_2 \leq t < \tau_3$, the equation (1) has the form

$$dx_1(t) = dw_1(t) - \gamma_2 \frac{\sin(\xi_1 + \theta_2)}{\cos \theta_2} dL_x^{c_2}(t), \quad (5)$$

$$dx_2(t) = dw_2(t) + \gamma_2 \frac{\cos(\xi_1 + \theta_2)}{\cos \theta_2} dL_x^{c_2}(t). \quad (6)$$

We do the clockwise rotation of coordinates defined by the formulas

$$\begin{aligned} x'_1 &= x_1 \cos \xi_1 + x_2 \sin \xi_1, \\ x'_2 &= -x_1 \sin \xi_1 + x_2 \cos \xi_1. \end{aligned} \quad (7)$$

This map takes the ray c_2 to the ray c_1 , i.e. $c'_2 = c_1$. Moreover, it preserves the angles. Using the Ito formula we get the stochastic differential equations

$$dx'_1(t) = dw'_1(t) - \gamma_2 \tan \theta_2 dL_{x'}^{c'_2}(t), \quad (8)$$

$$dx'_2(t) = dw'_2(t) + \gamma_2 dL_{x'}^{c'_2}(t), \quad (9)$$

which holds true for $\tau_2 \leq t < \tau_3$. Here $w'_1(t) = \cos \xi_1 dw_1(t) + \sin \xi_1 dw_2(t)$, $t \geq 0$, and $w'_2(t) = -\sin \xi_1 dw_1(t) + \cos \xi_1 dw_2(t)$, $t \geq 0$, are independent one-dimensional Wiener processes as a square integrable martingales with characteristics t . By the arguments similar to those of (i) there exists a unique strong solution to equations (8) and (9) for $\tau_2 \leq t < \tau_3$. It means, because the inverse transformation exists, that the equations (5) and (6) has a unique strong solution for $\tau_2 \leq t < \tau_3$. The case of $x(\tau_2) \in c_n$ can be treated analogously. Thus we have built a strong solution to equation (1) up to the time τ_3 and proved the uniqueness of it.

(iii) Repeating this procedure of localization of equation (1) we build a solution to (1) up to the time $\zeta = \lim_{m \rightarrow \infty} \tau_m$. It is easily

seen that the process $(x(t))_{t \geq 0}$ does not reach infinity in finite time a.s. and does not intersect any ring $a < |x| < b$, $0 < a < b$, $x \in \mathbb{R}^2$, infinitely many times on $[0, T]$. So, $\zeta = +\infty$ or $x(\zeta-) = 0$. \square

2. Coordinate transformation and change of time

As above, we assume that $\xi_k \leq \pi/2$, $k = 1, \dots, n$.

To investigate the value of probability $\mathbb{P}_{x_0}\{\zeta < \infty\}$ we make certain transformations of the process.

Let us change coordinates by the formulas

$$\begin{aligned}\tilde{x}_1 &= -\ln \sqrt{x_1^2 + x_2^2} = -\ln r, \\ \tilde{x}_2 &= \varphi,\end{aligned}$$

where

$$\begin{aligned}r \cos \varphi &= x_1, \\ r \sin \varphi &= x_2.\end{aligned}$$

Therefore,

$$x_1 = e^{-\tilde{x}_1} \cos \tilde{x}_2, \quad (10)$$

$$x_2 = e^{-\tilde{x}_1} \sin \tilde{x}_2. \quad (11)$$

The Jacobian matrix of this transformation is

$$\frac{\partial(\tilde{x}_1, \tilde{x}_2)}{\partial(x_1, x_2)} = \left(\frac{\partial(x_1, x_2)}{\partial(\tilde{x}_1, \tilde{x}_2)} \right)^{-1} = \begin{bmatrix} -e^{\tilde{x}_1} \cos \tilde{x}_2 & -e^{\tilde{x}_1} \sin \tilde{x}_2 \\ -e^{\tilde{x}_1} \sin \tilde{x}_2 & e^{\tilde{x}_1} \cos \tilde{x}_2 \end{bmatrix}.$$

Moreover, $\Delta \tilde{x}_1 = \Delta \tilde{x}_2 = 0$.

The image of the ray c_k , $k = 1, \dots, n$, under this map is the straight-line

$$\tilde{c}_k = \{(\tilde{x}_1, \tilde{x}_2) : \tilde{x}_1 \in \mathbb{R}, \tilde{x}_2 = \varphi_k\}.$$

Clearly, the image of S_k is the strip $\tilde{S}_k = \{(\tilde{x}_1, \tilde{x}_2) : \tilde{x}_1 \in \mathbb{R}, \varphi_k \leq \tilde{x}_2 \leq \varphi_{k+1}\}$. As above we put $\varphi_{nl+k} = \varphi_k$, $\tilde{c}_{nl+k} = \tilde{c}_k$, $k = 1, \dots, n$, $l \in \mathbb{N}$.

Note that if the function f is twice continuously differentiable on $\mathbb{R}^2 \setminus \{0\}$, then by the Ito formula

$$df(x(t)) = (\nabla f, dw(t)) + \sum_{k=1}^n \gamma_k (\nabla f, v_k) dL_x^{c_k}(t) + \frac{1}{2} \Delta f(x(t)) dt, \quad t < \zeta. \quad (12)$$

So

$$\begin{aligned} d\tilde{x}_1(t) &= -e^{\tilde{x}_1(t)}(\cos \tilde{x}_2(t)dw_1(t) + \sin \tilde{x}_2(t)dw_2(t)) - \\ &- e^{\tilde{x}_1(t)} \sum_{k=1}^n \gamma_k (v_{k1} \cos \tilde{x}_2(t) + v_{k2} \sin \tilde{x}_2(t)) dL_x^{c_k}(t), \quad t < \zeta, \end{aligned} \quad (13)$$

where (v_{k1}, v_{k2}) are the cartesian coordinates of the vector v_k . As easily seen, $v_{k1} = -\frac{\sin(\varphi_k + \theta_k)}{\cos \theta_k}$, $v_{k2} = \frac{\cos(\varphi_k + \theta_k)}{\cos \theta_k}$. The process $(L_x^{c_k}(t))_{t \geq 0}$ increases only at those moments of time when $\tilde{x} \in \tilde{c}_k$ and, consequently, $\tilde{x}_2 = \varphi_k$. Besides, the process $\tilde{w}_1(t) = -\cos \tilde{x}_2(t)dw_1(t) - \sin \tilde{x}_2(t)dw_2(t)$, $t \geq 0$, is a Wiener process as a square integrable martingale with its characteristic being equal to t . Thus, we can rewrite the equation (13) as follows

$$d\tilde{x}_1(t) = e^{\tilde{x}_1(t)} d\tilde{w}_1(t) + e^{\tilde{x}_1(t)} \sum_{k=1}^n \gamma_k \tan \theta_k dL_x^{c_k}(t). \quad (14)$$

By analogy,

$$d\tilde{x}_2(t) = e^{\tilde{x}_1(t)} d\tilde{w}_2(t) + e^{\tilde{x}_1(t)} \sum_{k=1}^n \gamma_k dL_x^{c_k}(t), \quad (15)$$

where $\tilde{w}_2(t) = \cos \tilde{x}_2(t)dw_2(t) - \sin \tilde{x}_2(t)dw_1(t)$, $t \geq 0$, is a one-dimensional Wiener process independent of the process $(\tilde{w}_1(t))_{t \geq 0}$.

From (14), (15) we get the equation

$$d\tilde{x}(t) = e^{\tilde{x}_1(t)} d\tilde{w}(t) + e^{\tilde{x}_1(t)} \sum_{k=1}^n \gamma_k \tilde{v}_k dL_x^{c_k}(t), \quad (16)$$

where $\tilde{w}(t) = (\tilde{w}_1(t), \tilde{w}_2(t))$, $t \geq 0$; $\tilde{v}_k = (\tan \theta_k, 1)$, $k = 1, \dots, n$.

We emphasize that $L_x^{c_k}$ is not a local time of the process \tilde{x} but that of the process x . Let us give a form of equation (16) that involves $L_{\tilde{x}}^{\tilde{c}_k}$ being a local time of the process \tilde{x} at the line \tilde{c}_k instead of $L_x^{c_k}$, $k = 1, \dots, n$.

To do this we make use of the following lemma.

Lemma 2. *Let $u(t)$, $t \in [0, T]$, be a continuous function on $[0, T]$ such that for all $t \in [0, T]$ there exists a limit*

$$\lim_{\varepsilon \downarrow 0} \frac{1}{2\varepsilon} \int_0^t \mathbb{I}_{[-\varepsilon, \varepsilon]}(u(s)) ds =: L(t).$$

Suppose f is a continuous function on $[0, T]$ with positive values, $g(s, \varepsilon)$ is a function on $[0, T] \times [0, T]$ with non-negative values such that $g(s, \varepsilon)/\varepsilon \rightarrow f(s)$ as $\varepsilon \downarrow 0$ uniformly in s . Then for all $t \in [0, T]$

$$\lim_{\varepsilon \downarrow 0} \frac{1}{2\varepsilon} \int_0^t \mathbb{I}_{[-g(s, \varepsilon), g(s, \varepsilon)]}(u(s)) ds = \int_0^t f(s) dL(s).$$

Proof. (i) The statement of Lemma is easily justified for $g(s, \varepsilon) = \varepsilon \mathbb{I}_{[a, b]}(s)$, where a, b are constant, $0 \leq a < b$, and, consequently, it is true for linear combinations of such functions.

(ii) Let $\underline{f}_n, \bar{f}_n$ be sequences of piecewise constant functions on $[0, T]$ such that $\underline{f}_n \leq f \leq \bar{f}_n$ and $\underline{f}_n \rightarrow f, \bar{f}_n \rightarrow f$ as $n \rightarrow \infty$ uniformly. By (i) we have

$$\begin{aligned} \int_0^t \underline{f}_n(s) dL(s) &= \lim_{\varepsilon \downarrow 0} \frac{1}{2\varepsilon} \int_0^t \mathbb{I}_{[-\varepsilon \underline{f}_n(s), \varepsilon \underline{f}_n(s)]}(u(s)) ds \leq \\ &\leq \lim_{\varepsilon \downarrow 0} \frac{1}{2\varepsilon} \int_0^t \mathbb{I}_{[-\varepsilon f(s), \varepsilon f(s)]}(u(s)) ds \leq \overline{\lim}_{\varepsilon \downarrow 0} \frac{1}{2\varepsilon} \int_0^t \mathbb{I}_{[-\varepsilon f(s), \varepsilon f(s)]}(u(s)) ds \leq \\ &\leq \lim_{\varepsilon \downarrow 0} \frac{1}{2\varepsilon} \int_0^t \mathbb{I}_{[-\varepsilon \bar{f}_n(s), \varepsilon \bar{f}_n(s)]}(u(s)) ds = \int_0^t \bar{f}_n(s) dL(s). \end{aligned}$$

Passing to the limit as $n \rightarrow \infty$ we get

$$\lim_{\varepsilon \downarrow 0} \frac{1}{2\varepsilon} \int_0^t \mathbb{I}_{[-\varepsilon f(s), \varepsilon f(s)]}(u(s)) ds = \int_0^t f(s) dL(s).$$

(iii) Since $\frac{g(s, \varepsilon)}{\varepsilon} \rightarrow f(s)$ as $\varepsilon \downarrow 0$ uniformly in $s \in [0, T]$, then for each $t \in [0, T]$, $\delta > 0$ there exists $\varepsilon_\delta > 0$ such that for all $\varepsilon < \varepsilon_\delta$ the inequalities

$$\varepsilon(1 - \delta)f(s) \leq g(s, \varepsilon) \leq \varepsilon(1 + \delta)f(s)$$

hold for all $s \in [0, t]$. Similarly to (ii) we have

$$\begin{aligned} (1 - \delta) \int_0^t f(s) dL(s) &\leq \lim_{\varepsilon \downarrow 0} \frac{1}{2\varepsilon} \int_0^t \mathbb{I}_{[-g(s, \varepsilon), g(s, \varepsilon)]}(u(s)) ds \leq \\ &\leq \overline{\lim}_{\varepsilon \downarrow 0} \frac{1}{2\varepsilon} \int_0^t \mathbb{I}_{[-g(s, \varepsilon), g(s, \varepsilon)]}(u(s)) ds \leq (1 + \delta) \int_0^t f(s) dL(s). \end{aligned}$$

Letting $\delta \downarrow 0$ we obtain the statement of the Lemma. \square

Now let us express $L_{\tilde{x}}^{\tilde{c}_k}$ in term of $L_x^{c_k}$.

Lemma 3. *The process*

$$L_{\tilde{x}}^{\tilde{c}_k}(t) = \int_0^t e^{-\tilde{x}_1(s)} dL_x^{c_k}(s), t \leq \zeta,$$

is a local time of the process $(\tilde{x}(t))_{t \geq 0}$ on the straight-line \tilde{c}_k .

Proof. Let $k = 1$. According to (11)

$$\begin{aligned} L_{\tilde{x}}^{\tilde{c}_1}(t) &= \lim_{\varepsilon \downarrow 0} \frac{1}{2\varepsilon} \int_0^t \mathbb{I}_{[-\varepsilon, \varepsilon]}(\tilde{x}_2(s)) ds = \\ &= \lim_{\varepsilon \downarrow 0} \frac{1}{2\varepsilon} \int_0^t \mathbb{I}_{[-e^{-\tilde{x}_1(s)} \sin \varepsilon, e^{-\tilde{x}_1(s)} \sin \varepsilon]} \mathbb{I}_{\{\cos \tilde{x}_2(s) > 0\}} (e^{-\tilde{x}_1(s)} \sin \tilde{x}_2(s)) ds = \\ &= \lim_{\varepsilon \downarrow 0} \frac{1}{2\varepsilon} \int_0^t \mathbb{I}_{[-e^{-\tilde{x}_1(s)} \sin \varepsilon, e^{-\tilde{x}_1(s)} \sin \varepsilon]} (x_2(s)) ds = \\ &= \int_0^t e^{-\tilde{x}_1(s)} dL_x^{c_1}(s). \end{aligned}$$

The last equality follows from Lemma 2. This proves Lemma 3 in the case of $k = 1$. The statement of Lemma for $k = 2, \dots, n$, can be obtained analogously. \square

Using this Lemma we get the form of equation (16) involving only the process \tilde{x} as follows

$$d\tilde{x}(t) = e^{\tilde{x}_1(t)} d\tilde{w}(t) + e^{2\tilde{x}_1(t)} \sum_{k=1}^n \gamma_k \tilde{v}_k dL_{\tilde{x}}^{\tilde{c}_k}(t). \quad (17)$$

Let us make a change of time. Put $A(t) = \int_0^t e^{2\tilde{x}_1(s)} ds$, $t < \zeta$, and denote $\eta = \int_0^\zeta e^{2\tilde{x}_1(s)} ds$. Define the process

$$\hat{x}(t) = \tilde{x}(A^{-1}(t)), \quad t \leq \eta,$$

where

$$A^{-1}(t) = \inf \left\{ s : \int_0^s e^{2\tilde{x}_1(u)} du \geq t \right\}.$$

The equation (17) implies the stochastic integral equation

$$\begin{aligned} \tilde{x}(A^{-1}(t)) &= \tilde{x}(0) + \int_0^{A^{-1}(t)} e^{\tilde{x}_1(s)} d\tilde{w}(t) + \\ &+ \sum_{k=1}^n \gamma_k \tilde{v}_k \int_0^{A^{-1}(t)} e^{2\tilde{x}_1(s)} dL_{\tilde{x}}^{\tilde{c}_k}(s), \quad t \leq \eta. \end{aligned} \quad (18)$$

Similarly to [5], Theorem 7.2, it can be proved that the process $\hat{w}(t) = \int_0^{A^{-1}(t)} e^{\tilde{x}_1(s)} d\tilde{w}(s)$ is a Wiener process in \mathbb{R}^2 up to the time η .

Lemma 4. *The process*

$$L_{\hat{x}}^{\tilde{c}_k}(t) = \int_0^{A^{-1}(t)} e^{2\tilde{x}_1(s)} dL_{\tilde{x}}^{\tilde{c}_k}(s), \quad t \leq \eta,$$

is a local time of the process $(\hat{x}(t))_{t < \eta}$ on the straight-line \tilde{c}_k .

Proof. For $k = 1$ we have

$$\begin{aligned} L_{\hat{x}}^{\tilde{c}_1}(t) &= \lim_{\varepsilon \downarrow 0} \frac{1}{2\varepsilon} \int_0^t \mathbb{I}_{[-\varepsilon, \varepsilon]}(\hat{x}_2(s)) ds = \\ &= \lim_{\varepsilon \downarrow 0} \frac{1}{2\varepsilon} \int_0^t \mathbb{I}_{[-\varepsilon, \varepsilon]}(\tilde{x}_2(A^{-1}(s))) ds = \\ &= \lim_{\varepsilon \downarrow 0} \frac{1}{2\varepsilon} \int_0^{A^{-1}(t)} \mathbb{I}_{[-\varepsilon, \varepsilon]}(\tilde{x}_2(s)) e^{2\tilde{x}_1(s)} ds = \\ &= \int_0^{A^{-1}(t)} e^{2\tilde{x}_1(s)} dL_{\tilde{x}}^{\tilde{c}_1}(s). \end{aligned}$$

The cases of $k = 2, \dots, n$ can be treated analogously. \square

Thus using (18) we get the stochastic differential equation for the process $(\hat{x}(t))_{t < \eta}$ as follows

$$d\hat{x}(t) = d\hat{w}(t) + \sum_{k=1}^n \gamma_k \tilde{v}_k dL_{\hat{x}}^{\tilde{c}_k}(t). \quad (19)$$

Remark 2. The process $(\hat{x}(t))_{t < \eta}$ is a continuous Markov one as a result of the random change of time for the Markov process $(\tilde{x}(t))_{t \geq 0}$ which is continuous (cf. [2]). So, either $\hat{x}(t)$ is defined for all $t \geq 0$ or

$\overline{\lim}_{t \rightarrow \eta^-} |\hat{x}(t)| = \infty$. It is easy to see that any solution to SDE (19) does not blow up in finite time a.s. Moreover, in the next Section we will show that $\mathbb{E}L_{\hat{x}_2}^{\tilde{c}_k}(t) < \infty$ for each $t \geq 0$ (see (22)). Then, by (19), the trajectories of $\hat{x}(t)$ do not reach infinity in finite time a.s. Thus the process $\hat{x}(t)$ is defined for all $t \geq 0$, i.e. $\mathbb{P}\{\eta = \infty\} = 1$.

3. Ergodic behavior of some Markov chain related to the Brownian motion with membranes.

For $k \in \mathbb{N}$, let $p_k = \frac{1+\gamma_k}{2}, q_k = \frac{1-\gamma_k}{2}$. Recall that $\xi_k = \varphi_{k+1} - \varphi_k$, $\tau_m, m \in \mathbb{N}$, is defined in the proof of Lemma 1. Put $\xi_0 := \xi_n$.

The process $\hat{x}_2(t)$ behaves on (τ_m, τ_{m+1}) , $m = 1, 2, \dots$, as an ordinary skew Brownian motion with skewing at the point $\hat{x}_2(\tau_m) \in \{\varphi_1, \dots, \varphi_n\}$ (Fig. 2). The skew Brownian motion is a strong Markov process as a continuous Feller process (cf. [2], Theorem 3.10). This yields $(\hat{x}_2(\tau_m))_{m \geq 1}$ form a homogeneous Markov chain with the phase space $\{\varphi_1, \dots, \varphi_n\}$ and the transition matrix

$$\begin{bmatrix} 0 & \tilde{p}_1 & 0 & \dots & 0 & \tilde{q}_1 \\ \tilde{q}_2 & 0 & \tilde{p}_2 & 0 & \dots & 0 \\ \dots & \dots & \dots & \dots & \dots & \dots \\ 0 & \dots & 0 & \tilde{q}_{n-1} & 0 & \tilde{p}_{n-1} \\ \tilde{p}_n & 0 & \dots & 0 & \tilde{q}_n & 0 \end{bmatrix},$$

where

$$\begin{aligned} \tilde{p}_k &= \mathbb{P}_{\varphi_k} \{ \hat{x}_2(\sigma_{\varphi_{k-1}} \wedge \sigma_{\varphi_{k+1}}) = \varphi_{k+1} \}, \tilde{q}_k = 1 - \tilde{p}_k, \\ \sigma_a &= \inf\{t \geq 0 : \hat{x}_2(t) = a\}. \end{aligned}$$

One can see (cf. [3], Ch. 3) that, for $k = 1, \dots, n$,

$$\tilde{p}_k = \frac{p_k \xi_{k-1}}{p_k \xi_{k-1} + q_k \xi_k}, \quad (20)$$

$$\tilde{q}_k = \frac{q_k \xi_k}{p_k \xi_{k-1} + q_k \xi_k}. \quad (21)$$

Now we find $M_k := \mathbb{E}_{\varphi_k} L_{\hat{x}_2}^{\varphi_k}(\sigma_{\varphi_{k-1}} \wedge \sigma_{\varphi_{k+1}})$. Equation (19) implies

$$\mathbb{E}_{\varphi_k} \hat{x}_2(\sigma_{\varphi_{k-1}} \wedge \sigma_{\varphi_{k+1}}) = \varphi_k + \gamma_k \mathbb{E}_{\varphi_k} L_{\hat{x}_2}^{\tilde{c}_k}(\sigma_{\varphi_{k-1}} \wedge \sigma_{\varphi_{k+1}}).$$




Figure 2.

Using (20),(21) we get

$$M_k = \frac{\xi_{k-1}\xi_k}{p_k\xi_{k-1} + q_k\xi_k}. \quad (22)$$

3.1. Assume $0 < p_k < 1$, $k = 1, \dots, n$. The Markov chain $(\hat{x}_2(\tau_m))_{m \geq 1}$ is irreducible and all of its states are positive-recurrent. This implies the existence of a unique invariant distribution (π_1, \dots, π_n) which is the solution to the following system of equations

$$\pi_k = \pi_{k-1}\tilde{p}_{k-1} + \pi_{k+1}\tilde{q}_{k+1}, \quad k = 2, \dots, n-1, \quad (23)$$

$$\pi_1 = \pi_n\tilde{p}_n + \pi_2\tilde{q}_2, \quad (24)$$

$$\sum_{k=1}^n \pi_k = 1. \quad (25)$$

Let us find it out. We can rewrite the equation (23) in the form

$$\pi_k(\tilde{p}_k + \tilde{q}_k) = \pi_{k-1}\tilde{p}_{k-1} + \pi_{k+1}\tilde{q}_{k+1}, \quad k = 2, \dots, n-1.$$

This implies the equality

$$\pi_{k+1}\tilde{q}_{k+1} - \pi_k\tilde{p}_k = \pi_k\tilde{q}_k - \pi_{k-1}\tilde{p}_{k-1}$$

valid for $k = 2, \dots, n-1$. Put

$$C := \pi_{k+1}\tilde{q}_{k+1} - \pi_k\tilde{p}_k, \quad k = 1, \dots, n-1. \quad (26)$$

Let $k = 1$. Then from (26) we have

$$\pi_2 = C \frac{1}{\tilde{q}_2} + \pi_1 \frac{\tilde{p}_1}{\tilde{q}_2}. \quad (27)$$

Put $k = 2$ in (26). Taking into account (27) we obtain

$$\pi_3 = C \left(\frac{1}{\tilde{q}_3} + \frac{\tilde{p}_2}{\tilde{q}_2 \tilde{q}_3} \right) + \pi_1 \frac{\tilde{p}_1 \tilde{p}_2}{\tilde{q}_2 \tilde{q}_3}.$$

Finally by recursion we arrive at the formula

$$\pi_k = C \left(\frac{1}{\tilde{q}_k} + \frac{\tilde{p}_{k-1}}{\tilde{q}_{k-1} \tilde{q}_k} + \cdots + \frac{\tilde{p}_2 \cdots \tilde{p}_{k-1}}{\tilde{q}_2 \cdots \tilde{q}_k} \right) + \pi_1 \frac{\tilde{p}_1 \cdots \tilde{p}_{k-1}}{\tilde{q}_2 \cdots \tilde{q}_k}$$

justified for $k = 2, \dots, n$.

Substituting this expression for π_n in the equality

$$\pi_1 \tilde{q}_1 - \pi_n \tilde{p}_n = C,$$

which is a simple consequence of (24), we get

$$\pi_1 = \frac{1 + \tilde{r}_n + \tilde{r}_{n-1} \tilde{r}_n + \cdots + \tilde{r}_2 \cdots \tilde{r}_n}{\tilde{q}_1 (1 - \tilde{r}_1 \cdots \tilde{r}_n)} C, \text{ if } \tilde{p}_1 \cdots \tilde{p}_n \neq \tilde{q}_1 \cdots \tilde{q}_n,$$

where $\tilde{r}_k = \frac{\tilde{p}_k}{\tilde{q}_k}$, $k = 1, 2, \dots, n$. Note that $\tilde{p}_1 \cdots \tilde{p}_n \neq \tilde{q}_1 \cdots \tilde{q}_n$ if and only if $p_1 \cdots p_n \neq q_1 \cdots q_n$. Taking into account that we can chose the ray \tilde{c}_1 in an arbitrary way, we get for $k = 1, 2, \dots, n$

$$\pi_k = \frac{1 + \tilde{r}_{k-1} + \tilde{r}_{k-2} \tilde{r}_{k-1} + \cdots + \tilde{r}_1 \cdots \tilde{r}_{k-1} \tilde{r}_{k+1} \cdots \tilde{r}_n}{\tilde{q}_k (1 - \tilde{r}_1 \cdots \tilde{r}_n)} C,$$

if $p_1 \cdots p_n \neq q_1 \cdots q_n$,

One can find the value of C by substitution of the last formula into (25). Computations lead to the formula

$$\pi_k = C_1 \beta_k, \quad (28)$$

where

$$\beta_k = \frac{\sum_{j=0}^{n-1} q_{k+1} \cdots q_{k+j} p_{k+j+1} \cdots p_{n+k-1} \xi_{k+1} \cdots \xi_{k+j} \xi_{k+j} \cdots \xi_{n+k-2}}{\prod_{i=1, i \neq k}^n (p_i \xi_{i-1} + q_i \xi_i)},$$

$$C_1 = 1 / \sum_{k=1}^n \beta_k.$$

If $p_1 \cdot \dots \cdot p_n = q_1 \cdot \dots \cdot q_n$, it is easy to see that

$$\begin{aligned} \pi_1 &= \frac{1}{1 + \frac{\tilde{p}_1}{\tilde{q}_2} + \frac{\tilde{p}_1 \tilde{p}_2}{\tilde{q}_2 \tilde{q}_3} + \dots + \frac{\tilde{p}_1 \dots \tilde{p}_{n-1}}{\tilde{q}_2 \dots \tilde{q}_n}}, \\ \pi_k &= \pi_1 \frac{\tilde{p}_1 \cdot \dots \cdot \tilde{p}_{k-1}}{\tilde{q}_2 \cdot \dots \cdot \tilde{q}_k}, \quad k = 2, \dots, n. \end{aligned} \tag{29}$$

Now suppose $0 < p_k \leq 1$, $k = 1, \dots, n$, or $0 \leq p_k < 1$, $k = 1, \dots, n$. It is easy to check that formula (28) has no singularities in this case.

Thus we have proved the following statement

Lemma 5. *Let, for $k = 1, \dots, n$, $0 \leq p_k < 1$ or let, for $k = 1, \dots, n$, $0 < p_k \leq 1$. Then there exists a unique invariant distribution $(\pi_k)_{k=1}^n$ of the Markov chain $(\hat{x}_2(\tau_m))_{m \geq 1}$.*

If $p_1 \cdot \dots \cdot p_n \neq q_1 \cdot \dots \cdot q_n$ this invariant distribution is given by formula (28), $k = 1, \dots, n$.

If $p_1 \cdot \dots \cdot p_n = q_1 \cdot \dots \cdot q_n$ the invariant distribution is given by formula (29), $k = 1, \dots, n$.

Remark 3. The method of finding the invariant distribution was proposed by Professor Alexey Kulik.

Now partial cases when some of p_k are equal to 0 and some of them are equal to 1 are left unattended. We consider them in Subsections 3.3, 3.4.

3.2. Further on we make use of the following well-known result. Given a homogeneous Markov chain with state space $\{0, 1, \dots, N\}$ and transition matrix

$$\begin{bmatrix} 0 & 1 & 0 & \dots & \dots & 0 \\ x_1 & 0 & y_1 & 0 & \dots & 0 \\ \dots & \dots & \dots & \dots & \dots & \dots \\ 0 & \dots & 0 & x_{N-1} & 0 & y_{N-1} \\ 0 & \dots & \dots & 0 & 1 & 0 \end{bmatrix},$$

where $0 < x_i < 1$, $0 < y_i < 1$, $i = 1, \dots, N - 1$.

As is known there exists a unique invariant distribution

$$\{f_i\}_{i=0}^N = \{f(x_1, \dots, x_{N-1}, y_1, \dots, y_{N-1}, N)\}_{i=0}^N$$

where

$$f_0 = \frac{1}{1 + \frac{1}{x_1} + \frac{y_1}{x_1 x_2} + \dots + \frac{y_1 \dots y_{N-2}}{x_1 \dots x_{N-1}} + \frac{y_1 \dots y_{N-1}}{x_1 \dots x_{N-1}}}, \quad (30)$$

$$f_1 = \frac{1}{x_1} f_0, \quad (31)$$

$$f_i = \frac{y_1 \dots y_{i-1}}{x_1 \dots x_i} f_0, \quad i = 2, \dots, N-1, \quad (32)$$

$$f_N = y_{N-1} f_{N-1}. \quad (33)$$

3.3. Suppose $p_{k_0} = 1$, $p_{k_1} = \dots = p_{k_l} = 0$, $k_0, k_1, \dots, k_l \in \{1, \dots, n\}$, and $0 < p_k < 1$, $k \in \{1, \dots, n\} \setminus \{k_0, k_1, \dots, k_l\}$. Without loss of generality we may assume that $k_0 < k_1 < \dots < k_l$. Then the Markov chain $(\hat{x}_2(\tau_m))_{m \geq 1}$ has a unique communicating positive-recurrent class $\{\varphi_{k_0}, \varphi_{k_0+1}, \dots, \varphi_{k_1}\}$. All other states are transient. So there exists a unique invariant distribution $(\pi_k)_{k=1}^n$. Besides, it is not difficult to verify that

$$\begin{aligned} \pi_k &= 0, \quad 1 \leq k < k_0 \text{ or } k_1 < k \leq n, \\ \pi_{k_0+i} &= f_i(\tilde{q}_{k_0+1}, \dots, \tilde{q}_{k_1-1}, \tilde{p}_{k_0+1}, \dots, \tilde{p}_{k_1-1}, k_1 - k_0), \\ & \quad i = 0, 1, \dots, k_1 - k_0. \end{aligned}$$

where $f_i, i = 0, 1, \dots, k_1 - k_0$, is defined by formulas (30) – (33).

The case of $p_{k_0} = 0$ and $0 < p_k \leq 1$, $k \in \{1, \dots, n\} \setminus \{k_0\}$ can be treated analogously.

3.4. Suppose there exist $k_1, k_2, \dots, k_{2l} \in \{1, \dots, n\}$, $k_1 < k_2 < \dots < k_{2l}$, such that $p_{k_1} = p_{k_3} = \dots = p_{k_{2l-1}} = 1$, $p_{k_2} = p_{k_4} = \dots = p_{k_{2l}} = 0$, and $0 < p_k < 1$, $k_{2j-1} < k < k_{2j}$, $j = 1, \dots, l$. Then each of the classes $\{\varphi_{k_1}, \varphi_{k_1+1}, \dots, \varphi_{k_2}\}, \dots, \{\varphi_{k_{2l-1}}, \varphi_{k_{2l-1}+1}, \dots, \varphi_{k_{2l}}\}$ is a communicating one for $(\hat{x}_2(\tau_m))_{m \geq 1}$. (Elements of different classes are not communicate.) For each class, there exists a unique invariant distribution which can be found by formulas (30)–(33).

Assume $\hat{x}_2(\tau_1) = \varphi_{k_0}$, $k_{2j} < k_0 < k_{2j+1}$ for some $j \in \{1, \dots, l\}$. ($k_{2l+1} := k_1$). Define

$$\alpha(k_0) = \mathbb{P}_{\varphi_{k_0}} \{ \hat{x}_2(\sigma_{\varphi_{2j}} \wedge \sigma_{\varphi_{2j+1}}) = \varphi_{2j} \}.$$

Note that

$$\mathbb{P}_{\varphi_{k_0}} \{ \sigma_{\varphi_{2j}} \wedge \sigma_{\varphi_{2j+1}} < \infty \} = 1.$$

If $0 < p_k < 1$, $k_{2j} < k < 2j + 1$, computations lead to the formula

$$\alpha(k_0) = \frac{\sum_{i=k_0}^{k_{2j+1}} \frac{\tilde{q}_{k_{2j+1}} \cdots \tilde{q}_i}{\tilde{p}_{k_{2j+1}} \cdots \tilde{p}_i}}{1 + \sum_{i=k_{2j+1}}^{k_{2j+1}} \frac{\tilde{q}_{k_{2j+1}} \cdots \tilde{q}_i}{\tilde{p}_{k_{2j+1}} \cdots \tilde{p}_i}}.$$

If $0 \leq p_k < 1$, $k_{2j} < k < k_{2j+1}$, and equality is reached, $\alpha(k_0) = 1$. Similarly, if $0 < p_k \leq 1$, $k_{2j} < k < k_{2j+1}$, and equality is reached, $\alpha(k_0) = 0$.

If there exist $i_1, \dots, i_s, i_{s+1}, \dots, i_r$ such that $k_{2j} < i_1 < \dots < i_s < k_0 < i_{s+1} < \dots < i_r < k_{2j+1}$, $p_{i_1} = \dots = p_{i_s} = 1$, $p_{i_{s+1}} = \dots = p_{i_r} = 0$. Then

$$\alpha(k_0) = \mathbb{P}_{\varphi_{k_0}} \{ \hat{x}_2(\sigma_{\varphi_s} \wedge \sigma_{\varphi_{s+1}}) = \varphi_s \}.$$

4. The main result.

The behavior of the process $(x(t))_{t \geq 0}$ is uniquely determined by that of the process $(\hat{x}(t))_{t \geq 0}$. As we know (see Remark 2) $\eta = \infty$ a.s. So $x(t) \rightarrow 0$ as $t \rightarrow \zeta -$ if and only if $\hat{x}(t) \rightarrow +\infty$ as $t \rightarrow +\infty$.

Let us obtain an estimate of the local time $L_{\hat{x}}^{\tilde{c}_k}(t)$, $k = 1, \dots, n$. Fix $k \in \{1, \dots, n\}$. Then the functional $L_{\hat{x}}^{\tilde{c}_k}(t)$ increases at those and only those intervals of time $[\tau_m, \tau_{m+1})$, $m \geq 1$, for which $\hat{x}(\tau_m) \in \tilde{c}_k$. Moreover,

$$L_{\hat{x}}^{\tilde{c}_k}(t) = L_{\hat{x}_2}^{\varphi_k}(t) =: \lim_{\varepsilon \downarrow 0} \frac{1}{2\varepsilon} \int_0^t \mathbb{1}_{[\varphi_k - \varepsilon, \varphi_k + \varepsilon]}(\hat{x}_2(s)) ds.$$

Let $\nu_k(t)$ be the number of such intervals on $[0, t]$. According to the strong Law of large numbers

$$\frac{L_{\hat{x}}^{\tilde{c}_k}(t)}{\nu_k(t)} \rightarrow \mathbb{E}_{\varphi_k} L_{\hat{x}_2}^{\varphi_k}(\sigma_{\varphi_{k-1}} \wedge \sigma_{\varphi_{k+1}}) \text{ as } t \rightarrow +\infty \text{ a.s.}$$

Fix $\varepsilon > 0$. Then there exists $t_{k1}(\omega) > 0$ such that for all $t > t_{k1}$ the inequality

$$\nu_k(t) M_k(1 - \varepsilon) \leq L_{\hat{x}}^{\tilde{c}_k}(t) \leq \nu_k(t) M_k(1 + \varepsilon) \quad (34)$$

holds. Recall that $M_k := \mathbb{E}_{\varphi_k} L_{\hat{x}_2}^{\varphi_k}(\sigma_{\varphi_{k-1}} \wedge \sigma_{\varphi_{k+1}})$ and it is defined by formula (22).

Further, by the renewal theorem (cf. [6], Ch. XI),

$$\frac{\mathbb{E}\nu_k(t)}{t} \rightarrow K\pi_k \text{ as } t \rightarrow \infty,$$

where K is some positive constant. Using the strong law of large numbers we get

$$\frac{\nu_k(t)}{t} \rightarrow K\pi_k \text{ as } t \rightarrow \infty \text{ a.s.}$$

Thus there exists $t_{k2} \geq t_{k1}$ such that for all $t > t_{k2}$

$$K\pi_k t(1 - \varepsilon) \leq \nu_k(t) \leq K\pi_k t(1 + \varepsilon) \text{ a.s.} \quad (35)$$

Finally from (34) and (35) we obtain the inequality

$$\begin{aligned} (1 - \varepsilon)^2 K\pi_k t M_k &\leq L_{\hat{x}}^{\tilde{c}_k}(t) \leq \\ &\leq (1 + \varepsilon)^2 K\pi_k t M_k, \quad t > t_{k2}, \text{ a.s.} \end{aligned} \quad (36)$$

We use the inequality (36) to get an estimation of the process $(\hat{x}_1(t))_{t \geq 0}$.

Let $\sum_{k=1}^n \gamma_k \pi_k M_k \tan \theta_k \neq 0$. Integrating (19) over $[0, t]$ and applying (36) we come to the inequality

$$\begin{aligned} \hat{x}_1(0) + \hat{w}_1(t) + Kt \sum_{k=1}^n \gamma_k (1 - \varepsilon_k)^2 \pi_k M_k \tan \theta_k &\leq \\ &\leq \hat{x}_1(t) \leq \\ &\leq \hat{x}_1(0) + \hat{w}_1(t) + Kt \sum_{k=1}^n \gamma_k (1 + \varepsilon_k)^2 \pi_k M_k \tan \theta_k \end{aligned} \quad (37)$$

valid for all $t > t^* = \max_{k=1, \dots, n} \{t_{12}, \dots, t_{n2}\}$, where

$$\varepsilon_k = \begin{cases} \varepsilon & \text{if } \gamma_k \tan \theta_k > 0, \\ -\varepsilon & \text{if } \gamma_k \tan \theta_k < 0. \end{cases}$$

So if $\sum_{k=1}^n \gamma_k \pi_k M_k \tan \theta_k \neq 0$ we can choose ε so small that the expressions

$$\sum_{k=1}^n \gamma_k \pi_k M_k \tan \theta_k,$$

$$\sum_{k=1}^n \gamma_k (1 + \varepsilon_k)^2 \pi_k M_k \tan \theta_k,$$

and

$$\sum_{k=1}^n \gamma_k (1 - \varepsilon_k)^2 \pi_k M_k \tan \theta_k$$

have the same sign. By the law of iterated logarithm there exists $t_* > 0$ such that for all $t > t_*$ the inequality

$$|\hat{w}_1(t)| \leq (1 + \varepsilon) \sqrt{2t \ln \ln t} \quad (38)$$

fulfilled. Combining (37) and (38) we see that

$$\begin{aligned} \hat{x}_1(t) &\rightarrow +\infty \quad \text{as } t \rightarrow +\infty \quad \text{if } \sum_{k=1}^n \gamma_k \pi_k M_k \tan \theta_k > 0 \\ \hat{x}_1(t) &\rightarrow -\infty \quad \text{as } t \rightarrow +\infty \quad \text{if } \sum_{k=1}^n \gamma_k \pi_k M_k \tan \theta_k < 0. \end{aligned}$$

Now we can return from the process $(\hat{x}(t))_{t \geq 0}$ to $(x(t))_{t \geq 0}$ making the inverse change of time and doing the rotation of coordinate system inverse to that defined by (7). Then we see that $x(t) \rightarrow 0$, as $t \rightarrow \zeta^-$, if $\sum_{k=1}^n (p_k - q_k) \pi_k \frac{\xi_{k-1} \xi_k}{p_k \xi_{k-1} + q_k \xi_k} \tan \theta_k > 0$. In this case, as a consequence of (37), there exist $t_0, K_1, K_2 > 0$ such that for all $t > t_0$ the inequality $\hat{x}_1(t) > K_1 + K_2 t$ holds true a.s. This implies convergence of the integral $\int_0^\infty e^{-2\hat{x}_1(s)} ds$. Therefore, $\zeta < \infty$ a.s. Similarly, if $\sum_{k=1}^n (p_k - q_k) \pi_k \frac{\xi_{k-1} \xi_k}{p_k \xi_{k-1} + q_k \xi_k} \tan \theta_k < 0$, then $\zeta = +\infty$ a.s. and $x(t) \rightarrow \infty$ as $t \rightarrow \infty$ a.s.

Let $\sum_{k=1}^n \gamma_k \pi_k M_k \tan \theta_k = 0$. Define

$$\mu_1 = \inf\{t \geq 0 : \hat{x}(t) \in \tilde{c}_1\},$$

and, for $j \geq 2$,

$$\mu_j = \inf\{t > \tilde{\mu}_{j-1} : \hat{x}(t) \in \tilde{c}_1\},$$

where

$$\tilde{\mu}_j = \inf\{t > \mu_j : \hat{x}(t) \in \{\tilde{c}_n, \tilde{c}_2\}\}$$

(see Fig. 3).




Figure 3.

Denote

$$\psi_j = \sum_{k=1}^n \gamma_k \tan \theta_k (L_{\hat{x}}^{\tilde{c}_k}(\mu_{j+1}) - L_{\hat{x}}^{\tilde{c}_k}(\mu_j)), \quad j \geq 1.$$

It is easily seen that if $\sum_{k=1}^n \gamma_k \pi_k M_k \tan \theta_k = 0$ then $\mathbb{E}\psi_j = 0$, $j \geq 1$. From (19) we get

$$\hat{x}_1(\mu_j) = \hat{x}_1(\mu_1) + \sum_{l=1}^j (\hat{w}_1(\mu_{l+1}) - \hat{w}_1(\mu_l) + \psi_l).$$

Note that $\chi_l = \hat{w}_1(\mu_{l+1}) - \hat{w}_1(\mu_l) + \psi_l$, $l \geq 1$, are independent identically distributed random variables (i.i.d.r.v.), $\mathbb{E}\chi_l = 0$, and all the moments are finite. Then by law of iterated logarithm for sum of i.i.d.r.v. (cf. [7]) we have

$$\overline{\lim}_{j \rightarrow \infty} \hat{x}_1(\mu_j) = +\infty, \quad \underline{\lim}_{j \rightarrow \infty} \hat{x}_1(\mu_j) = -\infty \quad \text{a.s.}$$

It is easily seen that in this case the integral $\int_0^\infty e^{-2\hat{x}_1(s)} ds$ diverges a.s. Therefore $\zeta = +\infty$ a.s.

Recall that conventionally $c_{nl+k} = c_k$, $\xi_{nl+k} = \xi_k$ and put additionally $\gamma_{nl+k} = \gamma_k$, $\pi_{nl+k} = \pi_k$, $k = 1, \dots, n$, $l \in \mathbb{N}$.

Definition 1. We call $c_i 0 c_j$, $1 \leq i < j$, a characteristic wedge if $\varphi_i, \varphi_{i+1}, \dots, \varphi_j$ is a communicating class for the Markov chain $(\hat{x}_2(\tau_m))_{m \geq 1}$.

Theorem 1. Let $x(0) = x^0$, $x^0 \neq 0$. Then

- 1) $\mathbb{P}_{x^0}\{\zeta < \infty\} + \mathbb{P}_{x^0}\{\overline{\lim}_{t \rightarrow +\infty} |x(t)| = \infty\} = 1$.
- 2) The probability $\mathbb{P}_{x^0}\{\zeta < \infty\}$ is equal to the probability of hitting some characteristic wedge $c_{j_1}0c_{j_2}$, $1 \leq j_1 < j_2$, for which

$$\sum_{k=j_1}^{k=j_2} (p_k - q_k) \pi_k \frac{\xi_{k-1} \xi_k}{p_k \xi_{k-1} + q_k \xi_k} \tan \theta_k > 0$$

where π_k , $j_1 \leq k \leq j_2$, is defined in Section 3.

In particular, suppose $0 \leq p_k < 1$ or $0 < p_k \leq 1$, $k = 1, \dots, n$. Then, if $\sum_{k=1}^n (p_k - q_k) \pi_k \frac{\xi_{k-1} \xi_k}{p_k \xi_{k-1} + q_k \xi_k} \tan \theta_k > 0$, the process $(x(t))_{t \geq 0}$ hits the origin a.s.; if $\sum_{k=1}^n (p_k - q_k) \pi_k \frac{\xi_{k-1} \xi_k}{p_k \xi_{k-1} + q_k \xi_k} \tan \theta_k < 0$, the process $(x(t))_{t \geq 0}$ does not hit the origin a.s. and $|x(t)| \rightarrow \infty$ as $t \rightarrow +\infty$ a.s.; if $\sum_{k=1}^n (p_k - q_k) \pi_k \frac{\xi_{k-1} \xi_k}{p_k \xi_{k-1} + q_k \xi_k} \tan \theta_k = 0$ the process does not hit the origin a.s. and $\overline{\lim}_{t \rightarrow +\infty} |x(t)| = \infty$ a.s.

Remark 4. It is not difficult to calculate that if $0 \leq p_k < 1, p_1 \cdot \dots \cdot p_n \neq q_1 \cdot \dots \cdot q_n$, the process $(x(t))_{t \geq 0}$ hits the origin if and only if

$$\sum_{k=1}^n \xi_k \sum_{j=1}^n r_{k+1} \cdot \dots \cdot r_{k+j-1} (r_{k+j} - 1) \tan \theta_{k+j} > 0$$

where $r_j = p_j/q_j$, $j = 1, \dots, n$.

5. Examples

Example 1. Let $n = 1$. This is a degenerate case in which the process reaches the origin if and only if $\tan \theta_1 > 0$, i.e. $\theta_1 > 0$.

Example 2. Let $n = 2, \xi_1 = \xi, 0 \leq p_1, p_2 \leq 1$. Then $\pi_1 = \pi_2 = 1/2$ and the process hits the origin if and only if

$$(p_1 - q_1)(p_2 - q_2)\xi(\tan \theta_1 - \tan \theta_2) + 2\pi((p_1 - q_1)q_2 \tan \theta_1 + (p_2 - q_2)p_1 \tan \theta_2) > 0.$$

Consider the particular case when $p_1 = 1, p_2 = 0$. Then the part of the process in the interior of the wedge c_10c_2 is a Brownian motion with instantaneous reflection at the boundary of the wedge (Fig. 4). Let the starting point x^0 be inside the wedge and away from the origin. In this case the process hits the origin if and only if




Figure 4.

$\tan \theta_1 - \tan \theta_2 > 0$, i.e. $\theta_1 - \theta_2 > 0$. Thus the result we obtained coincides with that of [1].

Example 3. Let $n = 3, 0 < p_k \leq 1$ or $0 \leq p_k \leq 1, k = 1, 2, 3$. If $p_1 p_2 p_3 \neq q_1 q_2 q_3$, the invariant distribution has the form

$$\begin{aligned}\pi_1 &= \frac{\tilde{q}_2 + \tilde{p}_2 \tilde{p}_3}{D}, \\ \pi_2 &= \frac{\tilde{q}_3 + \tilde{p}_1 \tilde{p}_3}{D}, \\ \pi_3 &= \frac{\tilde{q}_1 + \tilde{p}_1 \tilde{p}_2}{D},\end{aligned}$$

where $D = \tilde{q}_1 + \tilde{q}_2 + \tilde{q}_3 + \tilde{p}_1 \tilde{p}_2 + \tilde{p}_2 \tilde{p}_3 + \tilde{p}_1 \tilde{p}_3$. In this case the process hits the origin if and only if

$$\begin{aligned}(p_1 - q_1) \tan \theta_1 (q_2 p_3 \xi_2 + q_2 q_3 \xi_3 + p_2 p_3 \xi_1) + \\ + (p_2 - q_2) \tan \theta_2 (p_1 q_3 \xi_3 + q_1 q_3 \xi_1 + p_1 p_3 \xi_2) + \\ + (p_3 - q_3) \tan \theta_3 (q_1 p_2 \xi_1 + q_1 q_2 \xi_2 + p_1 p_2 \xi_3) > 0.\end{aligned}$$

If $p_1 p_2 p_3 = q_1 q_2 q_3$ we get

$$\begin{aligned}\pi_1 &= \frac{\tilde{q}_2 \tilde{q}_3}{\tilde{q}_2 \tilde{q}_3 + \tilde{p}_1 \tilde{q}_3 + \tilde{p}_1 \tilde{p}_2}, \\ \pi_2 &= \frac{\tilde{p}_1 \tilde{q}_3}{\tilde{q}_2 \tilde{q}_3 + \tilde{p}_1 \tilde{q}_3 + \tilde{p}_1 \tilde{p}_2}, \\ \pi_3 &= \frac{\tilde{p}_1 \tilde{p}_2}{\tilde{q}_2 \tilde{q}_3 + \tilde{p}_1 \tilde{q}_3 + \tilde{p}_1 \tilde{p}_2}\end{aligned}$$

and the process hits the origin if and only if

$$q_2 q_3 (p_1 - q_1) \tan \theta_1 + p_1 q_3 (p_2 - q_2) \tan \theta_2 + p_1 p_2 (p_3 - q_3) \tan \theta_3 > 0.$$

References

- [1] Varadhan S.R.S., Williams R.J., *Brownian motion in a wedge with oblique reflection*, Comm. Pure Appl. Math. 38, 1985, pp. 405–443.
- [2] Dynkin E.B. *Markov processes*, Fizmatgiz, Moscow, 1963; English transl., vol I, II, New York: Acad.Press., Berlin: Springer, 1965.
- [3] Portenko N.I., *Generalized diffusion processes*, Naukova Dumka, Kiev, 1982; English transl. Amer. Math. Soc., Providence, RI, 1990
- [4] Harrison J.M., Shepp L.A., *On skew brownian motion*, Annals Prob., 9, 1981, pp. 309–313.
- [5] Ikeda N., Watanabe S., *Stochastic differential equations and diffusion processes*, North-Holland, Amsterdam (Kodansha Ltd., Tokyo), 1981.
- [6] Feller W., *An introduction to probability theory and its applications, Vol.II*, J. Wiley & Sons, New York, 1971.
- [7] Petrov V.V., *Sum of Independent Random Variables*, Springer, Berlin, 1975.
- [8] Aryasova O.V. *Diffusion in a plane with semipermeable membranes on two straight lines*, Nonlinear Oscillations 2, 1999, pp. 439-447.

Institute of Geophysics, National Academy of Sciences of Ukraine,
Palladina pr. 32, 03680, Kiev-142, Ukraine,
oaryasova@mail.ru

Institute of Mathematics, National Academy of Sciences of Ukraine,
Tereshchenkivska str. 3, 01601, Kiev, Ukraine
apilip@imath.kiev.ua