

Quasitoric Manifolds with Invariant Almost Complex Structure

Andrei Kustarev*

April 20, 2019

Abstract

We prove that any quasitoric manifold M^{2n} admits a T^n -invariant almost complex structure if and only if M^{2n} admits a positive omniorientation. In particular, we show that all obstructions to existence of T^n -invariant almost complex structure on M^{2n} arise from cohomology of underlying polytope - and hence are trivial.

1 Introduction

Let M^{2n} be an oriented closed compact manifold. We say that M^{2n} is *quasitoric* over simple polytope P ([6],[10]) if:

1. there is locally standart action of torus T^n on M^{2n} and
2. there exists a map $\pi : M^{2n} \rightarrow P^n$ which is a superposition of projection $M^{2n} \rightarrow M^{2n}/T^n$ and diffeomorphism of manifolds with corners $M^{2n}/T^n \rightarrow P$.

Any smooth projective toric variety gives the example of quasitoric manifold, but not vice versa. Any quasitoric manifold admits a canonical smooth structure ([11]).

Submanifolds of the form $\pi^{-1}(F_j)$, $j = 1 \dots m$, where F_j is a codimension one facet, are called *characteristic submanifolds*. An *omniorientation* of M^{2n} is the orientation of M^{2n} and all of its characteristic submanifolds. Every omniorientation determines a canonical T^n -invariant weakly complex structure on M^{2n} ([11]), which may also be constructed in terms of *characteristic map* $\lambda : \mathbb{Z}^m \rightarrow \mathbb{Z}^n$.

*Dep. of Mechanics and Mathematics, MSU. E-mail: kustarev@gmail.com

Every smooth projective toric variety admits canonical T^n -invariant complex structure. One may ask the following question ([6], Prob. 7.6): find the criterion for existence of T^n -invariant almost complex structure on quasitoric manifold M^{2n} in terms of λ . In this paper we present the solution of this problem.

T^n -invariant weakly complex structure on manifold M^{2n} with torus action α is an isomorphism c_τ of bundle $\tau(M^{2n}) \oplus \mathbb{C}^{m-n}$ with complex vector bundle ξ . The map

$$\xi \xrightarrow{c_\tau^{-1}} \tau(M^{2n}) \oplus \mathbb{C}^{m-n} \xrightarrow{d\alpha(t) \oplus I} \tau(M^{2n}) \oplus \mathbb{C}^{m-n} \xrightarrow{c_\tau} \xi$$

is complex-linear for every t .

If v is a fixed point for α , then there is a map $\tau_v(M^{2n}) \xrightarrow{i} \xi_v \xrightarrow{\pi'} \mathbb{C}^n$, where i is an injection and π' is a projection map along fixed subspace $\mathbb{C}^{m-n} \subset \xi_v$. The *sign* of v is the sign of determinant of the map $\pi' \circ i$.

An equivalent definition of sign is as follows: we set $\text{sign}(v) = +1$, if the orientations of $\tau(M)_v$ determined by orientation of M and orientations of characteristic submanifolds coincide, and $\text{sign}(v) = -1$, if not.

We call omniorientation *positive*, if signs of all fixed points are positive.

Theorem 1. Quasitoric manifold M^{2n} admits a T^n -invariant almost complex structure if and only if it admits a positive omniorientation.

2 Remarks and examples

As follows from [8], quasitoric manifold M with positive omniorientation and corresponding weakly complex structure J satisfies $c_n(J) = \chi(M)$. So by [2], the structure J is equivalent to some almost complex structure on M , not necessarily T^n -invariant.

Every smooth projective toric variety M admits canonical T^n -invariant complex structure and symplectic form. So by [4], there's a *moment map* $\pi : M \rightarrow P$, where P is a simple convex polytope. Since all characteristic submanifolds of M are complex subvarieties, M has positive omniorientation.

Note that in real dimension 6 there exist examples of non-projective smooth toric varieties (see [10]). Nevertheless, the quotient by the action of T^3 is still isomorphic to simple convex polytope.

Now we turn to examples of non-algebraic quasitoric manifolds. Denote by $\mathbb{C}P_k^2$ a connected sum of k copies of $\mathbb{C}P^2$ with standart orientation and smooth structure. As follows from equivariant connected sum constuction

([11]), all $\mathbb{C}P_k^2$'s are quasitoric. Also, if k is odd, then $\mathbb{C}P_k^2$ admits a positive omniorientation. If k is even, then $\mathbb{C}P_k^2$ doesn't admit an almost complex structure.

Theorem 2.1 ([1]). Let M^4 be an oriented smooth manifold with no boundary. Then M admits an almost complex structure if and only if the number $td(M) = \frac{1}{4}(\chi(M) + \text{sign}(M))$ is integer. Here $\text{sign}(M) = (b_2^+(M) - b_2^-(M))$ is the signature of a cohomology intersection form on M .

We have $td(\mathbb{C}P_k^2) = \frac{k+1}{2}$ and this implies non-existence of almost complex structure on $\mathbb{C}P_k^2$ for even k .

From Theorem 1 we obtain that the manifold $\mathbb{C}P_k^2$ admits T^2 -invariant almost complex structure, if k is odd. But if $k \geq 3$, $\mathbb{C}P_k^2$ can't be a toric variety since all toric varieties are rational and their Todd genus is equal to 1.

As follows from [9], the manifold $\mathbb{C}P_k^2$ doesn't admit T^2 -invariant symplectic structure, if $k \geq 2$.

Theorem 2.2 ([9]). Let M be a manifold equipped with a symplectic circle action with only isolated fixed points. Then $td(M) = 1$ if the action is Hamiltonian, and $td(M) = 0$ if not.

The proof of following result is based on Seiberg-Witten theory of four-dimensional manifolds.

Theorem 2.3 ([7]). The manifold $\mathbb{C}P_k^2$ doesn't admit a symplectic structure, if $k \geq 2$.

Hence, manifolds $\mathbb{C}P_k^2$ with $k \geq 3$, k odd, are neither symplectic manifolds nor toric varieties – but they admit T^2 -invariant almost complex structure by Theorem 1.

3 Proof of Theorem 1

The “only if” part of Theorem 1 is obvious. If J is an invariant almost complex structure on M , then all characteristic submanifolds of M are J -invariant and all signs of fixed points are positive. Therefore, it remains to show that existence of positive omniorientation implies the existence of invariant almost complex structure.

3.1 Notations

Denote by $sk_i(P)$ the set of all i -dimensional facets of P . Let $\iota : P \rightarrow M$ be an embedding of P to M satisfying the following conditions:

1. $\pi \circ \iota = id$;
2. restriction of ι on $Int G$ is smooth for any facet $G \subset P$.

An example of such ι is given by composition $P \rightarrow P \times T^n \rightarrow M$, where the last map is factorization map ([10]).

We assume that M admits T^n -invariant metric g (see [3]). Then our problem is equivalent to constructing an operator J on $\tau(M)$ with following properties:

1. $J^2 = -1$;
2. J commutes with $d\alpha(t)$ for every $t \in T^n$;
3. J is orthogonal with respect to the metric g .

Recall that the space of orthogonal complex structures on oriented vector space \mathbb{R}^{2i} is homeomorphic to $SO(2i)/U(i)$.

The words “structure on X ” will mean “ T^n -invariant complex structure on vector bundle $\tau(M)$ restricted on X ”.

If $M_G = \pi^{-1}(G)$ is a $2i$ -dimensional quasitoric submanifold in M , then we denote by $\xi_1 \dots \xi_{n-i}$ two-dimensional vector bundles over M corresponding to $(n-i)$ one-dimensional stationary toric subgroups $T_1 \dots T_{n-i}$ of M_G . The fibers of $\xi_1 \dots \xi_{n-i}$ over M_G are J -invariant since J commutes with $d\alpha$.

The structure J on $\pi^{-1}(Int G)$ is called *positive* with respect to omniorientation o if it agrees with orientations of ξ_1, \dots, ξ_{n-i} determined by o .

3.2 Positivity and one-dimensional facets

Lemma 3.1 If $M_G \subset M$ is a quasitoric submanifold, then $\xi_j \perp M_G$ and $\xi_j \perp \xi_k$, if $j \neq k$,

□ Let $v \in \tau(M_G)$, $v_j \in \xi_j$ and $v_k \in \xi_k$ be nonzero vectors in the point $x \in M_G$, and $t_\pi \in T_j$ be an element of toric subgroup T_j corresponding to multiplication by -1 . Since T_j acts trivially on ξ_k and $\tau(M_G)$, we have $g(v, v_j) = g(v, t_\pi v_j) = g(v, -v_j) = 0$ and $g(v_j, v_k) = g(t_\pi v_j, v_k) = g(-v_j, v_k) = 0$. □

Let o be any omniorientation of M , not necessarily positive. Then o determines structure J on $\pi^{-1}(sk_0(P))$ in the following way: J is rotation

by angle $\pi/2$ in the direction specified by o on every fiber $\xi_1|_x, \dots, \xi_n|_x$, where $x \in \pi^{-1}(sk_0(P))$.

Proposition 3.2. Let J be a structure on $\pi^{-1}(sk_0(P))$ determined by o . Then J may be extended over $\pi^{-1}(sk_1(P))$ if and only if o is positive.

□ Let us prove the “if” part first. Consider $I \subset P$ – an edge connecting vertices x_0 and x_1 , $T_1 \dots T_{n-1}$ – toric subgroups corresponding to I . Then $\tau(M) \simeq \tau(\pi^{-1}(I)) \oplus \xi_1 \oplus \dots \oplus \xi_{n-1}$ over $\pi^{-1}(I)$. The orientation of bundles $\xi_1 \dots \xi_{n-1}$ over $\pi^{-1}(I)$ is determined by o .

Denote by W_0 and W_1 orthogonal complements to $\xi_1 \dots \xi_{n-1}$ at the points $\pi^{-1}(x_0)$ and $\pi^{-1}(x_1)$ respectively. Since J was constructed by o , J is a rotation by $\pi/2$ in spaces W_0 and W_1 . Note that W_0 and W_1 are tangent spaces to $\pi^{-1}(I)$ at $\pi^{-1}(x_0)$ and $\pi^{-1}(x_1)$. The positivity of o implies that orientations of W_0 and W_1 agree as orientations of $\tau(\pi^{-1}(I)) = \tau(S^2)$. So we can define J on $\tau(\pi^{-1}(I))$ as rotation by $\pi/2$ in the direction specified by orientations of W_0 and W_1 . On the bundles ξ_1, \dots, ξ_{n-1} structure J is defined as rotation by $\pi/2$ in the direction specified by o .

The “only if” part of Prop.3.2 is proved in similar fashion. If J is defined on entire $\pi^{-1}(I)$, then the orientations of M at $\pi^{-1}(x_0)$ and $\pi^{-1}(x_1)$ agree – and that holds for every pair of adjacent vertices (x_0, x_1) . □

3.3 Triviality of higher obstructions

Now we consider the case $i > 1$. We suppose that J is defined on $\pi^{-1}(sk_{i-1}(P))$ and try to extend it over $\pi^{-1}(sk_i(P))$.

Lemma 3.3. Let $G \subset P$ be an i -dimensional facet. The space of positive structures J over $\pi^{-1}(Int G)$ is homeomorphic to space of continuous mappings $\text{Map}(Int G, SO(2i)/U(i))$.

□ The positivity of J implies that J is uniquely defined on bundle $\tau(\pi^{-1}(Int G))^\perp \simeq \xi_1 \oplus \dots \oplus \xi_{n-i}$. So it suffices to define J on the tangent vector bundle $\tau(\pi^{-1}(Int G)) \simeq \tau(M_G)|_{Int G}$.

Denote by T^i any i -dimensional toric subgroup complementary to T^{n-i} in T^n . Then T^i acts freely on $\pi^{-1}(Int G)$ and the quotient by the action is homeomorphic to $Int G$ itself. Since J is T^n -invariant structure, it suffices to define J only on $\iota(Int G)$. The bundle $\tau(M_G)$ is trivial oriented $2i$ -dimensional real vector bundle over $\iota(Int G)$ – and so J may be chosen in an arbitrary way over it (the only condition is that J must be orthogonal with respect to the metric g). □

Note that the bundle $\tau(M_G)$ over $\iota(G)$ is also trivial. Since J is already defined over $\pi^{-1}(sk_{i-1}(P))$ and $\iota(\partial G) \subset \pi^{-1}(sk_{i-1}(P))$, there is a continuous map $\partial G \simeq S^{i-1} \rightarrow SO(2i)/U(i)$ defined by J . We denote by C_G the corresponding homotopy class. Recall that the polytope P and all of its facets are oriented.

We summarize all this in following statement.

Lemma 3.4. Let J be a positive structure on $\pi^{-1}(sk_{i-1}(P))$. Then J may be extended over $\pi^{-1}(sk_{i-1}(P)) \cup M_G$ if and only if $C_G = 0$ in the group $\pi_{i-1}(SO(2i)/U(i))$.

We define cochain $\sigma_J^i \in C^i(P, \pi_{i-1}(SO(2i)/U(i)))$ by the rule $\sigma_J^i(G) = C_G$. By definition, σ_J^i is zero if and only if J may be extended from $\pi^{-1}(sk_{i-1}(P))$ to $\pi^{-1}(sk_i(P))$. Our next aim is to prove that σ_J^i is a cocycle.

Lemma 3.5. Suppose that J is a positive structure defined on $\pi^{-1}(sk_{i-1}(P))$, Q is an $(i+1)$ -dimensional facet of P . Then

$$\sum_{G_j \subset \partial Q} \sigma_J^i(G_j) = 0.$$

□ We'll need some preliminary results before the proof.

There is a canonical embedding of spaces of complex structures $c : SO(2i)/U(i) \rightarrow SO(2i+2)/U(i+1)$, defined by formula $J \rightarrow J \oplus t_{\pi/2}$, where $t_{\pi/2}$ is rotation by $\pi/2$ in added two-dimensional subspace.

Lemma 3.6. The map

$$c_* : \pi_j(SO(2i)/U(i)) \rightarrow \pi_j(SO(2i+2)/U(i+1))$$

is an isomorphism for every $j \leq 2i-2$.

□ The embedding c may be viewed as the composition

$$SO(2i)/U(i) \rightarrow SO(2i+2)/U(i) \rightarrow SO(2i+2)/U(i+1),$$

where first map is an embedding of fiber to bundle space over $SO(2i+2)/SO(2i)$ and the second map is a projection map of bundle space with fiber S^{2i+1} . An exact homotopy sequence now implies that c induces isomorphism of corresponding homotopy groups up to dimension $2i-2$. □

If $i > 0$, then $i-1 \leq 2i-2$ and $c_* : \pi_{i-1}(SO(2i)/U(i)) \rightarrow \pi_{i-1}(SO(2i+2)/U(i+1))$ is an isomorphism.

Now we can prove Lemma 3.5. It suffices to show that $\sum_{j=1}^k c_* \sigma_J^i(G_j) = 0$ in the group $\pi_{i-1}(SO(2i+2)/U(i+1))$.

We can say nothing on the sum of $\sigma_J^i(G_j)$'s itself, since the trivializations of bundles $\tau(\pi^{-1}(G_j))$ over different $\iota(G_j)$'s don't agree.

But the structure J on $\tau(\pi^{-1}(G_j))|_{\iota(G_j)}$ automatically defines structure $c(J)$ on $\tau(\pi^{-1}(Q))|_{\iota(G_j)}$ by the formula $J \rightarrow J \oplus t_{\pi/2}$. The bundle $\tau(\pi^{-1}(Q))$ is trivial over Q . Since $\iota(G_j) \subset \iota(Q)$ for every j , the map $f_J : Q \cap sk_{i-1}(P) \rightarrow SO(2i+2)/U(i+1)$ such that $f_J|_{\partial G_j} = c_* \sigma_J^i(G_j)$ is well-defined.

Let U_ε be closed i -dimensional tubular neighbourhood of $Q \cap sk_{i-1}(P)$ in ∂Q (such that $G_j \cap U_\varepsilon \neq \emptyset$ for every j). The set $Q \cap sk_{i-1}(P)$ is a deformational retract of U_ε , and the retraction determines a map $\tilde{f}_J : U_\varepsilon \rightarrow SO(2i+2)/U(i+1)$. The space U_ε is homeomorphic to i -dimensional sphere with k holes; let $S_1 \subset G_1, \dots, S_k \subset G_k$ be the boundary spheres of the holes. Then any homotopy class $\tilde{f}_J|_{S_j}$ coincides with $c_* \sigma_J^i(G_j)$. Since the sum of spheroids $\tilde{f}_J|_{S_j}$ must be zero in $\pi_{i-1}(SO(2i+2)/U(i+1))$, this completes the proof. \square

Lemma 3.7. Suppose that J is a positive structure on $\pi^{-1}(sk_{i-1}(P))$ and $\sigma_J^i = \partial\beta$. Then one can change J on $\pi^{-1}(sk_{i-1}(P) \setminus sk_{i-2}(P))$ and obtain new positive structure J' on $\pi^{-1}(sk_{i-1}(P))$ such that $\sigma_{J'}^i = 0$.

\square Let H be some $(i-1)$ -dimensional facet. Then the space of positive structures on $\pi^{-1}(Int H)$ is homeomorphic to $Int H \rightarrow SO(2i-2)/U(i-1)$ by Lemma 3.3. Denote by f_H the map $f_H : Int H \rightarrow SO(2i-2)/U(i-1)$ corresponding to J .

By Lemma 3.6 the map $c : SO(2i-2)/U(i-1) \rightarrow SO(2i)/U(i)$ induces an isomorphism of homotopy groups up to dimension $2i-4$. If $i > 2$, then $i-1 \leq 2i-4$, and if $i=2$, then $SO(4)/U(2)$ and $SO(2)/U(1)$ are simply connected – so the map $c_* : \pi_{i-1}(SO(2i-2)/U(i-1)) \rightarrow \pi_{i-1}(SO(2i)/U(i))$ is an isomorphism for $i > 1$.

Let us identify $Int H$ with open i -dimensional ball in \mathbb{R}^i of radius 1. Consider the map $\tilde{f}_H : Int H \rightarrow SO(2i-2)/U(i-1)$ satisfying following conditions:

- $\tilde{f}_H(x) = f_H((2|x| - 1) \cdot x)$, if $1/2 \leq |x| \leq 1$;
- the homotopy class of spheroid $\tilde{f}_H|_{\{|x| \leq 1/2\}}$ is $(-c_*^{-1}\beta(H))$.

If we replace f_H with \tilde{f}_H , we'll obtain new structure J' on $\pi^{-1}(sk_i(P))$. Then $\sigma_{J'}^i(G) = \sigma_J^i(G) - \beta(H)$ for every i -dimensional facet $G, H \subset G$. So if

we replace f_H with \tilde{f}_H for every $(i-1)$ -dimensional facet $H \subset P$, we'll have $\sigma_{J'}^i = 0$, since $\sigma_J^i = \partial\beta$. \square

3.4 Proof of the main theorem

Now we can prove Theorem 1. Proposition 3.2 and the positivity of o guarantee that J may be extended from $\pi^{-1}(sk_0(P))$ to $\pi^{-1}(sk_1(P))$. From Lemmas 3.4 and 3.5 it follows that for $i > 1$ obstruction to extending J from $\pi^{-1}(sk_{i-1}(P))$ to $\pi^{-1}(sk_i(P))$ is a cocycle $\sigma_J^i \in C^i(P, \pi_{i-1}(SO(2i)/U(i)))$. Since P has trivial homology, σ_J^i is a coboundary, so by Lemma 3.7, there exists structure J' on $\pi^{-1}(sk_{i-1}(P))$ that may be extended to $\pi^{-1}(sk_i(P))$.

4 Acknowledgements

The author is grateful to his advisor, Prof.V.M.Buchstaber, and to T.E.Panov for constant attention to this work, and also to N.E.Dobrinskaya for useful e-mail correspondence. The statement of Theorem 1 was proved earlier by N.E.Dobrinskaya for $n \leq 7$.

References

- [1] W.-T. Wu. Sur le classes caractéristique des structures fibrées sphériques // Actualites Sci. Industr. 1183 (1952).
- [2] E. Thomas. Complex Structures on Real Vector Bundles // Amer. J. Math. 1967. V.89. P.887-908.
- [3] Glen E. Bredon. Introduction to Compact Transformation Groups. Pure and Applied Math. 46, Academic Press, 1972.
- [4] M. F. Atiyah. Convexity and Commuting Hamiltonians // Bull. London Math. Soc. 1982. V.14, N1. P. 1-15.
- [5] V. Guilemin, S. Sternberg. Convexity properties of the moment mapping // Inv. Math. 67 (1982), N3, P. 491-513.
- [6] M. Davis, T. Januskiewicz. Convex Polytopes, Coxeter Orbifolds and Torus Actions // Duke Math J. 1991. V.62 N2 P. 417-451.
- [7] C.H.Taubes. The Seiberg-Witten Invariants and Symplectic Forms // Math. Res. Lett. 1994. V.1. P.809-822.
- [8] T. E. Panov. Hirzebruch Genera of Manifolds with Torus Action // Izv. RAN. Ser. Mat., 65:3 (2001), 123138
- [9] K.E.Feldman. Hirzebruch Genera of Manifolds Equipped with a Hamiltonian Circle Action. arXiv:math/0110028v2
- [10] Victor M. Buchstaber, Taras E. Panov. Torus Actions and Their Applications in Topology and Combinatorics. AMS Bookstore, 2002.
- [11] V. Buchstaber, T. Panov, N. Ray. Spaces of Polytopes and Cobordism of Quasitoric Manifolds // Moscow Math. J (2007) V.7 N2.