

PERVERSE COHERENT SHEAVES

DMITRY ARINKIN, ROMAN BEZRUKAVNIKOV

ABSTRACT. This note introduces an analogue of perverse t -structure [1] on the derived category of coherent sheaves on an algebraic stack (subject to some mild technical conditions). Under additional assumptions construction of coherent “intersection cohomology” sheaves is given. Those latter assumptions are rather restrictive but hold in some examples of interest in representation theory.

Similar results were obtained by Deligne (unpublished), Gabber [10] and Kashiwara [13].

To Pierre Deligne with admiration.

1. INTRODUCTION

Let X be a reasonable stratified topological space; or let X be a reasonable scheme, stratified by locally closed subschemes. Let D be the full subcategory in, respectively, derived category of sheaves on X , or in the derived category of étale sheaves on X , consisting of complexes smooth along the stratification.

For an integer-valued function p (perversity) on the set of strata Beilinson, Bernstein, and Deligne [1] defined a t -structure on the category D ; the objects of corresponding abelian category (core of the t -structure) are called perverse sheaves.

The question addressed in this note is whether an analogous construction can be carried out for the derived category of coherent sheaves on a reasonable scheme. Surprisingly, the answer is positive (with some modifications), easy, and not widely known (although it was known to Deligne for a long time, see [8]).

Let us summarize the difference between the coherent case considered here, and the constructible case treated in [1].

First, in the coherent case we can not work with complexes “smooth” along a given stratification, because the corresponding subcategory in the derived category is not a full triangulated subcategory. (If f is a function whose divisor intersects the open stratum, then the cone of the morphism $\mathcal{O} \xrightarrow{f} \mathcal{O}$ has singularity on the open stratum). This forces us to define perversity as a function on the set of generic points of all irreducible subvarieties, i.e. on the topological space of a scheme.

The second, more essential difference is that in the derived category of coherent sheaves the functor j^* of pull-back under an open embedding j does not have adjoint functors. Recall that in constructible situation the right adjoint to j^* is the functor j_* of direct image, and the left adjoint is the functor $j_!$ of extension by zero. In coherent set-up the functor j_* is defined in the larger category of quasi-coherent sheaves (Ind-coherent sheaves), while $j_!$ is defined in the Grothendieck dual category (consisting of Pro-coherent sheaves) introduced in Deligne’s appendix to [12].

It turns out, however, that in the proof of the existence of perverse t -structure one can use instead of the object $j_!(\mathcal{F})$ (where $j : U \hookrightarrow X$ is an open embedding)

any extension $\tilde{\mathcal{F}}$ of $\mathcal{F}$ to X such that the restriction of $\tilde{\mathcal{F}}$ to $X - U$ has no cohomology above certain degree (depending on the perversity function). If the perversity function is *monotone* (see Definition 3.9 below) it is very easy to construct such $\tilde{\mathcal{F}}$. Applying the Grothendieck-Serre duality to this construction, we get a substitute for $j_*(\mathcal{F})$, which exists if the perversity function is comonotone. Otherwise the proof is parallel to that in [1].

Thus the t -structure is constructed not for an arbitrary perversity function, but only for a monotone and comonotone one. (In the topological situation one also needs this condition to get a t -structure on the whole derived category of constructible sheaves rather than on the category corresponding to a fixed stratification.)

In [8] Deligne used the Grothendieck's Finiteness Theorem ([11], VIII.2.1) to show that the formulas for $\tau_{\leq 0}^p, \tau_{\geq 0}^p$ of [1], *a priori* making sense in a larger category containing $D^b(\text{Coh})$, give in fact objects of $D^b(\text{Coh})$, provided the perversity function is monotone and comonotone (see also Remark 3.13).

The results on the existence of a “perverse” t -structure carry over to the case of G -equivariant coherent sheaves, where G is a (reasonable) algebraic group acting on a (reasonable) scheme X . In this case perversity $p(x)$ is assigned only to orbits of G (including “generic orbits”), since an equivariant sheaf is automatically smooth along the orbits.

More generally, the perverse t -structure can be constructed for coherent sheaves on a (reasonable) algebraic stack $\mathcal{X}$, given a perversity defined on points of $\mathcal{X}$. The case of G -equivariant coherent sheaves on X corresponds to working with the stack $\mathcal{X} = X/G$.

The general formalism in all three situations: sheaves on a scheme, equivariant sheaves on a scheme, and sheaves on a stack — is very similar, to the extent that we found it easier to work with algebraic stacks and treat the other situations as special cases. However, one construction does not apply to (non-equivariant) sheaves on a scheme. Namely, the definition of the minimal (Goresky-MacPherson, or IC) extension functor $j_{!*}$ requires a *strictly* monotone and comonotone perversity. Such perversity exists only if the dimensions of adjacent points differ by at least two, which excludes schemes (other than finite schemes). On the other hand, in equivariant settings, it is possible that the dimensions of adjacent orbits differ by at least two, and a strictly monotone and comonotone perversity exists.

If the perversity is strictly monotone and comonotone, an analogue of the usual description of irreducible perverse sheaves as minimal extensions of local systems is valid, and the core of the perverse t -structure is Artinian and Noetherian (in contrast with the core of the standard t -structure). Some examples of this situation are given at the end of the paper.

A version of the main result in a restricted generality appeared in the preprint [3] by the second author. Later related constructions were published by Gabber [10] and Kashiwara [13]. A similar result was known to Deligne [8] long before the date of [3].

Organization. In Section 2, we prove some basic properties of coherent sheaves, and study dualizing complexes on stacks.

Section 3 contains the definition of the perverse t -structure on a stack. The main result of this section (Theorem 3.10) verifies axioms of a t -structure.

In Section 4, we define the minimal extension functor (Theorem 4.3). We then use it to study irreducible perverse coherent sheaves (Proposition 4.11), and prove that the category of perverse coherent sheaves is Artinian and Noetherian (Corollary 4.13). As we already mentioned, these results require additional assumptions; in particular, all results are empty in the case of (non-equivariant) coherent sheaves on a scheme.

Acknowledgements. We are much obliged to Pierre Deligne for valuable explanations, comments on the text and a kind permission to use his unpublished results.

We thank Alexander Beilinson, Victor Ginzburg, and Dmitry Panyushev for discussions and references.

This paper grew out of the preprint [3] which was started in the spring of 1999 when the second author was a member at the Institute for Advanced Study. He thanks IAS for excellent work conditions and Leonid Positselski for his participation in the early stages of the work on [3].

The first author is a Sloan Research Fellow, and he is grateful to the Alfred P. Sloan Foundation for the support. The work of the second author was partly supported by DARPA grant HR0011-04-1-0031 and NSF grant DMS-0625234.

2. PRELIMINARIES

In this section we collect some results needed in the exposition. The reader familiar with algebraic geometry should feel free to skip our proofs.

2.1. Quasi-coherent sheaves on stacks. Let $\mathcal{X}$ be a Noetherian algebraic stack. In particular, $\mathcal{X}$ is quasi-compact. Assume also that $\mathcal{X}$ is semi-separated: that is, the diagonal morphism $\mathcal{X} \rightarrow \mathcal{X} \times \mathcal{X}$ is affine. Let us fix a presentation of $\mathcal{X}$, that is, a surjective smooth morphism $\pi : X \rightarrow \mathcal{X}$, where X is an algebraic space.

Lemma 2.1. *Let $\mathcal{X}$ be an algebraic stack. The following conditions are equivalent:*

- (a) $\mathcal{X}$ is quasi-compact and semi-separated.
- (b) $\mathcal{X}$ admits a presentation $\pi : X \rightarrow \mathcal{X}$ such that X is an affine scheme and π is an affine morphism.
- (c) $\mathcal{X}$ admits a presentation $\pi : X \rightarrow \mathcal{X}$ such that X is a quasi-compact semi-separated algebraic space and π is an affine morphism.

Proof. (a) $\Rightarrow$ (b). Let $\pi : X \rightarrow \mathcal{X}$ be a presentation of $\mathcal{X}$. Note that X is quasi-compact, so passing to its étale cover, we may assume that X is an affine scheme. But then

$$X \times_{\mathcal{X}} X = (X \times X) \times_{(X \times X)} \mathcal{X}$$

is an affine scheme; therefore, π is an affine morphism.

(b) $\Rightarrow$ (c) is obvious.

(c) $\Rightarrow$ (a). Clearly, $\mathcal{X}$ is quasi-compact; let us prove it is semi-separated. The morphism

$$X \times_{\mathcal{X}} X \rightarrow X$$

is affine, because it is obtained from π by a base change. Since X is semi-separated, so is $X \times_{\mathcal{X}} X$. Therefore, the composition

$$X \times_{\mathcal{X}} X \rightarrow (X \times_{\mathcal{X}} X) \times (X \times_{\mathcal{X}} X) \rightarrow X \times X$$

is affine. It remains to notice that the composition is obtained from the diagonal $\mathcal{X} \rightarrow \mathcal{X} \times \mathcal{X}$ by a base change. $\square$

Remark 2.2. To simplify the exposition, we only consider presentations $\pi : \mathcal{X} \rightarrow X$ where X is a scheme from now on. This assumption allows us to avoid a separate discussion of perverse coherent sheaves on algebraic spaces.

Also, we do not need smoothness of presentations: it is enough to assume that $\pi : X \rightarrow \mathcal{X}$ is a faithfully flat Gorenstein morphism of finite type. Recall that a flat morphism of finite type $\pi : X \rightarrow Y$ between locally Noetherian schemes is *Gorenstein* if its fibers are Gorenstein schemes; equivalently, $\pi^! \mathcal{O}_Y$ is an invertible sheaf (concentrated in a single cohomological dimension that need not be zero), see [12, Exercise V.9.7]. The class of Gorenstein morphisms is local in smooth topology; this allows to define Gorenstein morphisms between locally Noetherian stacks.

Let us agree that a *presentation* of a stack is a (representable) faithfully flat Gorenstein morphism of finite type $\pi : X \rightarrow \mathcal{X}$, where X is a scheme.

Denote by $\mathrm{Coh}(\mathcal{X}) \subset \mathrm{QCoh}(\mathcal{X})$ the categories of coherent and quasi-coherent sheaves on $\mathcal{X}$, respectively. The presentation $\pi : X \rightarrow \mathcal{X}$ defines a simplicial algebraic space $X_\bullet$ (the *coskeleton* of π): X_i is the fiber product of $i + 1$ copies of X over $\mathcal{X}$ ($i \geq 0$). We can interpret quasi-coherent sheaves on $\mathcal{X}$ as cartesian quasi-coherent sheaves on $X_\bullet$.

Example 2.3. An important example is the quotient stack $\mathcal{X} = X/G$. Let us assume that X is a semi-separated Noetherian scheme and G is a flat finitely presented affine group scheme (over some base scheme) acting on X . Then $\mathcal{X}$ is an algebraic stack by the Artin criterion (see Theorem 10.1, Corollary 10.6 in [16]); it is Noetherian and semi-separated. The natural morphism $X \rightarrow \mathcal{X}$ is a presentation of $\mathcal{X}$ if G is smooth (or at least Gorenstein, since we consider presentations in the sense of Remark 2.2). In this example, quasi-coherent sheaves on $\mathcal{X}$ are simply G -equivariant quasi-coherent sheaves on X .

Remark. Example 2.3 demonstrates our main reason for working with Gorenstein presentations, rather than smooth presentations as in [16]. Namely, there are interesting group schemes G that are Gorenstein, but not smooth (for example, G could be a non-reduced group scheme over a field of non-zero characteristic). By our definition, the morphism $X \rightarrow \mathcal{X}$ is still a presentation of $\mathcal{X} = X/G$ for such G .

Denote by $D_{qcoh}(\mathcal{X})$ the derived category of quasi-coherent sheaves on $\mathcal{X}$. Recall its definition ([16]).

Definition 2.4. Consider the lisse-étale topology on $\mathcal{X}$. (The underlying category is the category of morphisms $u : U \rightarrow \mathcal{X}$, where U is an algebraic space smooth over $\mathcal{X}$; coverings are surjective étale morphisms.) This site is equipped with the sheaf of rings $\mathcal{O}_{\mathcal{X}}$. Let $D(\mathcal{O}_{\mathcal{X}})$ be the derived category of complexes of $\mathcal{O}_{\mathcal{X}}$ -modules. Then $D_{qcoh}(\mathcal{X}) \subset D(\mathcal{O}_{\mathcal{X}})$ is the full subcategory formed by complexes whose cohomology sheaves are quasi-coherent, and $D_{coh}(\mathcal{X}) \subset D_{qcoh}(\mathcal{X})$ is the full subcategory of complexes with coherent cohomology. The notation $D_{coh}^+(\mathcal{X})$, $D_{qcoh}^b(\mathcal{X})$, and so on is self-explanatory.

$D_{qcoh}(\mathcal{X})$ can be described more explicitly in terms of a presentation $\pi : X \rightarrow \mathcal{X}$. Let $X_\bullet$ be the corresponding simplicial algebraic space. (More generally, one can consider flat hypercovers $X_\bullet \rightarrow \mathcal{X}$.)

Consider the derived category of quasi-coherent sheaves on $X_\bullet$ (cf. [2, Appendix B]). We denote by $D(\mathcal{O}_{X_\bullet})$ the derived category of complexes of simplicial $\mathcal{O}_{X_\bullet}$ -modules on $X_\bullet$. The natural functor from $\mathbf{QCoh}(\mathcal{X})$ to the category of simplicial $\mathcal{O}_{X_\bullet}$ -modules is fully faithful. It provides an equivalence between $\mathbf{QCoh}(\mathcal{X})$ and the category of cartesian quasi-coherent sheaves on $X_\bullet$. Define $D_{qcoh, cart}(X_\bullet) \subset D(\mathcal{O}_{X_\bullet})$ to be the full subcategory of complexes whose cohomology objects are cartesian quasicohherent sheaves.

Claim 2.5. (a) *In Definition 2.4, the lisse-étale topology can be replaced by the smooth topology or the fppf topology, or a ‘hybrid topology’ such as ‘fppf-Zariski’. If $\mathcal{X}$ is a Deligne-Mumford stack (resp. scheme), one can work with the étale topology (resp. Zariski topology) on $\mathcal{X}$. In all these cases, the corresponding versions of $D_{qcoh}(\mathcal{X})$ are naturally equivalent.*
 (b) *Similarly, in the definition of $D_{qcoh, cart}(X_\bullet)$, one can use on X_i ’s any of the usual topologies: smooth, flat, étale, or Zariski (assuming X_i ’s are schemes). The corresponding derived categories are naturally equivalent.*

Claim 2.6. *The natural functor*

$$D_{qcoh}(\mathcal{X}) \rightarrow D_{qcoh, cart}(X_\bullet)$$

is an equivalence of categories.

Claim 2.7. $D_{qcoh}^+(\mathcal{X})$ *is identified with the bounded below derived category of the abelian category $\mathbf{QCoh}(\mathcal{X})$.*

Claims 2.5 and 2.6 are well known for the bounded below derived category $D_{qcoh}^+(\mathcal{X})$. In this case, they fit into the framework of cohomological descent [18] (see also [9, Chapter 5]); see Theorem 13.5.5 in [16] for a proof of Claim 2.6 under this assumption. Laszlo and Olsson extended cohomological descent to unbounded derived categories in [15]; the claims easily follow.

Proof of Claims 2.5 and 2.6. Claim 2.6 immediately follows from Theorem 2.2.3 of [15] (see [15, Example 2.2.5]). To prove Claim 2.5(a), we first note that there are natural functors relating different versions of the derived category (the functors are the direct and inverse images under the continuous maps between the corresponding sites). We need to verify that the functors are equivalences. This also follows from [15, Theorem 2.2.3]. (Technically, [15, Theorem 2.3.3] only considers maps from a simplicial topos to a topos, but any site can be viewed as a simplicial site.) Same argument applies to sites associated with different topologies on $X_\bullet$, proving Claim 2.5(b). $\square$

Let us now prove Claim 2.7. Fix a presentation $\pi : X \rightarrow \mathcal{X}$ such that X is an affine scheme and π is an affine morphism; such presentation exists by Lemma 2.1. Let $X_\bullet$ be the corresponding simplicial scheme. Consider the derived categories $D_{qcoh}(\mathcal{X})$ and $D_{qcoh, cart}(X_\bullet)$.

Claim 2.6 provides an equivalence between the categories. The equivalence is given by functors $\pi_{\bullet*}$ and $\pi_\bullet^*$, where $\pi_\bullet$ stands for the morphism $X_\bullet \rightarrow \mathcal{X}$. (We often use the same notation for a functor on an abelian category and its derived functor on the derived category.)

Now consider the derived category $D_{cart}(\mathbf{QCoh}(X_\bullet))$. Its objects are complexes of quasi-coherent sheaves on $X_\bullet$ whose cohomology sheaves are cartesian.

Lemma 2.8. *The natural functor $D_{cart}(\mathrm{QCoh}(X_\bullet)) \rightarrow D_{cart,qcoh}(X_\bullet)$ is an equivalence of categories.*

Proof. Once again, we derive the statement from [15, Theorem 2.2.3]. Let $Y_\bullet$ be the final object in the category of simplicial topological spaces: Y_i is a singleton set for all i . Equip Y_i with a sheaf of rings $\mathcal{O}_{Y_i}$ by setting

$$H^0(Y_i, \mathcal{O}_{Y_i}) = H^0(X_i, \mathcal{O}_{X_i}) \quad (i \geq 0).$$

Clearly, this gives rise to a simplicial sheaf of rings $\mathcal{O}_{Y_\bullet}$ on $Y_\bullet$ and a morphism of simplicial ringed sites $\Gamma_\bullet : X_\bullet \rightarrow Y_\bullet$. (Of course, a simplicial sheaf of rings on $Y_\bullet$ is nothing but a cosimplicial ring.)

The inverse image $\Gamma_\bullet^*$ identifies the category of $\mathcal{O}_{Y_\bullet}$ -modules and the category of quasi-coherent $\mathcal{O}_{X_\bullet}$ -modules. By [15, Theorem 2.2.3], the functors $\Gamma_{\bullet*}$ and $\Gamma_\bullet^*$ induce mutually inverse equivalences between $D_{cart,qcoh}(X_\bullet)$ and $D_{cart}(Y_\bullet)$ (the derived category of complexes of $\mathcal{O}_{Y_\bullet}$ -modules whose cohomology objects are cartesian). $\square$

Remark. Lemma 2.8 is a simplicial version of [7, Theorem 5.1].

By Lemma 2.8, we can restate Claim 2.7 using the category $D_{cart}^+(\mathrm{QCoh}(X_\bullet))$ in place of $D_{cart,qcoh}^+(X_\bullet)$. The advantage of working with $D_{cart}^+(\mathrm{QCoh}(X_\bullet))$ is that the functor $\pi_{\bullet*}$ can be described more explicitly. Namely, for $\mathcal{F} \in \mathrm{QCoh}(X_\bullet)$, consider its Čech complex

$$\check{\pi}_{\bullet*}\mathcal{F} = (\pi_{0*}\mathcal{F}_0 \rightarrow \pi_{1*}\mathcal{F}_1 \rightarrow \dots).$$

Here we denote by $\mathcal{F}_i$ the component of $\mathcal{F}$ concentrated on X_i ; $\pi_i : X_i \rightarrow \mathcal{X}$ is the projection. The definition immediately extends to complexes of quasi-coherent sheaves on $X_\bullet$. This yields a functor

$$\check{\pi}_{\bullet*} : D_{cart}^+(\mathrm{QCoh}(X_\bullet)) \rightarrow D^+(\mathrm{QCoh}(\mathcal{X}))$$

such that the composition

$$D_{cart}^+(\mathrm{QCoh}(X_\bullet)) \rightarrow D^+(\mathrm{QCoh}(\mathcal{X})) \rightarrow D_{qcoh}^+(\mathcal{X})$$

is naturally isomorphic $\pi_{\bullet*}$.

Proof of Claim 2.7. Let us show that $\check{\pi}_{\bullet*}$ is an equivalence, with inverse equivalence given by $\pi_\bullet^*$. An isomorphism between $\pi_\bullet^* \circ \check{\pi}_{\bullet*} = \pi_\bullet^* \circ \pi_{\bullet*}$ and the identity functor is given by Claim 2.6. On the other hand, for any $\mathcal{F} \in \mathrm{QCoh}(\mathcal{X})$, we have a natural map

$$\mathcal{F} \rightarrow \check{\pi}_{\bullet*}(\pi_\bullet^*\mathcal{F}).$$

It leads to a functorial morphism from the identity functor to $\check{\pi}_{\bullet*} \circ \pi_\bullet^*$. Clearly, it is an isomorphism of functors. $\square$

Remark. Let us sketch another proof of Claim 2.7 for bounded derived categories. We need to show that the natural functor

$$D^b(\mathrm{QCoh}(\mathcal{X})) \rightarrow D_{qcoh}^b(\mathcal{X})$$

is an equivalence. It is enough to verify that it is fully faithful; then essential surjectivity follows because every $\mathcal{F} \in D_{qcoh}^b(\mathcal{X})$ is obtained from quasi-coherent sheaves by taking cones. For the same reason, we only need to check that the natural morphism of functors

$$(2.1) \quad \mathrm{Ext}_{\mathrm{QCoh}(\mathcal{X})}^i(\mathcal{F}, \mathcal{G}) \rightarrow \mathrm{Ext}_{\mathcal{O}_{\mathcal{X}}}^i(\mathcal{F}, \mathcal{G}) \quad (\mathcal{F}, \mathcal{G} \in \mathrm{QCoh}(\mathcal{X}))$$

is an isomorphism for all i . This is trivial for $i = 0$. For $i > 0$, it suffices to check that both sides of (2.1) are effaceable as functors of $\mathcal{G} \in \mathrm{QCoh}(\mathcal{X})$.

Let $\pi : X \rightarrow \mathcal{X}$ be an affine presentation. For any $\mathcal{G} \in \mathrm{QCoh}(\mathcal{X})$, pick an embedding $\pi^*\mathcal{G} \hookrightarrow \mathcal{I}$ into injective $\mathcal{I} \in \mathrm{QCoh}(X)$. Then $\mathcal{G} \rightarrow \pi_*\mathcal{I}$ is also an embedding. It remains to notice that

$$\mathrm{Ext}_{\mathrm{QCoh}(\mathcal{X})}^i(\mathcal{F}, \pi_*\mathcal{I}) \rightarrow \mathrm{Ext}_{\mathcal{O}_{\mathcal{X}}}^i(\mathcal{F}, \pi_*\mathcal{I}) = 0 \quad (\mathcal{F} \in \mathrm{QCoh}(\mathcal{X}), i > 0)$$

by adjunction.

Remark. Bökstedt and Neeman proved that for a quasi-compact separated scheme X , the unbounded derived categories $D_{qcoh}(X)$ and $D(\mathrm{QCoh}(X))$ are equivalent ([7, Corollary 5.5]). It is possible that Claim 2.7 also holds for unbounded derived categories.

Note however that a finite affine cover of a quasi-compact semi-separated scheme X gives rise to a ‘bounded’ affine semi-simplicial scheme $X_\bullet$ (that is, $X_i = \emptyset$ for $i \gg 0$). The corresponding Čech complex is bounded; this allows us to extend the proof of Claim 2.7 to unbounded derived categories (the argument is then almost identical to that of Bökstedt and Neeman, see [7, Section 6.7]). In the case of stacks, such simplification is no longer available.

2.2. Coherent sheaves on stacks. Consider now the derived category $D_{coh}(\mathcal{X})$.

Lemma 2.9 ([16, Proposition 15.4]). *Any $\mathcal{F} \in \mathrm{QCoh}(\mathcal{X})$ is the union of its coherent subsheaves.* $\square$

Corollary 2.10. *Let $\mathcal{F}^\bullet$ be a bounded above complex of quasi-coherent sheaves on $\mathcal{X}$ whose cohomology is coherent. Then $\mathcal{F}^\bullet$ contains a coherent subcomplex $\mathcal{F}_c^\bullet \subset \mathcal{F}^\bullet$ quasi-isomorphic to $\mathcal{F}^\bullet$. Moreover, for any coherent subcomplex $\mathcal{G}^\bullet \subset \mathcal{F}^\bullet$, we can choose such $\mathcal{F}_c^\bullet$ so that $\mathcal{F}_c^\bullet \supset \mathcal{G}^\bullet$.*

Proof. The argument is almost standard (similar to [12, Proposition I.4.8], for instance). We can construct subsheaves $\mathcal{F}_c^i \subset \mathcal{F}^i$ by descending induction in i . We require that $\mathcal{F}_c^i$ has the following properties:

$$\begin{aligned} d(\mathcal{F}_c^i) &= \mathcal{F}_c^{i+1} \cap \mathcal{B}^{i+1} \\ (\mathcal{F}_c^i \cap \mathcal{Z}^i) &\rightarrow \mathcal{Z}^i / \mathcal{B}^i = H^i(\mathcal{F}^\bullet) \\ \mathcal{F}_c^i &\supset \mathcal{G}^i. \end{aligned}$$

Here $\mathcal{Z}^i, \mathcal{B}^i \subset \mathcal{F}^i$ denote, respectively, the kernel and the image of the differential. $\square$

Corollary 2.11. *The category $D_{coh}^b(\mathcal{X})$ is naturally equivalent to the derived category $D^b(\mathrm{Coh}(\mathcal{X}))$ of complexes of coherent sheaves.*

Proof. Consider the category $D_{coh}^b(\mathrm{QCoh}(\mathcal{X}))$. By Claim 2.7, it is equivalent to $D_{coh}^b(\mathcal{X})$. Therefore, we need to show that the natural functor

$$D^b(\mathrm{Coh}(\mathcal{X})) \rightarrow D_{coh}^b(\mathrm{QCoh}(\mathcal{X}))$$

is an equivalence. Essential surjectivity follows from Corollary 2.10. Let us show that the functor is faithful. Suppose

$$f : \mathcal{F}^\bullet \rightarrow \mathcal{G}^\bullet$$

is a morphism between bounded complexes of coherent sheaves whose image in $D^b(\mathrm{QCoh}(\mathcal{X}))$ vanishes. Then there exists a quasi-isomorphism

$$\iota : \mathcal{F}_1^\bullet \rightarrow \mathcal{F}^\bullet$$

such that $f\iota$ is homotopic to zero. Here $\mathcal{F}_1^\bullet$ is a bounded complex of quasi-coherent sheaves. Using Corollary 2.10, we replace $\mathcal{F}_1^\bullet$ by a quasi-isomorphic coherent subcomplex. This shows that the image of f in $D^b(\mathrm{Coh}(\mathcal{X}))$ also vanishes.

Now let us prove the functor is full. Given bounded complexes of coherent sheaves $\mathcal{F}^\bullet, \mathcal{G}^\bullet$, a morphism between their images in $D^b(\mathrm{QCoh}(\mathcal{X}))$ is of the form $f\iota^{-1}$, where $f : \mathcal{F}_1^\bullet \rightarrow \mathcal{G}^\bullet$ is a morphism between complexes of quasi-coherent sheaves, and $\iota : \mathcal{F}_1^\bullet \rightarrow \mathcal{F}^\bullet$ is a quasi-isomorphism. By Corollary 2.10, we can replace $\mathcal{F}_1^\bullet$ by a quasi-isomorphic coherent subcomplex. Then $\iota^{-1}f$ is defined in $D^b(\mathrm{Coh}(\mathcal{X}))$. $\square$

We also need statements relating coherent sheaves on $\mathcal{X}$ to coherent sheaves on its substacks.

Lemma 2.12. *Let $\mathcal{U} \subset \mathcal{X}$ be an open substack.*

- (a) (cf. [16, Corollary 15.5]) *Any $\mathcal{F} \in D_{coh}^b(\mathcal{U})$ extends to $\tilde{\mathcal{F}} \in D_{coh}^b(\mathcal{X})$ such that $\tilde{\mathcal{F}}|_{\mathcal{U}} \simeq \mathcal{F}$.*
- (b) *For any $\mathcal{F}, \mathcal{G} \in D_{coh}^b(\mathcal{U})$ and a morphism $f : \mathcal{F} \rightarrow \mathcal{G}$, we can choose extensions $\tilde{\mathcal{F}}, \tilde{\mathcal{G}} \in D_{coh}^b(\mathcal{X})$ and a morphism $\tilde{f} : \tilde{\mathcal{F}} \rightarrow \tilde{\mathcal{G}}$ such that $\tilde{f}|_{\mathcal{U}} \cong f$.*
- (c) *For any $\tilde{\mathcal{F}}', \tilde{\mathcal{F}}'' \in D_{coh}^b(\mathcal{X})$ and an isomorphism $f : \tilde{\mathcal{F}}'|_{\mathcal{U}} \cong \tilde{\mathcal{F}}''|_{\mathcal{U}}$, there exists $\tilde{\mathcal{F}} \in D_{coh}^b(\mathcal{X})$ and morphisms $f' : \tilde{\mathcal{F}}' \rightarrow \tilde{\mathcal{F}}$, $f'' : \tilde{\mathcal{F}}'' \rightarrow \tilde{\mathcal{F}}$ such that $f'|_{\mathcal{U}}, f''|_{\mathcal{U}}$ are isomorphisms and $f = (f'|_{\mathcal{U}}) \circ (f''|_{\mathcal{U}})^{-1}$.*

Proof. The argument is parallel to the proof of Corollary 2.10. Indeed, $\mathcal{F} \in D_{coh}^b(\mathcal{U})$ can be represented by a complex of coherent $\mathcal{O}_{\mathcal{U}}$ -modules. The complex extends to a complex of quasi-coherent $\mathcal{O}_{\mathcal{X}}$ -modules $\overline{\mathcal{F}}^\bullet$. We can then replace $\overline{\mathcal{F}}^\bullet$ by a coherent subcomplex $\mathcal{F}_c^\bullet$ such that the restriction $\mathcal{F}_c^\bullet|_{\mathcal{U}} \hookrightarrow \overline{\mathcal{F}}^\bullet|_{\mathcal{U}}$ is a quasi-isomorphism. Moreover, any coherent subcomplex of $\overline{\mathcal{F}}^\bullet$ lies in such a subcomplex. The statement follows. $\square$

Remark. Lemma 2.12 implies that $D_{coh}^b(\mathcal{U})$ is equivalent to the localization of $D_{coh}^b(\mathcal{X})$ with respect to the class of morphisms that become isomorphisms when restricted to $\mathcal{U}$.

Lemma 2.13. *Let $\mathcal{X}^{top}$ be the set of points of $\mathcal{X}$ equipped with the Zariski topology. Given closed $\mathbf{Z} \subset \mathcal{X}^{top}$ and $\mathcal{F} \in D_{coh}^b(\mathcal{X})$ such that $\mathrm{supp} \mathcal{F} \subset \mathbf{Z}$, there exists a closed substack $i_{\mathbf{Z}} : \mathcal{Z} \hookrightarrow \mathcal{X}$ with $\mathcal{Z}^{top} \subset \mathbf{Z}$ such that*

$$\mathcal{F} \cong i_{\mathbf{Z}*}(\mathcal{F}_{\mathbf{Z}}) \text{ for some } \mathcal{F}_{\mathbf{Z}} \in D_{coh}^b(\mathcal{Z}).$$

Proof. Let $\mathrm{QCoh}_{\mathbf{Z}}(\mathcal{X}) \subset \mathrm{QCoh}(\mathcal{X})$ be the full subcategory of sheaves supported by $\mathbf{Z}$. It is easy to see that the $\mathrm{QCoh}_{\mathbf{Z}}(\mathcal{X})$ has enough objects that are injective in $\mathrm{QCoh}(\mathcal{X})$; therefore, $\mathcal{F}$ can be represented by a (bounded from below) complex of injective objects of $\mathrm{QCoh}_{\mathbf{Z}}(\mathcal{X})$. Truncation gives a bounded complex whose terms need not be injective. Finally, by Corollary 2.10 we see that $\mathcal{F}$ can be represented by a complex of coherent sheaves supported by $\mathbf{Z}$. This implies the statement. $\square$

2.3. Dualizing complexes of stacks. As before, $\mathcal{X}$ is a Noetherian stack. Recall ([12, §V.2]) that the dualizing complex on a Noetherian scheme X is an object $\mathbb{D}\mathbb{C}_X \in D_{coh}^b(X)$ of finite injective dimension (in $\mathbf{QCoh}(X)$) such that every $\mathcal{F} \in D_{coh}^b(X)$ is $\mathbb{D}\mathbb{C}_X$ -reflexive. That is, the natural transformation

$$\mathcal{F} \rightarrow \underline{Hom}(\underline{Hom}(\mathcal{F}, \mathbb{D}\mathbb{C}), \mathbb{D}\mathbb{C}) \quad (\mathcal{F} \in D_{coh}^b(X))$$

is an isomorphism.

The notion of a dualizing complex is local with respect to flat Gorenstein covers:

Lemma 2.14. *Let $\pi : Y \rightarrow X$ be a faithfully flat Gorenstein morphism of finite type between Noetherian schemes. Then $\mathcal{F} \in D_{coh}^b(X)$ is a dualizing complex on X if and only if $\pi^*\mathcal{F}$ is a dualizing complex on Y .* $\square$

Definition 2.15. Let $\mathcal{X}$ be a Noetherian stack. A *dualizing complex* on $\mathcal{X}$ is an object $\mathbb{D}\mathbb{C}_{\mathcal{X}} \in D_{coh}^b(\mathcal{X})$ such that for a presentation $\pi : X \rightarrow \mathcal{X}$, $\pi^*\mathbb{D}\mathbb{C}_{\mathcal{X}}$ is a dualizing complex on X .

By Lemma 2.14, Definition 2.15 does not depend on the presentation π . It is easy to see that $\mathbb{D}\mathbb{C}_{\mathcal{X}}$ has finite injective dimension in $\mathbf{QCoh}(\mathcal{X})$, and that all objects of $D_{coh}^b(\mathcal{X})$ are $\mathbb{D}\mathbb{C}_{\mathcal{X}}$ -reflexive.

Example 2.16. Suppose the ground scheme S is Noetherian and admits a dualizing complex. Let G be a separated Gorenstein S -group scheme of finite type acting on a S -scheme X of finite type. By Definition 2.15, a dualizing complex $\mathbb{D}\mathbb{C}_{\mathcal{X}}$ on $\mathcal{X} = X/G$ can be viewed as a G -equivariant dualizing complex on X : a G -equivariant complex is an equivariant dualizing complex if and only if it is a dualizing complex on X . Proposition 2.17 shows that $\mathbb{D}\mathbb{C}_{\mathcal{X}}$ exists.

Proposition 2.17. *Suppose $\mathcal{X}$ is of finite type over a Noetherian scheme S that admits a dualizing complex. Then $\mathcal{X}$ admits a dualizing complex.*

Proof. Let $\pi : X \rightarrow \mathcal{X}$ be a presentation of $\mathcal{X}$. Since π is Gorenstein, it is easy to define the relative dualizing sheaf $\omega_{X/\mathcal{X}}$, which is a line bundle on X . Let us include the appropriate cohomological shift by $\dim(X/\mathcal{X})$ in $\omega_{X/\mathcal{X}}$, so that

$$\pi^!(\mathcal{F}) = \pi^*(\mathcal{F}) \otimes \omega_{X/\mathcal{X}} \quad (\mathcal{F} \in D_{coh}^b(\mathcal{X})),$$

as in [12, Remark on pp. 143–144].

Now fix a dualizing complex $\mathbb{D}\mathbb{C}_S \in D_{coh}^b(S)$, and set

$$\mathcal{F} = f_X^!(\mathbb{D}\mathbb{C}_S) \otimes \omega_{X/\mathcal{X}}^{-1} \in D_{coh}^b(X),$$

where $f_X : X \rightarrow S$. Clearly, $\mathcal{F}$ is a dualizing complex on X . Let us show that it descends to $\mathcal{X}$.

By [1, Theorem 3.2.4] (see also [1, Section 3.2.5]), an object of the derived category of sheaves on a site can be given locally provided negative local Ext's from the object to itself vanish. Applying it to the flat covering $\pi : X \rightarrow \mathcal{X}$, we see that it is enough to provide an isomorphism

$$\pi_2^*\mathcal{X} \simeq \pi_1^*\mathcal{X} \in D_{coh}^b(X \times_{\mathcal{X}} X)$$

satisfying the associativity constraint on $X \times_{\mathcal{X}} X \times_{\mathcal{X}} X$. Here $\pi_{1,2} : X \times_{\mathcal{X}} X \rightarrow X$ are the projections. However, $\pi_{1,2}$ are Gorenstein, so

$$\pi_{1,2}^! = \pi_{1,2}^* \otimes \omega_{X \times_{\mathcal{X}} X/X},$$

and

$$\pi_{1,2}^* \mathcal{F} = f_{X \times_{\mathcal{X}} X}^! (\mathbb{D}\mathbb{C}_S) \otimes \omega_{X \times_{\mathcal{X}} X/\mathcal{X}}^{-1}.$$

This provides the desired isomorphism. The associativity constraint follows from functorial properties of $f^!$. $\square$

Remark. The dualizing complex on $\mathcal{X}$ can be constructed more explicitly if there exists a closed embedding $\iota : \mathcal{X} \hookrightarrow \mathcal{Y}$ into a smooth S -stack $\mathcal{Y}$. In this case, we can take

$$\mathbb{D}\mathbb{C}_{\mathcal{X}} = \iota^! (f_{\mathcal{Y}}^* (\mathbb{D}\mathbb{C}_S) \otimes \omega_{\mathcal{Y}/S})$$

for $f_{\mathcal{Y}} : \mathcal{Y} \rightarrow S$. (A less canonical choice is $\mathbb{D}\mathbb{C}_{\mathcal{X}} = \iota^! f_{\mathcal{Y}}^* (\mathbb{D}\mathbb{C}_S)$.)

In particular, suppose X is a normal quasiprojective variety over an algebraically closed field, and G is a connected linear algebraic group acting on X . Sumihiro's embedding theorem ([19], see also [14]) claims that there exists an action of G on $\mathbb{P}^n$ and a G -equivariant embedding $X \hookrightarrow \mathbb{P}^n$. Thus the stack $\mathcal{X} = X/G$ admits a closed embedding

$$\mathcal{X} \hookrightarrow \mathbb{P}^n/G.$$

2.4. *-restriction and !-restriction. We now summarize the properties of restriction to a (not necessarily) closed point. To simplify the exposition, we only use the functors on schemes. Let X be a Noetherian scheme.

Remark. We do not need to assume that X is semi-separated. This assumption is only used in Claim 2.7, but it is well known that the claim holds for locally Noetherian schemes ([12, Corollary II.7.19]).

We write X^{top} for the underlying topological space. Given a point $x \in X^{top}$ (resp. a locally closed subscheme $Z \subset X$), we write $i_x : \{x\} \hookrightarrow X^{top}$ (respectively $i_Z : Z \hookrightarrow X$) for the embedding.

For $x \in X^{top}$, consider the restriction functors

$$i_x^* : D_{qcoh}(X) \rightarrow D(\mathcal{O}_x - mod), \quad i_x^! : D_{qcoh}^+(X) \rightarrow D^+(\mathcal{O}_x - mod).$$

They are derived of an exact functor and a left exact functor, respectively. Note that the functor $i_x^!$ has finite homological dimension, because so does the functor j_* for an open embedding j .

Lemma 2.18. *Let $Z \subset X$ be a locally closed subscheme, and n be an integer. Let $x \in X^{top}$ be a generic point of Z . Then*

- (a) *For $\mathcal{F} \in D_{coh}^-(X)$ we have $i_x^*(\mathcal{F}) \in D^{\leq n}(\mathcal{O}_x - mod)$ if and only if there exists an open subscheme $Z^0 \subset Z$, $Z^0 \ni x$ such that $i_{Z^0}^*(\mathcal{F}) \in D_{coh}^{\leq n}(Z^0)$;*
- (b) *For $\mathcal{F} \in D_{coh}^+(X)$ we have $i_x^!(\mathcal{F}) \in D^{\geq n}(\mathcal{O}_x - mod)$ if and only if there exists an open subscheme $Z^0 \subset Z$, $Z^0 \ni x$ such that $i_{Z^0}^!(\mathcal{F}) \in D_{coh}^{\geq n}(Z^0)$.*

Proof. Existence of an open subscheme $Z^0 \subset Z$ as in (a) is equivalent to

$$i_x^* i_Z^*(\mathcal{F}) \in D^{\leq n}(\mathcal{O}_{Z,x} - mod).$$

Indeed, if this holds, we can let Z^0 be the complement in Z to support of $\mathcal{H}^k(i_Z^*(\mathcal{F}))$, $k > n$; the converse is obvious.

Let us rewrite

$$i_x^* i_Z^*(\mathcal{F}) = i_x^*(\mathcal{F}) \otimes_{\mathcal{O}_x}^L \mathcal{O}_{Z,x}.$$

Since the functor of tensor product with $\mathcal{O}_{Z,x}$ over $\mathcal{O}_x$ is right exact, and kills no finitely generated $\mathcal{O}_x$ modules by the Nakayama Lemma, we see that the top

cohomology of $i_x^*(\mathcal{F}) \otimes_{\mathcal{O}_x}^L \mathcal{O}_{Z,x}$ and of $i_x^*(\mathcal{F})$ occur in the same degree. This proves (a).

Similarly, the second condition in (b) says that

$$i_x^! i_Z^!(\mathcal{F}) = i_x^* i_Z^!(\mathcal{F}) \in D^{\geq n}((\mathcal{O}_Z)_x - \text{mod})$$

(the equality here is due to the fact x is generic in Z). We rewrite

$$i_x^! i_Z^!(\mathcal{F}) = R\text{Hom}_{\mathcal{O}_x}(\mathcal{O}_{Z,x}, i_x^!(\mathcal{F})),$$

and see that the lowest cohomology of $i_x^!(\mathcal{F})$ and of $R\text{Hom}_{\mathcal{O}_x}(\mathcal{O}_{Z,x}, i_x^!(\mathcal{F}))$ occur in the same degree. Indeed, the functor $\text{Hom}_{\mathcal{O}_x}(\mathcal{O}_{Z,x}, _)$ is left exact, and kills no torsion module, while cohomology of $i_x^!(\mathcal{F})$ is a torsion $\mathcal{O}_x$ -module. $\square$

Lemma 2.19 (cf. e.g. [12], Theorem V.4.1). *Let $i : \mathbf{Z} \hookrightarrow X^{\text{top}}$ be the embedding of a closed subspace. For any $\mathcal{F} \in D_{\text{coh}}^-(X)$, $\mathcal{G} \in D_{\text{coh}}^+(X)$ we have*

$$\text{Hom}(\mathcal{F}, i_* i^!(\mathcal{G})) = \varinjlim_Z \text{Hom}(\mathcal{F}, i_{Z*} i_Z^!(\mathcal{G})),$$

where Z runs over the set of closed subschemes of X with the underlying topological space $\mathbf{Z}$.

Proof. Let us represent $\mathcal{G}$ by a bounded below complex $I_{\mathcal{G}}$ of injective quasi-coherent sheaves. Also, we can represent $\mathcal{F}$ by a bounded above complex $P_{\mathcal{F}}$ whose terms are finite direct sums of sheaves of the form $(j_U)_! \mathcal{O}_U$ for open affine embeddings $j_U : U \hookrightarrow X$.

Then $R\text{Hom}(\mathcal{F}, i_{Z*} i_Z^!(\mathcal{G}))$ can be computed by the complex

$$\text{Hom}^\bullet(P_{\mathcal{F}}, i_{Z*} i_Z^!(I_{\mathcal{G}})).$$

On the other hand, $R\text{Hom}(\mathcal{F}, i_* i^!(\mathcal{G}))$ is computed by the complex

$$\text{Hom}^\bullet(P_{\mathcal{F}}, i_* i^!(I_{\mathcal{G}})).$$

Finally,

$$i_* i^!(I_{\mathcal{G}}) = \bigcup_Z i_{Z*} i_Z^!(I_{\mathcal{G}}),$$

and therefore

$$\text{Hom}^\bullet(P_{\mathcal{F}}, i_* i^!(I_{\mathcal{G}})) = \bigcup_Z \text{Hom}^\bullet(P_{\mathcal{F}}, i_{Z*} i_Z^!(I_{\mathcal{G}})).$$

This implies the lemma. $\square$

3. PERVERSE COHERENT SHEAVES

3.1. Perverse t -structure on schemes. Let us first consider the case of schemes. Suppose X is a Noetherian scheme that admits a dualizing complex.

Fix a dualizing complex

$$\mathbb{D}\mathbb{C}_X \in D_{\text{coh}}^b(X).$$

The choice defines the *codimension function* d on points $x \in X^{\text{top}}$ such that $i_x^!(\mathbb{D}\mathbb{C}_X)$ is concentrated in homological degree $d(x)$ (see [12], §V.7). Clearly, $d(x)$ is bounded, because $\mathbb{D}\mathbb{C}_X$ has finite injective dimension. We set $\dim(x) = -d(x)$.

Example 3.1. Suppose X is of finite type over a field. Then we can choose $\mathbb{D}\mathbb{C}_X$ so that $\dim(x)$ equals to the dimension of the closure of x .

Let p (the *perverseity*) be an integer-valued function $p : X^{\text{top}} \rightarrow \mathbb{Z}$. The *dual perverseity* (for given $\mathbb{D}\mathbb{C}_X$) is given by $\bar{p}(x) = -\dim(x) - p(x)$.

Definition 3.2. Define $D^{p,\leq 0}(X) \subset D_{coh}^-(X)$, $D^{p,\geq 0}(X) \subset D_{coh}^+(X)$ as follows:

$\mathcal{F} \in D^{p,\geq 0}(X)$ if for any $x \in X^{top}$ we have $i_x^!(\mathcal{F}) \in D^{\geq p(x)}(\mathcal{O}_x - mod)$.

$\mathcal{F} \in D^{p,\leq 0}(X)$ if for any $x \in X^{top}$ we have $i_x^*(\mathcal{F}) \in D^{\leq p(x)}(\mathcal{O}_x - mod)$.

Lemma 3.3. (a) $\mathbb{D}(D^{p,\leq 0}(X)) = D^{\bar{p},\geq 0}$, where $\mathbb{D} = \underline{Hom}(_, \mathbb{D}\mathbb{C}_X)$ is the duality functor.

(b) Let $i_Z : Z \hookrightarrow X$ be a locally closed subscheme. Define the induced perversity on Z by $p_Z = p \circ i_Z : Z^{top} \rightarrow \mathbb{Z}$. Then

$$i_Z^*(D^{p,\leq 0}(X)) \subset D^{p_Z,\leq 0}(Z) \quad \text{and} \quad i_Z^!(D^{p,\geq 0}(X)) \subset D^{p_Z,\geq 0}(Z).$$

(c) In the situation of (b) assume that Z is closed. Then

$$i_{Z*}(D^{p_Z,\leq 0}(Z)) \subset D^{p,\leq 0}(X) \quad \text{and} \quad i_{Z*}(D^{p_Z,\geq 0}(Z)) \subset D^{p,\geq 0}(X).$$

Proof. (a) One knows from [12], §V.6 that

$$i_x^!(\mathbb{D}(\mathcal{F})) = \mathrm{Hom}_{\mathcal{O}_x}(i_x^*(\mathcal{F}), I_{\mathcal{O}_x})[-\dim(x)] \quad (\mathcal{F} \in D_{coh}^b(X), x \in X^{top}),$$

where $I_{\mathcal{O}_x}$ is the injective hull of the residue field of $\mathcal{O}_x$. Since $\mathrm{Hom}_{\mathcal{O}_x}(_, I_{\mathcal{O}_x})$ is exact and kills no finitely generated $\mathcal{O}_x$ -module, (a) follows.

(b) follows from Lemma 2.18; in view of this lemma if $\mathcal{F} \in D^{p,\leq 0}$, then for any $x \in Z^{top} \subset X^{top}$ there exists a subscheme $Z' \subset Z$ with generic point x such that $i_{Z'}^*(\mathcal{F}) = i_{Z'}^*(i_Z^*(\mathcal{F})) \in D^{\leq p(x)}(Z)$. This implies $i_Z^*(\mathcal{F}) \in D^{p_Z,\leq 0}(Z)$. The argument for $i_Z^!(\mathcal{F})$ is similar.

(c) is obvious. $\square$

It is clear that the categories $D^{p,\leq 0}(X)$, $D^{p,\geq 0}(X)$ are local with respect to flat Gorenstein coverings.

Lemma 3.4. Suppose $\pi : Y \rightarrow X$ is a faithfully flat Gorenstein morphism of finite type. Consider on Y the perversity $\pi^*p = p \circ \pi$. Then for $\mathcal{F} \in D_{coh}(X)$,

$$\mathcal{F} \in D^{p,\geq 0}(X) \text{ if and only if } \pi^*\mathcal{F} \in D^{\pi^*p,\geq 0}(Y)$$

$$\mathcal{F} \in D^{p,\leq 0}(X) \text{ if and only if } \pi^*\mathcal{F} \in D^{\pi^*p,\leq 0}(Y).$$

$\square$

Remark. One could consider on Y the perversity $\pi^!p$ given by

$$\pi^!p(y) = p(\pi(y)) - \mathrm{codim}(\overline{\{y\}} \subset \overline{\pi^{-1}(\pi(y))}) \quad (y \in Y^{top}).$$

Then

$$\pi^!\overline{p} = \overline{\pi^*p},$$

assuming the dualizing complex on Y is the pull-back of the dualizing complex on X . It is easy to see that Lemma 3.4 remains true for perversity $\pi^!p$; this statement is dual to Lemma 3.4 in the sense of Grothendieck-Serre duality.

Proposition 3.5. For $\mathcal{F} \in D^{p,\leq 0}(X)$, $\mathcal{G} \in D^{p,>0}(X)$ we have $\mathrm{Hom}(\mathcal{F}, \mathcal{G}) = 0$.

Proof. Fix $\mathcal{F}$ and $\mathcal{G}$. We proceed by Noetherian induction in X ; thus we can assume that the statement with (X, p) replaced by (Z, p_Z) for all closed subschemes $Z \subsetneq X$ is known.

Let x be a generic point of X . By Lemma 2.18, there exists an open subscheme $j : U \hookrightarrow X$ containing x such that $j^*(\mathcal{F}) \in D_{coh}^{\leq p(x)}(U)$ and $j^*(\mathcal{G}) = j^!(\mathcal{G}) \in D_{coh}^{> p(x)}(U)$. Thus, of course, $\mathrm{Hom}(j^*(\mathcal{F}), j^*(\mathcal{G})) = 0$.

Let i denote the closed embedding of $X^{top} - U^{top}$ into X^{top} . It induces a distinguished triangle

$$i_* i^! \mathcal{G} \rightarrow \mathcal{G} \rightarrow j_* j^*(\mathcal{G}) \rightarrow i_* i^! \mathcal{G}[1]$$

in $D_{coh}^b(X)$. By Lemma 2.19,

$$\mathrm{Hom}(\mathcal{F}, i_* i^! \mathcal{G}) = \varinjlim_Z \mathrm{Hom}(\mathcal{F}, i_{Z*} i_Z^! (\mathcal{G})) = \varinjlim_Z \mathrm{Hom}(i_Z^*(\mathcal{F}), i_Z^! (\mathcal{G})),$$

where Z runs over closed subschemes of X whose underlying set is $X^{top} - U^{top}$. Notice finally that for any such Z ,

$$i_Z^*(\mathcal{F}) \in D^{pz, \leq 0}(Z) \quad \text{and} \quad i_Z^! (\mathcal{G}) \in D^{pz, > 0}(Z)$$

by Lemma 3.3(b), so

$$\mathrm{Hom}(i_Z^*(\mathcal{F}), i_Z^! (\mathcal{G})) = 0$$

by the induction hypotheses. This implies the desired equality $\mathrm{Hom}(\mathcal{F}, \mathcal{G}) = 0$, since $\mathrm{Hom}(\mathcal{F}, j_* j^*(\mathcal{G})) = \mathrm{Hom}(j^*(\mathcal{F}), j^*(\mathcal{G})) = 0$. $\square$

If perversity p is both monotone and comonotone (Definition 3.9), the categories $D^{p, \geq 0}(X)$ and $D^{p, \leq 0}(X)$ yield a t -structure on $D_{coh}^b(X)$. To avoid repeating the argument, we prove this for perversities on stacks in Theorem 3.10.

3.2. Perverse t -structure on stacks. Now let $\mathcal{X}$ be a Noetherian semi-separated stack that admits a dualizing complex. Fix a dualizing complex $\mathbb{D}\mathbb{C}_{\mathcal{X}} \in D_{coh}^b(\mathcal{X})$. We denote by $\mathcal{X}^{top}$ the set of points of $\mathcal{X}$, which we consider as a topological space with respect to the Zariski topology. By definition, a *perversity* on $\mathcal{X}$ is a function $p : \mathcal{X}^{top} \rightarrow \mathbb{Z}$.

Fix a presentation $\pi : X \rightarrow \mathcal{X}$ of $\mathcal{X}$. On X , we consider the induced perversity $\pi^* p = p \circ \pi : X^{top} \rightarrow \mathbb{Z}$ and the induced dualizing sheaf $\mathbb{D}\mathbb{C}_X = \pi^* \mathbb{D}\mathbb{C}_{\mathcal{X}}$. For a point $x \in \mathcal{X}^{top}$, take $y \in \pi^{-1}(x) \in X^{top}$, and define the codimension function

$$d(x) = d(y) - \mathrm{codim}(y/x),$$

where $\mathrm{codim}(y/x)$ is the codimension of the closure of y in the fiber of π over x . The definition is independent of presentation π and preimage y . Set $\dim(x) = -d(x)$, and define the dual perversity $\bar{p} : \mathcal{X}^{top} \rightarrow \mathbb{Z}$ by the same formula as in the case of schemes.

Example 3.6. Suppose $\mathcal{X}$ is of finite type over a field. Then for appropriate choice of $\mathbb{D}\mathbb{C}_{\mathcal{X}}$, we have

$$\dim(x) = \dim(\overline{\{x\}}) \quad (x \in \mathcal{X}^{top}).$$

Definition 3.7.

$$\begin{aligned} D^{p, \leq 0}(\mathcal{X}) &= \{\mathcal{F} \in D^-(\mathcal{X}) : \pi^* \mathcal{F} \in D^{\pi^* p, \leq 0}(X)\} \\ D^{p, \geq 0}(\mathcal{X}) &= \{\mathcal{F} \in D^+(\mathcal{X}) : \pi^* \mathcal{F} \in D^{\pi^* p, \geq 0}(X)\}. \end{aligned}$$

It follows from Lemma 3.4 that Definition 3.7 does not depend on the choice of presentation $\pi : X \rightarrow \mathcal{X}$. It is clear that Lemmas 3.3 and 3.4 hold for stacks. Let us now prove Proposition 3.5 in these settings.

Proposition 3.8. *For $\mathcal{F} \in D^{p, \leq 0}(\mathcal{X})$, $\mathcal{G} \in D^{p, > 0}(\mathcal{X})$ we have $\mathrm{Hom}(\mathcal{F}, \mathcal{G}) = 0$.*

Proof. Fix $\mathcal{F}$ and $\mathcal{G}$, and let us prove that $\text{Ext}^i(\mathcal{F}, \mathcal{G}) = 0$ for $i \leq 0$ by induction. The statement holds for $i \ll 0$, because $\mathcal{F} \in D^-(\mathcal{X})$ and $\mathcal{G} \in D^+(\mathcal{X})$. It remains to prove that if the statement holds for all $i < i_0$, it also holds for $i = i_0$, where $i_0 \leq 0$ is fixed. Now [1, Proposition 3.2.2] shows that homomorphisms from $\mathcal{F}$ to $\mathcal{G}[i_0]$ form a sheaf in the flat topology on $\mathcal{X}$. To complete the proof, note that $\text{Hom}(\pi^*\mathcal{F}, \pi^*\mathcal{G}[i_0]) = 0$ by Proposition 3.5. $\square$

Definition 3.9. A perversity function p is

monotone if $p(x') \geq p(x)$ whenever $x' \in \overline{\{x\}}$, $x, x' \in \mathcal{X}^{top}$;

strictly monotone if $p(x') > p(x)$ whenever $x' \in \overline{\{x\}}$, $x, x' \in \mathcal{X}^{top}$, $x \neq x'$;

(strictly) *comonotone* if the dual perversity $\overline{p}(x) = -\dim(x) - p(x)$ is (strictly) monotone.

Theorem 3.10. Suppose that a perversity p is monotone and comonotone. Then $(D^{p, \leq 0}(\mathcal{X}) \cap D^b(\mathcal{X}), D^{p, \geq 0}(\mathcal{X}) \cap D^b(\mathcal{X}))$ define a t -structure on $D_{coh}^b(\mathcal{X})$.

Proof. In view of Proposition 3.8 we have only to construct for any $\mathcal{F} \in D_{coh}^b(\mathcal{X})$ a distinguished triangle

$$\mathcal{F}' \rightarrow \mathcal{F} \rightarrow \mathcal{F}'' \rightarrow \mathcal{F}'[1]$$

with $\mathcal{F}' \in D^{p, \leq 0}(\mathcal{X})$, $\mathcal{F}'' \in D^{p, > 0}(\mathcal{X})$.

We abuse notation as follows. For a category A , we denote the set of isomorphism classes of $\text{Ob}(A)$ by the same letter A . In particular, for a functor $F : A \rightarrow B$, we denote by $F(A)$ the image of the induced map the set of isomorphism classes of $\text{Ob}(A)$ to that of $\text{Ob}(B)$.

Let us also adopt the following convention (see [1], 1.3.9). If D' , D'' are sets of (isomorphism classes) of objects of a triangulated category D , then $D' * D''$ is the set of (isomorphism classes) of objects of D , defined by the following condition: $B \in D' * D''$ if and only if there exists a distinguished triangle

$$A \rightarrow B \rightarrow C \rightarrow A[1] \quad (A \in D', C \in D'').$$

The octahedron axiom implies (see [1], Lemma 1.3.10) that the $*$ operation is associative:

$$(D' * D'') * D''' = D' * (D'' * D''').$$

Thus the meaning of the notation $D_1 * \dots * D_n$ is unambiguous.

In this notation, we need to prove that

$$D_{coh}^b(\mathcal{X}) \subset D^{p, \leq 0}(\mathcal{X}) * D^{p, > 0}(\mathcal{X}).$$

We proceed by Noetherian induction; thus we can assume that for any closed substack $\mathcal{Z} \subsetneq \mathcal{X}$,

$$D_{coh}^b(\mathcal{Z}) \subset D^{p_{\mathcal{Z}}, \leq 0}(\mathcal{Z}) * D^{p_{\mathcal{Z}}, > 0}(\mathcal{Z}),$$

where $p_{\mathcal{Z}} : \mathcal{Z}^{top} \rightarrow \mathbb{Z}$ is the induced perversity. By Lemma 3.3(c), it suffices to show that

$$(3.1) \quad D_{coh}^b(\mathcal{X}) \subset \bigcup_{\mathcal{Z}} D^{p, \leq 0}(\mathcal{X}) * i_{\mathcal{Z}*} D_{coh}^b(\mathcal{Z}) * D^{p, > 0}(\mathcal{X}),$$

where the union is over closed substacks $i_{\mathcal{Z}} : \mathcal{Z} \hookrightarrow \mathcal{X}$, $\mathcal{Z} \neq \mathcal{X}$.

Let us prove (3.1). Fix $\mathcal{F} \in D_{coh}^b(\mathcal{X})$. Choose a generic point $y \in \mathcal{X}^{top}$ and a closed irreducible substack $i_{\mathcal{Y}} : \mathcal{Y} \hookrightarrow \mathcal{X}$ that coincides with $\mathcal{X}$ in a neighborhood of y . Set

$$\mathcal{F}^- = \tau_{\leq p(y)}^{stand}(i_{\mathcal{Y}*} i_{\mathcal{Y}}^!(\mathcal{F})).$$

Here $\tau_{\leq n}^{stand} : D^b(\mathcal{X}) \rightarrow D^{\leq n}(\mathcal{X})$ is the truncation functor for the usual t -structure ($n \in \mathbb{Z}$). Clearly $\mathcal{F}^- \in D^{p, \leq 0}(\mathcal{X})$, because it is supported on $\mathcal{Y}$ and p is monotone. It is equipped with a canonical morphism $\mathcal{F}^- \rightarrow \mathcal{F}$. Set

$$\mathcal{F}_1 = \text{Cone}(\mathcal{F}^- \rightarrow \mathcal{F}).$$

Over an open subset of $\mathcal{Y}$, $\mathcal{F}_1$ is concentrated in cohomological degrees above $p(y)$.

The dual procedure (in the sense of Grothendieck-Serre duality) gives $\mathcal{F}^+ \in D^{p, > 0}(\mathcal{X})$ and a morphism $f : \mathcal{F}_1 \rightarrow \mathcal{F}^+$ that is an isomorphism over y . More precisely, set

$$\mathcal{F}^+ = \mathbb{D}(\tau_{< \overline{p}(y)}^{stand} i_{\mathcal{Y}*} i_{\mathcal{Y}}^! (\mathbb{D}(\mathcal{F}_1))).$$

Since p is comonotone, we see by Lemma 3.3(a) that $\mathcal{F}^+ \in D^{p, > 0}(\mathcal{X})$. Note that the duality is exact over the generic point y ([12], §V.6), which implies that f is an isomorphism at y .

Thus

$$\mathcal{F}^0 = \text{Cone}(\mathcal{F}_1 \rightarrow \mathcal{F}^+)$$

vanishes at y . By Lemma 2.13, there is a closed substack $i_{\mathcal{Z}} : \mathcal{Z} \hookrightarrow \mathcal{X}$, $\mathcal{Z} \neq \mathcal{X}$ such that

$$\mathcal{F}^0 \cong i_{\mathcal{Z}*}(\mathcal{F}_{\mathcal{Z}}) \quad (\mathcal{F}_{\mathcal{Z}} \in D_{coh}^b(\mathcal{Z})).$$

Therefore,

$$\mathcal{F} \in \{\mathcal{F}^-\} * \{\mathcal{F}^0[-1]\} * \{\mathcal{F}^+\} \subset D^{p, \leq 0}(\mathcal{X}) * i_{\mathcal{Z}*} D_{coh}^b(\mathcal{Z}) * D^{p, > 0}(\mathcal{X}),$$

which proves (3.1). $\square$

Remark. Construction of an object $\mathcal{F}^+ \in D^{p, > 0}(\mathcal{X})$ with given generic fiber (and with a morphism from a given object) is the only place where we use the duality formalism on stacks.

Corollary 3.11. *($D^{p, \leq 0}(\mathcal{X}), D^{p, \geq 0}(\mathcal{X})$) define a t -structure on $D_{coh}(\mathcal{X})$. In other words, Theorem 3.10 holds for the unbounded derived category.*

Proof. We need to prove that

$$D^b(\mathcal{X}) = D^{p, \leq 0}(\mathcal{X}) * D^{p, > 0}(\mathcal{X}),$$

where $*$ is defined in the proof of Theorem 3.10. Clearly,

$$D^{p, \leq 0}(\mathcal{X}) \supset D_{coh}^{< -N}(\mathcal{X}) \quad \text{and} \quad D^{p, > 0}(\mathcal{X}) \supset D_{coh}^{> N}(\mathcal{X}) \quad (N \gg 0).$$

Therefore,

$$\begin{aligned} D^{p, \leq 0}(\mathcal{X}) * D^{p, > 0}(\mathcal{X}) &= D_{coh}^{< -N}(\mathcal{X}) * D^{p, \leq 0}(\mathcal{X}) * D^{p, > 0}(\mathcal{X}) * D_{coh}^{> N}(\mathcal{X}) \supset \\ &D_{coh}^{< -N}(\mathcal{X}) * D_{coh}^b(\mathcal{X}) * D_{coh}^{> N}(\mathcal{X}) = D_{coh}(\mathcal{X}). \end{aligned}$$

$\square$

Corollary 3.12. *Let $j : \mathcal{U} \hookrightarrow \mathcal{X}$ be an open substack, $p : \mathcal{X}^{top} \rightarrow \mathbb{Z}$ be a monotone and comonotone perversity, and $\mathcal{F} \in D^{p, \geq 0}(\mathcal{U}) \cap D_{coh}^b(\mathcal{U})$. Consider $j_*(\mathcal{F}) \in D_{qcoh}^b(\mathcal{X})$, and let $n = \min_{x \notin \mathcal{U}^{top}} p(x)$. Then $\tau_{\leq n-2}^{stand}(j_*(\mathcal{F}))$ has coherent cohomology. Recall that $\tau_{\leq n-2}^{stand}$ is the truncation functor for the usual t -structure.*

Proof. Passing to a presentation of $\mathcal{X}$, we can assume that $X = \mathcal{X}$ is a scheme and $U = \mathcal{U}$ is an open subscheme of X . Let $\tilde{\mathcal{F}} \in D_{coh}^b(X)$ be any extension of $\mathcal{F}$ (see Lemma 2.12). Truncating $\tilde{\mathcal{F}}$ with respect to the perverse t-structure, we can achieve that $\tilde{\mathcal{F}} \in D^{p, \geq 0}(X)$. Consider the closed embedding $i : X^{top} - U^{top} \hookrightarrow X^{top}$. It yields a distinguished triangle

$$i_* i^!(\tilde{\mathcal{F}}) \rightarrow \tilde{\mathcal{F}} \rightarrow j_*(\mathcal{F}) \rightarrow i_* i^!(\tilde{\mathcal{F}})[1].$$

Since $\tilde{\mathcal{F}}$ has coherent cohomology, it is enough to show that

$$\tau_{\leq n-2}^{stand}(i_* i^!(\tilde{\mathcal{F}})[1]) \in D_{coh}^b(X).$$

However, the assumption $\tilde{\mathcal{F}} \in D^{p, \geq 0}(X)$ implies that for any closed subscheme $Z \subset X$ with $Z^{top} = X^{top} - U^{top}$, we have

$$i_Z^!(\tilde{\mathcal{F}}) \in D^{p, \geq 0}(Z) \subset D_{coh}^{\geq n}(Z).$$

Hence

$$i_* i^!(\tilde{\mathcal{F}}) \in D_{qcoh}^{\geq n}(X),$$

and $\tau_{\leq n-2}^{stand}(i_* i^!(\tilde{\mathcal{F}})[1]) = 0$. $\square$

Remark 3.13. This statement and idea of proof of Corollary 3.12 are copied from Deligne's message to the second author. (Possible mistakes belong to the authors.) It was pointed out to us by Deligne that Corollary 3.12 is equivalent to the Grothendieck Finiteness Theorem, [11], VIII.2.1.

4. COHERENT IC-SHEAVES

4.1. IC-extension. Suppose now that $p : \mathcal{X}^{top} \rightarrow \mathbb{Z}$ is a monotone and comonotone perversity function. As before, $\mathcal{X}$ is a semi-separated Noetherian stack admitting a dualizing sheaf. Denote by $\mathcal{P} = \mathcal{P}_{\mathcal{X}} = \mathcal{P}_{\mathcal{X}, p} \subset D_{coh}^b(\mathcal{X})$ the core of the t-structure constructed in the previous section.

Let $\mathbf{Z} \subset \mathcal{X}^{top}$ be a closed subset. Define auxiliary perversities $p^{\pm} = p_{(\mathbf{Z})}^{\pm} : \mathcal{X}^{top} \rightarrow \mathbb{Z}$ by

$$p^-(x) = \begin{cases} p(x), & x \notin \mathbf{Z} \\ p(x) - 1, & x \in \mathbf{Z}, \end{cases} \quad p^+(x) = \begin{cases} p(x), & x \notin \mathbf{Z} \\ p(x) + 1, & x \in \mathbf{Z}. \end{cases}$$

Lemma 4.1. *Let $\mathcal{F} \in \mathcal{P}_{\mathcal{X}}$.*

(a) *The following conditions are equivalent:*

- (i) $\mathcal{F} \in D^{p^-, \leq 0}(\mathcal{X})$.
- (ii) $i_{\mathcal{Z}}^*(\mathcal{F}) \in D^{p_{\mathcal{Z}}, < 0}(\mathcal{Z})$ for any closed substack $\mathcal{Z} \subset \mathcal{X}$ with $\mathcal{Z}^{top} \subset \mathbf{Z}$.
- (iii) $\text{Hom}(\mathcal{F}, \mathcal{G}) = 0$ for all $\mathcal{G} \in \mathcal{P}$ such that $\text{supp } \mathcal{G} \subset \mathbf{Z}$.

(b) *The following conditions are equivalent:*

- (i) $\mathcal{F} \in D^{p^+, \geq 0}(\mathcal{X})$.
- (ii) $i_{\mathcal{Z}}^!(\mathcal{F}) \in D^{p_{\mathcal{Z}}, > 0}(\mathcal{Z})$ for any closed substack $\mathcal{Z} \subset \mathcal{X}$ with $\mathcal{Z}^{top} \subset \mathbf{Z}$.
- (iii) $\text{Hom}(\mathcal{G}, \mathcal{F}) = 0$ for all $\mathcal{G} \in \mathcal{P}$ such that $\text{supp } \mathcal{G} \subset \mathbf{Z}$.

Proof. We prove only (a); the proof of (b) is completely analogous. The equivalence between (ai) and (aii) follows from Lemma 2.18. Let us show (aaii) $\iff$ (aiii).

Let $\mathcal{Z}$ be a closed substack with $\mathcal{Z}^{top} \subset \mathbf{Z}$. Since $\mathcal{F} \in \mathcal{P}_{\mathcal{X}}$,

$$i_{\mathcal{Z}}^*(\mathcal{F}) \in D^{p_{\mathcal{Z}}, \leq 0}(\mathcal{Z}).$$

On the other hand, given any triangulated category D with a t -structure and an object $A \in D^{\leq 0}$, it is clear that $A \in D^{< 0}$ if and only if $\text{Hom}(A, B) = 0$ for all B in the core of the t -structure. Therefore, $i_{\mathcal{Z}}^*(\mathcal{F}) \in D^{p_{\mathcal{Z}}, < 0}(\mathcal{Z})$ if and only if

$$\text{Hom}(\mathcal{F}, i_{\mathcal{Z}*}(\mathcal{G})) = \text{Hom}(i_{\mathcal{Z}}^*(\mathcal{F}), \mathcal{G}) = 0 \quad \text{for all } \mathcal{G} \in \mathcal{P}_{\mathcal{Z}}.$$

Now the statement follows from Lemma 2.13. $\square$

Suppose now that $j : \mathcal{U} \hookrightarrow \mathcal{X}$ is an open substack, and consider the perversities $p^{\pm} = p_{(\mathcal{X} - \mathcal{U})}^{\pm}$. Define a full subcategory $\mathcal{P}_{!*}(\mathcal{U}) \subset \mathcal{P}_{\mathcal{X}}$ by

$$\mathcal{P}_{!*}(\mathcal{U}) = D^{p^-, \leq 0}(\mathcal{X}) \cap D^{p^+, \geq 0}(\mathcal{X}).$$

Theorem 4.2. *Suppose that $p(x') > p(x)$, $\overline{p}(x') > \overline{p}(x)$ for any $x \in \mathcal{U}^{top}$, $x' \in \overline{\{x\}} - \mathcal{U}^{top}$. Then j^* induces an equivalence between $\mathcal{P}_{!*}(\mathcal{U})$ and $\mathcal{P}_{\mathcal{U}}$.*

The inverse equivalence is denoted by

$$j_{!*} : \mathcal{P}_{\mathcal{U}} \rightarrow \mathcal{P}_{!*}(\mathcal{U}) \subset \mathcal{P}_{\overline{\mathcal{U}}},$$

and is called the functor of minimal (or Goresky-MacPherson, or IC) extension.

Proof. By the hypotheses, both p^- and p^+ are monotone and comonotone perversities on $\mathcal{X}^{top}$. Hence they define t -structures on $D_{coh}^b(\mathcal{X})$; let τ^- , τ^+ be the corresponding truncation functors.

Define a functor $J_{!*}$ from $D_{coh}^b(\mathcal{X})$ to itself by

$$J_{!*} = \tau_{\leq 0}^- \circ \tau_{\geq 0}^+.$$

We need the following claim.

Lemma 4.3. (a) $J_{!*}$ takes values in $\mathcal{P}_{!*}(\mathcal{U})$.

(b) If a morphism $f : \mathcal{F} \rightarrow \mathcal{G}$ in $D_{coh}^b(\mathcal{X})$ is such that $f|_{\mathcal{U}}$ is an isomorphism, then $J_{!*}(f)$ is an isomorphism.

Proof. Fix $\mathcal{F} \in D_{coh}^b(\mathcal{X})$. By construction, $J_{!*}(\mathcal{F}) \in D^{p^-, \leq 0}(\mathcal{X})$. Set

$$\mathcal{F}_1 = \tau_{\geq 0}^+ \mathcal{F}.$$

Then $J_{!*}(\mathcal{F})$ fits in a distinguished triangle

$$(\tau_{> 0}^- \mathcal{F}_1)[-1] \rightarrow J_{!*}(\mathcal{F}) \rightarrow \mathcal{F}_1 \rightarrow \tau_{> 0}^- \mathcal{F}_1.$$

Since $\mathcal{F}_1 \in D^{p^+, \geq 0}(\mathcal{X})$ and $(\tau_{> 0}^- \mathcal{F}_1)[-1] \in D^{p^-, \geq 2}(\mathcal{X}) \subset D^{p^+, \geq 0}(\mathcal{X})$, part (a) follows.

Let f be as in (b). Then $J_{!*}(f)$ is a morphism in $\mathcal{P}_{!*}(\mathcal{U})$ that is an isomorphism over $\mathcal{U}$. Therefore, its kernel and cokernel are objects of $\mathcal{P}_{\mathcal{X}}$ supported by $\mathcal{X} - \mathcal{U}$. But by Lemma 4.1, $J_{!*}(\mathcal{F})$ and $J_{!*}(\mathcal{G})$ have neither subobjects nor quotients supported by $\mathcal{X} - \mathcal{U}$. $\square$

We now finish the proof of Theorem 4.2. By Lemmas 2.12 and 4.3, the functor $J_{!*}$ decomposes as

$$D_{coh}^b(\mathcal{X}) \xrightarrow{j^*} D_{coh}^b(\mathcal{U}) \rightarrow \mathcal{P}_{!*}(\mathcal{U})$$

for a canonically defined functor $\tilde{j}_{!*} : D_{coh}^b(\mathcal{U}) \rightarrow \mathcal{P}_{!*}(\mathcal{U})$. Set $j_{!*} = \tilde{j}_{!*}|_{\mathcal{P}_{\mathcal{U}}}$. By construction, $j^* \circ j_{!*} = id_{\mathcal{P}_{\mathcal{U}}}$ canonically. Also $j_{!*} \circ j^*|_{\mathcal{P}_{!*}(\mathcal{U})} = id$ canonically, because $J_{!*}|_{\mathcal{P}_{!*}(\mathcal{U})} = id$. Thus j^* and $j_{!*}$ are inverse equivalences between $\mathcal{P}_{!*}(\mathcal{U})$ and $\mathcal{P}_{\mathcal{U}}$. $\square$

Remark. Note that the hypotheses of Theorem 4.2 are always satisfied if p is strictly monotone and strictly comonotone.

We need some elementary properties of the minimal extension. Let us keep the hypotheses of Theorem 4.2.

Lemma 4.4. *For any $\mathcal{F} \in \mathcal{P}_{\mathcal{X}}$, $j_{!*}(\mathcal{F}|_{\mathcal{U}})$ is a subquotient of $\mathcal{F}$ in the abelian category $\mathcal{P}_{\mathcal{X}}$.*

Proof. Since $\mathcal{F} \in \mathcal{P}_{\mathcal{X}}$, we have

$$\begin{aligned}\tau_{\geq 0}^+ \mathcal{F} &\in D^{p^+, \geq 0}(\mathcal{X}) \subset D^{p, \geq 0}(\mathcal{X}) \\ \tau_{< 0}^+ \mathcal{F} &\in D^{p^+, < 0}(\mathcal{X}) \subset D^{p, \leq 0}(\mathcal{X}).\end{aligned}$$

Now the distinguished triangle

$$\tau_{< 0}^+ \mathcal{F} \rightarrow \mathcal{F} \rightarrow \tau_{\geq 0}^+ \mathcal{F} \rightarrow \tau_{< 0}^+ \mathcal{F}[1]$$

implies that $\tau_{\geq 0}^+(\mathcal{F}), \tau_{< 0}^+(\mathcal{F}) \in \mathcal{P}_{\mathcal{X}}$. In other words, $\tau_{\geq 0}^+(\mathcal{F})$ is a quotient of $\mathcal{F}$.

Repeating the argument, we see that

$$J_{!*}(\mathcal{F}) = \tau_{\leq 0}^- \tau_{\geq 0}^+ \mathcal{F} = j_{!*}(\mathcal{F}|_{\mathcal{U}}) \in \mathcal{P}_{\mathcal{X}}$$

is a subobject of $\tau_{\geq 0}^+(\mathcal{F})$, as required. $\square$

Lemma 4.5. *Let $\bar{i} : \mathcal{Y} \hookrightarrow \mathcal{X}$ be a closed substack. Set $\mathcal{V} = \mathcal{U} \cap \mathcal{Y}$. Then*

$$j_{!*} \circ i_* \simeq \bar{i}_* \circ j'_{!*}.$$

Here i is the closed embedding $\mathcal{V} \hookrightarrow \mathcal{U}$, and j' is the open embedding $\mathcal{V} \hookrightarrow \mathcal{Y}$.

Proof. Indeed, $\bar{i}_*(\mathcal{P}_{!*}(\mathcal{V})) \subset \mathcal{P}_{!*}(\mathcal{U})$ by Lemma 3.3(c). $\square$

Definition 4.6. Let $j : \mathcal{U} \hookrightarrow \mathcal{X}$ be a locally closed embedding that satisfies the hypotheses of Theorem 4.2:

$$p(x') > p(x) \quad \text{and} \quad \bar{p}(x') > \bar{p}(x) \quad (x \in \mathcal{U}^{top}, x' \in \overline{\{x\}} - \mathcal{U}^{top}).$$

The *minimal (or Goresky-MacPherson, or IC) extension functor*

$$j_{!*} : \mathcal{P}_{\mathcal{U}} \rightarrow \mathcal{P}_{\mathcal{X}}$$

is given by $i_* \circ j'_{!*}$, where $i : \mathcal{Z} \hookrightarrow \mathcal{X}$ is a closed substack such that $j' : \mathcal{U} \hookrightarrow \mathcal{Z}$ is an open embedding. By Lemma 4.5, the functor is independent of the choice of $\mathcal{Z}$.

4.2. Irreducible perverse sheaves. For the rest of the paper, we assume that $\mathcal{X}$ is a semi-separated stack of finite type over a field k .

Let x be a point of $\mathcal{X}$. Let us represent it by a morphism $\xi : \text{Spec } K \rightarrow \mathcal{X}$ for a field extension $K \supset k$. Denote by G_{ξ} the automorphism group of ξ ; it is a K -group scheme of finite type. Note that $\dim(G_{\xi})$ depends only on $x \in \mathcal{X}^{top}$, and that it is an upper semi-continuous function of x (because G_{ξ} can be thought of as the fiber of the diagonal morphism $\mathcal{X} \rightarrow \mathcal{X} \times \mathcal{X}$ over (ξ, ξ)).

Example 4.7. Suppose $\mathcal{X} = X/G$, where X is a semi-separated scheme of finite type over k and G is an affine group scheme of finite type acting on X . A point $\xi : \text{Spec } K \rightarrow X$ defines a point of $\mathcal{X}$; the corresponding automorphism group is the stabilizer of ξ .

Denote by $\mathcal{X}_{lc}^{top} \subset \mathcal{X}^{top}$ the set of points $x \in \mathcal{X}^{top}$ that are defined over the algebraic closure $\bar{k} \supset k$. For any presentation $\pi : X \rightarrow \mathcal{X}$, $x \in \mathcal{X}^{top}$ is defined over $\bar{k}$ if and only if it can be lifted to a $\bar{k}$ -point of X , because π has finite type.

Lemma 4.8. *For $x \in \mathcal{X}^{top}$, the set $\{x\}$ is locally closed if and only if $x \in \mathcal{X}_{lc}^{top}$.*

Proof. Suppose $\{x\}$ is locally closed. Then there is a locally closed substack $\mathcal{Y} \subset \mathcal{X}$ such that $\mathcal{Y}^{top} = \{x\}$. However, $\mathcal{Y}_{lc}^{top} \neq \emptyset$ (because its presentation has $\bar{k}$ -points), and therefore x is defined over $\bar{k}$.

Suppose x is defined over $\bar{k}$. Passing to a locally closed subset of $\mathcal{X}$, we may assume that $\dim(G_\eta)$ is constant for all $\eta : \text{Spec } K \rightarrow \mathcal{X}$. But then $x \in \mathcal{X}^{top}$ has minimal dimension (equal to $-\dim(G_\eta)$), and therefore $\{x\} \subset \mathcal{X}^{top}$ is closed. (For another proof, note that $\{x\} \subset \mathcal{X}^{top}$ is the image of a morphism $\text{Spec } K \rightarrow \mathcal{X}$, where $K \supset k$ is a finite extension. This morphism is of finite type, therefore $\{x\}$ is a constructible set. This implies $\{x\}$ is locally closed.) $\square$

Lemma 4.9. *Suppose $x \in \mathcal{X}^{top} - \mathcal{X}_{lc}^{top}$. Then there is a point $x' \in \overline{\{x\}}$ of dimension $\dim(x') = \dim(x) - 1$.*

Proof. For a presentation $\pi : X \rightarrow \mathcal{X}$, consider preimage $\pi^{-1}(x) \subset X^{top}$. Choose a point $y \in \pi^{-1}(x)$ that is closed in the induced topology, and set $Y = \overline{\{y\}} \subset X$. Note that $\dim Y > 0$, otherwise x would be defined over $\bar{k}$.

The restriction $\pi|_Y : Y \rightarrow \mathcal{X}$ is of finite type; therefore, we can replace Y with its open subset such that the dimension of fibers of $\pi|_Y$ is constant and equal to $\dim Y - \dim x$. (Actually, it is easy to see that $\dim Y - \dim x = \dim G_\xi$, where ξ represents x .) Take a point $y' \in Y$ such that $\text{codim } \overline{\{y'\}} = 1$, and set $x' = \pi(y')$. We claim that $\dim x' = \dim x - 1$.

Indeed, $\dim x' < \dim x$, because $x' \in \overline{\{x\}}$ and $x' \neq x$. On the other hand,

$$\dim x' = \dim \overline{\{x'\}} \geq \dim(\pi^{-1}(\overline{\{x'\}}) \cap Y) - (\dim Y - \dim x) = \dim x - 1.$$

$\square$

Remark. Lemmas 4.8 and 4.9 are also easy to derive from [16, Corollary 10.8]. If $\mathcal{X}$ is of the form X/G , Lemmas 4.8 and 4.9 follow from Rosenlicht's Theorem ([17], see also exposition in [20]).

By Lemma 4.8, a point $x \in \mathcal{X}_{lc}^{top}$ determines a reduced locally closed substack $\mathcal{G}_x \subset \mathcal{X}$: the *residue gerbe* of x . We denote the embedding $\mathcal{G}_x \hookrightarrow \mathcal{X}$ by $j^{(x)}$.

Remark 4.10. Let k_x be the residue field of x (see [16, Chapter 11]); since $x \in \mathcal{X}_{lc}^{top}$, $k_x \supset k$ is a finite extension. Suppose that x is defined over k_x so that x is the image of a morphism

$$\xi : \text{Spec } k_x \rightarrow \mathcal{X}.$$

Then ξ provides a presentation of $\mathcal{G}_x$, and $\mathcal{G}_x$ is isomorphic to the classifying stack of G_ξ . In particular, quasi-coherent sheaves on $\mathcal{G}_x$ identify with representations of G_ξ .

In general, we can always represent x as the image of $\xi : \text{Spec } K \rightarrow \mathcal{X}$, where K is a finite extension of k_x . Then the extension of scalars

$$\mathcal{G}_x \times_{\text{Spec } k_x} \text{Spec } K$$

identifies with the classifying stack of G_ξ , so in a sense $\mathcal{G}_x$ is a k_x -form of this classifying stack.

Let us fix a monotone and comonotone perversity $p : \mathcal{X}^{top} \rightarrow \mathbb{Z}$. Recall that

$$\mathcal{P}_{\mathcal{X}} \subset D_{coh}^b(\mathcal{X})$$

is the kernel of the associated t-structure.

Proposition 4.11. *For $\mathcal{F} \in D_{coh}^b(\mathcal{X}^{top})$ the following statements are equivalent:*

- (a) $\mathcal{F}$ is an irreducible object of $\mathcal{P}_{\mathcal{X}}$.
- (b) There exists a point $x \in \mathcal{X}_{lc}^{top}$ such that $p(x) < p(y)$, $\overline{p}(x) < \overline{p}(y)$ for any $y \in \overline{\{x\}} - \{x\}$ and an irreducible coherent sheaf L on $\mathcal{G}_x$ such that $\mathcal{F} = j_{!*}^{(x)}(L[p(x)])$.

Proof. (b) $\Rightarrow$ (a). Suppose that $\mathcal{F}$ has a proper subobject $\mathcal{F}'$. Then $\text{supp}(\mathcal{F}') \subset \overline{\{x\}}$, so by Lemma 2.13, $\mathcal{F}'$ can be obtained by a direct image from a closed substack $\mathcal{Y} \subset \mathcal{X}$ with generic point x . Replacing $\mathcal{X}$ with $\mathcal{Y}$, we may assume that $x \in \mathcal{X}^{top}$ is dense, and therefore open.

Let $j : \mathcal{U} \hookrightarrow \mathcal{X}$ be the open substack with $\mathcal{U}^{top} = \{x\}$; its maximal reduced substack is $\mathcal{G}_x$. By Lemma 4.5,

$$\mathcal{F} = j_{!*}i_*(L[p(x)]),$$

where $i : \mathcal{G}_x \hookrightarrow \mathcal{U}$ is the natural closed embedding. Clearly, $L[p(x)] \in \mathcal{P}_{\mathcal{U}}$ is irreducible (note that $\mathcal{P}_{\mathcal{U}} = \text{Coh}(\mathcal{U})[p(x)]$). Now irreducibility of $\mathcal{F}$ follows from Lemma 4.1.

(a) $\Rightarrow$ (b). Let x be a generic point of $\text{supp}(\mathcal{F})$. Since $\mathcal{F}$ is irreducible, we have

$$\text{Hom}(\mathcal{F}, i_{\mathcal{Z}*}(\mathcal{G})) = 0, \quad \text{Hom}(i_{\mathcal{Z}*}(\mathcal{G}), \mathcal{F}) = 0$$

for any closed substack $i_{\mathcal{Z}} : \mathcal{Z} \hookrightarrow \mathcal{X}$ not containing x and any $\mathcal{G} \in \mathcal{P}_{\mathcal{Z}}$.

Thus Lemma 4.1 implies that

$$i_{\mathcal{Z}}^*(\mathcal{F}) \in D^{p_{\mathcal{Z}}, < 0}(\mathcal{Z}), \quad i_{\mathcal{Z}}^!(\mathcal{F}) \in D^{p_{\mathcal{Z}}, > 0}(\mathcal{Z}).$$

In particular, this shows that $\text{supp } \mathcal{F}$ is irreducible (otherwise we can take $\mathcal{Z}$ to be an irreducible component of $\text{supp } \mathcal{F}$ that does not contain x).

Now take $y \in \text{supp } \mathcal{F} = \overline{\{x\}}$, $y \neq x$. Then $p(x) < p(y)$: otherwise, the coherent sheaf $\mathcal{H}^{p(x)}(\mathcal{F})$ has a nonzero fiber at x , but has zero fiber at y , which contradicts the Nakayama Lemma. A dual argument shows that $\overline{p}(x) < \overline{p}(y)$. In particular, $\dim(y) < \dim(x) - 1$. By Lemma 4.9, $x \in \mathcal{X}_{lc}^{top}$. By Lemma 2.13, $\mathcal{F}$ can be obtained by a direct image from a closed substack $\mathcal{Y} \subset \mathcal{X}$ with generic point x .

Consider the open substack $j : \mathcal{U} \hookrightarrow \mathcal{Y}$ such that $\mathcal{U}^{top} = \{x\}$. Then Theorem 4.2 implies that

$$\mathcal{F} = j_{!*}(L'[p(x)]) \quad (L' \in \text{Coh}(\mathcal{U})).$$

However, L' has to be irreducible, and, in particular, it is supported by the reduced substack $\mathcal{G}_x \subset \mathcal{U}$. $\square$

Example 4.12. Oftentimes, Proposition 4.11 can be made more explicit using Remark 4.10. For instance, suppose $k = \overline{k}$. Then every point $x \in \mathcal{X}_{lc}^{top}$ is defined over k , and an irreducible coherent sheaf L on $\mathcal{G}_x$ is simply an irreducible representation of G_{ξ} . Here $\xi : \text{Spec } k \rightarrow \mathcal{X}$ is the k -point representing x .

Corollary 4.13. *Suppose that $\mathcal{X}$ has finitely many points. If the perversity $p : \mathcal{X}^{top} \rightarrow \mathbb{Z}$ is strictly monotone and strictly comonotone, then the category $\mathcal{P}_{\mathcal{X}}$ is Artinian and Noetherian.*

Proof. In this case, every point $x \in \mathcal{X}^{top}$ is locally closed, so $x \in \mathcal{X}_{lc}^{top}$, and $j_{!*}^{(x)}$ is defined (by the hypotheses). By induction in the number of points, one can deduce that the irreducible objects generate the triangulated category $D_{coh}^b(\mathcal{X})$. This implies that $\mathcal{P}_{\mathcal{X}}$ is Artinian and Noetherian. $\square$

Corollary 4.14. *Suppose that $\mathcal{X}$ has finitely many points and that the perversity $p : \mathcal{X}^{top} \rightarrow \mathbb{Z}$ is strictly monotone and strictly comonotone. Then classes of irreducible objects $\mathcal{F} \in \mathcal{P}_{\mathcal{X}}$ (described in Proposition 4.11) form a basis in $K^0(\text{Coh}(\mathcal{X}))$.*

Proof. The derived category $D_{coh}^b(\mathcal{X})$ has two bounded t-structures: the standard t-structure and the perverse t-structure. For the hearts of these t-structures, we get an identification

$$K^0(\text{Coh}(\mathcal{X})) = K^0(\mathcal{P}_{\mathcal{X}}).$$

Now the claim follows from Corollary 4.13. $\square$

Example 4.15. Let G be a semisimple group over a field of characteristic 0, or of large finite characteristic, and let $\mathcal{N} \subset G$ be the subvariety of unipotent elements. Then G acts on $\mathcal{N}$ by conjugation, and this action has a finite number of orbits. Moreover, dimensions of orbits are known to be even. Points x of the stack $\mathcal{N}/G$ correspond to G -orbits $O \subset \mathcal{N}$, and we can define the *middle perversity* $p : (\mathcal{N}/G)^{top}$ by

$$p(x) = -\frac{\dim(O)}{2}.$$

Obviously, p is strictly monotone and comonotone, hence by Proposition 4.13 the heart of the corresponding t-structure is Artinian and Noetherian. See [4], [5] for more information on this example.

Example 4.16. Let G be as above and let Gr denote the affine Grassmannian of G . This is an ind-scheme acted upon by the (infinite dimensional) group scheme G_O . The orbits are in bijection with the set of dominant coweights of G . Moreover, it is known that orbits in a given connected component of Gr have dimension of a fixed parity. Thus the perversity function $p(x) = -[\frac{\dim(O)}{2}]$ is monotone and comonotone, and (a straightforward generalization of) Corollary 4.13 applies. See [6] for more information on this example.

REFERENCES

- [1] A. A. Beilinson, J. Bernstein, and P. Deligne. Faisceaux pervers. In *Analysis and topology on singular spaces, I (Luminy, 1981)*, volume 100 of *Astérisque*, pages 5–171. Soc. Math. France, Paris, 1982.
- [2] J. Bernstein and V. Lunts. *Equivariant sheaves and functors*, volume 1578 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin, 1994.
- [3] R. Bezrukavnikov. Perverse coherent sheaves (after Deligne). [arXiv:math/0010089](https://arxiv.org/abs/math/0010089).
- [4] R. Bezrukavnikov. Quasi-exceptional sets and equivariant coherent sheaves on the nilpotent cone. *Represent. Theory*, 7:1–18, 2003.
- [5] R. Bezrukavnikov. Cohomology of tilting modules over quantum groups and t-structures on derived categories of coherent sheaves. *Invent. Math.*, 166(2):327–357, 2006.
- [6] R. Bezrukavnikov, M. Finkelberg, and I. Mirković. Equivariant homology and K-theory of affine Grassmannians and Toda lattices. *Compos. Math.*, 141(3):746–768, 2005.
- [7] M. Bökstedt and A. Neeman. Homotopy limits in triangulated categories. *Compositio Math.*, 86(2):209–234, 1993.
- [8] P. Deligne. Perversité cohérente. Handwritten notes.
- [9] P. Deligne. Théorie de Hodge. III. *Inst. Hautes Études Sci. Publ. Math.*, (44):5–77, 1974.

- [10] O. Gabber. Notes on some t -structures. In *Geometric aspects of Dwork theory. Vol. I, II*, pages 711–734. Walter de Gruyter GmbH & Co. KG, Berlin, 2004.
- [11] A. Grothendieck. *Cohomologie locale des faisceaux cohérents et théorèmes de Lefschetz locaux et globaux (SGA 2)*. North-Holland Publishing Co., Amsterdam, 1968. Séminaire de Géométrie Algébrique du Bois-Marie, 1962, Advanced Studies in Pure Mathematics, Vol. 2.
- [12] R. Hartshorne. *Residues and duality*. Lecture notes of a seminar on the work of A. Grothendieck, given at Harvard 1963/64. With an appendix by P. Deligne. Lecture Notes in Mathematics, No. 20. Springer-Verlag, Berlin, 1966.
- [13] M. Kashiwara. t -structures on the derived categories of holonomic D -modules and coherent O -modules. *Mosc. Math. J.*, 4(4):847–868, 981, 2004.
- [14] F. Knop, H. Kraft, D. Luna, and T. Vust. Local properties of algebraic group actions. In *Algebraische Transformationsgruppen und Invariantentheorie*, volume 13 of *DMV Sem.*, pages 63–75. Birkhäuser, Basel, 1989.
- [15] Y. Laszlo and M. Olsson. The six operations for sheaves on Artin stacks. I. Finite coefficients. *Publ. Math. Inst. Hautes Études Sci.*, (107):109–168, 2008.
- [16] G. Laumon and L. Moret-Bailly. *Champs algébriques*, volume 39 of *Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics [Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics]*. Springer-Verlag, Berlin, 2000.
- [17] M. Rosenlicht. A remark on quotient spaces. *An. Acad. Brasil. Ci.*, 35:487–489, 1963.
- [18] B. Saint-Donat. Techniques de descente cohomologique. In *Théorie des topos et cohomologie étale des schémas. Tome 2*, Lecture Notes in Mathematics, Vol. 270. Springer-Verlag, Berlin, 1972. Séminaire de Géométrie Algébrique du Bois-Marie 1963–1964 (SGA 4), Dirigé par M. Artin, A. Grothendieck et J. L. Verdier. Exposé V^{bis} .
- [19] H. Sumihiro. Equivariant completion. *J. Math. Kyoto Univ.*, 14:1–28, 1974.
- [20] È. B. Vinberg and V. L. Popov. Invariant theory. In *Algebraic geometry, 4 (Russian)*, Itogi Nauki i Tekhniki, pages 137–314, 315. Akad. Nauk SSSR Vsesoyuz. Inst. Nauchn. i Tekhn. Inform., Moscow, 1989.