

On the geometry of the connection with totally skew-symmetric torsion on almost complex manifolds with Norden metric

Dimitar Mekerov

November 20, 2018

University of Plovdiv, Faculty of Mathematics and Informatics, 236 Bulgaria Blvd., 4003 Plovdiv, Bulgaria, mircho@uni-plovdiv.bg

Abstract

We consider an almost complex manifold with Norden metric (i. e. a metric with respect to which the almost complex structure is an anti-isometry). On such a manifold we study a linear connection preserving the almost complex structure and the metric and having a totally skew symmetric torsion tensor (i. e. a 3-form). We prove that if a non-Kähler almost complex manifold with Norden metric admits such connection then the manifold is quasi-Kählerian (i. e. has non-integrable almost complex structure). We prove that this connection is unique, determine its form, and construct an example of it on a Lie group. We consider the case when the manifold admits a connection with parallel totally skew-symmetric torsion and the case when such connection has a Kähler curvature tensor. We get necessary and sufficient conditions for an isotropic Kähler manifold with Norden metric.

Mathematics Subject Classification (2000): Primary 53C15, 53B05; Secondary 53C50, 22E60.

Key words: Norden metric, almost complex manifold, indefinite metric, linear connection, Bismut connection, KT connection, skew-symmetric torsion, parallel torsion.

1 Introduction

There is a strong interest in the metric connections with totally skew-symmetric torsion tensor (3-form). These connections arise in a natural way in theoretical and mathematical physics. For example, such a connection is of particular interest in string theory [1]. In mathematics this connection was used by Bismut to prove the local index theorem for non-Kähler Hermitian manifolds [2]. A connection with totally skew-symmetric torsion tensor is called a KT connection

by physicists, and among mathematicians this connection is known as a Bismut connection.

According to Gauduchon [3], on any non-Kähler Hermitian manifold, there exists a unique Hermitian connection (i. e. one preserving the almost complex structure and the metric), whose torsion tensor is totally skew-symmetric.

The goal of the present work is to solve the analogous problem for almost complex manifolds with Norden metric. We build upon the results in [4], where we have considered a connection with totally skew-symmetric torsion tensor in the case of quasi-Kähler manifolds with Norden metric.

The main results in the present paper are contained in Theorem (3.1). We prove that if a non-Kähler almost complex manifold with Norden metric admits a linear connection ∇' preserving the almost complex structure and the metric and having totally skew-symmetric tensor, then the manifold is quasi-Kählerian and the connection ∇' is unique. We get the form of ∇' .

We construct a 4-parametric family of 4-dimensional quasi-Kähler manifolds with Norden metric by a Lie group, and derive the connection ∇' for an arbitrary manifold of this family.

We consider the case when ∇' has a parallel torsion and obtain a relation between the scalar curvatures of this connection and the Levi-Civita connection. We show that the manifold is isotropic Kählerian iff these curvatures are equal.

We prove that if the manifold admits a parallel connection with a Kähler curvature tensor, then the manifold is isotropic Kählerian.

2 Preliminaries

Let (M, J, g) be a $2n$ -dimensional *almost complex manifold with Norden metric*, i. e.

$$J^2x = -x, \quad g(Jx, Jy) = -g(x, y),$$

for all differentiable vector fields x, y on M . The *associated metric* \tilde{g} of g on M , given by $\tilde{g}(x, y) = g(x, Jy)$, is a Norden metric, too. The signature of both metrics is necessarily (n, n) .

Further, x, y, z, w will stand for arbitrary differentiable vector fields on M (or vectors in the tangent space of M at an arbitrary point $p \in M$).

The Levi-Civita connection of g is denoted by ∇ . The tensor filed F of type $(0, 3)$ on M is defined by

$$F(x, y, z) = g((\nabla_x J)y, z). \quad (2.1)$$

It has the following properties [5]:

$$F(x, y, z) = F(x, z, y) = F(x, Jy, Jz), \quad F(x, Jy, z) = -F(x, y, Jz).$$

In [6], the considered manifolds are classified into eight classes with respect to F . The class \mathcal{W}_0 of the *Kähler manifolds with Norden metric* is contained in each of the other seven classes. It is determined by the condition $F(x, y, z) = 0$, which is equivalent to $\nabla J = 0$.

The condition

$$\mathfrak{S}_{x,y,z} F(x, y, z) = 0, \quad (2.2)$$

where $\mathfrak{S}_{x,y,z}$ is the cyclic sum over x, y, z , characterizes the class \mathcal{W}_3 of the *quasi-Kähler manifolds with Norden metric*. This is the only class of manifolds with non-integrable almost complex structure J .

Let $\{e_i\}$ ($i = 1, 2, \dots, 2n$) be an arbitrary basis of the tangent space of M at a point $p \in M$. The components of the inverse matrix of g , with respect to this basis, are denoted by g^{ij} .

Following [7], the *square norm* $\|\nabla J\|^2$ of ∇J is defined in [8] by

$$\|\nabla J\|^2 = g^{ij} g^{ks} g((\nabla_{e_i} J) e_k, (\nabla_{e_j} J) e_s), \quad (2.3)$$

where it is proven that

$$\|\nabla J\|^2 = -2g^{ij} g^{ks} g((\nabla_{e_i} J) e_k, (\nabla_{e_s} J) e_j). \quad (2.4)$$

There, the manifold with $\|\nabla J\|^2 = 0$ is called an *isotropic-Kähler manifold with Norden metric*. It is clear that every Kähler manifold with Norden metric is isotropic-Kähler, but the inverse implication is not always true.

Let R be the curvature tensor of ∇ , i. e. $R(x, y)z = \nabla_x \nabla_y z - \nabla_y \nabla_x z - \nabla_{[x,y]} z$. The corresponding (0,4)-tensor is determined by $R(x, y, z, w) = g(R(x, y)z, w)$. The Ricci tensor ρ and the scalar curvature τ with respect to ∇ are defined by

$$\rho(y, z) = g^{ij} R(e_i, y, z, e_j), \quad \tau = g^{ij} \rho(e_i, e_j).$$

A tensor L of type (0,4) with the properties

$$L(x, y, z, w) = -L(y, x, z, w) = -L(x, y, w, z), \quad (2.5)$$

$$\mathfrak{S}_{x,y,z} L(x, y, z, w) = 0 \quad (\text{the first Bianchi identity}) \quad (2.6)$$

is called a *curvature-like tensor*. Moreover, if the curvature-like tensor L has the property

$$L(x, y, Jz, Jw) = -L(x, y, z, w), \quad (2.7)$$

it is called a *Kähler tensor* [9].

Let ∇' be a linear connection with a tensor Q of the transformation $\nabla \rightarrow \nabla'$ and a torsion tensor T , i. e.

$$\nabla'_x y = \nabla_x y + Q(x, y), \quad T(x, y) = \nabla'_x y - \nabla'_y x - [x, y]. \quad (2.8)$$

The corresponding (0,3)-tensors are defined by

$$Q(x, y, z) = g(Q(x, y), z), \quad T(x, y, z) = g(T(x, y), z). \quad (2.9)$$

The symmetry of the Levi-Civita connection implies

$$T(x, y) = Q(x, y) - Q(y, x), \quad T(x, y) = -T(y, x). \quad (2.10)$$

A partial decomposition of the space \mathcal{T} of the torsion (0,3)-tensors T (i. e. $T(x, y, z) = -T(y, x, z)$) is valid on an almost complex manifold with Norden metric (M, J, g) : $\mathcal{T} = \mathcal{T}_1 \oplus \mathcal{T}_2 \oplus \mathcal{T}_3 \oplus \mathcal{T}_4$, where \mathcal{T}_i ($i = 1, 2, 3, 4$) are invariant orthogonal subspaces [10]. For the projection operators p_i of \mathcal{T} in \mathcal{T}_i it is established that:

$$\begin{aligned}
4p_1(x, y, z) &= T(x, y, z) - T(Jx, Jy, z) - T(Jx, y, Jz) - T(x, Jy, Jz), \\
4p_2(x, y, z) &= T(x, y, z) - T(Jx, Jy, z) + T(Jx, y, Jz) + T(x, Jy, Jz), \\
8p_3(x, y, z) &= 2T(x, y, z) - T(y, z, x) - T(z, x, y) - T(Jy, z, Jx) \\
&\quad - T(z, Jx, Jy) + 2T(Jx, Jy, z) - T(Jy, Jz, x) \\
&\quad - T(Jz, Jx, y) + T(y, Jz, Jx) + T(Jz, x, Jy), \\
8p_4(x, y, z) &= 2T(x, y, z) + T(y, z, x) + T(z, x, y) + T(Jy, z, Jx) \\
&\quad + T(z, Jx, Jy) + 2T(Jx, Jy, z) + T(Jy, Jz, x) \\
&\quad + T(Jz, Jx, y) - T(y, Jz, Jx) - T(Jz, x, Jy).
\end{aligned} \tag{2.11}$$

A linear connection ∇' on an almost complex manifold with Norden metric (M, J, g) is called a *natural connection* if $\nabla'J = \nabla'g = 0$. The last conditions are equivalent to $\nabla'g = \nabla'\tilde{g} = 0$. If ∇' is a linear connection with a tensor Q of the transformation $\nabla \rightarrow \nabla'$ on an almost complex manifold with Norden metric, then it is a natural connection iff the following conditions are valid:

$$F(x, y, z) = Q(x, y, Jz) - Q(x, Jy, z), \tag{2.12}$$

$$Q(x, y, z) = -Q(x, z, y). \tag{2.13}$$

Since $\nabla'g = 0$, equalities (2.10) and (2.11) imply

$$Q(x, y, z) = \frac{1}{2}\{T(x, y, z) - T(y, z, x) + T(z, x, y)\}, \tag{2.14}$$

which is the Hayden theorem ([11]).

Theorem 2.1 ([12]). *For a natural connection with a torsion tensor T on a quasi-Kähler manifold with Norden metric (M, J, g) , which is non-Kählerian, the following properties are valid*

$$p_2 \neq 0, \quad p_3 = 0. \tag{2.15}$$

□

3 Connection with totally skew-symmetric torsion

3.1 Main results

The main results in the present work are consist in the following

Theorem 3.1. *Let ∇' be a natural connection with a totally skew-symmetric torsion tensor on a non-Kähler almost complex manifold with Norden metric (M, J, g) . Then*

- (i) (M, J, g) is quasi-Kählerian;
- (ii) ∇' has a torsion tensor T in the class $\mathcal{T}_2 \oplus \mathcal{T}_4$;
- (iii) ∇' is unique and the tensor Q of the transformation $\nabla \rightarrow \nabla'$ is determined by

$$Q(x, y, z) = \frac{1}{4} \{F(x, Jy, z) - F(Jx, y, z) - 2F(y, Jx, z)\}. \quad (3.1)$$

Proof. Since ∇' has a totally skew-symmetric torsion tensor T , then we have

$$T(x, y, z) = -T(y, x, z) = -T(x, z, y) = -T(z, y, x). \quad (3.2)$$

From (2.14) and (3.2) it follows that for the tensor Q of the transformation $\nabla \rightarrow \nabla'$ it is valid

$$Q(x, y, z) = \frac{1}{2}T(x, y, z). \quad (3.3)$$

Since ∇' is a natural connection, then (2.12) holds and consequently

$$\mathfrak{S}_{x,y,z} F(x, y, z) = \mathfrak{S}_{x,y,z} \{Q(x, y, Jz) - Q(x, Jy, z)\}. \quad (3.4)$$

According to (3.3), Q is also 3-form and then we have

$$Q(x, y, Jz) = Q(y, Jz, x). \quad (3.5)$$

Equalities (3.4) and (3.5) imply (2.2), which completes the proposition (i).

According to Theorem (2.1), for the tensor T we have $p_2 \neq 0$ and $p_3 = 0$, i. e. $T \in \mathcal{T}_1 \oplus \mathcal{T}_2 \oplus \mathcal{T}_4$. Let us suppose that $T \in \mathcal{T}_1$. Then $T = p_1$ and from (2.11) it follows

$$3T(x, y, z) + T(Jx, Jy, z) + T(Jx, y, Jz) + T(x, Jy, Jz) = 0. \quad (3.6)$$

We substitute Jy for y and Jz for z in (3.6) and the obtained equality we add to (3.6). In the result we substitute Jz for z and according to (3.3) we get $Q(x, y, Jz) - Q(x, Jy, z) = 0$. Then, by (2.12) we obtain $F(x, y, z) = 0$, i. e. (M, J, g) is Kählerian, which is a contradiction. Therefore $T \notin \mathcal{T}_1$, i. e. $p_1 = 0$.

Combining (2.11), (3.2) and (3.3), we find that

$$p_4(x, y, z) = Q(x, y, z) + Q(Jx, Jy, z). \quad (3.7)$$

Let us suppose that $p_4 = 0$. Then from (3.7) we have

$$Q(x, y, z) + Q(Jx, Jy, z) = 0. \quad (3.8)$$

We substitute $y \leftrightarrow z$ in (3.8), and subtract the obtained equality from (3.8). In the result we apply (3.8) and the fact that Q is a 3-form. In such a way we obtain

$$Q(x, y, z) + Q(Jx, y, Jz) = 0. \quad (3.9)$$

The equalities (3.8) and (3.9) imply $Q(x, y, Jz) - Q(x, Jy, z) = 0$. Thus, according to (2.12), we obtain $F(x, y, z) = 0$, i. e. (M, J, g) is Kählerian, which is impossible. Therefore $p_4 \neq 0$.

Thereby we establish the following conditions for T : $p_1 = 0$, $p_2 \neq 0$, $p_3 = 0$, $p_4 \neq 0$. Hence $T \in \mathcal{T}_2 \oplus \mathcal{T}_4$, i. e. (ii) holds.

From (2.12), having in mind that Q is a 3-form, we obtain

$$\begin{aligned} & F(x, Jy, z) - F(Jx, y, z) - 2F(y, Jx, z) \\ &= Q(Jx, Jy, z) + Q(Jx, y, Jz) + Q(x, Jy, Jz) + 3Q(x, y, z). \end{aligned} \quad (3.10)$$

Since $p_1 = 0$, from (2.11) and (3.3) we have

$$Q(Jx, Jy, z) + Q(Jx, y, Jz) + Q(x, Jy, Jz) = Q(x, y, z). \quad (3.11)$$

The equalities (3.10) and (3.11) imply (3.1), which completes the proof of (iii). \square

According to (2.1), equality (3.1) is equivalent to

$$Q(x, y) = \frac{1}{4} \{ (\nabla_x J) Jy - (\nabla_{Jx} J) y - 2(\nabla_y J) Jx \}. \quad (3.12)$$

Remark 3.1. *In [13] it is found a 2-parametric family of natural connections on almost complex manifolds with Norden metric which contains the connection ∇' with totally skew-symmetric torsion tensor.*

3.2 An example

Let V be a real 4-dimensional vector space with a basis $\{E_i\}$. Let us consider a structure of a Lie algebra determined by the commutators $[E_i, E_j] = C_{ij}^k E_k$, where C_{ij}^k are structure constants satisfying the anti-commutativity condition $C_{ij}^k = -C_{ji}^k$ and the Jacobi identity $C_{ij}^k C_{ks}^l + C_{js}^k C_{ki}^l + C_{si}^k C_{kj}^l = 0$.

Let G be the associated connected Lie group and $\{X_i\}$ be a global basis for the left invariant vector fields that is induced by the basis $\{E_i\}$ of V . Then we have the decomposition

$$[X_i, X_j] = C_{ij}^k X_k. \quad (3.13)$$

Let us consider the almost complex manifold with Norden metric (G, J, g) , where

$$JX_1 = X_3, \quad JX_2 = X_4, \quad JX_3 = -X_1, \quad JX_4 = -X_2 \quad (3.14)$$

and

$$\begin{aligned} g(X_1, X_1) &= g(X_2, X_2) = -g(X_3, X_3) = -g(X_4, X_4) = 1, \\ g(X_i, X_j) &= 0 \quad \text{for } i \neq j. \end{aligned} \quad (3.15)$$

Because of (3.15) the following equality is valid

$$2g(\nabla_{X_i} X_j, X_k) = g([X_i, X_j], X_k) + g([X_k, X_i], X_j) + g([X_k, X_j], X_i). \quad (3.16)$$

We add the condition for the associated metric \tilde{g} of g to be a Killing metric [14]. Then

$$g([X_i, X_j], JX_k) + g([X_i, X_k], JX_j) = 0. \quad (3.17)$$

Combining (2.1), (3.14), (3.15), (3.16) and (3.17), we obtain

$$2F(X_i, X_j, X_k) = g([JX_i, X_j], X_k) + g([JX_i, X_k], X_j). \quad (3.18)$$

From (3.18) it follows immediately that $\sum_{X_i, X_j, X_k} F(X_i, X_j, X_k) = 0$, i. e.

(G, J, g) is a quasi-Kähler manifold with Norden metric.

According to (3.13) and (3.17), we get

$$\begin{aligned} [X_1, X_2] &= \lambda_1 X_1 + \lambda_2 X_2, & [X_1, X_3] &= \lambda_3 X_2 - \lambda_1 X_4, \\ [X_1, X_4] &= -\lambda_3 X_1 - \lambda_2 X_4, & [X_2, X_3] &= \lambda_4 X_2 + \lambda_1 X_3, \\ [X_2, X_4] &= -\lambda_4 X_1 + \lambda_2 X_3, & [X_3, X_4] &= \lambda_3 X_3 + \lambda_4 X_4, \end{aligned} \quad (3.19)$$

where

$$\begin{aligned} \lambda_1 &= C_{12}^1 = C_{23}^3 = -C_{13}^4, & \lambda_2 &= C_{12}^2 = C_{24}^3 = -C_{14}^4, \\ \lambda_3 &= C_{13}^2 = C_{34}^3 = -C_{14}^1, & \lambda_4 &= C_{23}^2 = C_{34}^4 = -C_{24}^1. \end{aligned}$$

Let equalities (3.19) are valid for an almost complex manifold with Norden metric (G, J, g) , where J and g are determined by (3.14) and (3.15). Then we verify directly that the Jacobi identity for the commutators $[X_i, X_j]$ is satisfied and the associated metric \tilde{g} of g is a Killing metric.

Therefore, the following theorem is valid.

Theorem 3.2. *Let (G, J, g) be a 4-dimensional almost complex manifold with Norden metric, where G is the connected Lie group with an associated Lie algebra determined by a global basis $\{X_i\}$ of left invariant vector fields, and J and g are the almost complex structure and the Norden metric determined by (3.14) and (3.15), respectively. Then (G, J, g) is a quasi-Kähler manifold with a Killing associated metric \tilde{g} iff G belongs to the 4-parametric family of Lie groups, defined by (3.19). \square*

Let (G, J, g) be the quasi-Kähler manifold determined by the conditions of Theorem (3.2).

By (3.18) and (3.19) we get the non-trivial components $F_{ijk} = F(X_i, X_j, X_k)$ of the tensor F :

$$\begin{aligned} -2F_{114} &= 2F_{123} = -2F_{312} = 2F_{334} = 2F_{411} = 2F_{433} = \lambda_1, \\ 2F_{223} &= -2F_{241} = -2F_{322} = -2F_{344} = 2F_{412} = 2F_{434} = \lambda_2, \\ -2F_{112} &= -2F_{134} = F_{211} = F_{233} = 2F_{314} = -2F_{332} = \lambda_3, \\ -F_{122} &= F_{144} = 2F_{221} = 2F_{234} = 2F_{414} = -2F_{432} = \lambda_4. \end{aligned}$$

This leads to the following

Proposition 3.3. *(G, J, g) is a non-Kähler manifold with Norden metric.* \square

Using (3.16) and (3.17), we obtain

$$2g(\nabla_{X_i} X_j, X_k) = g([X_i, X_j], X_k) - g(J[X_i, JX_j], X_k) + g(J[X_i, X_j], X_k).$$

Then we have

$$2\nabla_{X_i} X_j = [X_i, X_j] - J[X_i, JX_j] + J[JX_i, X_j]. \quad (3.20)$$

The equality (3.20) implies

$$\begin{aligned} 2\nabla_{X_i} JX_j &= [X_i, JX_j] + J[X_i, X_j] + J[JX_i, JX_j], \\ 2J\nabla_{X_i} X_j &= J[X_i, X_j] + [X_i, JX_j] - [JX_i, X_j]. \end{aligned}$$

We subtract the last two equalities, apply the formula for covariant derivation and obtain

$$2(\nabla_{X_i} J) X_j = J[JX_i, JX_j] + [JX_i, X_j]. \quad (3.21)$$

By (3.19) and (3.21) we obtain the components of ∇J :

$$\begin{aligned} 2(\nabla_{X_1} J) X_1 &= 2(\nabla_{X_3} J) X_3 = -\lambda_3 X_2 + \lambda_1 X_4, \\ 2(\nabla_{X_2} J) X_2 &= 2(\nabla_{X_4} J) X_4 = \lambda_4 X_1 - \lambda_2 X_3, \\ 2(\nabla_{X_1} J) X_3 &= -2(\nabla_{X_3} J) X_1 = \lambda_1 X_2 + \lambda_3 X_4, \\ 2(\nabla_{X_2} J) X_4 &= -2(\nabla_{X_4} J) X_2 = -\lambda_4 X_3 - \lambda_2 X_1, \\ 2(\nabla_{X_1} J) X_2 &= -\lambda_3 X_1 - 2\lambda_4 X_2 - \lambda_1 X_3, \\ 2(\nabla_{X_1} J) X_4 &= -\lambda_1 X_1 + \lambda_3 X_3 + 2\lambda_4 X_4, \\ 2(\nabla_{X_2} J) X_1 &= 2\lambda_3 X_1 + \lambda_4 X_2 + \lambda_2 X_4, \\ 2(\nabla_{X_2} J) X_3 &= \lambda_2 X_2 - 2\lambda_3 X_3 - \lambda_4 X_4, \\ 2(\nabla_{X_3} J) X_2 &= -\lambda_1 X_1 - 2\lambda_2 X_2 + \lambda_3 X_3, \\ 2(\nabla_{X_3} J) X_4 &= \lambda_3 X_1 + \lambda_1 X_3 + 2\lambda_2 X_4, \\ 2(\nabla_{X_4} J) X_1 &= 2\lambda_1 X_1 + \lambda_2 X_2 - \lambda_4 X_4, \\ 2(\nabla_{X_4} J) X_3 &= -\lambda_4 X_2 - 2\lambda_1 X_3 - \lambda_2 X_4. \end{aligned} \quad (3.22)$$

From (2.3) and (3.22) it follows that $\|\nabla J\|^2 = -4(\lambda_1^2 + \lambda_2^2 - \lambda_3^2 - \lambda_4^2)$, and therefore the following is valid.

Proposition 3.4. *(G, J, g) is an isotropic Kähler manifold with Norden metric iff $\lambda_1^2 + \lambda_2^2 - \lambda_3^2 - \lambda_4^2 = 0$.* \square

Let ∇' be the connection with totally skew-symmetric torsion tensor T on (G, J, g) .

By virtue of (3.3), (3.12) and (3.21), we have

$$4T(X_i, X_j) = 3[JX_i, JX_j] - J[X_i, JX_j] - J[JX_i, X_j] + [X_i, X_j], \quad (3.23)$$

which implies

$$\begin{aligned}
 T(X_1, X_2) &= \lambda_3 X_3 + \lambda_4 X_4, & T(X_1, X_3) &= \lambda_3 X_2 - \lambda_1 X_4, \\
 T(X_1, X_4) &= \lambda_4 X_2 + \lambda_1 X_3, & T(X_2, X_3) &= -\lambda_3 X_1 - \lambda_2 X_4, \\
 T(X_2, X_4) &= -\lambda_4 X_1 + \lambda_2 X_3, & T(X_3, X_4) &= \lambda_1 X_1 + \lambda_2 X_2.
 \end{aligned} \tag{3.24}$$

Then the non-trivial components $T_{ijk} = T(X_i, X_j, X_k)$ of the corresponding (0,3)-tensor T are:

$$T_{134} = \lambda_1, \quad T_{234} = \lambda_2, \quad T_{123} = -\lambda_3, \quad T_{124} = -\lambda_4. \tag{3.25}$$

According to (3.19) and (3.24), we have

$$T(X_i, X_j) = [JX_i, JX_j]$$

and then, from (3.23) it follows that

$$[JX_i, JX_j] + J[JX_i, X_j] = [X_i, X_j] - J[X_i, JX_j]. \tag{3.26}$$

Since we have $2Q = T$ for ∇' , from (2.8), (3.23) and (3.26), we obtain

$$\nabla'_{X_i} X_j = [X_i, X_j] - J[X_i, JX_j]. \tag{3.27}$$

Combining (3.19) and (3.27), we get the components of ∇' :

$$\begin{aligned}
 \nabla'_{X_1} X_1 &= \nabla'_{X_3} X_3 = -\lambda_1 X_2 - \lambda_3 X_4, \\
 \nabla'_{X_2} X_2 &= \nabla'_{X_4} X_4 = \lambda_2 X_1 + \lambda_4 X_3, \\
 \nabla'_{X_1} X_2 &= \nabla'_{X_3} X_4 = \lambda_1 X_1 + \lambda_3 X_3, \\
 \nabla'_{X_1} X_3 &= -\nabla'_{X_3} X_1 = \lambda_3 X_2 - \lambda_1 X_4, \\
 \nabla'_{X_1} X_4 &= -\nabla'_{X_3} X_2 = -\lambda_3 X_1 + \lambda_1 X_3, \\
 \nabla'_{X_2} X_1 &= \nabla'_{X_4} X_3 = -\lambda_2 X_2 - \lambda_4 X_4, \\
 \nabla'_{X_2} X_3 &= -\nabla'_{X_4} X_1 = \lambda_4 X_2 - \lambda_2 X_4, \\
 \nabla'_{X_2} X_4 &= -\nabla'_{X_4} X_2 = -\lambda_4 X_1 + \lambda_2 X_3.
 \end{aligned} \tag{3.28}$$

Thus we arrive at the following

Proposition 3.5. *Let ∇' be the connection with totally skew-symmetric torsion tensor T on (G, J, g) . Then the components of ∇' and T with respect to the basis $\{X_i\}$ are (3.28) and (3.25), respectively. \square*

In [12] and [15] we have considered the canonical connection ∇^C and the B-connection ∇^B on quasi-Kähler manifolds with Norden metric. Now we will show how these two connections relate to the connection ∇' with totally skew-symmetric torsion tensor on (G, J, g) .

The connection ∇^B is defined in [15] by

$$\nabla^B_{X_i} X_j = \nabla_{X_i} X_j + \frac{1}{2} (\nabla_{X_i} J) JX_j.$$

Hence, applying (3.14) and (3.22), we obtain directly $\nabla^B = \frac{3}{4}\nabla'$. On the other hand, according to [12], we have $\nabla^B = \frac{1}{2}(\nabla^C + \nabla')$. Thus we proved the following

Proposition 3.6. *Let ∇' , ∇^B and ∇^C be the connection with totally skew-symmetric torsion, the B-connection and the canonical connection on (G, J, g) , respectively. Then*

$$\nabla' = \frac{4}{3}\nabla^B = 2\nabla^C.$$

□

In [4] we have proved that ∇' has a Kähler curvature tensor on any quasi-Kähler manifolds with Norden metric iff the following identity holds

$$\mathfrak{S}_{x,y,z} \{g((\nabla_x J) Jy + (\nabla_{Jx} J) y, (\nabla_z J) Jw + (\nabla_{Jz} J) w)\} = 0. \quad (3.29)$$

Equality (3.21) implies

$$\begin{aligned} 2(\nabla_{X_i} J) JX_j &= -J[JX_i, X_j] + [JX_i, JX_j], \\ 2(\nabla_{JX_i} J) X_j &= -J[X_i, JX_j] - [X_i, X_j]. \end{aligned}$$

We add the above, and according to (3.26) obtain

$$(\nabla_{X_i} J) JX_j + (\nabla_{JX_i} J) X_j = [JX_i, JX_j] - [X_i, X_j].$$

Then, the condition (3.29) takes the following form on (G, J, g) :

$$\mathfrak{S}_{x,y,z} \{g([JX_i, JX_j] - [X_i, X_j], [JX_k, JX_s] - [X_k, X_s])\} = 0.$$

The last equality and equalities (3.14), (3.15) and (3.19) imply

Proposition 3.7. *The connection ∇' with totally skew-symmetric torsion on (G, J, g) has a Kähler curvature tensor iff $\lambda_1^2 + \lambda_2^2 = \lambda_3^2 + \lambda_4^2$. □*

4 Connection with parallel totally skew-symmetric torsion

Let ∇' be the connection with totally skew-symmetric torsion tensor T on the quasi-Kähler manifold with Norden metric (M, J, g) .

Now we consider the case when ∇' has a parallel torsion, i. e. $\nabla' T = 0$.

It is known that the curvature tensors R' and R of ∇' and ∇ , respectively, satisfy:

$$\begin{aligned} R'(x, y, z, w) &= R(x, y, z, w) + (\nabla_x Q)(y, z, w) - (\nabla_y Q)(x, z, w) \\ &\quad + Q(x, Q(y, z), w) - Q(y, Q(x, z), w). \end{aligned} \quad (4.1)$$

Equality (3.3) implies $\nabla'Q = 0$ in the considered case. Then from the formula for covariant derivation with respect to ∇' it follows that

$$xQ(y, z, w) - Q(\nabla'_x y, z, w) - Q(y, \nabla'_x z, w) - Q(y, z, \nabla'_x w) = 0. \quad (4.2)$$

According to the first equality of (2.8) we have

$$\begin{aligned} Q(\nabla'_x y, z, w) &= Q(\nabla_x y, z, w) + Q(Q(x, y), z, w), \\ Q(y, \nabla'_x z, w) &= Q(y, \nabla_x z, w) + Q(y, Q(x, z), w), \\ Q(y, z, \nabla'_x w) &= Q(y, z, \nabla_x w) + Q(y, z, Q(x, w)). \end{aligned} \quad (4.3)$$

Combining (4.2), (4.3), the first equality of (2.9) and having in mind the formula for covariant derivation with respect to ∇ , we obtain

$$\begin{aligned} (\nabla_x Q)(y, z, w) &= Q(Q(x, y), z, w) \\ &\quad - g(Q(x, z), Q(y, w)) - g(Q(y, z), Q(x, w)). \end{aligned} \quad (4.4)$$

From (4.4) and the first equality of (2.10) we have

$$\begin{aligned} (\nabla_x Q)(y, z, w) - (\nabla_y Q)(x, z, w) &= Q(T(x, y), z, w) \\ &\quad - 2g(Q(x, z), Q(y, w)) + 2g(Q(y, z), Q(x, w)). \end{aligned} \quad (4.5)$$

Because of (4.5), equality (4.1) can be rewritten as

$$\begin{aligned} R'(x, y, z, w) &= R(x, y, z, w) + Q(T(x, y), z, w) \\ &\quad - g(Q(x, z), Q(y, w)) + g(Q(y, z), Q(x, w)). \end{aligned} \quad (4.6)$$

Since $Q(e_i, e_j) = -Q(e_j, e_i)$ it follows that $g^{ij}Q(e_i, e_j) = 0$. Then, from (4.6) after contraction by $x = e_i, w = e_j$, we obtain the following equality for the Ricci tensor ρ' of ∇' :

$$\begin{aligned} \rho'(y, z) &= \rho(y, z) + 2g^{ij}g(Q(e_i, y), Q(z, e_j)) \\ &\quad - g^{ij}g(Q(e_i, z), Q(y, e_j)). \end{aligned} \quad (4.7)$$

Contracting by $y = e_k, z = e_s$ in (4.7), we get

$$\tau' = \tau + g^{ij}g^{ks}g(Q(e_i, e_k), Q(e_s, e_j)), \quad (4.8)$$

where τ' is the scalar curvature of ∇' .

By virtue of (4.8), (3.12), (2.3) and (2.4) we have

$$\tau' = \tau - \frac{1}{8} \|\nabla J\|^2. \quad (4.9)$$

Thus we arrive at the following

Theorem 4.1. *Let ∇' be the connection with parallel totally skew-symmetric torsion on the quasi-Kähler manifold with Norden metric (M, J, g) . Then for the Ricci tensor ρ' and the scalar curvature τ' of ∇' are valid (4.7) and (4.9), respectively. \square*

Equality (4.9) leads to the following

Corollary 4.2. *Let ∇' be the connection with parallel totally skew-symmetric torsion on the quasi-Kähler manifold with Norden metric (M, J, g) . Then the manifold (M, J, g) is isotropic Kählerian iff ∇' and ∇ have equal scalar curvatures. \square*

Remark 4.1. *The 4-parametric family of 4-dimensional quasi-Kähler manifolds (G, J, g) considered in subsection 3.2 does not admit any connection with parallel totally skew-symmetric torsion.*

5 Connection with parallel totally skew-symmetric torsion and Kähler curvature tensor

Let ∇' be a connection with parallel totally skew-symmetric torsion on the quasi-Kähler manifold with Norden metric (M, J, g) .

We will find conditions for the curvature tensor R' of ∇' to be Kählerian.

From (3.3), having in mind that Q is a 3-form, we have

$$\begin{aligned} Q(T(x, y), z, w) &= Q(z, w, T(x, y)) = g(Q(z, w), T(x, y)) \\ &= g(T(x, y), Q(z, w)) = 2g(Q(x, y), Q(z, w)). \end{aligned}$$

Then (4.6) obtains the form

$$\begin{aligned} R'(x, y, z, w) &= R(x, y, z, w) + 2g(Q(x, y), Q(z, w)) \\ &\quad - g(Q(x, z), Q(y, w)) + g(Q(y, z), Q(x, w)). \end{aligned} \tag{5.1}$$

From (5.1), identities (2.5) and (2.7) for R' follow immediately. Therefore R' is a Kähler tensor iff the first Bianchi identity (2.6) for R' is satisfied. Since this identity is valid for R , then (5.1) implies that R' is Kählerian iff

$$\mathfrak{S}_{x, y, z} \{2g(Q(x, y), Q(z, w)) - g(Q(x, z), Q(y, w)) + g(Q(y, z), Q(x, w))\} = 0.$$

Thus, using that Q is a skew-symmetric tensor, we arrive the following

Theorem 5.1. *Let ∇' be the connection with parallel totally skew-symmetric torsion on the quasi-Kähler manifold with Norden metric (M, J, g) . Then the curvature tensor for ∇' is a Kähler tensor iff*

$$\mathfrak{S}_{x, y, z} g(Q(x, y), Q(z, w)) = 0. \tag{5.2}$$

\square

Because of the skew-symmetry of Q , (5.2) implies

$$g(Q(y, z), Q(x, w)) - g(Q(x, z), Q(y, w)) = -g(Q(x, y), Q(z, w)).$$

The last equality and (5.1) lead to the following

Corollary 5.2. *Let ∇' be the connection with parallel totally skew-symmetric torsion and Kähler curvature tensor on the quasi-Kähler manifold with Norden metric (M, J, g) . Then*

$$R'(x, y, z, w) = R(x, y, z, w) + g(Q(x, y), Q(z, w)). \quad (5.3)$$

□

If R' is a Kähler tensor then $R'(x, y, Jz, Jw) = -R'(x, y, z, w)$, and because of (5.3) we have

$$\begin{aligned} R(x, y, Jz, Jw) + R(x, y, z, w) &= -g(Q(x, y), Q(Jz, Jw)) \\ &\quad - g(Q(x, y), Q(z, w)). \end{aligned} \quad (5.4)$$

From (3.12) we get

$$Q(x, Jy) = JQ(x, y) - (\nabla_x J) y.$$

Then we have

$$Q(Jx, Jy) = -Q(x, y) - (\nabla_{Jx} J) y - (\nabla_y J) Jx$$

and consequently

$$\begin{aligned} g(Q(x, y), Q(Jz, Jw)) &= -g(Q(x, y), Q(z, w)) \\ &\quad - g(Q(x, y), (\nabla_{Jz} J) w + (\nabla_w J) Jz). \end{aligned}$$

The last equality and (5.4) imply the following

Corollary 5.3. *Let ∇' be the connection with parallel totally skew-symmetric torsion and Kähler curvature tensor on the quasi-Kähler manifold with Norden metric (M, J, g) . Then*

$$R(x, y, Jz, Jw) + R(x, y, z, w) = g(Q(x, y), (\nabla_{Jz} J) w + (\nabla_w J) Jz). \quad (5.5)$$

□

Contracting by $x = e_i, w = e_j$ in (5.5), we obtain

$$g^{ij} R(e_i, y, Jz, J e_j) + \rho(y, z) = g^{ij} g(Q(e_i, y), (\nabla_{Jz} J) e_j + (\nabla_{e_j} J) Jz).$$

Then, after a contraction by $y = e_k, z = e_s$, it follows

$$\tau^{**} + \tau = g^{ij} g^{ks} g(Q(e_i, e_k), (\nabla_{J e_s} J) e_j + (\nabla_{e_j} J) J e_s), \quad (5.6)$$

where $\tau^{**} = g^{ij}g^{ks}R(e_i, e_k, J e_s, J e_j)$.

From (3.12), (2.3) and (2.4) we have

$$g^{ij}g^{ks}g(Q(e_i, e_k), (\nabla_{J e_s} J) e_j + (\nabla_{e_j} J) J e_s) = -\frac{1}{8} \|\nabla J\|^2.$$

Then (5.6) can be rewritten as

$$\tau^{**} + \tau = -\frac{1}{8} \|\nabla J\|^2.$$

On the other hand, according to [8], we have

$$\tau^{**} + \tau = -\frac{1}{2} \|\nabla J\|^2.$$

Then $\|\nabla J\|^2 = 0$ and therefore the following is valid.

Theorem 5.4. *Let (M, J, g) be a quasi-Kähler manifold with Norden metric which admit a connection with parallel totally skew-symmetric torsion and Kähler curvature tensor. Then (M, J, g) is a isotropic Kähler manifold with Norden metric. \square*

References

- [1] A. Strominger, Superstrings with torsion. *Nuclear Phys. B*, 274:253-284 (1986).
- [2] J.-M. Bismut, A local index theorem for a non-Kähler manifolds. *Math. Ann.* 284: 681-699 (1989).
- [3] P. Gauduchon, Hermitian connections and Dirac operators. *Boll. Unione Mat. Ital. Sez. B Artic. Ric. Mat. (8)* 11: 257-289 (1997).
- [4] D. Mekerov, A connection with skew symmetric torsion and Kähler curvature tensor on a quasi-Kähler manifolds with Norden metric. *C. R. Acad. Bulgare Sci.* 61: 1249-1256 (2008).
- [5] K. I. Gribachev, D. G. Mekerov and G. D. Djelepov, Generalized B-manifold. *C. R. Acad. Bulgare Sci.* 38: 299-302 (1985).
- [6] G. T. Ganchev and A. V. Borisov, Note on the almost complex manifolds with a Norden metric. *C. R. Acad. Bulgare Sci.* 39: 31-34 (1986).
- [7] E. Garcia-Rio and Y. Matsushita, Isotropic Kähler structures on Engel 4-manifolds. *J. Geom. Phys.* 33: 288-294 (2000).
- [8] D. Mekerov and M. Manev, On the geometry of quasi-Kähler manifolds with Norden metric. *Nihonkai Math. J.* 16: 89-93 (2005).

- [9] G. Ganchev, K. Gribachev and V. Mihova, B-connection and their conformal invariants on conformally Kahler manifolds with B-metric. *Publ. Inst. Math. (Beograd) (N.S.)* 42: 107-121 (1987).
- [10] G. Ganchev and V. Mihova, Canonical connection and the canonical conformal group on an almost complex manifold with B-metric. *Annuaire Univ. Sofia Fac. Math. Inform.* 81: 195-206 (1987).
- [11] H. Hayden, Subspaces of a space with torsion. *Proc. London Math. Soc.* (3) 34: 27-50 (1934).
- [12] D. Mekerov, Canonical connection on quasi-Kahler manifolds with Norden metric (to appear). *arXiv:0812.3516* (December 2008).
- [13] M. Teofilova, On the geometry of almost complex manifolds with a Norden metric. Ph. D. Thesis, Univ. of Plovdiv, Sofia, 2008.
- [14] S. Helgason, *Differential geometry, Lie groups and symmetric spaces*. Academic Press, NY, 1978.
- [15] D. Mekerov, On the geometry of the B-connection on quasi-Kahler manifolds with Norden metric. *C. R. Acad. Bulgare Sci.* 61: 1105-1110 (2008).