

On the minimal norm of a non-regular generalized character of an arbitrary finite group

*Geoffrey R. Robinson
Department of Mathematics
University of Aberdeen
Aberdeen AB24 3UE
Scotland
United Kingdom
e-mail:g.r.robinson@abdn.ac.uk*

INTRODUCTION AND NOTATION: As usual, for G a finite group, we denote the set of non-identity elements of G by $G^\#$, the number of conjugacy classes of G by $k(G)$ and the set of complex irreducible characters of G by $\text{Irr}(G)$. When H is a non-trivial subgroup of G , we let $k_+(G, H)$ denote the number of irreducible characters of G which do not vanish identically on $H^\#$ (and we remark that $k_+(G, H) = k(G)$ when $|H| \geq \sqrt{|G|}$).

For θ a generalized character of a finite group G , we set

$$\sigma(G^\#, \theta) = \sum_{g \in G^\#} |\theta(g)|^2.$$

Notice that

$$\sigma(G^\#, \theta) = |G| \langle \theta, \theta \rangle - \theta(1)^2,$$

where \langle, \rangle is the usual inner product on the space of complex-valued class functions of G . We denote the regular character of G by ρ . We let $\text{cf}(G)$ denote the space of complex-valued class functions of G , and for $\alpha, \beta \in \text{cf}(G)$, we set

$$(\alpha, \beta) = |G| \langle \alpha, \beta \rangle - \alpha(1)\beta(1).$$

Then $(,)$ induces an inner product on the quotient space $\text{cf}(G)/\mathbb{C}\rho$. We label the irreducible characters of G as $\mu_1 (= 1), \mu_2, \dots, \mu_k$ where $\mu_1(1) \leq \dots \leq \mu_k(1)$.

In the earlier papers [1],[2] and [3], we considered the minimum possible value of $\sigma(G^\#, \theta)$ for θ a generalized character (other than a multiple of the regular character) of a non-trivial finite group G . We denote this minimum possible value by $m(G)$ (which is clearly a positive integer). Note that with the above notation, this is the minimum non-zero value of (θ, θ) for $\theta \in \mathbb{Z}\text{Irr}(G) \setminus \mathbb{Z}\rho$.

We proved in [1] that $m(G) \geq k(G) - 1$ for every non-trivial finite group G . In the papers [2],[3] we proved that $m(G) = |G| - \mu_k(1)^2$ whenever G is a non-trivial nilpotent group. In particular, in that case,

$$m(G) \geq |G| - [G : Z(G)]$$

whenever G since $\mu_k(1)^2$ divides $[G : Z(G)]$.

We asked in [3] whether it was the case that for every non-trivial finite group G we have

$$m(G) \geq \frac{|G|}{\mu_k(1)} - 1$$

(which would yield $m(G) > \sqrt{|G|} - 1$). We answer that question in the affirmative here. Notice that if G is a Frobenius group with Abelian kernel K and complement H , then $m(G) = |K| - 1$. In that case, the maximal degree of an irreducible character of G is $|H|$, and we do have

$$m(G) = \frac{|G|}{\mu_k(1)} - 1.$$

For if θ is a generalized character of G which does not vanish identically on $H^\#$, we see that $\sigma(G^\#, \theta) \geq |K|\sigma(H^\#, \text{Res}_H^G(\theta)) \geq |K|$, as there are $|K|$ conjugates of H which intersect trivially with each other. On the other hand, if the generalized character θ does not vanish identically on $K^\#$, then

$$\sigma(G^\#, \theta) \geq m(K) = |K| - 1,$$

as K is Abelian. Finally, let θ be the sum of the distinct irreducible characters of G which do not contain K in their kernels. By the orthogonality relations, we see that $\theta(x) = -1$ for all $x \in K^\#$, while also θ vanishes identically outside K , so that $\sigma(G^\#, \theta) = |K| - 1$, as claimed.

In view of these examples, our main theorem may be regarded as optimal in this generality. We prove:

THEOREM 1 : *Let G be a non-trivial finite group. Then*

$$m(G) \geq \frac{|G|}{\mu_k(1)} - 1.$$

Furthermore, equality is attained if and only if G is either Abelian, or else is a Frobenius group with an Abelian Frobenius kernel.

As noted above, Theorem 1 has an interpretation in terms of the minimal non-zero value of a certain semi-definite integral quadratic form.

COROLLARY 2: *Let G be a non-trivial finite group and let a_1, a_2, \dots, a_k be any k integers such that for some $i \neq j$, we have $a_i\mu_j(1) \neq a_j\mu_i(1)$. Then*

$$\sum_{i < j} (a_i\mu_j(1) - a_j\mu_i(1))^2 \geq \frac{|G|}{\mu_k(1)} - 1.$$

COROLLARY 3: *Let H be a non-trivial subgroup of the finite group G and let μ be a complex irreducible character of maximal degree of H . Then we have*

$$k_+(G, H) \leq \frac{\mu(1)|H| - \mu(1)}{|H| - \mu(1)} \max_{h \in H^\#} |C_G(h)|.$$

PROOF OF THEOREM 1

Suppose that $m(G) < \frac{|G|}{\mu_k(1)}$. To prove the Theorem, by the cases discussed above, we may suppose that G is non-Abelian. Let us choose a generalized character θ of G such that $m(G) = \sigma(G^\#, \theta)$. We will prove that $\theta(x) \in \{0, -1\}$ for all $x \in G^\#$, and that we may suppose that θ is a multiplicity free irreducible character of G whose irreducible constituents all have degree $\mu_k(1)$, and that if there are r such constituents, then $|G| = \mu_k(1) + r\mu_k(1)^2$.

If θ has the form $t\rho + \delta\mu_i$ for some integer t , some sign δ and some i , then we may replace θ by μ_i . Then we have

$$\sigma(G^\#, \theta) = |G| - \mu_i(1)^2,$$

so we may suppose that $i = k$, as μ_i must also have maximal degree by the choice of θ . We then have

$$\sigma(G^\#, \theta) = |G| - \mu_k(1)^2 \geq \frac{|G|}{\mu_k(1)} - 1.$$

Furthermore, equality forces either $\mu_k(1) = 1$ (in which case G is Abelian, contrary to hypothesis), or else $|G| = \mu_k(1)^2 + \mu_k(1)$. In the latter case, we have $\sum_{i=1}^{k-1} \mu_i(1)^2 = \mu_k(1)$. Now $\langle \theta, \mu_1 \rangle = 0$, so that $\sum_{g \in G^\#} \mu_k(g) = -\mu_k(1)$. Under the current assumptions, we also have

$$\sigma(G^\#, \theta) = \mu_k(1).$$

Now μ_k is the unique irreducible character of G of its degree, so it is rational-valued. Hence we must have $\mu_k(g) = -1$ whenever $\mu_k(g) \neq 0$. In order to establish the initial claim, then, we may now suppose that θ does not have the form $t\rho + \delta\mu_i$ for some integer t , some sign δ and some i .

By the arguments of [2] and [3], we may suppose that θ is a rational-valued character of G with $0 < \theta(1) < \frac{|G|}{2}$ (in which irreducible characters of the same degree occur with equal multiplicity), and that $\langle \theta, \mu_i \rangle$ is the closest integer to $\frac{\theta(1)\mu_i(1)}{|G|}$ for each irreducible character μ_i of G . For the convenience of the reader, we review the necessary results using the notation currently adopted. After subtracting integer multiples of ρ from θ and replacing θ by $-\theta$ if necessary, we may suppose that $0 < \theta(1) \leq \frac{|G|}{2}$.

For (by the choice of θ) we have

$$(\theta - \delta\mu_i, \theta - \delta\mu_i) \geq (\theta, \theta)$$

for each i and any choice of sign δ . This leads easily to

$$|(\theta, \mu_i)| \leq \frac{|G| - \mu_i(1)^2}{2}$$

and then

$$\left| \langle \theta, \mu_i \rangle - \frac{\theta(1)\mu_i(1)}{|G|} \right| \leq \frac{1}{2} \left(1 - \frac{\mu_i(1)^2}{|G|} \right).$$

Hence $\langle \theta, \mu_i \rangle$ is the (unique under current assumptions) closest integer to $\frac{\theta(1)\mu_i(1)}{|G|}$, and only depends on $\mu_i(1)$. Notice that we can't now have $\theta(1) = \frac{|G|}{2}$, for if we did, we would have the contradiction

$$\left| \langle \theta, \mu_1 \rangle - \frac{1}{2} \right| < \frac{1}{2}.$$

Thus we have $0 < \theta(1) < \frac{|G|}{2}$. Also, θ is a rational (integer) valued character, since it contains algebraically conjugate irreducible characters with the same non-negative multiplicity.

Notice that the trivial character does not now occur as an irreducible constituent of θ . Hence we have $\sum_{g \in G^\#} \theta(g) = -\theta(1)$, so that

$$\sum_{g \in G^\#} |\theta(g)| \geq \theta(1)$$

and certainly

$$\sigma(G^\#, \theta) \geq \theta(1)$$

(as θ is integer-valued).

Now let us write

$$\theta = \sum_{i=1}^k a_i \mu_i,$$

and set

$$\epsilon_i = a_i - \frac{\theta(1)\mu_i(1)}{|G|}.$$

Then we have

$$\sum_{i=1}^k \epsilon_i \mu_i(1) = 0$$

and

$$\sigma(G^\#, \theta) = |G| \left(\sum_{i=1}^k |\epsilon_i|^2 \right)$$

(for notice that θ agrees with $\sum_{i=1}^k \epsilon_i \mu_i$ on $G^\#$). We have noted already that $|\epsilon_i| < \frac{1}{2}$ for each i .

Suppose that $a_i \geq 2$ for some i . Then we have

$$\frac{\theta(1)\mu_i(1)}{|G|} > \frac{3}{2},$$

so that

$$\sigma(G^\#, \theta) \geq \theta(1) > \frac{G}{\mu_i(1)} \geq \frac{|G|}{\mu_k(1)},$$

contrary to hypothesis. Hence we have $a_i \in \{0, 1\}$ for each i .

Now suppose that $\epsilon_i < 0$ for $i = 1, \dots, k-r$, and that $\epsilon_i > 0$ for $i = k-r+1, \dots, k$. Then

$$\theta = \sum_{i=1}^r \mu_{k-r+i}.$$

Since we have dealt with the case that θ is irreducible, we may, and do, suppose that $r > 1$.

By assumption, we have

$$\frac{|G|}{\mu_k(1)} - 1 \geq m(G) = \sigma(G^\#, \theta) \geq \theta(1),$$

while we clearly also have

$$\theta(1) \geq \frac{|G| - \sum_{i=1}^{k-r} \mu_i(1)^2}{\mu_k(1)},$$

so that

$$\sum_{i=1}^{k-r} \mu_i(1)^2 \geq \mu_k(1)$$

(and hence

$$|G| \geq \mu_k(1) + \sum_{i=k-r+1}^k \mu_i(1)^2.$$

Now we have

$$\sigma(G^\#, \theta) = \sum_{i < j} (a_i \mu_j(1) - a_j \mu_i(1))^2 \geq r \left(\sum_{i=1}^{k-r} \mu_i(1)^2 \right),$$

using the fact that $a_j = 1$ for $j > k-r$ and $a_i = 0$ for $i \leq k-r$.

Thus we have (by assumption)

$$r \left[|G| - \left(\sum_{i=k-r+1}^k \mu_i(1)^2 \right) \right] \leq \sigma(G^\#, \theta) \leq \frac{|G|}{\mu_k(1)} - 1,$$

so that

$$|G| \left(1 - \frac{1}{r\mu_k(1)} \right) < \left(\sum_{i=k-r+1}^k \mu_i(1)^2 \right).$$

Now for $n \geq 2$, we have

$$\frac{\sum_{i=k-r+1}^k \mu_i(1)^2}{r^n \mu_k(1)^n} \leq \frac{1}{r^{n-1} \mu_k(1)^{n-2}}.$$

Hence

$$\begin{aligned} |G| &< \sum_{i=k-r+1}^k \mu_i(1)^2 + \frac{\sum_{i=k-r+1}^k \mu_i(1)^2}{r \mu_k(1)} + \frac{1}{r} \left(1 - \frac{1}{r \mu_k(1)}\right)^{-1} \\ &= \sum_{i=k-r+1}^k \mu_i(1)^2 + \frac{\sum_{i=k-r+1}^k \mu_i(1)^2}{r \mu_k(1)} + \frac{1}{r - \frac{1}{\mu_k(1)}}. \end{aligned}$$

Now $r > 1$ and also $\mu_k(1) > 1$ (as G is non-Abelian).

Thus

$$|G| < 1 + \mu_k(1) + \sum_{i=k-r+1}^k \mu_i(1)^2,$$

so that we now have

$$|G| = \mu_k(1) + \sum_{i=k-r+1}^k \mu_i(1)^2,$$

as we already have

$$|G| \geq \mu_k(1) + \sum_{i=k-r+1}^k \mu_i(1)^2.$$

We also now have

$$\frac{|G|}{\mu_k(1)} - 1 \leq \theta(1) \leq \sigma(G^\#, \theta) \leq \frac{|G|}{\mu_k(1)} - 1,$$

so that

$$\theta(1) = \sigma(G^\#, \theta) = \frac{|G|}{\mu_k(1)} - 1.$$

Furthermore, we have $\mu_i(1) = \mu_k(1)$ whenever $i > k - r$. Hence

$$|G| = \mu_k(1) + r \mu_k(1)^2.$$

Furthermore, we also have $|\theta(x)|^2 = 1 = -\theta(x)$ for all $x \in G^\#$ such that $\theta(x) \neq 0$. Thus the claim of the first paragraph of the proof is complete, and in the discussion which follows, we allow the possibility that $r = 1$.

The orthogonality relations (and the fact that $\mu_k(1) = \sum_{i=1}^{k-r} \mu_i(1)^2$) now yield that $x \in \bigcap_{i=1}^{k-r} \ker \mu_i$ whenever $\theta(x) \neq 0$. Conversely, if $\theta(x) = 0$, we have $\sum_{i=1}^{k-r} \mu_i(1) \mu_i(x) = 0$.

Set $K = \bigcap_{i=1}^{k-r} \ker \mu_i$. Now

$$|G| = \mu_k(1) + r\mu_k(1)^2$$

and

$$|K| = 1 + \theta(1) = 1 + r\mu_k(1).$$

Thus K is a Hall subgroup of G , and there is a complement, H say, to K in G . Since $\mu_k(1) = |H|$, we see that, for all $i > k - r$, we have $\mu_i(y) = 0$ whenever $\gcd(|\langle y \rangle|, |H|) > 1$. In particular (using a well-known observation of Feit and Thompson),

$$|C_G(y)| = |C_{G/K}(yK)| = |C_H(y)|$$

for all $y \in H^\#$. Thus $C_G(y) \leq H$ for all $y \in H^\#$ and G is a Frobenius group with kernel K , complement H . Furthermore, every irreducible character of G which does not contain K in its kernel has degree $\mu_k(1) = [G : K]$, so that every non-trivial irreducible character of K is linear. Hence K is Abelian.

PROOF OF COROLLARY 2: As noted in [1], if $\theta = \sum_{i=1}^k a_i \mu_i$, then $(\theta, \theta) = \sum_{i < j} (a_i \mu_j(1) - a_j \mu_i(1))^2$.

PROOF OF COROLLARY 3: By the orthogonality relations for G and Theorem 1 (applied to H), and noting the presence of the trivial character, we have

$$\sum_{h \in H^\#} |C_G(h)| \geq (|H| - 1) + (k_+(G, H) - 1) \left(\frac{|H|}{\mu(1)} - 1 \right).$$

REFERENCES

- [1] Robinson, G.R., *A bound on norms of generalized characters with applications*, Journal of Algebra, **212**, (1999), 660-668.
- [2] Robinson, G.R., *More bounds on norms of generalized characters with applications to p -local bounds and blocks*, Bulletin of London Mathematical Society, **37**, (4), (2005), 555-565.
- [3] Robinson, G.R., *On generalized characters of nilpotent groups*, Journal of Algebra, **308**, (2007), 822-827.