

On linear forms containing the Euler constant*

A.I.Aptekarev

1 Introduction

We study the arithmetical nature of two numbers

$$\gamma := - \int_0^\infty \ln x e^{-x} dx \quad \text{and} \quad \delta := \int_0^\infty \ln(x+1) e^{-x} dx .$$

The first number is the famous Euler constant

$$\gamma := \lim_{n \rightarrow \infty} \left(\sum_{k=1}^n \frac{1}{k} - \ln(n) \right) \approx 0.577 \dots$$

The second number

$$\delta := \int_0^\infty \frac{e^{-x} dx}{x+1} \approx 0.596 \dots$$

is related to the Laguerre polynomials, which give a sequence of rational approximants for δ

$$\frac{\tilde{p}_n}{\tilde{q}_n} \rightarrow \delta, \quad n \rightarrow \infty, \tag{1}$$

generated by the recurrence relations

$$\tilde{q}_{n+1} = 2(n+1)\tilde{q}_n - n^2\tilde{q}_{n-1},$$

with initial condition

$$\begin{aligned} \tilde{p}_0 &= 0, & \tilde{p}_1 &= 1, \\ \tilde{q}_0 &= 1, & \tilde{q}_1 &= 2. \end{aligned}$$

The Perron asymptotics for the Laguerre polynomials

$$\begin{aligned} \tilde{q}_n &= n! \frac{e^{2\sqrt{n}}}{\sqrt[4]{n}} \left(\frac{1}{2\sqrt{\pi e}} + O(n^{-1/2}) \right), \\ \tilde{p}_n - \delta \tilde{q}_n &= O \left(n! \frac{e^{-2\sqrt{n}}}{\sqrt[4]{n}} \right), \end{aligned}$$

confirm (1). However, the rational approximants (1) do not imply that δ is an irrational number. As far as we know, irrationality of δ is still an open problem, as well as the famous open problem about irrationality of the Euler constant γ .

*The work was supported by Programm 1 DMS RAS and grants N-Sh-3906.2008.1, RFBR-08-01-00179.

2 Result

The following fact holds true

Theorem 1. *One of two numbers γ or δ is an irrational number.*

This result is a corollary of

Theorem 2. *Given sequences of numbers*

$$u := \{u_n\}, \quad v := \{v_n\}, \quad w := \{w_n\},$$

generated by the recurrence relations

$$\begin{aligned} (16n+1)(16n-15)u_{n+1} = & (16n-15)(256n^3 + 528n^2 + 352n + 73)u_n \\ & - (16n+17)(128n^3 + 40n^2 - 82n - 45)u_{n-1} \\ & + n^2(16n+17)(16n+1)u_{n-2}, \end{aligned} \quad (2)$$

with initial conditions

$$\begin{aligned} u_0 = -2, u_1 = 7, u_2 = 558, \\ v_0 = -1, v_1 = -22, v_2 = -1518, \\ w_0 = 0, w_1 = -17, w_2 = -1209, \end{aligned}$$

Then

- 1) $u_n, v_n, w_n \in \mathbb{Z}$;
- 2) *difference equation (2) has solutions with three different asymptotics as $n \rightarrow \infty$*

$$u_n, v_n, w_n = O\left(\frac{(2n)!4^n}{n^{3/2}}\right),$$

$$[w_n + (e\gamma + \delta)u_n], [v_n + eu_n] = O\left(\frac{e^{\sqrt{2n}}n^{5/4}}{4^n}\right), \quad (3)$$

$$l_n := u_n\delta - \gamma q_n + w_n = \frac{n^{5/4}}{e^{\sqrt{2n}}4^n} \left(\frac{2\sqrt[4]{2}}{e^{3/8}} + O(n^{-1/2}) \right).$$

Remark 1. Validity of Theorem 1 follows from $l_n \rightarrow 0$ as $n \rightarrow \infty$. However, the growth of the coefficients u_n, v_n, w_n is too fast. It does not allow one to conclude the rational independence of the numbers $(\gamma, \delta, 1)$.

Remark 2. Theorem 2 also implies that

$$\varepsilon := \gamma + \frac{\delta}{e} = - \int_0^1 \ln x e^{-x} dx \quad \text{and} \quad (e\varepsilon),$$

are irrational numbers (in view of $w_n + (\gamma + \frac{\delta}{e})v_n \rightarrow 0$ as $n \rightarrow \infty$).

3 A rational approximation of Euler constant

The numbers generated by (2) are intimately related with rational approximants of the Euler constant (studied in [1]). These approximants are produced by the functional Hermite-Pade rational approximants for a system of four functions $\{\hat{\mu}_1, \hat{\mu}_2, \hat{s}_1, \hat{s}_2\}$:

$$\hat{\mu}_k(z) := \int_0^1 \frac{w_k(z)dx}{z-x}, \quad \hat{s}_k(z) := \int_1^\infty \frac{w_k(z)dx}{z-x}, \quad (4)$$

where

$$w_k(x) := x^{\alpha_k} (1-x)^\alpha e^{-\beta x}, \quad k = 1, 2. \quad (5)$$

For the denominators Q_n of the Hermite-Pade rational approximants of this system (4)-(5) the generalized Rodrigues formula was known (see the example of subsection 2.2 in [2]; for similar systems, see also [3] and [4])

$$Q_n(z) = \frac{1}{(n!)^2} w_2^{-1} \frac{d^n}{dz^n} \left[w_2 z^n w_1^{-1} \frac{d^n}{dz^n} [w_1 z^n (1-z)^{2n}] \right]. \quad (6)$$

Then the above mentioned rational approximants $\frac{p_n}{q_n}$ of the Euler constant γ are defined as

$$p_n - \gamma q_n := \int_0^\infty Q_n(x) \ln(x) e^{-x} dx =: f_n, \quad (7)$$

where Q_n is taken from (6) – (5) with parameters $\alpha_1 = \alpha_2 = 0$ $\alpha = -\beta = 1$.

In [5]–[6] recurrence relations (of six, seven and eight terms) for polynomials (6) were studied, which in [7] eventually brought a four-term recurrence relation for sequences of numbers $p := \{p_n\}$ and $q := \{q_n\}$

$$(16n - 15) q_{n+1} = (128n^3 + 40n^2 - 82n - 45) q_n - n^2(256n^3 - 240n^2 + 64n - 7) q_{n-1} + n^2(n-1)^2(16n+1) q_{n-2}, \quad (8)$$

with initial conditions

$$\begin{aligned} p_0 &= 0, \quad p_1 = 2, \quad p_2 = 31, \\ q_0 &= 1, \quad q_1 = 3, \quad q_2 = 50. \end{aligned}$$

We also highlight a solution $r := \{r_n\}$ of difference equation (8) with initial conditions

$$r_0 = 0, \quad r_1 = 1, \quad r_2 = 24.$$

The fact that numbers $q_n, p_n, r_n \in \mathbb{Z}$ are integers for $n \in \mathbb{N}$ was proven in [10]. Finally in [8] and [9], the following asymptotics for $\{q_n, p_n, r_n\}$ and the linear forms with these coefficients were obtained

$$q_n, p_n, r_n = O\left(\frac{(2n)! e^{\sqrt{2n}}}{\sqrt[4]{n}}\right),$$

$$f_n := p_n - \gamma q_n = (2n)! \frac{e^{-\sqrt{2n}}}{\sqrt[4]{n}} \left(\frac{2\sqrt{\pi}}{(4e)^{3/8}} + O(n^{-1/2}) \right), \quad (9)$$

$$g_n := ep_n - (e\gamma + \delta)q_n + r_n = \frac{1}{16^n} \left(\frac{1}{8} + O(n^{-1}) \right).$$

4 Proof of Theorem 2

A) We define

$$u_n := \frac{\Delta_n^{(qp)}}{(n!)^2}, \quad v_n := \frac{\Delta_n^{(qr)}}{(n!)^2}, \quad w_n := \frac{\Delta_n^{(pr)}}{(n!)^2}, \quad (10)$$

where we use a notation

$$\Delta_n^{(ab)} := \det \begin{pmatrix} a_{n+1} & a_n \\ b_{n+1} & b_n \end{pmatrix}, \quad a := \{a_n\}, \quad b := \{b_n\}. \quad (11)$$

B) Substituting the recurrence relations

$$\begin{pmatrix} a_{n+1} \\ b_{n+1} \end{pmatrix} = A_n \begin{pmatrix} a_n \\ b_n \end{pmatrix} + B_n \begin{pmatrix} a_{n-1} \\ b_{n-1} \end{pmatrix} + C_n \begin{pmatrix} a_{n-2} \\ b_{n-2} \end{pmatrix}$$

into determinant (11), we get

$$\begin{vmatrix} a_{n+1} & a_n \\ b_{n+1} & b_n \end{vmatrix} = -B_n \begin{vmatrix} a_n & a_{n-1} \\ b_n & b_{n-1} \end{vmatrix} - C_n A_{n-1} \begin{vmatrix} a_{n-1} & a_{n-2} \\ b_{n-1} & b_{n-2} \end{vmatrix} + C_n C_{n-1} \begin{vmatrix} a_{n-2} & a_{n-3} \\ b_{n-2} & b_{n-3} \end{vmatrix},$$

that from (8) leads to (2).

C) The fact that $\frac{\Delta_n^{(qp)}}{(n!)^2} \in \mathbb{Z}$ is proven in [11]. We get that $v_n, w_n \in \mathbb{Z}$ analogously.

D) The asymptotics of

$$\Delta_n^{(qp)} = \begin{vmatrix} q_{n+1} & q_n \\ p_{n+1} - \gamma q_{n+1} & p_n - \gamma q_n \end{vmatrix} = O\left(\frac{(2n)!^2}{n^{3/2}}\right)$$

was computed in [11]. The same way, from (9) we deduce

$$\Delta_n^{(qr)} = \begin{vmatrix} q_{n+1} & q_n \\ r_{n+1} - \gamma q_{n+1} & r_n - \gamma q_n \end{vmatrix} = O\left(\frac{(2n)!^2}{n^{3/2}}\right).$$

Noticing that

$$\Delta_n^{(pr)} \frac{1}{e} - \Delta_n^{(pq)} \left(\gamma + \frac{\delta}{e} \right) = \Delta_n^{(pg)},$$

we get from (9) asymptotics for $[w_n + (e\gamma + \delta)u_n]$ and w_n . Analogously, the identity

$$\Delta_n^{(qp)} + \Delta_n^{(qr)} \frac{1}{e} = \Delta_n^{(qg)}$$

brings the asymptotics for $(v_n + e u_n)$. Finally,

$$-\Delta_n^{(pq)} \frac{\delta}{e} + \Delta_n^{(pr)} \frac{1}{e} - \Delta_n^{(qr)} \frac{\gamma}{e} = \Delta_n^{(fg)}$$

from (9) brings the asymptotics for l_n in (3).

The Theorem is proved.

References

- [1] A. I. Aptekarev (ed.), *Rational approximation of Euler's constant and recurrence relations*, Collected papers, Sovrem. Probl. Mat., 9, Steklov Math. Inst., RAS, Moscow, 2007, 84 p.(in Russian)
- [2] A. I. Aptekarev, A. Branquinho, and W. Van Assche, *Multiple orthogonal polynomials for classical weights*, Trans. Amer. Math. Soc., 2003, 355, 3887–3914.
- [3] V. N. Sorokin, *Simultaneous Pade approximants for finite and infinite intervals*, Izv. Vyssh. Uchebn. Zaved. Mat. 8 (1984) 45-52; English transl, in Sov. Math. Iz. VUZ 28 (1984).
- [4] V.N. Sorokin, *On simultaneous Pade approximants for functions of Stieltjes type*, Sibirsk. Math. Zh. 31 (N5) (1990); English transl, in Siberian Math. J. 31 (1990).
- [5] D. V. Khristoforov, *Recurrence relations for the Hermite-Pade approximants for a system of four functions of Markov and Stieltjes type*, Rational approximation of Euler's constant and recurrence relations, Collected papers, Sovrem. Probl. Mat., 9, Steklov Math. Inst., RAS, Moscow, 2007, 11–26, (in Russian)
- [6] A. I. Bogolyubskii, *Recurrence relations with rational coefficients for some jointly orthogonal polynomials defined by Rodriguess formula*, Rational approximation of Euler's constant and recurrence relations, Collected papers, Sovrem. Probl. Mat., 9, Steklov Math. Inst., RAS, Moscow, 2007, 27–35, (in Russian)
- [7] A. I. Aptekarev, D. N. Tulyakov, *Four-termed recurrence relations for γ -forms*, Rational approximation of Euler's constant and recurrence relations, Collected papers, Sovrem. Probl. Mat., 9, Steklov Math. Inst., RAS, Moscow, 2007, 37–43, (in Russian)
- [8] A. I. Aptekarev, V. G. Lysov, *Asymptotics of γ -forms jointly generated by orthogonal polynomials*, Rational approximation of Euler's constant and recurrence relations, Collected papers, Sovrem. Probl. Mat., 9, Steklov Math. Inst., RAS, Moscow, 2007, 55–62, (in Russian)
- [9] D. N. Tulyakov, *On some procedures for finding asymptotic expansions of solutions of difference equations*, Rational approximation of Euler's constant and recurrence relations, Collected papers, Sovrem. Probl. Mat., 9, Steklov Math. Inst., RAS, Moscow, 2007, 45–53, (in Russian)
- [10] D. N. Tulyakov, *System of recurrence relations for rational approximants of Euler constant*, Mat. Zametki (in press); English translation in Math. Notes (in press)
- [11] A. I. Aptekarev, D. N. Tulyakov, *On determinant of a lattice generated by rational approximants of the Euler constant*, Trudy Moskov. Mat. Obshch. (in press); English translation in Trans. Moscow Math. Soc (in press)