

Some integrals involving the Stieltjes constants

Donal F. Connon

dconnon@btopenworld.com

12 February 2009

Abstract

Some new integrals involving the Stieltjes constants are developed in this paper.

1. Introduction

Using the binomial theorem we obtain

$$\begin{aligned}\int_0^{\infty} e^{-ux} (1 - e^{-x})^n x^{p-1} dx &= \int_0^{\infty} e^{-ux} \sum_{j=0}^n \binom{n}{j} (-1)^j e^{-jx} x^{p-1} dx \\ &= \sum_{j=0}^n \binom{n}{j} (-1)^j \int_0^{\infty} e^{-(u+j)x} x^{p-1} dx\end{aligned}$$

and employing the definition of the gamma function for $\text{Re}(p) > 0$

$$\Gamma(p) = \int_0^{\infty} e^{-x} x^{p-1} dx$$

we see for $\text{Re}(u) > 0$ that

$$\frac{\Gamma(p)}{u^p} = \int_0^{\infty} e^{-ux} x^{p-1} dx$$

Hence for $\text{Re}(u) > 0$ and $\text{Re}(p) > 0$ we have

$$(1.1) \quad \int_0^{\infty} e^{-ux} (1 - e^{-x})^n x^{p-1} dx = \Gamma(p) \sum_{j=0}^n \binom{n}{j} \frac{(-1)^j}{(u+j)^p}$$

With $n = 0$ in (1.1) we have $\int_0^{\infty} e^{-ux} x^{p-1} dx = \frac{\Gamma(p)}{u^p}$ and therefore (1.1) is also valid for

$n = 0$. The integral (1.1) appears in Gradshteyn and Ryzhik [16, p.357, Eq. 3.432.1] where it is attributed to Bierens de Haan.

As $p \rightarrow 0$ the right-hand side of (1.1) is indeterminate but we may use L'Hôpital's rule when considering the ratio

$$\frac{\sum_{j=0}^n \binom{n}{j} \frac{(-1)^j}{(u+j)^p}}{1/\Gamma(p)}$$

(since $\sum_{j=0}^n \binom{n}{j} (-1)^j = \delta_{n,0}$ we require that $n \neq 0$ in order to ensure that the numerator is equal to zero as $p \rightarrow 0$).

As regards the numerator, it is easily seen that

$$\frac{d}{dp} \sum_{j=0}^n \binom{n}{j} \frac{(-1)^j}{(u+j)^p} = - \sum_{j=0}^n \binom{n}{j} \frac{(-1)^j \log(u+j)}{(u+j)^p}$$

We have using Euler's reflection formula for the gamma function

$$\begin{aligned} \frac{d}{dp} [1/\Gamma(p)] &= \frac{1}{\pi} \frac{d}{dp} [\Gamma(1-p) \sin \pi p] \\ &= \Gamma(1-p) \cos \pi p - \frac{1}{\pi} \Gamma'(1-p) \sin \pi p \end{aligned}$$

and therefore we see that where r is a positive integer

$$\lim_{p \rightarrow 1-r} \frac{d}{dp} [1/\Gamma(p)] = (-1)^{r+1} \Gamma(r)$$

This gives us

$$(1.2) \quad \int_0^{\infty} \frac{e^{-ux} (1-e^{-x})^n}{x^r} dx = \frac{(-1)^r}{\Gamma(r)} \sum_{j=0}^n \binom{n}{j} (-1)^j (u+j)^{r-1} \log(u+j)$$

The above integral was originally given by Anglesio [3] in 1997. [Could one adopt a Eulerian approach at this stage and assume that r was a continuous variable with the representation $(-1)^r = \cos \pi r$?]

We now let $u=1$ and $r=1$ to obtain

$$(1.3) \quad \int_0^{\infty} \frac{e^{-x}(1-e^{-x})^n}{x} dx = -\sum_{j=0}^n \binom{n}{j} (-1)^j \log(1+j)$$

Making the substitution $y = e^{-x}$, equation (1.3) becomes for $n \geq 1$

$$(1.4) \quad \int_0^1 \frac{(1-y)^n}{\log y} dy = \sum_{j=0}^n \binom{n}{j} (-1)^j \log(1+j)$$

and this is in agreement with G&R [16, p.541, Eqn. 4.267.1] as corrected by Boros and Moll (see the Errata to G&R [16] dated 26 April 2005).

Making the same substitution $y = e^{-x}$ in (1.2) we see that for $n \geq 1$

$$(1.5) \quad \int_0^1 \frac{y^{u-1}(1-y)^n}{\log^r y} dy = \frac{1}{\Gamma(r)} \sum_{j=0}^n \binom{n}{j} (-1)^j (u+j)^{r-1} \log(u+j)$$

and we have the summation (where we are required to start at $n = 1$)

$$\sum_{n=1}^{\infty} \frac{1}{n+1} \int_0^1 \frac{y^{u-1}(1-y)^n}{\log^r y} dy = \frac{1}{\Gamma(r)} \sum_{n=1}^{\infty} \frac{1}{n+1} \sum_{j=0}^n \binom{n}{j} (-1)^j (u+j)^{r-1} \log(u+j)$$

Using the logarithmic expansion

$$\sum_{n=1}^{\infty} \frac{(1-y)^n}{n+1} = -\frac{1}{1-y} [\log y + 1 - y]$$

we therefore obtain

$$(1.6) \quad \int_0^1 \left[\frac{1}{1-y} + \frac{1}{\log y} \right] \frac{y^{u-1}}{\log^{r-1} y} dy = -\frac{1}{\Gamma(r)} \sum_{n=1}^{\infty} \frac{1}{n+1} \sum_{j=0}^n \binom{n}{j} (-1)^j (u+j)^{r-1} \log(u+j)$$

Starting the summation at $n = 0$ gives us

(1.7)

$$\int_0^1 \left[\frac{1}{1-y} + \frac{1}{\log y} \right] \frac{y^{u-1}}{\log^{r-1} y} dy = \frac{1}{\Gamma(r)} u^{r-1} \log u - \frac{1}{\Gamma(r)} \sum_{n=0}^{\infty} \frac{1}{n+1} \sum_{j=0}^n \binom{n}{j} (-1)^j (u+j)^{r-1} \log(u+j)$$

and reference to the Hasse identity [18]

$$(s-1)\zeta(s) = \sum_{n=0}^{\infty} \frac{1}{n+1} \sum_{j=0}^n \binom{n}{j} \frac{(-1)^j}{(u+j)^{s-1}}$$

shows that the right-hand side of (1.7) is related to $\zeta'(2-r, u)$.

With $r=1$ we get

$$(1.8) \quad \int_0^1 \left[\frac{1}{1-y} + \frac{1}{\log y} \right] y^{u-1} dy = \log u - \sum_{n=0}^{\infty} \frac{1}{n+1} \sum_{j=0}^n \binom{n}{j} (-1)^j \log(u+j)$$

and it is known from [17] and [12] that the digamma function $\psi(u)$ may be represented by

$$(1.9) \quad \psi(u) = \sum_{n=0}^{\infty} \frac{1}{n+1} \sum_{j=0}^n \binom{n}{j} (-1)^j \log(u+j)$$

Equation (1.8) then becomes

$$(1.10) \quad \int_0^1 \left[\frac{1}{1-y} + \frac{1}{\log y} \right] y^{u-1} dy = \log u - \psi(u)$$

This integral is due to Binet [22, p.175] and with $u=1$ in (1.10) we obtain the well-known integral for Euler's constant [6, p.178]

$$(1.11) \quad \gamma = \int_0^1 \left[\frac{1}{1-y} + \frac{1}{\log y} \right] dy$$

The identity (1.9) may also be obtained in the following way by employing a method due to Dirichlet [25]. We see that

$$\begin{aligned} \frac{\Gamma(u+h) - \Gamma(u)}{h} &= \frac{\Gamma(u+h)}{\Gamma(1+h)} \Gamma(h) - \frac{\Gamma(u+h)}{\Gamma(1+h)} \frac{\Gamma(u)\Gamma(h)}{\Gamma(u+h)} \\ &= \frac{\Gamma(u+h)}{\Gamma(1+h)} [\Gamma(h) - B(u, h)] \\ &= \frac{\Gamma(u+h)}{\Gamma(1+h)} \int_0^1 \left[\frac{1}{\log^{1-h}(1/y)} - \frac{y^{u-1}}{(1-y)^{1-h}} \right] dy \end{aligned}$$

and in the limit as $h \rightarrow 0$ we have

$$\Gamma'(u) = -\Gamma(u) \int_0^1 \left[\frac{1}{\log y} + \frac{y^{u-1}}{1-y} \right] dy$$

Therefore we have

$$(1.12) \quad \psi(u) = -\int_0^1 \left[\frac{1}{\log y} + \frac{y^{u-1}}{1-y} \right] dy$$

We see that

$$\frac{1}{u} = \int_0^1 y^{u-1} dy$$

and integration of this results in the well-known Frullani integral [8, p.472]

$$(1.13) \quad \log u = \int_0^1 \left[\frac{y^{u-1}}{\log y} - \frac{1}{\log y} \right] dy$$

We then have

$$\log u - \psi(u) = \int_0^1 \left[\frac{1}{1-y} + \frac{1}{\log y} \right] y^{u-1} dy$$

and, comparing this with (1.8), results in (1.9).

Making the substitution $y = e^{-x}$ in (1.10) we see that

$$I = \int_0^1 \left[\frac{1}{1-y} + \frac{1}{\log y} \right] y^{u-1} dy = \int_0^\infty \left[\frac{1}{1-e^{-x}} - \frac{1}{x} \right] e^{-ux} dx$$

and noting that

$$\frac{1}{1-e^{-x}} = \frac{e^x}{e^x - 1} = 1 + \frac{1}{e^x - 1}$$

we obtain

$$I = \frac{1}{u} + \int_0^\infty \left[\frac{1}{e^x - 1} - \frac{1}{x} \right] e^{-ux} dx$$

Therefore we have from (1.10) [26, p.15]

$$(1.14) \quad \int_0^{\infty} \left[\frac{1}{e^x - 1} - \frac{1}{x} \right] e^{-ux} dx = -\frac{1}{u} + \log u - \psi(u) = \log u - \psi(1+u)$$

Integrating (1.14) with respect to u over the interval $[0, t]$ gives us a Binet-type integral representation for $\log \Gamma(u)$

$$(1.15) \quad \int_0^{\infty} \left[\frac{1}{e^x - 1} - \frac{1}{x} \right] \frac{1 - e^{-tx}}{x} dx = t \log t - t - \log \Gamma(1+t)$$

Alternatively, integrating (1.14) with respect to u over the interval $[1, t]$ gives us

$$(1.16) \quad \int_0^{\infty} \left[\frac{1}{e^x - 1} - \frac{1}{x} \right] \frac{e^{-x} - e^{-tx}}{x} dx = t \log t - t + 1 - \log \Gamma(1+t)$$

which recently appeared in [15].

Further integrations would in turn develop integrals for the Barnes multiple gamma functions defined in [26, p.24]. This is considered further in [12].

Further consideration of integrals such as (1.10) is contained in Appendix B to this paper.

2. Some integrals involving the Stieltjes constants

We now boldly assume that (1.6) is also valid in the case where r is a continuous variable (perhaps Carlson's Theorem [27, p.185] could be utilised to provide a rigorous proof: further advice on this aspect would be appreciated). In this regard, it may be noted that Adamchik [1] made a similar assumption in making the transition from equation (23) to equation (24) in his 1997 paper, "A class of logarithmic integrals".

Differentiation of (1.6) with respect to r results in

$$(2.1) \quad \int_0^1 \left[\frac{1}{1-y} + \frac{1}{\log y} \right] \frac{y^{u-1} \log |\log y|}{\log^{r-1} y} dy$$

$$= \frac{1}{\Gamma(r)} \sum_{n=1}^{\infty} \frac{1}{n+1} \sum_{j=0}^n \binom{n}{j} (-1)^j (u+j)^{r-1} \log^2(u+j) - \frac{\psi(r)}{\Gamma(r)} \sum_{n=1}^{\infty} \frac{1}{n+1} \sum_{j=0}^n \binom{n}{j} (-1)^j (u+j)^{r-1} \log(u+j)$$

Letting $r = 1$ results in

$$\int_0^1 \left[\frac{1}{1-y} + \frac{1}{\log y} \right] y^{u-1} \log |\log y| dy$$

$$= \sum_{n=1}^{\infty} \frac{1}{n+1} \sum_{j=0}^n \binom{n}{j} (-1)^j \log^2(u+j) + \gamma \sum_{n=1}^{\infty} \frac{1}{n+1} \sum_{j=0}^n \binom{n}{j} (-1)^j \log(u+j)$$

and starting the summation at $n = 0$ gives us

$$\int_0^1 \left[\frac{1}{1-y} + \frac{1}{\log y} \right] y^{u-1} \log |\log y| dy = \sum_{n=0}^{\infty} \frac{1}{n+1} \sum_{j=0}^n \binom{n}{j} (-1)^j \log^2(u+j) \\ + \gamma \sum_{n=0}^{\infty} \frac{1}{n+1} \sum_{j=0}^n \binom{n}{j} (-1)^j \log(u+j) - \log^2 u - \gamma \log u$$

We have previously shown in [13] that for integer $p \geq 0$

$$(2.2) \quad \gamma_p(u) = -\frac{1}{p+1} \sum_{n=0}^{\infty} \frac{1}{n+1} \sum_{j=0}^n \binom{n}{j} (-1)^j \log^{p+1}(u+j)$$

where the Stieltjes constants $\gamma_n(u)$ are the coefficients in the Laurent expansion of the Hurwitz zeta function $\zeta(s, u)$ about $s = 1$

$$(2.3) \quad \zeta(s, u) = \sum_{n=0}^{\infty} \frac{1}{(n+u)^s} = \frac{1}{s-1} + \sum_{n=0}^{\infty} \frac{(-1)^n}{n!} \gamma_n(u) (s-1)^n$$

and $\gamma_0(u) = -\psi(u)$, where $\psi(u)$ is the digamma function which is the logarithmic derivative of the gamma function $\psi(u) = \frac{d}{du} \log \Gamma(u)$. It is easily seen from the definition of the Hurwitz zeta function that $\zeta(s, 1) = \zeta(s)$ and accordingly that $\gamma_n(1) = \gamma_n$.

Hence we obtain

$$(2.4) \quad \int_0^1 \left[\frac{1}{1-y} + \frac{1}{\log y} \right] y^{u-1} \log |\log y| dy = -2\gamma_1(u) - \gamma\gamma_0(u) - \log^2 u - \gamma \log u$$

Letting $u = 1$ results in (however see (B.30) in Appendix B)

$$(2.5) \quad \int_0^1 \left[\frac{1}{1-y} + \frac{1}{\log y} \right] \log |\log y| dy = -(2\gamma_1 + \gamma^2)$$

We define $f(y)$ as

$$f(y) = \frac{1}{1-y} + \frac{1}{\log y} = \frac{\log y + 1 - y}{(1-y)\log y}$$

Using L'Hôpital's rule we can easily deduce that $\lim_{y \rightarrow 0} f(y) = 1$ and $\lim_{y \rightarrow 1} f(y) = -1/2$ and, since $f'(y) < 0$, we note that $f(y)$ is monotonically decreasing on the interval $[0, 1]$. However, the range of $\log|\log y|$ is $(-\infty, \infty)$ and hence it is not straightforward to determine the sign of the integral in (2.5). Another approach is therefore necessary.

With the substitution $y = 1/t$ in (2.5) we get

$$\int_0^1 \left[\frac{1}{1-y} + \frac{1}{\log y} \right] \log|\log y| dy = \int_1^\infty \left[\frac{1}{t-1} - \frac{1}{t \log t} \right] \frac{\log|\log(1/t)|}{t} dt$$

We denote $g(t)$ by

$$g(t) = \frac{1}{t-1} - \frac{1}{t \log t} = \frac{t \log t - (t-1)}{t(t-1) \log t} = \frac{\log t - (1-1/t)}{(t-1) \log t}$$

and using L'Hôpital's rule we obtain

$$\lim_{t \rightarrow 1} g(t) = 1/2 \quad \lim_{t \rightarrow \infty} g(t) = 0$$

Since $\log t > 1 - 1/t$ we see that

$$g(t) > 0 \text{ for } t \in [1, \infty)$$

We see that $h(t) = \frac{\log|\log(1/t)|}{t} \rightarrow -\infty$ as $t \rightarrow 1$ and using L'Hôpital's rule we have

$$\frac{\log|\log(1/t)|}{t} \rightarrow 0 \text{ as } t \rightarrow \infty$$

We have the derivative

$$h'(t) = \frac{-\frac{1}{|\log(1/t)|} - \log|\log(1/t)|}{t^2}$$

Since $\log t > 1 - 1/t$ for $t > 0$ we also have

$$\log|\log(1/t)| > 1 - \frac{1}{|\log(1/t)|} \quad \text{for } |\log(1/t)| > 0$$

and therefore $h'(t)$ is negative with the result that $h(t)$ is monotonic decreasing and negative for $t \in [1, \infty)$. We accordingly deduce that the integrand in (2.5) is negative and that the integral is also negative. This then enables us to conclude that

$$2\gamma_1 + \gamma^2 > 0$$

The constants η_k are defined by the following Maclaurin expansion

$$\log[(s-1)\zeta(s)] = -\sum_{k=0}^{\infty} \frac{\eta_k}{k+1} (s-1)^{k+1} = -\sum_{k=1}^{\infty} \frac{\eta_{k-1}}{k} (s-1)^k$$

where

$$\eta_0 = -\gamma$$

$$\eta_1 = 2\gamma_1 + \gamma^2$$

We note the approximate numerical values of the first three Stieltjes constants from [2]

$$\gamma = 0.5772 \dots \quad \gamma_1 = -0.0728 \dots \quad \gamma_2 = -0.0096 \dots$$

Coffey [9a] has shown that the sequence (η_n) has strict sign alteration, i.e.

$$\eta_n = (-1)^{n+1} \varepsilon_n$$

where ε_n are positive constants. It may be noted that the above analysis has confirmed, without performing any numerical calculations, that η_1 is indeed positive.

We see that $\frac{\log^{2n} |\log(1/t)|}{t} > 0$ and with $j(t) = \frac{\log^{2n+1} |\log(1/t)|}{t}$ we see that

$$\begin{aligned} j'(t) &= -\frac{(2n+1) \log^{2n} |\log(1/t)| \frac{1}{|\log(1/t)|} + \log^{2n+1} |\log(1/t)|}{t^2} \\ &= -\frac{\log^{2n} |\log(1/t)|}{t^2} \left(\frac{(2n+1)}{|\log(1/t)|} + \log |\log(1/t)| \right) \end{aligned}$$

Since $\log t > 1 - 1/t$ for $t > 0$ we also have

$$\log^{2n+1} t > (1 - 1/t)^{2n+1} > 1 - \frac{2n+1}{t}$$

where, in the last step, we have employed Bernoulli's inequality. We then have

$$\log^{2n+1} |\log(1/t)| > 1 - \frac{2n+1}{|\log(1/t)|} \quad \text{for } |\log(1/t)| > 0$$

and conclude that $j'(t)$ is negative.

We have therefore determined that

$$(2.6) \quad \int_0^1 \left[\frac{1}{1-y} + \frac{1}{\log y} \right] \log^n |\log y| dy = \int_1^\infty \left[\frac{1}{t-1} - \frac{1}{t \log t} \right] \frac{\log^n |\log(1/t)|}{t} dt = (-1)^n d_n$$

where d_n are positive constants. This relationship is used in (3.15) below.

□

For convenience, we now write (2.1) as

$$(2.7) \quad \int_0^1 \left[\frac{1}{1-y} + \frac{1}{\log y} \right] \frac{y^{u-1} \log |\log y|}{\log^{r-1} y} dy = \frac{1}{\Gamma(r)} S(r, u, 2) - \frac{\psi(r)}{\Gamma(r)} S(r, u, 1)$$

where $S(r, u, p)$ is defined as

$$S(r, u, p) = \sum_{n=1}^{\infty} \frac{1}{n+1} \sum_{j=0}^n \binom{n}{j} (-1)^j (u+j)^{r-1} \log^p (u+j)$$

and note that $\frac{d}{dr} S(r, u, p) = S(r, u, p+1)$.

Differentiation of (2.7) with respect to r gives us

$$(2.8) \quad -\int_0^1 \left[\frac{1}{1-y} + \frac{1}{\log y} \right] \frac{y^{u-1} \log^2 |\log y|}{\log^{r-1} y} dy = \frac{1}{\Gamma(r)} S(r, u, 3) - 2 \frac{\psi(r)}{\Gamma(r)} S(r, u, 2) \\ - \frac{\psi'(r) - [\psi(r)]^2}{\Gamma(r)} S(r, u, 1)$$

We additionally define $S_0(r, u, p)$ by

$$S_0(r, u, p) = \sum_{n=0}^{\infty} \frac{1}{n+1} \sum_{j=0}^n \binom{n}{j} (-1)^j (u+j)^{r-1} \log^p(u+j)$$

where the summation starts at $n = 0$ and we note that

$$S_0(r, u, p) = S(r, u, p) + u^{r-1} \log^p u$$

and

$$S_0(r, 1, p) = S(r, 1, p)$$

We also see from (2.2) with $r = 1$ that

$$S(1, u, p+1) = -(p+1)\gamma_p(u) - \log^{p+1} u$$

and

$$S(1, 1, p+1) = -(p+1)\gamma_p$$

We then have with $r = 1$ in (2.8)

$$-\int_0^1 \left[\frac{1}{1-y} + \frac{1}{\log y} \right] y^{u-1} \log^2 |\log y| dy = S(1, u, 3) - 2\psi(1)S(1, u, 2) - (\psi'(1) - [\psi(1)]^2) S(1, u, 1)$$

which is equivalent to

$$(2.9) \quad \int_0^1 \left[\frac{1}{1-y} + \frac{1}{\log y} \right] y^{u-1} \log^2 |\log y| dy \\ = 3\gamma_2(u) + \log^3 u + 2\gamma[2\gamma_1(u) + \log^2 u] - [\zeta(2) - \gamma^2][\gamma(u) + \log u]$$

Therefore we get with $u = 1$

$$(2.10) \quad \int_0^1 \left[\frac{1}{1-y} + \frac{1}{\log y} \right] \log^2 |\log y| dy = 3\gamma_2 + 4\gamma\gamma_1 - [\zeta(2) - \gamma^2]\gamma$$

Therefore we see that

$$(2.11) \quad 3\gamma_2 + 4\gamma\gamma_1 - [\zeta(2) - \gamma^2]\gamma > 0$$

Coffey [9b] has given the following recurrence relation for the η_n constants

$$\eta_n = (-1)^{n+1} \left[\frac{n+1}{n!} \gamma_n + \sum_{k=0}^{n-1} \frac{(-1)^{k+1}}{(n-k-1)!} \gamma_{n-k-1} \eta_k \right]$$

from which we obtain

$$(2.12) \quad \eta_2 = - \left[\frac{3}{2} \gamma_2 - \eta_0 \gamma_1 + \eta_1 \gamma \right] = - \left[\frac{3}{2} \gamma_2 + 3\gamma \gamma_1 + \gamma^3 \right]$$

We then see that

$$\left[\frac{3}{2} \gamma_2 + 3\gamma \gamma_1 + \gamma^3 \right] - \gamma \left[\gamma_1 + \frac{1}{2} \gamma + \frac{1}{2} \zeta(2) \right] > 0$$

but it is not clear if such inequalities are germane in connection with the Riemann Hypothesis via the Li/Keiper criterion.

Differentiation of (2.8) with respect to r gives us

$$(2.13) \quad \int_0^1 \left[\frac{1}{1-y} + \frac{1}{\log y} \right] \frac{y^{u-1} \log^3 |\log y|}{\log^{r-1} y} dy = \frac{1}{\Gamma(r)} S(r, u, 4) - 3 \frac{\psi(r)}{\Gamma(r)} S(r, u, 3) \\ - 3 \frac{\psi'(r) - [\psi(r)]^2}{\Gamma(r)} S(r, u, 2) - \frac{\psi''(r) - 2\psi(r)\psi'(r) + [\psi(r)]^3}{\Gamma(r)} S(r, u, 1)$$

Further differentiations will result in integrals $\int_0^1 \left[\frac{1}{1-y} + \frac{1}{\log y} \right] y^{u-1} \log^n |\log y| dy$ involving the Stieltjes constants $\gamma_n(u)$ (together with the resulting inequalities). □

Differentiation of (2.4) with respect to u results in

$$\int_0^1 \left[\frac{1}{1-y} + \frac{1}{\log y} \right] y^{u-1} \log y \cdot \log |\log y| dy = -2\gamma'_1(u) - \gamma'_0(u) - \frac{\gamma + 2 \log u}{u}$$

and with $u = 1$ we get using $\gamma'_0(u) = \psi'(u)$

$$- \int_0^1 \left[\frac{1}{1-y} + \frac{1}{\log y} \right] \log y \cdot \log |\log y| dy = 2\gamma'_1(1) - \gamma \zeta(2) + \gamma$$

Using equation (4.3.244) in [13]

$$\gamma_1'(1) = 2\pi^2 \zeta'(-1) + \zeta(2)(\gamma + \log 2\pi)$$

we may write this as

$$(2.14) \quad -\int_0^1 \left[\frac{1}{1-y} + \frac{1}{\log y} \right] \log y \cdot \log |\log y| dy = 4\pi^2 \zeta'(-1) + \zeta(2)(\gamma + 2 \log 2\pi) + \gamma$$

Using (C.1) this may be written as

$$(2.15) \quad -\int_0^1 \frac{\log y \cdot \log |\log y|}{1-y} dy = 4\pi^2 \zeta'(-1) + \zeta(2)(\gamma + 2 \log 2\pi)$$

3. An application of the (exponential) complete Bell polynomials

Alternatively, we may write (1.6) as

$$(3.1) \quad \Gamma(r) \int_0^1 \left[\frac{1}{1-y} + \frac{1}{\log y} \right] \frac{y^{u-1}}{\log^{r-1} y} dy = -\sum_{n=1}^{\infty} \frac{1}{n+1} \sum_{j=0}^n \binom{n}{j} (-1)^j (u+j)^{r-1} \log(u+j)$$

Letting $h(r)$ be defined as

$$h(r) = \frac{\Gamma(r)}{\log^{r-1} y}$$

we have

$$\log h(r) = \log \Gamma(r) - (r-1) \log |\log y|$$

and differentiation results in

$$(3.2) \quad h'(r) = h(r)[\psi(r) - \log |\log y|]$$

Using (3.2) we now differentiate (3.1) with respect to r to obtain

$$\Gamma(r) \int_0^1 \left[\frac{1}{1-y} + \frac{1}{\log y} \right] \frac{y^{u-1} [\psi(r) - \log |\log y|]}{\log^{r-1} y} dy = -\sum_{n=1}^{\infty} \frac{1}{n+1} \sum_{j=0}^n \binom{n}{j} (-1)^j (u+j)^{r-1} \log^2(u+j)$$

which may be expressed as

$$\Gamma(r) \int_0^1 \left[\frac{1}{1-y} + \frac{1}{\log y} \right] \frac{y^{u-1} [\psi(r) - \log |\log y|]}{\log^{r-1} y} dy = 2\gamma_1(u) + u^{r-1} \log^2 u$$

Letting $r = 1$ this becomes

$$\int_0^1 \left[\frac{1}{1-y} + \frac{1}{\log y} \right] y^{u-1} [\gamma + \log |\log y|] dy = -2\gamma_1(u) - \log^2 u$$

and with $u = 1$ we have

$$\int_0^1 \left[\frac{1}{1-y} + \frac{1}{\log y} \right] [\gamma + \log |\log y|] dy = -2\gamma_1$$

which of course is consistent with our previous results.

We now wish to obtain a formula for the m th derivative of (3.1).

It is shown in Appendix A that

$$(3.3) \quad \frac{d^m}{dx^m} e^{f(x)} = e^{f(x)} Y_m(f^{(1)}(x), f^{(2)}(x), \dots, f^{(m)}(x))$$

where the (exponential) complete Bell polynomials $Y_n(x_1, \dots, x_n)$ are defined by $Y_0 = 1$ and for $n \geq 1$

$$(3.4) \quad Y_n(x_1, \dots, x_n) = \sum_{\pi(n)} \frac{n!}{k_1! k_2! \dots k_n!} \left(\frac{x_1}{1!} \right)^{k_1} \left(\frac{x_2}{2!} \right)^{k_2} \dots \left(\frac{x_n}{n!} \right)^{k_n}$$

where the sum is taken over all partitions $\pi(n)$ of n , i.e. over all sets of integers k_j such that

$$k_1 + 2k_2 + 3k_3 + \dots + nk_n = n$$

Suppose that $h'(x) = h(x)g(x)$ and let $f(x) = \log h(x)$. We see that

$$f'(x) = \frac{h'(x)}{h(x)} = g(x)$$

and then using (3.3) above we have

$$(3.5) \quad \frac{d^m}{dx^m} h(x) = \frac{d^m}{dx^m} e^{\log h(x)} = h(x) Y_m \left(g(x), g^{(1)}(x), \dots, g^{(m-1)}(x) \right)$$

As an example, letting $h(x) = \Gamma(x)$ in (3.5) we obtain

$$(3.6) \quad \begin{aligned} \frac{d^m}{dx^m} e^{\log \Gamma(x)} &= \Gamma^{(m)}(x) = \Gamma(x) Y_m \left(\psi(x), \psi^{(1)}(x), \dots, \psi^{(m-1)}(x) \right) \\ &= \int_0^{\infty} t^{x-1} e^{-t} \log^m t \, dt \end{aligned}$$

and since [26, p.22]

$$(3.7) \quad \psi^{(p)}(x) = (-1)^{p+1} p! \zeta(p+1, x)$$

we may express $\Gamma^{(m)}(x)$ in terms of $\psi(x)$ and the Hurwitz zeta functions. In particular, Kölbig [20] noted that

$$(3.8) \quad \Gamma^{(m)}(1) = Y_m(-\gamma, x_1, \dots, x_{m-1})$$

where $x_p = (-1)^{p+1} p! \zeta(p+1)$. Values of $\Gamma^{(m)}(1)$ for $m \leq 10$ are reported in [26, p.265] and the first three are

$$\Gamma^{(1)}(1) = -\gamma$$

$$\Gamma^{(2)}(1) = \zeta(2) + \gamma^2$$

$$\Gamma^{(3)}(1) = -[2\zeta(3) + 3\gamma\zeta(2) + \gamma^3]$$

As shown in Appendix C, we note that $\Gamma^{(n)}(1)$ has the same sign as $(-1)^n$. This was also reported as an exercise in Apostol's book [4, p.303].

As a variation of (3.5) above, suppose that $j'(x) = j(x)[g(x) + \alpha]$ where α is independent of x and let $f(x) = \log j(x)$. We see that

$$f'(x) = \frac{j'(x)}{j(x)} = g(x) + \alpha \quad \text{and} \quad f^{(k+1)}(x) = g^{(k+1)}(x) \quad \text{for } k \geq 1$$

and therefore we obtain

$$(3.9) \quad \frac{d^m}{dx^m} j(x) = \frac{d^m}{dx^m} e^{\log j(x)} = j(x) Y_m \left(g(x) + \alpha, g^{(1)}(x), \dots, g^{(m-1)}(x) \right)$$

It is also shown in Appendix A that

$$Y_n(x_1 + \alpha, \dots, x_n) = \sum_{k=0}^n \binom{n}{k} \alpha^{n-k} Y_k(x_1, \dots, x_k)$$

and we then determine that

$$(3.10) \quad \frac{d^m}{dx^m} j(x) = j(x) \sum_{k=0}^m \binom{m}{k} \alpha^{m-k} Y_k \left(g(x), g^{(1)}(x), \dots, g^{(k-1)}(x) \right)$$

Now, referring back to (3.2), we see that with $g(r) = \psi(r) - \log |\log y|$

$$\frac{d^m}{dr^m} \frac{\Gamma(r)}{\log^{r-1} y} = \frac{\Gamma(r)}{\log^{r-1} y} \sum_{k=0}^m \binom{m}{k} (-1)^{m-k} \log^{m-k} |\log y| \cdot Y_k \left(\psi(r), \psi^{(1)}(r), \dots, \psi^{(k-1)}(r) \right)$$

and since we have from (3.5)

$$\Gamma^{(k)}(r) = \Gamma(r) Y_k \left(\psi(r), \psi^{(1)}(r), \dots, \psi^{(k-1)}(r) \right)$$

we obtain

$$\frac{d^m}{dr^m} \frac{\Gamma(r)}{\log^{r-1} y} = \frac{\Gamma(r)}{\log^{r-1} y} \sum_{k=0}^m \binom{m}{k} (-1)^{m-k} \Gamma^{(k)}(r) \log^{m-k} |\log y|$$

Hence, differentiating (3.1) m times with respect to r results in

$$(3.11) \quad \Gamma(r) \sum_{k=0}^m \binom{m}{k} (-1)^{m-k} \Gamma^{(k)}(r) \int_0^1 \left[\frac{1}{1-y} + \frac{1}{\log y} \right] \frac{y^{u-1} \log^{m-k} |\log y|}{\log^{r-1} y} dy$$

$$= - \sum_{n=1}^{\infty} \frac{1}{n+1} \sum_{j=0}^n \binom{n}{j} (-1)^j (u+j)^{r-1} \log^{m+1}(u+j)$$

Letting $r = 1$ this becomes

$$\sum_{k=0}^m \binom{m}{k} (-1)^{m-k} \Gamma^{(k)}(1) \int_0^1 \left[\frac{1}{1-y} + \frac{1}{\log y} \right] y^{u-1} \log^{m-k} |\log y| dy$$

$$\begin{aligned}
&= -\sum_{n=1}^{\infty} \frac{1}{n+1} \sum_{j=0}^n \binom{n}{j} (-1)^j \log^{m+1}(u+j) \\
&= (m+1)\gamma_m(u) + \log^{m+1} u
\end{aligned}$$

With $u = 1$ we obtain

$$(3.12) \quad (m+1)\gamma_m = \sum_{k=0}^m \binom{m}{k} (-1)^{m-k} \Gamma^{(k)}(1) \int_0^1 \left[\frac{1}{1-y} + \frac{1}{\log y} \right] \log^{m-k} |\log y| dy$$

For example, with $m = 1$ we get

$$2\gamma_1 = -\int_0^1 \left[\frac{1}{1-y} + \frac{1}{\log y} \right] \log |\log y| dy - \gamma \int_0^1 \left[\frac{1}{1-y} + \frac{1}{\log y} \right] dy$$

which is in agreement with our previous results.

Equation (3.12) sheds a little light upon the complexity involved in the alteration in the signs of γ_m because

$$(3.13) \quad \Gamma^{(k)}(1) = (-1)^k c_k$$

$$(3.14) \quad \int_0^1 \left[\frac{1}{1-y} + \frac{1}{\log y} \right] \log^{m-k} |\log y| dy = (-1)^{m-k} d_{m-k}$$

where c_k and d_k are positive constants. The relationship (3.13) is derived in Appendix C and (3.14) was shown in (2.6) above. We then have

$$(3.15) \quad (m+1)\gamma_m = \sum_{k=0}^m \binom{m}{k} (-1)^k c_k d_{m-k}$$

□

Referring back to (1.1) we make the summation

$$\sum_{n=0}^{\infty} \frac{1}{n+1} \int_0^{\infty} e^{-ax} (1-e^{-x})^n x^{p-1} dx = \Gamma(p) \sum_{n=0}^{\infty} \frac{1}{n+1} \sum_{k=0}^n \binom{n}{k} \frac{(-1)^k}{(a+k)^p}$$

and we have

$$\begin{aligned}
\sum_{n=0}^{\infty} \frac{1}{n+1} \int_0^{\infty} e^{-ax} (1-e^{-x})^n x^{p-1} dx &= \int_0^{\infty} e^{-ax} x^{p-1} \sum_{n=0}^{\infty} \frac{(1-e^{-x})^n}{n+1} dx \\
&= \int_0^{\infty} \frac{e^{-ax} x^{p-1}}{1-e^{-x}} \sum_{n=0}^{\infty} \frac{(1-e^{-x})^{n+1}}{n+1} dx \\
&= \int_0^{\infty} \frac{e^{-ax} x^{p-1} \log(1-e^{-x})}{1-e^{-x}} dx
\end{aligned}$$

Therefore we obtain

$$\int_0^{\infty} \frac{e^{-ax} x^{p-1} \log(1-e^{-x})}{1-e^{-x}} dx = \Gamma(p) \sum_{n=0}^{\infty} \frac{1}{n+1} \sum_{k=0}^n \binom{n}{k} \frac{(-1)^k}{(a+k)^p}$$

and with $a = 1$ we have

$$\int_0^{\infty} \frac{e^{-x} x^{p-1} \log(1-e^{-x})}{1-e^{-x}} dx = \Gamma(p) \sum_{n=0}^{\infty} \frac{1}{n+1} \sum_{k=0}^n \binom{n}{k} \frac{(-1)^k}{(1+k)^p}$$

We note that the Bernoulli polynomials may be represented by [17]

$$B_k(a) = \sum_{n=0}^{\infty} \frac{1}{n+1} \sum_{k=0}^n \binom{n}{k} (-1)^k (a+k)^k$$

and hence for $m \geq 1$

$$B_{2m+1}(1) = \sum_{n=0}^{\infty} \frac{1}{n+1} \sum_{k=0}^n \binom{n}{k} (-1)^k (1+k)^{2m+1} = 0$$

We now consider the limit as $p \rightarrow -(2m+1)$ where m is a positive integer and $m \geq 1$; the right-hand side of () is indeterminate but we may use L'Hôpital's rule when considering the ratio

$$\frac{\sum_{n=0}^{\infty} \frac{1}{n+1} \sum_{k=0}^n \binom{n}{k} \frac{(-1)^k}{(a+k)^p}}{1/\Gamma(p)}$$

As shown previously, using Euler's reflection formula we have

$$\frac{d}{dp} [1/\Gamma(p)] = \Gamma(1-p) \cos \pi p - \frac{1}{\pi} \Gamma'(1-p) \sin \pi p$$

and therefore we see that where $p \rightarrow -(2m+1)$

$$\lim_{p \rightarrow -(2m+1)} \frac{d}{dp} [1/\Gamma(p)] = (-1)^{2m+1} \Gamma(2m)$$

We thereby obtain

$$\int_0^{\infty} \frac{e^{-x} \log(1-e^{-x})}{(1-e^{-x})x^{2m}} dx = (-1)^{2m+1} \Gamma(2m) \sum_{n=0}^{\infty} \frac{1}{n+1} \sum_{k=0}^n \binom{n}{k} (-1)^k (1+k)^{2m+1} \log(1+k)$$

Appendix A

A brief survey of the (exponential) complete Bell polynomials

The (exponential) complete Bell polynomials are defined by $Y_0 = 1$ and for $n \geq 1$

$$(A1) \quad Y_n(x_1, \dots, x_n) = \sum_{\pi(n)} \frac{n!}{k_1! k_2! \dots k_n!} \left(\frac{x_1}{1!}\right)^{k_1} \left(\frac{x_2}{2!}\right)^{k_2} \dots \left(\frac{x_n}{n!}\right)^{k_n}$$

where the sum is taken over all partitions $\pi(n)$ of n , i.e. over all sets of integers k_j such that

$$k_1 + 2k_2 + 3k_3 + \dots + nk_n = n$$

The complete Bell polynomials have integer coefficients and the first six are set out below [11, p.307]

$$(A.2) \quad Y_1(x_1) = x_1$$

$$Y_2(x_1, x_2) = x_1^2 + x_2$$

$$Y_3(x_1, x_2, x_3) = x_1^3 + 3x_1x_2 + x_3$$

$$Y_4(x_1, x_2, x_3, x_4) = x_1^4 + 6x_1^2x_2 + 4x_1x_3 + 3x_2^2 + x_4$$

$$Y_5(x_1, x_2, x_3, x_4, x_5) = x_1^5 + 10x_1^3x_2 + 10x_1^2x_3 + 15x_1x_2^2 + 5x_1x_4 + 10x_2x_3 + x_5$$

$$Y_6(x_1, x_2, x_3, x_4, x_5, x_6) = x_1^6 + 6x_1x_5 + 15x_2x_4 + 10x_2^3 + 15x_1^2x_4 + 15x_2^3 + 60x_1x_2x_3$$

$$+20x_1^3x_3 + 45x_1^2x_2^2 + 15x_1^4x_1 + x_6$$

The total number of terms $\pi(n)$ increases rapidly; for example, as reported by Bell [5] in 1934, we have $\pi(22) = 1002$ terms.

The modulus operandi of the summation in (A.1) is easily illustrated by the following example: if $n = 4$, then $k_1 + 2k_2 + 3k_3 + 4k_4 = 4$ is satisfied by the integers in the following array

$$\begin{bmatrix} k_1 & k_2 & k_3 & k_4 \\ 4 & 0 & 0 & 0 \\ 2 & 1 & 0 & 0 \\ 1 & 0 & 1 & 0 \\ 0 & 2 & 0 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

The complete Bell polynomials are also given by the exponential generating function (Comtet [11, p.134])

$$(A.3) \quad \exp\left(\sum_{j=1}^{\infty} x_j \frac{t^j}{j!}\right) = 1 + \sum_{n=1}^{\infty} Y_n(x_1, \dots, x_n) \frac{t^n}{n!} = \sum_{n=0}^{\infty} Y_n(x_1, \dots, x_n) \frac{t^n}{n!}$$

Let us now consider a function $f(x)$ which has a Taylor series expansion around x : we have

$$\begin{aligned} e^{f(x+t)} &= \exp\left(\sum_{j=0}^{\infty} f^{(j)}(x) \frac{t^j}{j!}\right) = e^{f(x)} \exp\left(\sum_{j=1}^{\infty} f^{(j)}(x) \frac{t^j}{j!}\right) \\ &= e^{f(x)} \left\{ 1 + \sum_{n=1}^{\infty} Y_n\left(f^{(1)}(x), f^{(2)}(x), \dots, f^{(n)}(x)\right) \frac{t^n}{n!} \right\} \end{aligned}$$

We see that

$$\frac{d^m}{dx^m} e^{f(x)} = \frac{\partial^m}{\partial x^m} e^{f(x+t)} \Big|_{t=0} = \frac{\partial^m}{\partial t^m} e^{f(x+t)} \Big|_{t=0}$$

and we therefore obtain (as noted by Kölbig [20])

$$(A.4) \quad \frac{d^m}{dx^m} e^{f(x)} = e^{f(x)} Y_m \left(f^{(1)}(x), f^{(2)}(x), \dots, f^{(m)}(x) \right)$$

Differentiating (A.4) we see that

$$Y_{m+1} \left(f^{(1)}(x), f^{(2)}(x), \dots, f^{(m+1)}(x) \right) = \left(f^{(1)}(x) + \frac{d}{dx} \right) Y_m \left(f^{(1)}(x), f^{(2)}(x), \dots, f^{(m)}(x) \right)$$

Suppose that $h'(x) = h(x)g(x)$ and let $f(x) = \log h(x)$. We see that

$$f'(x) = \frac{h'(x)}{h(x)} = g(x)$$

and then using (A.4) above we have

$$(A.5) \quad \frac{d^m}{dx^m} h(x) = \frac{d^m}{dx^m} e^{\log h(x)} = h(x) Y_m \left(g(x), g^{(1)}(x), \dots, g^{(m-1)}(x) \right)$$

As an example, letting $f(x) = \log \Gamma(x)$ in (A.4) we obtain

$$(A.6) \quad \frac{d^m}{dx^m} e^{\log \Gamma(x)} = \Gamma^{(m)}(x) = \Gamma(x) Y_m \left(\psi(x), \psi^{(1)}(x), \dots, \psi^{(m-1)}(x) \right)$$

$$= \int_0^{\infty} t^{x-1} e^{-t} \log^m t \, dt$$

and since [26, p.22]

$$(A.7) \quad \psi^{(p)}(x) = (-1)^{p+1} p! \zeta(p+1, x)$$

we may express $\Gamma^{(m)}(x)$ in terms of $\psi(x)$ and the Hurwitz zeta functions. In particular, Kölbig [20] notes that

$$(A.8) \quad \Gamma^{(m)}(1) = Y_m(-\gamma, x_1, \dots, x_{m-1})$$

where $x_p = (-1)^{p+1} p! \zeta(p+1)$. Values of $\Gamma^{(m)}(1)$ are reported in [26, p.265] for $m \leq 10$ and the first three are

$$\Gamma^{(1)}(1) = -\gamma$$

$$\Gamma^{(2)}(1) = \zeta(2) + \gamma^2$$

$$\Gamma^{(3)}(1) = -[2\zeta(3) + 3\gamma\zeta(2) + \gamma^3]$$

The general form is

$$\Gamma^{(m)}(1) = (-1)^m \sum_{j=1}^m \varepsilon_{mj}$$

where ε_{mj} are positive constants.

We could also, for example, let $f(x) = \log \sin(\pi x)$ in (A.4) to obtain complete Bell polynomials involving the derivatives of $\cot(\pi x)$.

We now consider a minor modification of the arguments of the exponential generating function

$$\exp\left(\sum_{j=1}^{\infty} x_j \frac{t^j}{j!}\right) = \sum_{n=0}^{\infty} Y_n(x_1, \dots, x_n) \frac{t^n}{n!}$$

where we let $x_1 \rightarrow x_1 + \alpha$ and the other arguments remain unchanged. We then see that

$$\exp\left(\alpha t + \sum_{j=1}^{\infty} x_j \frac{t^j}{j!}\right) = \sum_{n=0}^{\infty} Y_n(x_1 + \alpha, \dots, x_n) \frac{t^n}{n!}$$

Since

$$\exp\left(\alpha t + \sum_{j=1}^{\infty} x_j \frac{t^j}{j!}\right) = \exp(\alpha t) \exp\left(\sum_{j=1}^{\infty} x_j \frac{t^j}{j!}\right)$$

we easily determine that

$$(A.9) \quad \exp(\alpha t) \sum_{n=0}^{\infty} Y_n(x_1, \dots, x_n) \frac{t^n}{n!} = \sum_{n=0}^{\infty} Y_n(x_1 + \alpha, \dots, x_n) \frac{t^n}{n!}$$

Using the Cauchy series product formula we find that

$$\sum_{n=0}^{\infty} \alpha^n \frac{t^n}{n!} \sum_{n=0}^{\infty} Y_n(x_1, \dots, x_n) \frac{t^n}{n!} = \sum_{n=0}^{\infty} \sum_{k=0}^n \frac{\alpha^{n-k}}{(n-k)!} \frac{Y_k(x_1, \dots, x_k)}{k!} t^n$$

and equating coefficients we see that

$$(A.10) \quad Y_n(x_1 + \alpha, \dots, x_n) = \sum_{k=0}^n \binom{n}{k} \alpha^{n-k} Y_k(x_1, \dots, x_k)$$

□

As a variation of (A.5) above, suppose that $j'(x) = j(x)[g(x) + \alpha]$ where α is independent of x and let $f(x) = \log j(x)$. We see that

$$f'(x) = \frac{j'(x)}{j(x)} = g(x) + \alpha \quad \text{and} \quad f^{(k+1)}(x) = g^{(k)}(x) \quad \text{for } k \geq 1$$

and therefore we obtain

$$(A.11) \quad \frac{d^m}{dx^m} j(x) = \frac{d^m}{dx^m} e^{\log j(x)} = j(x) Y_m(g(x) + \alpha, g^{(1)}(x), \dots, g^{(m-1)}(x))$$

Using (A.10) we then determine that

$$(A.12) \quad \frac{d^m}{dx^m} j(x) = j(x) \sum_{k=0}^m \binom{m}{k} \alpha^{m-k} Y_k(g(x), g^{(1)}(x), \dots, g^{(k-1)}(x))$$

The relation (A.10) may be generalised as follows to

$$(A.13) \quad Y_n(x_1 + y_1, \dots, x_n + y_n) = \sum_{k=0}^n \binom{n}{k} Y_{n-k}(x_1, \dots, x_{n-k}) Y_k(y_1, \dots, y_k)$$

and we note that

$$Y_k(\alpha, 0, \dots, 0) = \alpha^k$$

Equation (A.13) follows by noting that

$$\begin{aligned} Y_n(x_1 + y_1, \dots, x_n + y_n) &= \exp\left(\sum_{j=1}^{\infty} (x_j + y_j) \frac{t^j}{j!}\right) = \exp\left(\sum_{j=1}^{\infty} x_j \frac{t^j}{j!}\right) \exp\left(\sum_{j=1}^{\infty} y_j \frac{t^j}{j!}\right) \\ &= \sum_{n=0}^{\infty} Y_n(x_1, \dots, x_n) \frac{t^n}{n!} \cdot \sum_{n=0}^{\infty} Y_n(y_1, \dots, y_n) \frac{t^n}{n!} \end{aligned}$$

and, as before, we apply the Cauchy series product formula.

Appendix B

Miscellaneous integral identities

In [24] Rivoal recently gave an elementary proof of the following lemma:

$$(B.1) \quad \int_0^1 x^n \Omega(x) dx = \gamma - [H_n - \log(n+1)]$$

where, for convenience $\Omega(x)$, is defined as $\Omega(x) = \frac{1}{1-x} + \frac{1}{\log x}$ and H_n is the harmonic number defined by

$$H_n = \sum_{k=1}^n \frac{1}{k}$$

Using L'Hôpital's rule it is easily seen that $\Omega(x)$ is continuous on $[0,1]$. For $x \in (0,1)$ we also have by a straightforward integration

$$\Omega(x) = \int_0^1 \frac{1-x^t}{1-x} dt$$

We see that

$$\int_0^1 \Omega(x) dx = \int_0^1 \left(\frac{1}{1-x} + \frac{1}{\log x} \right) dx = \int_0^{\infty} \left[\frac{1}{1-e^{-u}} - \frac{1}{u} \right] e^{-u} du = \gamma$$

Using the well-known integral representation of the Riemann zeta function

$$\zeta(s)\Gamma(s) = \int_0^{\infty} \frac{u^{s-1}}{e^u - 1} du$$

we also see that $\Gamma(s) = (s-1)\Gamma(s-1) = (s-1) \int_0^{\infty} e^{-u} u^{s-2} du$. Therefore we get

$$\left[\zeta(s) - \frac{1}{s-1} \right] \Gamma(s) = \int_0^{\infty} u^{s-1} \left[\frac{1}{e^u - 1} - \frac{1}{ue^u} \right] du$$

Hence, in the limit as $s \rightarrow 1$, we obtain the familiar limit

$$\lim_{s \rightarrow 1} \left[\zeta(s) - \frac{1}{s-1} \right] \Gamma(s) = \lim_{s \rightarrow 1} \left[\zeta(s) - \frac{1}{s-1} \right] = \int_0^{\infty} \left[\frac{1}{1-e^{-u}} - \frac{1}{u} \right] e^{-u} du = \gamma$$

We now continue with Rivoal's lemma.

We have by simple algebra

$$\int_0^1 x^n \Omega(x) dx = \int_0^1 \Omega(x) dx - \int_0^1 \frac{x^n - 1}{x - 1} dx + \int_0^1 \frac{x^n - 1}{\log x} dx$$

and this gives us

$$(B.2) \quad \int_0^1 x^n \left(\frac{1}{1-x} + \frac{1}{\log x} \right) dx = \gamma - H_n + \int_0^1 \frac{x^n - 1}{\log x} dx = \gamma - [H_n - \log(n+1)]$$

It may be noted that this is equivalent to letting $u = n+1$ in (1.10) because

$$\psi(n+1) = -\gamma + H_n$$

Hence by summation of (B.2) we obtain for $\text{Re}(s) > 1$

$$(B.3) \quad \gamma \zeta(s) - \sum_{n=1}^{\infty} \frac{H_n}{n^s} + \sum_{n=1}^{\infty} \frac{\log(n+1)}{n^s} = \int_0^1 Li_s(x) \left(\frac{1}{1-x} + \frac{1}{\log x} \right) dx$$

where $Li_s(x)$ is the polylogarithm function defined by

$$Li_s(x) = \sum_{n=1}^{\infty} \frac{x^n}{n^s}$$

In 1997 Candelpergher et al. [9] produced a somewhat similar result in the case where $s = 2$ by reference to "Ramanujan summation" but unfortunately I am not au fait with the underlying analysis contained in that paper. □

We have from [23]

$$(B.4) \quad H_n - \log n - \gamma = \int_0^{\infty} e^{-nx} \left[\frac{1}{x} - \frac{1}{e^x - 1} \right] dx$$

and on summation we obtain

$$\sum_{n=1}^{\infty} \frac{H_n}{n^s} + \zeta'(s) - \gamma \zeta(s) = \sum_{n=1}^{\infty} \frac{1}{n^s} \int_0^{\infty} e^{-nx} \left[\frac{1}{x} - \frac{1}{e^x - 1} \right] dx$$

In particular we have

$$\begin{aligned}
\sum_{n=1}^{\infty} \frac{1}{n^2} \int_0^{\infty} e^{-nx} \left[\frac{1}{x} - \frac{1}{e^x - 1} \right] dx &= \sum_{n=1}^{\infty} \int_0^{\infty} e^{-nu} du \int_0^{\infty} e^{-nv} dv \int_0^{\infty} e^{-nx} \left[\frac{1}{x} - \frac{1}{e^x - 1} \right] dx \\
&= \sum_{n=1}^{\infty} \int_0^{\infty} du \int_0^{\infty} dv \int_0^{\infty} e^{-n(x+u+v)} \left[\frac{1}{x} - \frac{1}{e^x - 1} \right] dx \\
&= \int_0^{\infty} du \int_0^{\infty} dv \int_0^{\infty} \frac{1}{e^{x+u+v} - 1} \left[\frac{1}{x} - \frac{1}{e^x - 1} \right] dx
\end{aligned}$$

We have

$$\int_0^{\infty} \frac{1}{e^{x+u+v} - 1} du = \int_0^{\infty} \frac{e^{-(x+u+v)}}{1 - e^{-(x+u+v)}} du = \log[1 - e^{-(x+u+v)}] \Big|_0^{\infty} = -\log[1 - e^{-(x+v)}]$$

and the Wolfram Integrator tells us that

$$\int \log[1 - e^{-(x+v)}] dv = \frac{1}{2} v^2 + v \log \frac{[1 - e^{-(x+v)}]}{[1 - e^{(x+v)}]} - Li_2[e^{(x+v)}]$$

With the substitution $y = e^{-v}$ we can easily find that

$$\int_0^{\infty} \log[1 - e^{-(x+v)}] dv = -Li_2[e^{-x}]$$

and hence we obtain

$$\sum_{n=1}^{\infty} \frac{H_n}{n^2} + \zeta'(2) - \gamma \zeta(2) = \int_0^{\infty} Li_2[e^{-x}] \left[\frac{1}{x} - \frac{1}{e^x - 1} \right] dx$$

or equivalently

$$\text{(B.5)} \quad 2\zeta(3) + \zeta'(2) - \gamma \zeta(2) = \int_0^{\infty} Li_2[e^{-x}] \left[\frac{1}{x} - \frac{1}{e^x - 1} \right] dx$$

where we have employed the well-known Euler sum [26, p.103] $\sum_{n=1}^{\infty} \frac{H_n}{n^2} = 2\zeta(3)$.

With the substitution $t = e^{-x}$ this may be written as

$$(B.6) \quad 2\zeta(3) + \zeta'(2) - \gamma\zeta(2) = \int_0^1 Li_2(t) \left[\frac{1}{t-1} - \frac{1}{t \log t} \right] dt$$

We see from (1.13) that

$$\sum_{n=1}^{\infty} \frac{\log n}{n^s} = \sum_{n=1}^{\infty} \frac{1}{n^s} \int_0^{\infty} \frac{e^{-x} - e^{-nx}}{x} dx$$

which may be written as

$$-\zeta'(s) = \int_0^{\infty} \frac{\zeta(s)e^{-x} - Li_s[e^{-x}]}{x} dx$$

With $t = e^{-x}$ this becomes

$$(B.7) \quad \zeta'(s) = \int_0^1 \frac{\zeta(s)t - Li_s(t)}{t \log t} dt$$

and with $s = 2$ we get

$$(B.8) \quad \zeta'(2) = \int_0^1 \frac{\zeta(2)t - Li_2(t)}{t \log t} dt$$

We note from (B.6) that

$$(B.9) \quad \begin{aligned} 2\zeta(3) + \zeta'(2) - \gamma\zeta(2) &= \int_0^1 Li_2(t) \left[\frac{1}{t-1} - \frac{1}{t \log t} \right] dt \\ &= \int_0^1 \left[\frac{Li_2(t)}{t-1} - \frac{Li_2(t)}{t \log t} \right] dt \\ &= \int_0^1 \left[\frac{Li_2(t)}{t-1} - \frac{\zeta(2)}{\log t} + \frac{\zeta(2)}{\log t} - \frac{Li_2(t)}{t \log t} \right] dt \\ &= \int_0^1 \left[\frac{Li_2(t)}{t-1} - \frac{\zeta(2)}{\log t} \right] dt + \int_0^1 \left[\frac{\zeta(2)}{\log t} - \frac{Li_2(t)}{t \log t} \right] dt \end{aligned}$$

and using (B.8) this becomes

$$= \int_0^1 \left[\frac{Li_2(t)}{t-1} - \frac{\zeta(2)}{\log t} \right] dt + \zeta'(2)$$

We therefore see that

$$(B.10) \quad 2\zeta(3) - \gamma\zeta(2) = \int_0^1 \left[\frac{Li_2(t)}{t-1} - \frac{\zeta(2)}{\log t} \right] dt$$

We note from (B.3) with $s = 2$ that

$$2\zeta(3) - \sum_{n=1}^{\infty} \frac{\log(n+1)}{n^2} - \gamma\zeta(2) = \int_0^1 Li_2(t) \left(\frac{1}{t-1} - \frac{1}{\log t} \right) dt$$

Subtracting (B.6) results in

$$(B.11) \quad \zeta'(2) + \sum_{n=1}^{\infty} \frac{\log(n+1)}{n^2} = \int_0^1 \frac{(t-1)Li_2(t)}{t \log t} dt$$

or equivalently

$$\sum_{n=1}^{\infty} \frac{\log(n+1)}{n^2} - \sum_{n=1}^{\infty} \frac{\log n}{n^2} = \int_0^1 \frac{(t-1)Li_2(t)}{t \log t} dt$$

□

We may write (B.10) as the limit

$$2\zeta(3) - \gamma\zeta(2) = \lim_{x \rightarrow 1} \int_0^x \left[\frac{Li_2(t)}{t-1} - \frac{\zeta(2)}{\log t} \right] dt$$

Using integration by parts we can easily determine that

$$\int \frac{Li_2(t)}{t-1} dt = Li_2(t) \log(1-t) + \log t \log^2(1-t) + 2Li_2(1-t) \log(1-t) - 2Li_3(1-t)$$

and we therefore have

$$\int_0^x \frac{Li_2(t)}{t-1} dt = Li_2(x) \log(1-x) + \log x \log^2(1-x) + 2Li_2(1-x) \log(1-x) - 2Li_3(1-x) + 2\zeta(3)$$

Using L'Hôpital's rule we find that

$$\lim_{x \rightarrow 1} [(1-x) \log(1-x)] = 0$$

and hence we have

$$\lim_{x \rightarrow 1} [Li_2(1-x) \log(1-x)] = 0$$

We therefore have the limit

$$\lim_{x \rightarrow 1} \int_0^x \frac{Li_2(t)}{t-1} dt = \lim_{x \rightarrow 1} [Li_2(x) \log(1-x) + \log x \log^2(1-x)] + 2\zeta(3)$$

which then implies that

$$-\gamma\zeta(2) = \lim_{x \rightarrow 1} \left[Li_2(x) \log(1-x) + \log x \log^2(1-x) - \int_0^x \frac{\zeta(2)}{\log t} dt \right]$$

We therefore have

$$(B.12) \quad \gamma\zeta(2) = \lim_{x \rightarrow 1} \left[-Li_2(x) \log(1-x) - \log x \log^2(1-x) + \zeta(2) li(x) \right]$$

where $li(x)$ is the logarithmic integral. Nielsen [21, p.3] has shown that for $0 < x < 1$

$$(B.13) \quad li(x) = \gamma + \log(-\log x) + \sum_{n=1}^{\infty} \frac{\log^n x}{n!n}$$

and hence we have

$$\gamma\zeta(2) = \lim_{x \rightarrow 1} \left[-Li_2(x) \log(1-x) - \log x \log^2(1-x) + \zeta(2) [\gamma + \log(-\log x)] \right]$$

which gives us

$$\lim_{x \rightarrow 1} \left[-Li_2(x) \log(1-x) - \log x \log^2(1-x) + \zeta(2) \log(-\log x) \right] = 0$$

Using Euler's identity

$$\zeta(2) = \log x \log(1-x) + Li_2(x) + Li_2(1-x)$$

we may write this as

$$\lim_{x \rightarrow 1} [Li_2(1-x) \log(1-x) + \zeta(2) \log(1-x) + \zeta(2) \log(-\log x)] = 0$$

and hence we have

$$(B.14) \quad \lim_{x \rightarrow 1} [\log(1-x) + \log(-\log x)] = 0$$

Since

$$\frac{d}{dt} \log[\log(1/t)] = -\frac{1}{t \log(1/t)} = \frac{1}{t \log t}$$

integration by parts gives us

$$\int_0^x \log[\log(1/t)] dt = t \log[\log(1/t)] \Big|_0^x - \int_0^x \frac{dt}{\log t}$$

Using L'Hôpital's rule we see that

$$\lim_{t \rightarrow 0} t \log[\log(1/t)] = \lim_{t \rightarrow 0} \frac{\frac{d}{dt} \log[\log(1/t)]}{-1/t^2} = 0$$

and therefore we obtain

$$(B.15) \quad \int_0^x \log[\log(1/t)] dt = x \log[\log(1/x)] - li(x)$$

Since $\int_0^1 \log[\log(1/t)] dt = -\gamma$ we find that

$$(B.16) \quad \lim_{x \rightarrow 1} [x \log[\log(1/x)] - li(x)] = -\gamma$$

Integration gives us

$$\begin{aligned} \int_0^x \left[\frac{1}{1-t} + \frac{1}{\log t} \right] dt &= -\log(1-t) + li(t) \Big|_0^x \\ &= -\log(1-x) + li(x) \end{aligned}$$

We note from (B.2) that

$$\gamma = \int_0^1 \left[\frac{1}{1-y} + \frac{1}{\log y} \right] dy$$

and therefore we have

$$(B.17) \quad \lim_{x \rightarrow 1} [-\log(1-x) + li(x)] = \gamma$$

Therefore we see that

$$(B.18) \quad \lim_{x \rightarrow 1} [x \log[\log(1/x)] - \log(1-x)] = 0$$

Multiplying (B.12) by x , upon taking the limit, we see that

$$\lim_{x \rightarrow 1} [xli(x) - x \log(-\log x)] = \gamma$$

Combining this with (B.18) we obtain

$$\lim_{x \rightarrow 1} [xli(x) - li(x) + \log(1-x) - x \log(-\log x)] = 0$$

and hence

$$(B.19) \quad \lim_{x \rightarrow 1} [(x-1)li(x)] = 0$$

This may also be verified using L'Hôpital's rule

$$\lim_{x \rightarrow 1} [(x-1)li(x)] = \lim_{x \rightarrow 1} \frac{li(x)}{1/(x-1)} = -\lim_{x \rightarrow 1} \frac{(x-1)^2}{\log x} = -\lim_{x \rightarrow 1} 2x(x-1) = 0$$

The limit (B.19) in conjunction with (B.12) implies that

$$(B.20) \quad \lim_{x \rightarrow 1} [(x-1) \log(-\log x)] = 0$$

□

We refer back to the Frullani integral (1.13)

$$\log u = \int_0^1 \left[\frac{y^{u-1}}{\log y} - \frac{1}{\log y} \right] dy$$

where the third derivative results in

$$\frac{2}{u^3} = \int_0^1 y^{u-1} \log y^2 dy$$

We now let $u = n$ and make a summation to obtain the well-known integral

$$2\zeta(3) = 2 \sum_{n=1}^{\infty} \frac{1}{n^3} = \int_0^1 \frac{\log y^2}{1-y} dy$$

□

The following is extracted from Brede's dissertation [7]. In 1999 Coppo showed that there exists a polynomial $p_n(z)$ such that

$$x^n = \int_0^{\infty} p_n(x - \log z) e^{-z} dz$$

With $p_n(z) = \sum_{k=0}^n a_k z^k$ we have

$$\begin{aligned} \int_0^{\infty} p_n(x - \log z) e^{-z} dz &= \int_0^{\infty} \sum_{k=0}^n a_k (x - \log z)^k e^{-z} dz \\ &= \sum_{k=0}^n a_k \int_0^{\infty} \sum_{l=0}^k \binom{k}{l} x^l (-\log z)^{k-l} e^{-z} dz \\ &= \sum_{k=0}^n a_k \sum_{l=0}^k \binom{k}{l} (-1)^{k-l} \Gamma^{(k-l)}(1) x^l \end{aligned}$$

Hence we have

$$\int_0^{\infty} p_n(x - \log z) e^{-z} dz = \sum_{k=0}^n \left[\sum_{l=0}^{n-k} (-1)^l \binom{k+l}{l} \Gamma^{(l)}(1) a_{k+l} \right] x^k$$

and we then select the coefficients a_k such that

$$\sum_{l=0}^{n-k} (-1)^l \binom{k+l}{l} \Gamma^{(l)}(1) a_{k+l} = \begin{cases} 1 & \text{for } k = n \\ 0 & \text{for } k \text{ less than } n \end{cases}$$

resulting in

$$x^n = \int_0^{\infty} p_n(x - \log z) e^{-z} dz$$

For example we have

$$p_0(z) = 1$$

$$p_1(z) = z - \gamma$$

$$p_2(z) = z^2 - 2\gamma z + \gamma^2 - \zeta(2)$$

Letting $x \rightarrow \log x$ we obtain

$$\log^n x = \int_0^{\infty} p_n(\log x - \log z) e^{-z} dz$$

and, with the substitution $z/x = -\log t$, we have

$$\frac{\log^n x}{x} = \int_0^1 p_n[-\log \log(1/t)] t^{x-1} dt$$

Referring to the well-known expression for the Stieltjes constants [19, p.4]

$$\begin{aligned} \gamma_n &= \sum_{k=1}^{\infty} \left[\frac{\log^n k}{k} - \frac{\log^n(k+1) - \log^n k}{n+1} \right] = \sum_{k=1}^{\infty} \left[\frac{\log^n k}{k} - \int_k^{k+1} \frac{\log^n t}{t} dt \right] \\ &= \lim_{N \rightarrow \infty} \left[\sum_{k=1}^N \frac{\log^n k}{k} - \int_1^{N+1} \frac{\log^n t}{t} dt \right] \end{aligned}$$

we then see that

$$\begin{aligned} \gamma_n &= \lim_{N \rightarrow \infty} \left[\sum_{k=1}^N \int_0^1 p_n[-\log \log(1/t)] t^{k-1} dt - \int_1^{N+1} \int_0^1 p_n[-\log \log(1/t)] t^{x-1} dt dx \right] \\ &= \lim_{N \rightarrow \infty} \int_0^1 p_n[-\log \log(1/t)] \left[\sum_{k=0}^{N-1} t^k - \int_1^{N+1} t^x dx \right] dt \\ &= \lim_{N \rightarrow \infty} \int_0^1 p_n[-\log \log(1/t)] \left[\frac{t^N - 1}{t - 1} - \frac{t^N - 1}{\log t} \right] dt \end{aligned}$$

Hence, as shown more completely in Brede's dissertation, we obtain as $N \rightarrow \infty$

$$(B.21) \quad \gamma_n = \int_0^1 p_n[-\log \log(1/t)] \left[\frac{1}{\log t} - \frac{1}{t-1} \right] dt$$

Since $\frac{1}{\log t} - \frac{1}{t-1} = \int_0^1 \frac{1-u^t}{1-u} du$ we may write this as a double integral

$$\gamma_n = \int_0^1 \int_0^1 p_n[-\log \log(1/t)] \frac{1-u^t}{1-u} du dt$$

□

The following is extracted from Coppo's paper [14]. With the substitution $u = xt$ we see that

$$\begin{aligned} \int_0^\infty e^{-xt} \log^n t dt &= \frac{1}{x} \int_0^\infty e^{-u} (\log u - \log x)^n du \\ &= \sum_{k=0}^n \binom{n}{k} (-1)^k \frac{\log^k x}{x} \int_0^\infty e^{-u} \log^{n-k} u du \\ &= \sum_{k=0}^n \binom{n}{k} (-1)^k \frac{\log^k x}{x} \Gamma^{(n-k)}(1) \end{aligned}$$

Let us assume that

$$(B.22) \quad \sum_{k=0}^n \binom{n}{k} (-1)^k \Gamma^{(n-k)}(1) \hat{a}_k(t) = \log^n t$$

where $\hat{a}_k(t)$ are to be determined. With $n = 0$ we have

$$\hat{a}_0(t) = 1$$

and $n = 1$ gives us

$$\Gamma^{(1)}(1) - \hat{a}_1(t) = \log t$$

We then multiply (B.22) by e^{-xt} and integrate to obtain

$$\sum_{k=0}^n \binom{n}{k} (-1)^k \Gamma^{(n-k)}(1) \int_0^{\infty} e^{-xt} \hat{a}_k(t) dt = \int_0^{\infty} e^{-xt} \log^n t dt$$

We then have

$$(B.23) \quad \sum_{k=0}^n \binom{n}{k} (-1)^k \Gamma^{(n-k)}(1) \left[\frac{\log^k x}{x} - \int_0^{\infty} e^{-xt} \hat{a}_k(t) dt \right] = 0$$

and Coppo deduces that

$$\frac{\log^k x}{x} = \int_0^{\infty} e^{-xt} \hat{a}_k(t) dt$$

It seems to me that this solution is only unique provided the coefficients appearing in (B.23) are linearly independent.

The associated Stieltjes functions $\hat{\gamma}_k(x)$ are defined by

$$\hat{\gamma}_k(x) = \int_0^{\infty} e^{-xt} \left(\frac{1}{1-e^{-t}} - \frac{1}{t} \right) \hat{a}_k(t) dt$$

and we have in particular

$$\hat{\gamma}_0(x) = \int_0^{\infty} e^{-xt} \left(\frac{1}{1-e^{-t}} - \frac{1}{t} \right) dt = \log x - \psi(x)$$

and with $x=1$ we have

$$\hat{\gamma}_0(1) = \int_0^{\infty} e^{-t} \left(\frac{1}{1-e^{-t}} - \frac{1}{t} \right) dt = \gamma$$

It is easily seen that

$$(B.24) \quad \sum_{k=0}^n \binom{n}{k} (-1)^k \Gamma^{(n-k)}(1) \hat{\gamma}_k(x) = \int_0^{\infty} e^{-xt} \left(\frac{1}{1-e^{-t}} - \frac{1}{t} \right) \log^n t dt$$

We have the well-known integral representation for the Hurwitz zeta function [26, p.92]

$$\zeta(s, x) = \frac{1}{\Gamma(s)} \int_0^{\infty} \frac{t^{s-1} e^{-(x-1)t}}{e^t - 1} dt = \frac{1}{\Gamma(s)} \int_0^{\infty} \frac{t^{s-1} e^{-xt}}{1 - e^{-t}} dt$$

and from the definition of the gamma function we see that

$$\frac{x^{1-s}}{s-1} = \frac{1}{\Gamma(s)} \int_0^{\infty} t^{s-2} e^{-xt} dt$$

Hence we have for $\text{Re}(s) > 1$

$$\zeta(s, x) - \frac{x^{1-s}}{s-1} = \frac{1}{\Gamma(s)} \int_0^{\infty} e^{-xt} \left(\frac{1}{1-e^{-t}} - \frac{1}{t} \right) t^{s-1} dt$$

which we write as

$$(B.25) \quad \left(\zeta(s, x) - \frac{x^{1-s}}{s-1} \right) \Gamma(s) = \int_0^{\infty} e^{-xt} \left(\frac{1}{1-e^{-t}} - \frac{1}{t} \right) t^{s-1} dt$$

We have

$$\zeta(s, x) - \frac{x^{1-s}}{s-1} = \zeta(s, x) - \frac{1}{s-1} + \frac{1}{s-1} - \frac{x^{1-s}}{s-1}$$

and we denote $f(x)$ as

$$f(x) = \frac{1-x^{1-s}}{s-1}$$

We can represent $f(x)$ by the following integral

$$f(x) = \frac{1-x^{1-s}}{s-1} = \int_1^x t^{-s} dt$$

so that

$$f^{(r)}(x) = (-1)^r \int_1^x t^{-s} \log^r t dt$$

and thus

$$f^{(r)}(1) = (-1)^r \int_1^x \frac{\log^r t}{t} dt = (-1)^r \frac{\log^{r+1} x}{r+1}$$

Differentiating (B.25) n times gives us

$$\frac{d^n}{ds^n} \left(\zeta(s, x) - \frac{x^{1-s}}{s-1} \right) \Gamma(s) = \int_0^{\infty} e^{-xt} \left(\frac{1}{1-e^{-t}} - \frac{1}{t} \right) t^{s-1} \log^n t dt$$

The Stieltjes constants $\gamma_n(u)$ are the coefficients in the Laurent expansion of the Hurwitz zeta function $\zeta(s, u)$ about $s = 1$

$$\zeta(s, x) = \frac{1}{s-1} + \sum_{k=0}^{\infty} \frac{(-1)^k}{k!} \gamma_k(x) (s-1)^k$$

and we therefore have

$$\frac{d^r}{ds^r} \left[\zeta(s, x) - \frac{1}{s-1} \right] = \sum_{k=0}^{\infty} \frac{(-1)^k}{k!} \gamma_k(x) k(k-1) \cdots (k-r+1) (s-1)^{k-r}$$

and thus

$$\lim_{s \rightarrow 1} \frac{d^r}{ds^r} \left[\zeta(s, x) - \frac{1}{s-1} \right] = (-1)^r \gamma_r(x)$$

$$\lim_{s \rightarrow 1} \frac{d^r}{ds^r} \left[\zeta(s, x) - \frac{x^{1-s}}{s-1} \right] = (-1)^r \gamma_r(x) + (-1)^r \frac{\log^{r+1} x}{r+1}$$

Applying the Leibniz rule we have

$$\lim_{s \rightarrow 1} \frac{d^n}{ds^n} \left(\zeta(s, x) - \frac{x^{1-s}}{s-1} \right) \Gamma(s) = \sum_{k=0}^n \binom{n}{k} (-1)^k \Gamma^{(n-k)}(1) \left[\gamma_k(x) + \frac{\log^{k+1} x}{k+1} \right]$$

and hence using (B.24) we see that

$$(B.26) \quad \sum_{k=0}^n \binom{n}{k} (-1)^k \Gamma^{(n-k)}(1) \left[\gamma_k(x) + \frac{\log^{k+1} x}{k+1} \right] = \int_0^{\infty} e^{-xt} \left(\frac{1}{1-e^{-t}} - \frac{1}{t} \right) \log^n t \, dt$$

We then see from (B.24) that (subject to the previous comment regarding the coefficients being linearly independent)

$$\hat{\gamma}_k(x) = \gamma_k(x) + \frac{\log^{k+1} x}{k+1}$$

and accordingly that

$$\hat{\gamma}_k(1) = \gamma_k(1) = \gamma_k$$

With $x = 1$ we have

$$(B.27) \quad \sum_{k=0}^n \binom{n}{k} (-1)^k \Gamma^{(n-k)}(1) \gamma_k = \int_0^\infty e^{-t} \left(\frac{1}{1-e^{-t}} - \frac{1}{t} \right) \log^n t \, dt$$

We also have

$$\zeta(s, x) - \frac{x^{1-s}}{s-1} = \sum_{k=0}^{\infty} \frac{(-1)^k}{k!} \hat{\gamma}_k(x) (s-1)^k$$

With the substitution $t = \log(1/u)$ we obtain

$$(B.28) \quad \sum_{k=0}^n \binom{n}{k} (-1)^k \Gamma^{(n-k)}(1) \hat{\gamma}_k(x) = \int_0^1 u^{x-1} \left(\frac{1}{1-u} + \frac{1}{\log u} \right) \log^n \log \left(\frac{1}{u} \right) du$$

and

$$(B.29) \quad \sum_{k=0}^n \binom{n}{k} (-1)^k \Gamma^{(n-k)}(1) \gamma_k = \int_0^1 \left(\frac{1}{1-u} + \frac{1}{\log u} \right) \log^n \log \left(\frac{1}{u} \right) du$$

and in particular we have

$$A_0(x) = \hat{\gamma}_0(x) = \log x - \psi(x)$$

$$A_1(x) = \Gamma^{(1)}(1) \hat{\gamma}_0(x) - \hat{\gamma}_1(x)$$

This gives us

$$\hat{\gamma}_1(x) = -\gamma [\log x - \psi(x)] - \int_0^1 u^{x-1} \left(\frac{1}{1-u} + \frac{1}{\log u} \right) \log \log \left(\frac{1}{u} \right) du$$

$$\gamma_1 = -\gamma^2 - \int_0^1 \left(\frac{1}{1-u} + \frac{1}{\log u} \right) \log \log \left(\frac{1}{u} \right) du$$

For $u \in [0, 1]$ we have

$$\log \left(\frac{1}{u} \right) = \left| \log \left(\frac{1}{u} \right) \right| = |-\log u| = |\log u|$$

and hence we obtain

$$(B.30) \quad \gamma_1 = -\gamma^2 - \int_0^1 \left(\frac{1}{1-u} + \frac{1}{\log u} \right) \log |\log u| du$$

which unfortunately differs from (2.5).

Appendix C

Derivatives of the gamma function

The gamma function is defined as

$$\Gamma(x) = \int_0^{\infty} y^{x-1} e^{-y} dy$$

and, with the substitution $y = -\log t$, this becomes

$$\Gamma(x) = \int_0^1 (\log(1/t))^{x-1} dt$$

We therefore have

$$\Gamma'(x) = \int_0^1 (\log(1/t))^{x-1} \log \log(1/t) dt$$

and thus

$$(C.1) \quad \Gamma'(1) = \int_0^1 \log \log(1/t) dt = -\gamma$$

More generally we see that

$$\Gamma^{(n)}(x) = \int_0^1 \log^{x-1}(1/t) (\log \log(1/t))^n dt$$

and as we shall see below

$$\Gamma^{(n)}(1) = \int_0^1 (\log \log(1/t))^n dt = (-1)^n c_n$$

where c_n are positive constants.

We have

$$\Gamma^{(n)}(x) = \int_0^{\infty} e^{-t} t^{x-1} \log^n t dt$$

and thus

$$\Gamma^{(n)}(1) = \int_0^{\infty} e^{-t} \log^n t dt$$

We see that

$$\int_0^{\infty} e^{-t} \log^n t dt = \int_0^1 e^{-t} \log^n t dt + \int_1^{\infty} e^{-t} \log^n t dt$$

and with the substitution $t = 1/y$ we have

$$\int_1^{\infty} e^{-t} \log^n t dt = (-1)^n \int_0^1 e^{-1/y} y^{-2} \log^n y dy$$

We therefore obtain

$$\Gamma^{(n)}(1) = \int_0^1 \left[t^2 + (-1)^n e^{t-1/t} \right] e^{-t} t^{-2} \log^n t dt$$

and it is clear that the integrand is positive in the interval $[0,1]$ in the case where n is an even integer. We now consider the case where n is an odd integer and we want to prove that for $t \in [0,1]$

$$t^2 \geq e^{t-1/t}$$

and this is easily seen by considering the logarithmic equivalent

$$2 \log t \geq t - 1/t$$

This inequality is easily proved by considering

$$f(t) = 2t \log t - t^2 + 1$$

where $f(0) = 1$ and $f'(t) = 2[\log t + 1 - t]$ and it is known that $\log t > 1 - t$ for $t > 0$.

Therefore we see that $\Gamma^{(n)}(1)$ has the same sign as $(-1)^n$. This result was reported as an exercise in Apostol's book [4, p.303] and we used it in (3.13) above.

REFERENCES

- [1] V.S.Adamchik, A Class of Logarithmic Integrals. Proceedings of the 1997 International Symposium on Symbolic and Algebraic Computation. ACM, Academic Press, 1-8, 2001.
<http://www-2.cs.cmu.edu/~adamchik/articles/logs.htm>
- [2] O.R. Ainsworth and L.W. Howell, The generalized Euler-Mascheroni constants. NASA Centre for AeroSpace Information (CASI) NASA-TP-2264 ;NAS 1.60.2264, 1984. [View PDF File](#)
- [3] J. Anglesio, A fairly general family of integrals. Amer. Math. Monthly, 104, 665-666, 1997.
- [4] T.M. Apostol, Mathematical Analysis, Second Ed., Addison-Wesley Publishing Company, Menlo Park (California), London and Don Mills (Ontario), 1974.
- [5] E.T. Bell, Exponential polynomials. Ann. of Math., 35 (1934), 258-277.
- [6] G. Boros and V.H. Moll, Irresistible Integrals: Symbolics, Analysis and Experiments in the Evaluation of Integrals. Cambridge University Press, 2004.
- [7] M. Brede, Eine reihenentwicklung der vervollständigten und ergänzten Riemannschen zetafunktion und verwandtes
<http://www.mathematik.uni-kassel.de/~koepf/Diplome/brede.pdf>
- [8] T.J.I'a Bromwich, Introduction to the theory of infinite series.2nd edition Macmillan & Co Ltd, 1965.
- [9] B. Candelpergher, M.A. Coppo and E. Delabaere, La Sommation de Ramanujan. L'Enseignement Mathématique, 43, 93-132, 1997.
<http://math.univ-angers.fr/~delab/Ramanujan.ps>
- [9a] M.W. Coffey, New results concerning power series expansions of the Riemann xi function and the Li/Keiper constants. (preprint) 2007.
- [9b] M.W. Coffey, Relations and positivity results for the derivatives of the Riemann ξ function. J. Comput. Appl. Math., 166, 525-534 (2004)
- [10] C.B. Collins, The role of Bell polynomials in integration. J. Comput. Appl. Math. 131 (2001) 195-211.
- [11] L. Comtet, Advanced Combinatorics, Reidel, Dordrecht, 1974.
- [12] D.F. Connon, Some series and integrals involving the Riemann zeta function, binomial coefficients and the harmonic numbers. Volume II(a), 2007.

[arXiv:0710.4023](#) [[pdf](#)]

- [13] D.F. Connon, Some series and integrals involving the Riemann zeta function, binomial coefficients and the harmonic numbers. Volume II(b), 2007.
[arXiv:0710.4024](#) [[pdf](#)]
- [14] M.A. Coppo, Nouvelles expressions des constantes de Stieltjes. Expositiones Mathematicae 17, No. 4, 349-358 (1999).
- [15] B. Fuglede, A sharpening of Wielandt's characterization of the gamma function. Amer. Math. Monthly, 115, 845-850, 2008.
- [16] I.S. Gradshteyn and I.M. Ryzhik, Tables of Integrals, Series and Products. Sixth Ed., Academic Press, 2000.
Errata for Sixth Edition http://www.mathtable.com/errata/gr6_errata.pdf
- [17] J. Guillera and J. Sondow, Double integrals and infinite products for some classical constants via analytic continuations of Lerch's transcendent. 2005. math.NT/0506319 [[abs](#), [ps](#), [pdf](#), [other](#)]
- [18] H. Hasse, Ein Summierungsverfahren für Die Riemannsche ζ - Reihe. Math.Z.32, 458-464, 1930.
<http://dz-srv1.sub.uni-goettingen.de/sub/digbib/loader?ht=VIEW&did=D23956&p=462>
- [19] A. Ivić, The Riemann Zeta- Function: Theory and Applications. Dover Publications Inc, 2003.
- [20] K.S. Kölbig, The complete Bell polynomials for certain arguments in terms of Stirling numbers of the first kind. J. Comput. Appl. Math. 51 (1994) 113-116.
Also available electronically at:
A relation between the Bell polynomials at certain arguments and a Pochhammer symbol. CERN/Computing and Networks Division, CN/93/2, 1993.
<http://doc.cern.ch/archive/electronic/other/preprints//CM-P/CM-P00065731.pdf>
- [21] N. Nielsen, Theorie des Integrallogarithmus und verwanter transzendenten. 1906.
http://www.math.uni-bielefeld.de/~rehmann/DML/dml_links_author_H.html
- [22] N. Nielsen, Die Gammafunktion. Chelsea Publishing Company, Bronx and New York, 1965.
- [23] S.K. Lakshamana Rao, On the Sequence for Euler's Constant. Amer. Math. Monthly, 63, 572-573, 1956.
- [24] T. Rivoal, Polynômes de type Legendre et approximations de la constant d'Euler. Note (2005). [DVI](#), [PS](#), [PDF](#)

<http://www-fourier.ujf-grenoble.fr/>

- [25] G.K. Srinivasan, The gamma function: An eclectic tour. Amer. Math. Monthly, 114, 297-315, 2007.
- [26] H.M. Srivastava and J. Choi, Series Associated with the Zeta and Related Functions. Kluwer Academic Publishers, Dordrecht, the Netherlands, 2001.
- [27] E.C. Titchmarsh, The Theory of Functions. 2nd Ed., Oxford University Press, 1932.

Donal F. Connon
Elmhurst
Dundle Road
Matfield
Kent TN12 7HD
England

dconnon@btopenworld.com