

LATTICE CONSTRUCTION OF LOGARITHMIC MODULES FOR CERTAIN VERTEX ALGEBRAS

DRAŽEN ADAMOVIĆ AND ANTUN MILAS

ABSTRACT. A general method for constructing logarithmic modules in vertex operator algebra theory is presented. By utilizing this approach, we give explicit vertex operator construction of certain indecomposable and logarithmic modules for the triplet vertex algebra $\mathcal{W}(p)$ and for other subalgebras of lattice vertex algebras and their $N = 1$ super extensions. We analyze in detail indecomposable modules obtained in this way, giving further evidence for the conjectural equivalence between the category of $\mathcal{W}(p)$ -modules and the category of modules for the restricted quantum group $\overline{\mathcal{U}}_q(\mathfrak{sl}_2)$, $q = e^{\pi i/p}$. We also construct logarithmic representations for a certain affine vertex operator algebra at admissible level realized in [4]. In this way we prove the existence of the logarithmic representations predicted in [23]. Our approach enlightens related logarithmic intertwining operators among indecomposable modules, which we also construct in the paper.

CONTENTS

1. Introduction	2
2. Construction of logarithmic modules	3
3. The vertex algebras $\mathcal{W}(p, p')$ and $\mathcal{W}(p)$	5
4. Extended vertex algebras $V_L \oplus V_{L-\alpha/p}$ and $V_L \oplus V_{L+\alpha/p'}$	6
5. Logarithmic and indecomposable modules for the triplet vertex algebra $\mathcal{W}(p)$	8
6. Indecomposable $\mathcal{W}(p)$ -modules and the quantum group $\overline{\mathcal{U}}_q(\mathfrak{sl}_2)$, $q = e^{\pi i/p}$	14
7. Logarithmic modules for $\mathcal{SW}(m)$ and $\mathcal{SW}(p, p')$	17
8. Logarithmic representations of the affine vertex operator algebra $L(-\frac{4}{3}\Lambda_0)$	20
9. Logarithmic intertwining operators	23
References	26

2000 *Mathematics Subject Classification.* Primary 17B69, Secondary 17B67, 17B68, 81R10.

Key words and phrases. vertex algebras, W-algebras, lattice vertex algebras; logarithmic representations; logarithmic intertwining operators; triplet vertex algebras; quantum groups; affine Lie algebras.

1. INTRODUCTION

An important problem in vertex algebra theory is to describe representations of irrational C_2 -cofinite vertex superalgebras, in particular the triplet vertex algebras $\mathcal{W}(p)$, $p \geq 2$. The triplet vertex algebra $\mathcal{W}(p)$ is interesting for several reasons: for instance, its representation theory is expected to be identical to the one of $\overline{\mathcal{U}}_q(\mathfrak{sl}_2)$, where q is root of unity (see [18], [19]). In fact, it was the triplet algebra (and especially early contributions [16] and [25]) that gave motivation for studying general C_2 -cofinite vertex algebras (see [26], [38], etc.).

The present work is a continuation of the series of papers [7]-[10] in which we studied the triplet and super-triplet vertex algebras $\mathcal{W}(p)$ and $S\mathcal{W}(m)$ and their representation theory. By using the theory of Zhu's algebra and modular invariance we proved that these vertex algebras admit logarithmic representations, i.e., the representation on which the Virasoro generator $L(0)$ does not act semisimply. On the other hand, irreducible and ordinary modules for $\mathcal{W}(p)$ and $S\mathcal{W}(m)$ can be constructed explicitly by using lattice vertex algebras and their modules. This raises a question whether logarithmic modules can be also constructed via lattice vertex algebras, even though lattice vertex algebras do not admit logarithmic modules. In contrast, for the Heisenberg and singlet vertex algebras (see [6] and [37]) logarithmic modules are easily constructed by deforming the action of the zero modes of the Heisenberg algebra. Thus, we believe it is important to present explicit construction of logarithmic representations for the triplet.

In this paper we present a general construction of logarithmic modules for vertex (super)algebras obtained as (intersections of) kernels of "screening" operators (subject to some additional properties). Vertex algebras obtained in this fashion are also known as \mathcal{W} -algebras, but we stress that \mathcal{W} -algebras considered in this paper do not arise from the usual quantum reduction as in [12], [14], [30], etc.

Although the idea of deforming CFTs with the use of screening operators has been considered in the literature on LCFT (especially [17]), our construction is new and covers models not previously considered. For instance, we are able to explicitly construct certain logarithmic representations for $\mathcal{W}(p)$ for every p , while in the literature the only explicit construction we are aware of is for $p = 2$ (cf. [1],[25]). Moreover, methods used to construct logarithmic modules are very useful to analyze general indecomposable $\mathcal{W}(p)$ -modules (see Section 5 and 6).

We also apply our method to other vertex superalgebras of interest, such as the supertriplet $S\mathcal{W}(m)$ introduced in [8]. Moreover, we construct certain logarithmic representations (cf. Theorem 4.1) of the vertex algebras $\mathcal{W}(p, p')$, which are extensions of the (p, p') -minimal models

for the Virasoro algebra introduced in [20]. We also present a construction of a new family of $N = 1$ vertex operator superalgebras, called $\mathcal{SW}(p, p')$, which are extensions of the $N = 1$ super minimal (p, p') -models, with $(p, \frac{p-p'}{2}) = 1$. This extension combines into the following diagram

$$V_{1/2}(c_{p,p'}, 0) \subset \mathcal{SW}(p, p') \subset V_L \otimes F,$$

where $V_{1/2}(c_{p,p'}, 0)$ is the (universal) Neveu-Schwarz vertex superalgebra of central charge $c_{p,p'}$, F is the free fermion vertex operator and V_L is the lattice vertex superalgebra based on the integral lattice $L = \mathbb{Z}\alpha$, $\langle \alpha, \alpha \rangle = pp'$.

The present work also contributes to better understanding of logarithmic representations of a certain affine vertex algebra on an admissible level. Previously, the first author [4] had obtain an explicit realization of the affine vertex operator algebra $L(-\frac{4}{3}\Lambda_0)$ (admissible module for $A_1^{(1)}$ [29], [30]) and related modules. In Section 8 we shall construct certain logarithmic $L(-\frac{4}{3}\Lambda_0)$ -representations. These logarithmic representations are important for description of extensions of certain weak modules and analysis of fusion product of admissible representations. In particular, we present the free field realization of the logarithmic module $\mathcal{R}_{-1/3}$ which appeared in [23] and [24]. It is also interesting that logarithmic representations for affine vertex operator algebras which we have constructed are not $\mathbb{Z}_{\geq 0}$ -graded and therefore they cannot be detected by using theory of Zhu's algebras.

It is well known now that logarithmic modules in vertex algebra theory are closely related to logarithmic intertwining operators [26], [27], [35]. As in the ordinary case, logarithmic intertwining operators are hard to construct explicitly. In the last part of the paper we show that already the module map for logarithmic modules considered earlier give rise to logarithmic intertwining operators. This is obtained by combining the theory of simple currents [34] with our method (cf. Theorems 9.1 and 9.2).

2. CONSTRUCTION OF LOGARITHMIC MODULES

In this part we assume some familiarity with vertex (super)algebras as in [28] and [32].

Assume that $(V, Y, \mathbf{1}, \omega)$ is a vertex operator superalgebra, with the parity decomposition $V = V_0 \oplus V_1$, together with the vertex operator map $Y(\cdot, x)$, $Y(u, x) = \sum_{m \in \mathbb{Z}} u_m x^{-m-1}$, such that

$$Y(\omega, x) = \sum_{m \in \mathbb{Z}} L(m) x^{-m-2},$$

where the operators $L(m)$, $m \in \mathbb{Z}$ close the Virasoro algebra.

Let $v \in V$ be an even vector such that:

$$(2.1) \quad [v_n, v_m] = 0 \quad \forall n, m \in \mathbb{Z},$$

$$(2.2) \quad L(n)v = \delta_{n,0}v \quad \forall n \in \mathbb{Z}_{\geq 0},$$

so that v is of conformal weight one.

Define

$$(2.3) \quad \Delta(v, x) = x^{v_0} \exp \left(\sum_{n=1}^{\infty} \frac{v_n}{-n} (-x)^{-n} \right).$$

If v_0 acts semisimply on V then the expression $x^{v_0}w$, where w is an eigenvector for v_0 corresponding to the eigenvalue λ , is defined as $x^\lambda w$. But (2.3) is ambiguous if v_0 does not act semisimply on V (however, see also Chapter 9).

The following easy but fundamental observation will be of great use in the paper.

Theorem 2.1. *Assume that V is a vertex operator superalgebra and that $v \in V$ is an even vector which satisfies conditions (2.1) and (2.2). Let \bar{V} be the vertex subalgebra of V such that $\bar{V} \subseteq \text{Ker}_V v_0$.*

Assume that (M, Y_M) is a weak V -module (resp. σ -twisted weak V -module). Define the pair $(\widetilde{M}, \widetilde{Y}_{\widetilde{M}})$ such that

$$\widetilde{M} = M \quad \text{as a vector space,}$$

$$\widetilde{Y}_{\widetilde{M}}(a, x) = Y_M(\Delta(v, x)a, x) \quad \text{for } a \in \bar{V}.$$

Then $(\widetilde{M}, \widetilde{Y}_{\widetilde{M}})$ is a weak \bar{V} -module (resp. σ -twisted weak \bar{V} -module).

Proof. The proof is essentially in [34] and [15]. There the same statement was proven in the case when v_0 is semisimple and without any restriction. But the same conclusion obviously holds for \bar{V} , because v_0 acts semisimply on \bar{V} . \square

Remark 2.1. *In the case when $v_n, n \in \mathbb{Z}$, satisfy the commutation relations for the Heisenberg algebra and v_0 acts semisimply on V with integral weights, $\Delta(v, x)$ can be used for construction of simple current modules (for details see [34], [15]).*

Among all (weak) V -modules, logarithmic modules are of great interest in this work. The next result gives a general method for their construction.

Theorem 2.2. *Assume that (M, Y_M) is a V -module such that $L(0)$ acts semisimply on M . Then $(\widetilde{M}, \widetilde{Y}_{\widetilde{M}})$ is logarithmic \bar{V} -module if and only if v_0 does not act semisimply on M .*

Proof. Let

$$\widetilde{L}(x) = \sum_{n \in \mathbb{Z}} \widetilde{L}(n) x^{-n-2} = \widetilde{Y}_{\widetilde{M}}(\omega, x).$$

Then

$$\widetilde{L}(x) = Y_M(\omega + x^{-1}v, x) = L(x) + x^{-1}Y(v, x),$$

which implies $\widetilde{L}(n) = L(n) + v_n$, and in particular

$$\widetilde{L}(0) = L(0) + v_0.$$

Since $L(0)$ and v_0 both preserve graded components and the operator $L(0)$ acts as a scalar on each graded component, it is clear that $\widetilde{L}(0)$ acts nonsemisimply if and only if v_0 acts nonsemisimply. The proof follows. \square

Remark 2.2. We should mention that in [17], the authors presented a related construction of logarithmic modules by using logarithmic deformation and the extension of the chiral algebra. We think that it will be interesting to relate these two constructions of logarithmic representations; we also refer the reader to Section 9, which is in the spirit of [17].

3. THE VERTEX ALGEBRAS $\mathcal{W}(p, p')$ AND $\mathcal{W}(p)$

Define the lattice

$$L = \mathbb{Z}\alpha, \quad \langle \alpha, \alpha \rangle = 2pp',$$

where $p, p' \in \mathbb{Z}_{>0}$, $p \neq p'$, $p \geq 2$. Let V_L be the associated (rank one) lattice vertex algebra with the vertex operator map Y (cf. [32]).

The vertex algebra V_L is a subalgebra of the generalized vertex algebra $V_{\widetilde{L}}$, where

$$\widetilde{L} = \mathbb{Z} \frac{\alpha}{2pp'}$$

is the dual lattice. Let us denote the (generalized) vertex operator map in $V_{\widetilde{L}}$ again by Y .

Define

$$\omega = \frac{1}{4pp'}\alpha(-1)^2 + \frac{p-p'}{2pp'}\alpha(-2).$$

Then ω is a conformal vector with central charge $c_{p,p'} = 1 - \frac{6(p-p')^2}{pp'}$. The associated screening operators are

$$Q = e_0^{\alpha/p'} = \text{Res}_x Y(e^{\alpha/p'}, x), \quad \widetilde{Q} = e_0^{-\alpha/p} = \text{Res}_x Y(e^{-\alpha/p}, x).$$

It is known (cf. also [3]) that we have

$$[Q, \widetilde{Q}] = 0.$$

The screening operators Q and \tilde{Q} enable us to define certain vertex subalgebras of V_L . Define

$$\overline{V}_L = \text{Ker}_{V_L} \tilde{Q}, \quad \overline{\overline{V}}_L = \text{Ker}_{V_L} Q, \quad \mathcal{W}(p, p') = \overline{V}_L \cap \overline{\overline{V}}_L.$$

Then \overline{V}_L , $\overline{\overline{V}}_L$ and $\mathcal{W}(p, p')$ are vertex operator algebras with central charge $c_{p, p'}$. The vertex algebra $\mathcal{W}(p, p')$ was introduced by Feigin et al. in [20].

It can be shown (by using Felder's complex or rationality of $L(c_{p, p'}, 0)$) that

$$V(c_{p, p'}, 0) \subset \mathcal{W}(p, p'),$$

where $V(c_{p, p'}, 0)$ is the universal Virasoro vertex algebra of central charge $c_{p, p'}$ (generalized Verma module) and $L(c_{p, p'}, 0)$ is its simple quotient.

It is not clear to us how to find a strongly generated set for $\mathcal{W}(p, p')$ in general, but for $p' = 2$ we recently made some progress [11]. Then, motivated mainly by [20] and [11], we do expect the following conjecture to be true (cf. also [11]).

Conjecture 3.1. *The vertex algebra $\mathcal{W}(p, p')$ is C_2 -cofinite with $2pp' + \frac{(p-1)(p'-1)}{2}$ inequivalent irreducible modules.*

Remark 3.1. *In the case $p' = 1$, $\overline{V}_L = \mathcal{W}(p)$ is the triplet vertex algebra with central charge $c_{p, 1}$. In this case*

$$\overline{\overline{V}}_L = \mathcal{W}(p, 1) = L^{Vir}(c_{p, 1}, 0).$$

4. EXTENDED VERTEX ALGEBRAS $V_L \oplus V_{L-\alpha/p}$ AND $V_L \oplus V_{L+\alpha/p'}$

Observe first that if V_L is of central charge $c_{p, p'}$, both $V_{L-\alpha/p}$ and $V_{L+\alpha/p'}$ are simple V_L -modules with integral weights. Then on $V_L \oplus V_{L-\alpha/p}$ (resp. $V_L \oplus V_{L+\alpha/p'}$) we have the structure of a vertex operator algebra (see [33]). In this section we shall investigate these extended vertex operator algebras and its modules. The associated vertex operators can be reconstructed from the generalized vertex algebra $(V_{\tilde{L}}, Y)$.

Define

$$\mathcal{V}_i = V_{L+\frac{i}{2p'}\alpha} \bigoplus V_{L+\frac{i}{2p'}\alpha-\frac{1}{p}\alpha}, \quad i = 0, \dots, 2p' - 1,$$

and

$$\mathcal{V}_j^o = V_{L+\frac{j}{2p}\alpha} \bigoplus V_{L+\frac{j}{2p}\alpha+\frac{1}{p'}\alpha}, \quad j = 0, \dots, 2p - 1.$$

Then $\mathcal{V} = \mathcal{V}_0$ (resp. $\mathcal{V}^o = \mathcal{V}_0^o$) is a vertex operator algebra and each \mathcal{V}_i (resp. \mathcal{V}_j^o) is an \mathcal{V} -module (resp. \mathcal{V}^o -module). The associated vertex operators $Y_{\mathcal{V}_i}$ are defined as follows:

$$Y_{\mathcal{V}_i}(u + v, x)(u_i + v_i) = Y(u + v, x)u_i + Y(u, x)v_i,$$

where $u \in V_L$, $v \in V_{L-\frac{1}{p}\alpha}$, $u_i \in V_{L+\frac{i}{2p'}\alpha}$, $v_i \in V_{L+\frac{i}{2p'}\alpha-\frac{1}{p}\alpha}$. (Note that we require $Y_{\mathcal{V}_i}(v, x)v_i = 0$.) Similarly we define the vertex operator $Y_{\mathcal{V}_j^o}$.

Remark 4.1. Assume that V is a vertex operator algebra and M is a V -module with integral weights. Then $V \oplus M$ carries the vertex operator algebra structure (cf. [33]). This structure in the case $V = V_L$ and $M = V_{L-\alpha/p}$ coincides with the vertex operator algebra structure on \mathcal{V} described above.

We fix $V = \mathcal{V}_0$. In this section we shall consider the case $v = e^{-\alpha/p}$. Then

$$[v_n, v_m] = 0 \quad \forall m, n \in \mathbb{Z}.$$

But $v_0 = \tilde{Q}$ is a screening operator and it does not acts semisimply on \mathcal{V}_0 . On the other hand \tilde{Q} acts trivially on \overline{V}_L and we can use $\Delta(v, x)$ to construct weak \overline{V}_L -modules. Similarly we can use $\Delta(e^{\alpha/p'}, x)$ to construct weak $\overline{\overline{V}_L}$ -modules.

Theorem 4.1.

(i) Assume that (M, Y_M) is a weak \mathcal{V} -module. Then the structure

$$(\tilde{M}, \tilde{Y}_{\tilde{M}}(\cdot, x)) := (M, Y_M(\Delta(e^{-\alpha/p}, x)\cdot, x))$$

is a weak \overline{V}_L -module. In particular,

$$(\tilde{\mathcal{V}}_i, \tilde{Y}_{\tilde{\mathcal{V}}_i}), \quad i = 0, \dots, 2p' - 1,$$

are weak \overline{V}_L -modules.

(ii) Assume that (M, Y_M) is a weak \mathcal{V}^o -module. Then the structure

$$(\tilde{M}, \tilde{Y}_{\tilde{M}}(\cdot, x)) := (M, Y_M(\Delta(e^{\alpha/p'}, x)\cdot, x))$$

is a weak $\overline{\overline{V}_L}$ -module. In particular,

$$(\tilde{\mathcal{V}}_j^o, \tilde{Y}_{\tilde{\mathcal{V}}_j^o}), \quad j = 0, \dots, 2p - 1,$$

are weak $\overline{\overline{V}_L}$ -modules.

We shall now investigate these \overline{V}_L -modules. For simplicity we set $\tilde{Y} = \tilde{Y}_{\tilde{V}_i}$. Let

$$L(x) = Y(\omega, x) = \sum_{n \in \mathbb{Z}} L(n)x^{-n-2}, \quad \widetilde{L}(x) = \tilde{Y}(\omega, x) = \sum_{n \in \mathbb{Z}} \widetilde{L}(n)x^{-n-2}.$$

Then

$$\begin{aligned} \widetilde{L}(x) &= L(x) + x^{-1}Y(e^{-\alpha/p}, x), \\ \widetilde{L}(n) &= L(n) + e_n^{-\alpha/p}. \end{aligned}$$

In particular,

$$\widetilde{L}(0) = L(0) + \tilde{Q}.$$

Similarly we see that on $\widetilde{\mathcal{V}}_i^o$ we have

$$\widetilde{L}(0) = L(0) + Q.$$

Now it is easy to see that modules $\tilde{\mathcal{V}}_i$ and $\widetilde{\mathcal{V}}_i^o$ are logarithmic.

Corollary 4.1.

- (i) For every $i = 0, \dots, 2p' - 1$, $\tilde{\mathcal{V}}_i$ are logarithmic \overline{V}_L -modules.
- (ii) For every $i = 0, \dots, 2p' - 1$, $\tilde{\mathcal{V}}_i$ are logarithmic $\mathcal{W}(p, p')$ -modules.
- (iii) For every $i = 0, \dots, 2p - 1$, $\widetilde{\mathcal{V}}_i^o$ are logarithmic $\mathcal{W}(p, p')$ -modules.

Remark 4.2. Observe that nothing had prevented us to work with more general operators $\Delta(\lambda e^{-\alpha/p}, x)$, where $\lambda \neq 0$. But this modification does not lead to new representations; it merely contributes to the endomorphism algebra of a (reducible) logarithmic module in question.

5. LOGARITHMIC AND INDECOMPOSABLE MODULES FOR THE TRIPLET VERTEX ALGEBRA

$$\mathcal{W}(p)$$

Now we shall consider the case $p' = 1$ and $p \geq 2$. Then \overline{V}_L is the triplet vertex algebra $\mathcal{W}(p)$ investigated in [7], [22], [16], [18], [19], etc. For more about the triplet algebra see [31], [21], [13], etc.

We know that $\mathcal{W}(p)$ is a vertex subalgebra of V_L , strongly generated by

$$\omega, \quad F = e^{-\alpha}, \quad H = QF, \quad E = Q^2F.$$

Recall also that $\mathcal{W}(p)$ has exactly $2p$ irreducible modules:

$$\Lambda(1), \dots, \Lambda(p), \Pi(1), \dots, \Pi(p).$$

The construction in the previous section provides two logarithmic $\mathcal{W}(p)$ -modules: $(\widetilde{\mathcal{V}}, \widetilde{Y})$ and $(\widetilde{\mathcal{V}}_1, \widetilde{Y})$.

For $a \in \mathcal{W}(p)$, let

$$\widetilde{Y}(a, x) = \sum_{n \in \mathbb{Z}} \widetilde{a}_n x^{-n-1}.$$

Define the following operator acting on $\widetilde{\mathcal{V}}$ and $\widetilde{\mathcal{V}}_1$:

$$(5.1) \quad G = e_0^\alpha + \nu_p e_{-1}^{\alpha-\alpha/p}, \quad \nu_p = \frac{p}{p-1}.$$

Then the definition of $\widetilde{Y}(a, x)$ implies that

$$(5.2) \quad \widetilde{a}_n = a_n + \sum_{m=1}^{\infty} \frac{(-1)^m}{-m} (e_m^{-\alpha/p} a)_{n-m}.$$

Lemma 5.1. *On $\mathcal{W}(p)$ -modules $\widetilde{\mathcal{V}}$ and $\widetilde{\mathcal{V}}_1$ we have*

- (i) $[G, \widetilde{a}_n] = \widetilde{(Qa)_n}$, for every $n \in \mathbb{Z}$ and $a \in \mathcal{W}(p)$.
- (ii) $[G, \widetilde{L}(n)] = 0$, i.e., G is a screening operator which commutes with the action of the Virasoro algebra on $\widetilde{\mathcal{V}}$ and $\widetilde{\mathcal{V}}_1$.

Proof. It is clear that (i) implies (ii). Let us prove assertion (i). By using formula (5.2) we get:

$$\begin{aligned} [G, \widetilde{a}_n] &= [Q + \nu_p e_{-1}^{\alpha-\alpha/p}, \widetilde{a}_n] \\ &= \widetilde{(Qa)_n} + \nu_p [e_{-1}^{\alpha-\alpha/p}, a_n] + \sum_{m=1}^{\infty} \frac{(-1)^m}{-m} ([Q, e_m^{-\alpha/p}] a)_{n-m} \\ &= \widetilde{(Qa)_n} + \nu_p \sum_{i=0}^{\infty} \binom{-1}{i} (e_i^{\alpha-\alpha/p} a)_{n-1-i} + \nu_p \sum_{m=1}^{\infty} (-1)^m (e_{m-1}^{\alpha-\alpha/p} a)_{n-m} \\ &= \widetilde{(Qa)_n}. \end{aligned}$$

The proof follows. □

For $X \in \{E, F, H\}$, we define

$$\widetilde{X}(n) := \widetilde{X}_{n+2p-2}.$$

We shall consider the following $\mathcal{W}(p)$ -module:

$$\mathcal{P}_{p-1}^+ = \mathcal{W}(p) \cdot e^{\frac{\alpha}{2}} \subseteq \widetilde{\mathcal{V}}_1.$$

Theorem 5.1.

(i) The module \mathcal{P}_{p-1}^+ is a $\mathbb{Z}_{\geq 0}$ -graded logarithmic $\mathcal{W}(p)$ -module

$$\mathcal{P}_{p-1}^+ = \bigoplus_{n \in \mathbb{Z}_{\geq 0}} \mathcal{P}_{p-1}^+(n),$$

with the top component $\mathcal{P}_{p-1}^+(0)$ of dimension two, spanned by $\text{span}_{\mathbb{C}}\{e^{\frac{\alpha}{2}}, e^{\frac{\alpha}{2} - \frac{\alpha}{p}}\}$. In this basis $\widetilde{L}(0)$ acts on $\mathcal{P}_{p-1}^+(0)$ as

$$\begin{bmatrix} \frac{2-p}{4} & 1 \\ 0 & \frac{2-p}{4} \end{bmatrix}.$$

(ii) The $\mathcal{W}(p)$ -module \mathcal{P}_{p-1}^+ is invariant under the action of G , and we have

$$\mathcal{P}_{p-1}^+ = \widetilde{\mathcal{V}}_1.$$

Proof. By using direct calculation, one can see that

$$\widetilde{X}(n)v = L(\widetilde{n+1})v = 0, \quad \text{for } n \in \mathbb{Z}_{\geq 0}$$

for $v \in \{e^{\frac{\alpha}{2}}, e^{\frac{\alpha}{2} - \frac{\alpha}{p}}\}$. This implies that \mathcal{P}_{p-1}^+ is a $\mathbb{Z}_{\geq 0}$ -graded module for $\mathcal{W}(p)$. Now the proof follows from:

$$\begin{aligned} \widetilde{L}(0)e^{\frac{\alpha}{2}} &= L(0)e^{\frac{\alpha}{2}} + \widetilde{Q}e^{\frac{\alpha}{2}} = \frac{2-p}{4}e^{\frac{\alpha}{2}} + e^{\frac{\alpha}{2} - \frac{\alpha}{p}}, \\ \widetilde{L}(0)e^{\frac{\alpha}{2} - \frac{\alpha}{p}} &= L(0)e^{\frac{\alpha}{2} - \frac{\alpha}{p}} = \frac{2-p}{4}e^{\frac{\alpha}{2} - \frac{\alpha}{p}}. \end{aligned}$$

Let us prove assertion (ii). First we recall the usual (undeformed) action of $\mathcal{W}(p)$ on $V_{L+\alpha/2}$. By using results from [7], one can easily see that under this action $V_{L+\alpha/2}$ is a cyclic module generated by $e^{\alpha/2}$. Moreover $V_{L+\alpha/2-\alpha/p}$ is a cyclic module generated by $e^{3/2\alpha-\alpha/p}$. On the other hand $\mathcal{W}(p) \cdot e^{\alpha/2-\alpha/p} = \Lambda(p-1) \subset \mathcal{P}_{p-1}^+$. Thus, it is sufficient to show $e^{3/2\alpha-\alpha/p} \in \mathcal{P}_{p-1}^+$. Our Lemma 5.1 directly implies that \mathcal{P}_{p-1}^+ is G -invariant.

Since $e^{-\alpha/2} \in \mathcal{P}_{p-1}^+$, we conclude that

$$G^2 e^{-\alpha/2} \in \mathcal{P}_{p-1}^+.$$

By direct calculation we have:

$$G^2 e^{-\alpha/2} = 2\nu_p Q e_{-1}^{\alpha-\alpha/p} e^{-\alpha/2} = C e^{3/2\alpha-\alpha/p}$$

for certain non-zero constant C . The constant C can be easily computed from the relation

$$Q e_{-1}^{\alpha-\alpha/p} e^{-\alpha/2} = \text{Res}_{x_1} \text{Res}_{x_2} x_2^{-1} Y(e^\alpha, x_1) Y(e^{\alpha-\alpha/p}, x_2) e^{-\alpha/2}$$

and the explicit formulas $Y(e^\alpha, x)$ and $Y(e^{\alpha-\alpha/p}, x)$. The proof follows. \square

Define the following submodules of \mathcal{P}_{p-1}^+ :

$$\begin{aligned}\mathcal{M}_1 &= \mathcal{W}(p).e^{-\alpha/2}, \\ \mathcal{N}_1 &= \mathcal{W}(p).e^{-\alpha/2} + \mathcal{W}(p).e^{3\alpha/2-\alpha/p} = \mathcal{M}_1 + V_{L+\alpha/2-\alpha/p}, \\ \mathcal{N}_2 &= \mathcal{W}(p).e^{\alpha/2-\alpha/p} \cong \Lambda(p-1).\end{aligned}$$

Lemma 5.2. *We have*

(i) *The $\mathcal{W}(p)$ -module $\mathcal{M}_1 = \mathcal{W}(p) \cdot e^{-\alpha/2}$ defines a non-split extension*

$$0 \rightarrow \Lambda(p-1) \rightarrow \mathcal{M}_1 \rightarrow \Pi(1) \rightarrow 0,$$

of modules $\Pi(1)$ and $\Lambda(p-1)$.

(ii) *The modules \mathcal{M}_1 and $V_{L+\alpha/2-\alpha/p}$ are non-isomorphic.*

Proof. We clearly have $\mathcal{N}_2 \subset V_{L+\alpha/2-\alpha/p}$. From Lemma 5.1, we compute

$$(5.3) \quad \widetilde{E(p-1)}e^{-\alpha/2} = (\widetilde{Q^2e^{-\alpha}})_{3p-3}e^{-\alpha/2} = (ad G)^2(e_{3p-3}^{-\alpha}) \cdot e^{-\alpha/2}.$$

But

$$e_{3p-3}^{-\alpha}e^{-\alpha/2} = e_{3p-3}^{-\alpha}Ge^{-\alpha/2} = 0,$$

and thus

$$\widetilde{E(p-1)}e^{-\alpha/2} = \widetilde{F(p-1)}G^2e^{-\alpha/2} = Ce^{\alpha/2-\alpha/p},$$

for some $C \neq 0$. This implies that $\mathcal{N}_2 \subset \mathcal{M}_1$.

To prove the assertion it is sufficient to show that:

$$(5.4) \quad V_{L+\alpha/2-\alpha/p} \cap \mathcal{M}_1 = \mathcal{N}_2.$$

By using Lemma 5.1 again we prove:

$$\widetilde{E(0)}e^{-\alpha/2} = -2 \binom{3p-2}{2p-1} Qe^{-\alpha/2} + w$$

where

$$w = -2\nu_p \binom{3p-2}{2p-1} e_{-1}^{\alpha-\alpha/p} e^{-\alpha/2} + \widetilde{F(0)}G^2e^{-\alpha/2}.$$

Since $Qw = 0$ we conclude that w is not a subsingular vector in $V_{L+\alpha/2-\alpha/p}$ which gives that $w \in \mathcal{N}_2$ (here we use the structure of $V_{L+\alpha/2-\alpha/p}$ as a module for the Virasoro algebra from [7]).

We also have:

$$\widetilde{H(0)}e^{-\alpha/2} = -\binom{3p-2}{2p-1} e^{-\alpha/2}, \quad \widetilde{H(0)}Qe^{-\alpha/2} = \binom{3p-2}{2p-1} Qe^{-\alpha/2} - \frac{1}{2}w.$$

Therefore the top component of the quotient module $\mathcal{M}_1/\mathcal{N}_2$ is of dimension two. Assume that $V_{L+\alpha/2-\alpha/p} \cap \mathcal{M}_1 \not\cong \mathcal{N}_2$. Then clearly

$$V_{L+\alpha/2-\alpha/p} \cap \mathcal{M}_1 = V_{L+\alpha/2-\alpha/p}$$

which implies that the top component of $\mathcal{M}_1/\mathcal{N}_2$ is 4-dimensional. A contradiction. Therefore (5.4) holds. This implies that the quotient module $\mathcal{M}_1/\mathcal{N}_2$ is isomorphic to $\Pi(1)$.

For (ii) we only have to show that $V_{L+\alpha/2-\alpha/p} \not\cong \mathcal{M}_1$. This is done by contradiction. Suppose there is an isomorphism from \mathcal{M}_1 to $V_{L+\alpha/2-\alpha/p}$. Since $e^{-\alpha/2}$ is a highest weight vector for the Virasoro algebra this vector has to be mapped to another highest weight vector in $V_{L+\alpha/2-\alpha/p}$ of the same conformal weight. Such a vector is unique up to a scalar and proportional to $e^{3\alpha/2-\alpha/p}$. But then $\tilde{F}_{3p-3}e^{3\alpha/2-\alpha/p} = e^{\alpha/2-\alpha/p}$, while $\tilde{F}_{3p-3}e^{-\alpha/2} = 0$, a contradiction. \square

The module \mathcal{M}_1 in the previous lemma was also predicted in [22] (see also [18]).

Proposition 5.1. *The socle part of the $\mathcal{W}(p)$ -module \mathcal{P}_{p-1}^+ is isomorphic to $\Lambda(p-1)$. The module \mathcal{P}_{p-1}^+ has semisimple length three with the following semisimple filtration:*

$$(5.5) \quad \mathcal{P}_{p-1}^+ = \mathcal{N}_0 \supset \mathcal{N}_1 \supset \mathcal{N}_2 \supset \mathcal{N}_3 = 0,$$

such that

$$\mathcal{N}_0/\mathcal{N}_1 \cong \Lambda(p-1), \quad \mathcal{N}_1/\mathcal{N}_2 \cong \Pi(1) \oplus \Pi(1), \quad \mathcal{N}_2/\mathcal{N}_3 \cong \Lambda(p-1).$$

Proof. Let W be a non-zero submodule of \mathcal{P}_{p-1}^+ . By using action of $\mathcal{W}(p)$ on vectors from W , one sees that

$$W \cap \mathcal{P}_{p-1}^+(0) \neq \{0\}.$$

Then action of $\widetilde{L}(0)$ on the top level $\mathcal{P}_{p-1}^+(0)$ implies that $e^{\alpha/2-\alpha/p} \in W$. In this way we have proved that W contains \mathcal{N}_2 . Since W is an arbitrary submodule, we conclude that the socle part of \mathcal{P}_{p-1}^+ is $\mathcal{N}_2 \cong \Lambda(p-1)$.

Since $\mathcal{P}_{p-1}^+/\mathcal{N}_2$ is not semisimple $\mathcal{W}(p)$ -module, there are no semisimple filtration of length two.

In order to prove that (5.5) is a semisimple filtration of length three, it suffices to show that $\mathcal{N}_1/\mathcal{N}_2$ is semisimple. First we notice that $V_{L+\alpha/2-\alpha/p}/\mathcal{N}_2 \cong \Pi(1)$ (cf. [7]) and $\mathcal{M}_1/\mathcal{N}_2 \cong \Pi(1)$ (cf. Lemma 5.2) which easily implies that

$$(V_{L+\alpha/2-\alpha/p}/\mathcal{N}_2) \cap \mathcal{M}_1/\mathcal{N}_2 = \{0\}.$$

In this way we have obtained that

$$\mathcal{N}_2/\mathcal{N}_1 = (\mathcal{M}_1 + V_{L+\alpha/2-\alpha/p})/\mathcal{N}_2 \cong \Pi(1) \oplus \Pi(1).$$

So filtration (5.5) is a semisimple filtration of minimal length, and therefore \mathcal{P}_{p-1}^+ has semisimple length three. \square

In the proof of Proposition 5.1 we have shown that every non-zero submodule of \mathcal{P}_{p-1}^+ intersects top component non-trivially. Thus, \mathcal{P}_{p-1}^+ is precisely the $\mathcal{W}(p)$ -module induced from the two-dimensional module $\mathcal{P}_{p-1}^+(0)$ for Zhu's algebra $A(\mathcal{W}(p))$.

Corollary 5.1. *Assume that M is a $\mathbb{Z}_{\geq 0}$ -graded $\mathcal{W}(p)$ -module such that:*

- (i) *The top component $M(0)$ is isomorphic to $\mathcal{P}_{p-1}^+(0)$ as a module for the Zhu's algebra $A(\mathcal{W}(p))$.*
- (ii) *If W is a non-zero submodule of M , then $W \cap M(0) \neq \{0\}$.*

Then we have $W \cong \mathcal{P}_{p-1}^+$. In particular, the module \mathcal{P}_{p-1}^+ is self-dual.

Proof. The first assertion follows from the theory of Zhu's algebras and the above discussion. Since the contragredient module for \mathcal{P}_{p-1}^+ also has the top component isomorphic to $\mathcal{P}_{p-1}^+(0)$, we easily conclude that \mathcal{P}_{p-1}^+ is a self-dual $\mathcal{W}(p)$ -module. \square

The other logarithmic $\mathcal{W}(p)$ -module $\widetilde{\mathcal{V}}$, which will be denoted by \mathcal{P}_{p-1}^- , has a slightly different structure as illustrated in our next result (the proof is analogous to the one in the previous proposition).

Proposition 5.2. *We have:*

- (i) *The module \mathcal{P}_{p-1}^- is of semisimple length three and its socle part is $\Pi(p-1)$. Moreover, \mathcal{P}_{p-1}^- admits the semisimple filtration*

$$(5.6) \quad \mathcal{P}_{p-1}^- = \mathcal{N}_0^- \supset \mathcal{N}_1^- \supset \mathcal{N}_2^- \supset \mathcal{N}_3^- = 0,$$

such that

$$\mathcal{N}_0^-/\mathcal{N}_1^- \cong \Pi(p-1), \quad \mathcal{N}_1^-/\mathcal{N}_2^- \cong \Lambda(1) \oplus \Lambda(1), \quad \mathcal{N}_2^-/\mathcal{N}_3^- \cong \Pi(p-1).$$

- (ii) *The first nontrivial $\widetilde{L}(0)$ -Jordan block of \mathcal{P}_{p-1}^- opens at the degree one (i.e., there exists $v \in \mathcal{P}_{p-1}^-(1)$ such that $\widetilde{L}(0)v = v + w$, where $w \neq 0$ and $\widetilde{L}(0)w = w$).*
- (iii) *If we let $\widetilde{Y}'(\cdot, x) = \widetilde{Y}(\Delta(\frac{\alpha(0)}{2}, x)\cdot, x)$, then $(\widetilde{\mathcal{V}}, \widetilde{Y}'(\cdot, x))$ is isomorphic to $\widetilde{\mathcal{V}}_1$.*

Remark 5.1. *Existence of certain logarithmic $\mathcal{W}(p)$ -modules was proven in [7] by using modular invariance and the theory of Zhu's algebras. In Theorem 5.1 we have presented an explicit realization of a logarithmic module from Proposition 6.6 of [7].*

Remark 5.2. *In [18],[19] (see also [22]) it was conjectured that $\mathcal{W}(p)$ admits $2(p-1)$ indecomposable (nonisomorphic) logarithmic modules. Although our method gives 2 logarithmic $\mathcal{W}(p)$ -modules for fixed p , there is a strong evidence that remaining logarithmic modules can be also constructed by using our method. The main difficulty here is to find a correct replacement for the Q operator in an appropriate extension of V_L . This is work in progress.*

6. INDECOMPOSABLE $\mathcal{W}(p)$ -MODULES AND THE QUANTUM GROUP $\overline{\mathcal{U}}_q(sl_2)$, $q = e^{\pi i/p}$

In this section discuss conjectural equivalence between the category of finitely generated $\mathcal{W}(p)$ -modules and the category of finite dimensional modules for the restricted quantum group $\overline{\mathcal{U}}_q(sl_2)$, where q is a primitive $2p$ -th root of unity (cf. [18], [19]).

We also discuss some additional indecomposable $\mathcal{W}(p)$ -modules.

In [7], we decomposed the category of ordinary $\mathcal{W}(p)$ -modules into $p+1$ blocks. Two blocks contain a single irreducible module, and remaining $p-1$ blocks contain two irreducible modules each. In fact, similar decomposition persists at the level of weak $\mathcal{W}(p)$ -modules, and a proof can be given along the lines of [7] by analyzing Virasoro algebra fusion. However, if p is prime, because of Lemma 6.2 [7] and the discussion after the lemma it is clear that there are no (logarithmic) extensions between different blocks, so no Virasoro fusion is needed. So let \mathcal{O} be the category of finitely generated weak $\mathcal{W}(p)$ -modules, and

$$\mathcal{O} = \bigoplus_{i=1}^{p+1} \mathcal{O}[h_{i,1}]$$

the corresponding block decomposition.


In the previous section we discussed in more details two logarithmic modules $\mathcal{P}_{p-1}^+ \in \text{Ob}(\mathcal{O}[h_{p-1,1}])$ and $\mathcal{P}_{p-1}^- \in \text{Ob}(\mathcal{O}[h_{1,1}])$.

Notice that inside the logarithmic module \mathcal{P}_{p-1}^+ we have realized the following non-trivial extensions of $\mathcal{W}(p)$ -modules:

$$\begin{aligned} 0 &\rightarrow \Lambda(p-1) \rightarrow \mathcal{M}_1 \rightarrow \Pi(1) \rightarrow 0; \\ 0 &\rightarrow \Lambda(p-1) \rightarrow V_{L+\alpha/2-\alpha/p} \rightarrow \Pi(1) \rightarrow 0; \\ 0 &\rightarrow \Lambda(p-1) \rightarrow \mathcal{N}_1 \rightarrow \Pi(1) \oplus \Pi(1) \rightarrow 0; \\ 0 &\rightarrow \mathcal{N}_1 \rightarrow \mathcal{P}_{p-1}^+ \rightarrow \Lambda(p-1) \rightarrow 0. \end{aligned}$$

From this we clearly obtain the following embedding structure for the module \mathcal{P}_{p-1}^+ (a similar diagram can be drawn for \mathcal{P}_{p-1}^-):


(6.1)



We find the structure of this logarithmic module identical to the structure of the corresponding projective modules in the case of the quantum group $\bar{U}_q(sl_2)$, $q = e^{\pi i/p}$ [18], [19].


From (6.1) we also obtain the following indecomposable module (the so-called "wedge"):

(6.2)



simply by taking the obvious quotient of \mathcal{P}_{p-1}^+ . But there is also another wedge of the form:

(6.3)



We can now "glue" together several wedges (6.2) to obtain:

$$(6.4) \quad \begin{array}{ccccccc} & \Lambda(p-1) & & \dots & & \Lambda(p-1) & \\ & \bullet & & & & \bullet & \\ & \swarrow & & & & \searrow & \\ \Pi(1) & & \Pi(1) & & \Pi(1) & & \Pi(1) \\ \bullet & & \bullet & & \bullet & & \bullet \end{array}$$

and similarly we get

$$(6.5) \quad \begin{array}{ccccccc} \Pi(1) & & \Pi(1) & & \dots & & \Pi(1) \\ \bullet & & \bullet & & & & \bullet \\ & \searrow & \swarrow & & \swarrow & & \searrow \\ & \Lambda(p-1) & & \dots & & \Lambda(p-1) & \\ & \bullet & & & & \bullet & \end{array}$$

All representations considered here can be constructed explicitly. For instance (6.3) is obtained by taking the $L(0)$ -semisimple part of \mathcal{P}_{p-1}^+ (cf. module \mathcal{N}_1 in the previous section). The representation (6.1) is the only logarithmic representation here and conjecturally this is a projective cover of the irreducible module $\Lambda(p-1)$. The other modules considered here are clearly indecomposable. We can of course obtain similar modules within the $\mathcal{O}[h_{1,1}]$ block, we only have to switch the roles of Λ and Π .

Another interesting class of representations is obtained as submodules of \mathcal{P}_{p-1}^+ , where there infinitely many nonisomorphic submodules parameterized by $[x : y] \in \mathbb{CP}^1$. These modules can be again constructed explicitly (we omit details here), and are in fact different Baer sums of extensions obtained from \mathcal{M}_1 and $V_{L+\alpha/2-\alpha/p}$. These modules have embedding structure

$$(6.6) \quad \begin{array}{c} \Pi(1) \\ \bullet \\ \begin{array}{c} \left. \begin{array}{l} \downarrow \\ \downarrow \end{array} \right\} x \\ \left. \begin{array}{l} \downarrow \\ \downarrow \end{array} \right\} y \end{array} \\ \Lambda(p-1) \\ \bullet \end{array}$$

This type of modules are known to exist on the quantum group side.

Remark 6.1. *The block $\mathcal{O}[h_{1,1}]$ is important for several reasons. Firstly, this is the principal block since it contains the "trivial" module $\mathcal{W}(p)$. But this block is also the "shortest" in the sense that two irreducible modules ($\mathcal{W}(p) = \Lambda(1)$ and $\Pi(p-1)$ are at the "conformal distance" one). Thus analysis of Zhu's algebra $A_1(\mathcal{W}(p))$ ought to be sufficient to classify all indecomposable modules in the block. We plan to study this problem in our future work.*

7. LOGARITHMIC MODULES FOR $\mathcal{SW}(m)$ AND $\mathcal{SW}(p, p')$

As in the previous section we can now construct logarithmic modules for the super-triplet vertex algebra $\mathcal{SW}(m)$ and related extensions of $N = 1$ superconformal minimal models $\mathcal{SW}(p, p')$. To the best of our knowledge, the algebra $\mathcal{SW}(p, p')$ has not been previously studied, so here we give its definition and a few basic properties omitting proofs that are identical as those in [8].

Let $p > p'$ be positive integers where $(p, \frac{p-p'}{2}) = 1$. Consider the rank one lattice $\mathbb{Z}\alpha$, with $\langle \alpha, \alpha \rangle = pp'$, and the vector space $V_L = M(1) \otimes \mathbb{C}[L]$, where $M(1)$ is as before. We have the parity decomposition $V_L = V_L^0 \oplus V_L^1$ in the standard way, and verify that this decomposition is compatible with the (super)bracket structure giving a vertex superalgebra structure on V_L . Then we form the vertex superalgebra $V_L \otimes F$, where F is the free fermion vertex superalgebra generated by the field $\phi(x) = \sum_{n \in \mathbb{Z}} \phi(n)x^{-n-1/2}$.

We introduce (super)conformal structure on $V_L \otimes F$ by choosing superconformal vector τ and conformal vector ω as follows:

$$\tau = \frac{\alpha(-1)\mathbf{1}}{\sqrt{pp'}} \otimes \phi(-1/2)\mathbf{1} + \frac{p-p'}{\sqrt{pp'}}\mathbf{1} \otimes \phi(-3/2)\mathbf{1},$$

and

$$\omega = \frac{\alpha(-1)^2\mathbf{1}}{2pp'} \otimes \mathbf{1} + (p-p')\frac{\alpha(-2)\mathbf{1}}{2pp'} \otimes \mathbf{1} + \frac{1}{2}(\mathbf{1} \otimes \phi(-3/2)\phi(-1/2)\mathbf{1}).$$

It is easy to check that $Y(\tau, x)$ and $Y(\omega, x)$ close the $N = 1$ Neveu-Schwarz superconformal algebra of central charge $\frac{3}{2}(1 - 2\frac{(p-p')^2}{pp'})$, and thus $V_L \otimes F$ is an $N = 1$ vertex operator superalgebra. If we denote the eigenvalue of $L(0)$ operator by $wt(v)$ then simple computation shows

$$wt(e^{l\alpha}) = \frac{l^2 pp' - l(p-p')}{2}.$$

Now, let L° denote the dual lattice and V_{L° generalized vertex operator superalgebra. Therefore $e^{\alpha/p'} \in \mathbb{C}[L^\circ]$ and $e^{-\alpha/p} \in \mathbb{C}[L^\circ]$ are odd vectors of weight $\frac{1}{2}$, and hence $e^{\alpha/p'} \otimes \phi(-1/2)\mathbf{1}$ and $e^{-\alpha/p} \otimes \phi(-1/2)\mathbf{1}$ are even vectors of weight one. It is not hard to show (see [8]) that the screening operators

$$Q = \text{Res}_x Y(e^{\alpha/p'} \otimes \phi(-1/2)\mathbf{1}, x), \quad \tilde{Q} = \text{Res}_x Y(e^{-\alpha/p} \otimes \phi(-1/2)\mathbf{1}, x)$$

commute with the action of the $N = 1$ superconformal algebra. We now define

$$\mathcal{SW}(p, p') := \text{Ker}_{V_L \otimes F}(Q) \cap \text{Ker}_{V_L \otimes F}(\tilde{Q}).$$

From the previous discussion we obtain

Proposition 7.1. *We have*

$$V_{1/2}(c_{p,p'}, 0) \subset \mathcal{SW}(p, p'),$$

thus $\mathcal{SW}(p, p')$ is an $N = 1$ superconformal vertex algebra.

We expect the vertex superalgebra $\mathcal{SW}(p, p')$ to be C_2 -cofinite. By using the construction analogous to the one for $\mathcal{W}(p, p')$ we can show that each $\mathcal{SW}(p, p')$ admits logarithmic modules.

Proposition 7.2.

- (i) For $i = 0, \dots, p' - 1$, let $\mathcal{SV}_i = \left(V_{L+\frac{i}{p'}\alpha} \oplus V_{L+\frac{i}{p'}\alpha-\frac{\alpha}{p}} \right) \otimes F$. Then $\mathcal{SV} = \mathcal{SV}_0$ is a vertex operator superalgebra and \mathcal{SV}_i is a \mathcal{SV} -module.
- (ii) For $j = 0, \dots, p - 1$, let $\mathcal{SV}_j^o = \left(V_{L+\frac{j}{p}\alpha} \oplus V_{L+\frac{j}{p}\alpha+\frac{\alpha}{p'}} \right) \otimes F$. Then $\mathcal{SV}^o = \mathcal{SV}_0^o$ is a vertex operator superalgebra and \mathcal{SV}_j^o is a \mathcal{SV}^o -module

Theorem 7.1.

- (i) Assume that (M, Y_M) is a weak \mathcal{SV} -module. Then the structure

$$(\widetilde{M}, \widetilde{Y}_{\widetilde{M}}(\cdot, x)) := (M, Y_M(\Delta(e^{-\alpha/p} \otimes \phi(-\frac{1}{2}), x) \cdot, x))$$

is a weak $\mathcal{SW}(p, p')$ -module. Moreover,

$$(\widetilde{\mathcal{SV}}_i, \widetilde{Y}_{\widetilde{\mathcal{SV}}_i}), \quad i = 0, \dots, p' - 1,$$

are logarithmic $\mathcal{SW}(p, p')$ -modules.

- (ii) Assume that (M, Y_M) is a weak \mathcal{SV}^o -module. Then the structure

$$(\widetilde{M}, \widetilde{Y}_{\widetilde{M}}(\cdot, x)) := (M, Y_M(\Delta(e^{\alpha/p'} \otimes \phi(-\frac{1}{2}), x) \cdot, x))$$

is a weak $\mathcal{SW}(p, p')$ -module. In particular,

$$(\widetilde{\mathcal{SV}}_j^o, \widetilde{Y}_{\widetilde{\mathcal{SV}}_j^o}), \quad j = 0, \dots, p - 1,$$

are logarithmic $\mathcal{SW}(p, p')$ -modules.

It is easy to modify the previous result to the case $p' = 1$ and $p = 2m + 1$ considered in [8], but now with only one screening operator. We shall use the same notation as in [8]. Let $L = \mathbb{Z}\alpha$ be a rank one lattice with nondegenerate form given by $\langle \alpha, \alpha \rangle = 2m + 1$, where $m \in \mathbb{Z}_{>0}$. Let V_L be the corresponding vertex superalgebra. Let F be the fermionic vertex algebra as above and

M its canonical twisted module as in [8] and [9]. We consider the vertex superalgebra $V_L \otimes F$. Let σ be the canonical \mathbb{Z}_2 automorphism of $V_L \otimes F$.

The super-triplet vertex algebra is defined as

$$\mathcal{SW}(m) = \text{Ker}_{V_L \otimes F} \tilde{Q}.$$

It was proved in [8] that $\mathcal{SW}(m)$ is an irrational, C_2 -cofinite vertex operator superalgebra. The irreducible and irreducible σ -twisted $\mathcal{SW}(m)$ -modules were classified in [8] and [9].

Now we shall construct certain logarithmic (σ -twisted) modules for $\mathcal{SW}(m)$.

Proposition 7.3.

- (i) The vector space $\mathcal{SV} = (V_L \oplus V_{L-\alpha/(2m+1)}) \otimes F$ admits a structure of vertex operator superalgebra.
- (ii) The vector space $\mathcal{RV} = (V_{L+\alpha/2} \oplus V_{L+\alpha/2-\alpha/(2m+1)}) \otimes M$ is a σ -twisted \mathcal{SV} -module.

Let now $v = e^{-\alpha/(2m+1)} \otimes \phi(-\frac{1}{2})$. Applying Theorem 2.1 in the case of the vertex operator superalgebra \mathcal{SV} we get construction of logarithmic modules for $\mathcal{SW}(m)$.

Theorem 7.2.

- (i) $(\widetilde{\mathcal{SV}}, \widetilde{Y}_{\widetilde{\mathcal{SV}}})$ is a logarithmic $\mathcal{SW}(m)$ -module of semisimple length three.
- (ii) $(\widetilde{\mathcal{RV}}, \widetilde{Y}_{\widetilde{\mathcal{RV}}})$ is logarithmic σ -twisted $\mathcal{SW}(m)$ -module of semisimple length three.

Let us describe the twisted logarithmic module $\widetilde{\mathcal{RV}}$ in more details. Let $(V_{\tilde{L}}, Y)$ be the generalized vertex algebra associated to the lattice $\tilde{L} = \mathbb{Z} \frac{\alpha}{2(2m+1)}$. As in [9], let

$$\phi(x) = \sum_{n \in \mathbb{Z}} \phi(n) x^{-n - \frac{3}{2}}$$

be the field which defines on M the structure of twisted F -module. Then one can show that

$$\tilde{Q} = \text{Res}_x Y(e^{-\alpha/(2m+1)}, x) \otimes \phi(z) = \sum_{n \in \mathbb{Z}} (e^{-\alpha/(2m+1)})_{-n - \frac{3}{2}} \otimes \phi(n)$$

is the (second) screening operator which commutes with the action of the Ramond algebra (see Remark 5.2 of [9]). By using similar analysis as in Section 5, one can see that on twisted module $\widetilde{\mathcal{RV}}$:

$$\widetilde{L}(0) = L(0) + \tilde{Q}.$$

Then $\widetilde{\mathcal{RV}}$ is a $\mathbb{Z}_{\geq 0}$ -graded and that on its top level $\widetilde{\mathcal{RV}}(0)$, the operator $\widetilde{L}(0)$ acts (in the obvious basis) as

$$\begin{bmatrix} h^{2m,1} & 1 \\ 0 & h^{2m,1} \end{bmatrix},$$

where $h^{2m,1} = \frac{1-2m}{8} + \frac{1}{16}$. This gives additional evidence for conjectures presented in [9].

8. LOGARITHMIC REPRESENTATIONS OF THE AFFINE VERTEX OPERATOR ALGEBRA $L(-\frac{4}{3}\Lambda_0)$

Admissible representations of affine Lie algebras were introduced by V. Kac and M. Wakimoto in [29]. These representations have been also studied in the framework of vertex operator algebras (cf. [2], [5], [30], [39]).

In this section we shall present a construction of logarithmic representations of the affine vertex operator algebra $L(-\frac{4}{3}\Lambda_0)$ which uses realization from [4]. This explicit realization enable us to apply Theorem 2.1 in the case of the vertex operator algebra $L(-\frac{4}{3}\Lambda_0)$.

First we shall briefly recall the construction from [4].

Define the following lattice

$$\tilde{L} = \mathbb{Z}\gamma + \mathbb{Z}\delta, \quad \langle \gamma, \gamma \rangle = -\langle \delta, \delta \rangle = \frac{1}{6}, \quad \langle \gamma, \delta \rangle = 0.$$

Let $V_{\tilde{L}}$ be the associated generalized vertex algebra with the vertex operator map Y .

Now we define the screening operators:

$$Q = \text{Res}_x Y(e^{-6\gamma}, x) = e_0^{-6\gamma},$$

$$\tilde{Q} = \text{Res}_x Y(e^{2\gamma}, x) = e_0^{2\gamma},$$

and the following Virasoro element

$$\omega = (3\gamma(-1)^2 - 2\gamma(-2) - 3\delta(-1)^2)\mathbf{1}.$$

Then ω generates the Virasoro vertex operator algebra $L^{Vir}(-6, 0)$. Set

$$L(x) = Y(\omega, x) = \sum_{n \in \mathbb{Z}} L(n)x^{-n-2}.$$

Define the following vectors in $V_{\tilde{L}}$:

$$e = e^{3(\gamma-\delta)},$$

$$h = 4\delta(-1)\mathbf{1},$$

$$f = -\frac{2}{9}Qe^{3(\gamma+\delta)}$$

$$= -(4\gamma(-1)^2 - \frac{2}{3}\gamma(-2))e^{-3(\gamma-\delta)}.$$

Then e, h, f are primary vectors of conformal weight 1 for the Virasoro algebra.

Define

$$D = \mathbb{Z}(3\gamma + 3\delta) + \mathbb{Z}(3\gamma - 3\delta).$$

Clearly, V_D is a subalgebra of $V_{\tilde{L}}$ which contains vectors e , f and h . Since D is an even lattice, V_D is a vertex algebra.

Theorem 8.1. [4] *The vectors e , f and h generate a subalgebra of the vertex algebra V_D isomorphic to the vertex operator algebra $L(-\frac{4}{3}\Lambda_0)$. Moreover,*

$$L(-\frac{4}{3}\Lambda_0) \subset \text{Ker}_{V_D} \tilde{Q}.$$

Now we shall consider the vertex operator algebra

$$\mathcal{V} = V_D \oplus V_{D+2\gamma}$$

and its simple module

$$\mathcal{M} = V_{D-3\gamma+\delta} \bigoplus V_{D-\gamma+\delta}.$$

(The associated vertex operators are restrictions of the vertex operator in the generalized vertex algebra $V_{\tilde{L}}$ and will be also denoted by Y).

We also note that $V_{D-2\gamma}$ and $V_{D-\gamma+\delta}$ are (irreducible) \mathcal{V} -modules (on which $V_{D+2\gamma}$ acts trivially) and that there is a non-trivial intertwining operator \mathcal{Y} of type

$$(8.1) \quad \begin{pmatrix} \mathcal{M} \\ V_{D-\gamma+\delta} & V_{D-2\gamma} \end{pmatrix}.$$

Let now $v = e^{2\gamma}$. Applying Theorem 2.1 in the case of the vertex operator superalgebra \mathcal{V} we get construction of logarithmic $L(-\frac{4}{3}\Lambda_0)$ -modules.

Theorem 8.2. *The pairs $(\tilde{\mathcal{V}}, \tilde{Y})$ and $(\tilde{\mathcal{M}}, \tilde{Y})$ are logarithmic $L(-\frac{4}{3}\Lambda_0)$ -modules. In particular, the components of the fields*

$$\begin{aligned} \widetilde{e}(x) &= \tilde{Y}(e, x) = Y(e, x), \\ \widetilde{h}(x) &= \tilde{Y}(h, x) = Y(h, x), \\ \widetilde{f}(x) &= \tilde{Y}(f, x) = Y(f, x) + \frac{4}{3}x^{-1}Y(\gamma(-1)e^{-\gamma+3\delta}, x) + \frac{1}{3}x^{-2}Y(e^{-\gamma+3\delta}, x), \end{aligned}$$

define on \mathcal{V} and \mathcal{M} structures of $A_1^{(1)}$ -modules of level $-\frac{4}{3}$.

Define now the following module

$$\mathcal{R}_{-1/3} = U(\widehat{sl_2}) \cdot e^{-3\gamma+\delta} \subset \tilde{\mathcal{M}}.$$

Recall that

$$\widetilde{L}(0) = L(0) + \tilde{Q}.$$

Then

$$\begin{aligned}\widetilde{L(0)}e^{-3\gamma+\delta} &= -\frac{1}{3}e^{-3\gamma+\delta} + e^{-\gamma+\delta}, \\ \widetilde{L(0)}e^{-\gamma+\delta} &= -\frac{1}{3}e^{-\gamma+\delta}.\end{aligned}$$

Therefore $\mathcal{R}_{-1/3}$ is an logarithmic $L(-\frac{4}{3}\Lambda_0)$ -module.

On the other hand, the results from [4] implies that

$$U(\widehat{sl_2}).e^{-\gamma+\delta} \cong L(-\frac{2}{3}\Lambda_0 - \frac{2}{3}\Lambda_1),$$

and that

$$E = U(\widehat{sl_2}).e^{-2\gamma}$$

is a $\mathbb{Z}_{\geq 0}$ -graded $L(-\frac{4}{3}\Lambda_0)$ -module.

Note that $\mathcal{R}_{-1/3}$ contains $e^{-\gamma+\delta}$ and therefore a submodule isomorphic to $L(-\frac{2}{3}\Lambda_0 - \frac{2}{3}\Lambda_1)$. But we shall now see that $\mathcal{R}_{-1/3}$ is not $\mathbb{Z}_{\geq 0}$ -graded. We have:

$$(8.2) \quad e(\widetilde{n+1})e^{-3\gamma+\delta} = f(\widetilde{n+2})e^{-3\gamma+\delta} = 0 \quad \text{for } n \geq 0,$$

$$(8.3) \quad \widetilde{e(0)}e^{-3\gamma+\delta} = e^{-2\delta}, \quad \widetilde{f(1)}e^{-3\gamma+\delta} = \nu e^{-4\gamma+4\delta} \quad (\nu \neq 0).$$

Thus $e^{-4\gamma+4\delta} \in \mathcal{R}_{-1/3}$.

By using explicit realization [4], one can see that

$$E_1 = U(\widehat{sl_2}).e^{-4\gamma+4\delta}$$

is a weak, ordinary module \mathbb{Z} -graded module, but gradation is not bounded. (Note that also $L(-\frac{2}{3}\Lambda_0 - \frac{2}{3}\Lambda_1)$ is isomorphic to a submodule of E_1). Now relations (8.2)-(8.3) imply that

$$E_2 = \mathcal{R}_{-1/3}/U(\widehat{sl_2}).e^{-4\gamma+4\delta}$$

is a non-logarithmic $\mathbb{Z}_{\geq 0}$ -graded $L(-\frac{4}{3}\Lambda_0)$ -module.

Proposition 8.1. *The logarithmic $L(-\frac{4}{3}\Lambda_0)$ -module $\mathcal{R}_{-1/3}$ appears in the extension*

$$0 \rightarrow E_1 \rightarrow \mathcal{R}_{-1/3} \rightarrow E_2 \rightarrow 0$$

of non-logarithmic weak module E_1 and non-logarithmic $\mathbb{Z}_{\geq 0}$ -graded module E_2 .

The results from this section imply the existence of the following intertwining operator (this intertwining operator can be also constructed from intertwining operator (8.1) and methods from Section 9 below).

Proposition 8.2. *There is a non-trivial (logarithmic) intertwining operator of type*

$$\begin{pmatrix} \mathcal{R}_{-1/3} & \\ L(-\frac{2}{3}\Lambda_0 - \frac{2}{3}\Lambda_1) & E \end{pmatrix}.$$

Remark 8.1. *We should mention that the existence of such logarithmic module $\mathcal{R}_{-1/3}$ was predicted in the fusion rules analysis of [23]. Gaberdiel found $\mathcal{R}_{-1/3}$ in the fusion product (which should correspond to tensor product) of non-logarithmic $\mathbb{Z}_{\geq 0}$ -graded modules $L(-\frac{2}{3}\Lambda_0 - \frac{2}{3}\Lambda_1)$ and E . Our intertwining operator from Proposition 8.2 is in agreement with this fusion product in [23].*

One can also check that the structure of our logarithmic module $\mathcal{R}_{-1/3}$ can be also illustrated by Figure 2 in [23]. So we believe that our construction present an explicit realization of this logarithmic module.

We can also make a similar analysis of other logarithmic submodules of $\tilde{\mathcal{V}}$ and $\tilde{\mathcal{M}}$. We plan to study these logarithmic modules and the corresponding fusion rules in our forthcoming papers.

9. LOGARITHMIC INTERTWINING OPERATORS

Let us recall the main ingredients in the definition of logarithmic intertwining operators following [35], [37] and [27]. These are simply intertwining-like operator maps

$$\mathcal{Y}(w, x) : W_1 \otimes W_2 \longrightarrow W_3\{x\}[\log(x)],$$

satisfying the $L(-1)$ -property and the Jacobi identity identical to the ordinary Jacobi identity in the definition of ordinary intertwining operator maps. These objects appear naturally if one studies differential equations satisfied by correlation functions [35]. First nontrivial constructions of intertwining operators were obtained in [37] and [6] (these structures were predicted earlier in [35]). In this part we present a general construction of intertwining operators among certain logarithmic modules. The construction here is heavily influenced by results in previous sections and especially [17], where closely related operators were considered.

We shall use the same framework as in Section 2. Let V be a vertex operator superalgebra. Let again $v \in V$ satisfies conditions (2.1) and (2.2). Denote by \bar{V} any vertex subalgebra of V contained in the kernel of v_0 . If M is a weak V -module, then the pair $(\tilde{Y}_{\bar{M}}, \tilde{M})$ defines a weak \bar{V} -module.

Since M is an V -module, we have an intertwining operator, which we denote by \mathcal{Y} , of type

$$\begin{pmatrix} M \\ V \quad M \end{pmatrix}.$$

We are aiming for a logarithmic intertwining operator of type

$$\begin{pmatrix} \widetilde{M} \\ V \widetilde{M} \end{pmatrix}.$$

Recall

$$\Delta(v, x) = x^{v_0} \exp\left(\sum_{n=1}^{\infty} \frac{v_n}{-n} (-x)^{-n}\right)$$

be as before. Observe (again!) that the expression

$$\Delta(v, x)w, \quad w \in V,$$

is ambiguous in general, but

$$\Delta(v, x)w, \quad w \in \overline{V},$$

is well-defined. To fix the problem we introduce a new operator (cf. [17] and [37] for closely related constructions):

$$(9.1) \quad \Delta_{\log}(v, x) = \exp\left(\log(x)v_0 + \sum_{n=1}^{\infty} \frac{v_n}{-n} (-x)^{-n}\right),$$

where as usual

$$\exp(\log(x)v_0) = \sum_{n=0}^{\infty} \frac{\log(x)^n v_0^n}{n!},$$

and $\log(x)$ is a formal variable satisfying $(\log(x))' = \frac{1}{x}$.

Notice that now $\Delta_{\log}(v, x)w$ involves logarithmic terms for $w \in V$, but if $w \in \overline{V}$ then

$$(9.2) \quad \Delta(v, x)w = \Delta_{\log}(v, x)w.$$

We first observe the following useful result:

Lemma 9.1. *With $u \in \overline{V}$ and v as above, then*

$$\Delta_{\log}(v, x_2)Y(u, y) = Y(\Delta(x_2 + y)u, y)\Delta_{\log}(v, x_2).$$

Proof. Because of $[v_0, v_m] = 0$ we can write

$$\Delta_{\log}(v, x) = \exp(\log(x)v_0)E^+(-v, -x),$$

where $E^+(v, x)$ is defined as usual.

As in Lemma 2.15 of [34] we get

$$E^+(v, x)Y(u, y)E^+(-v, x) = Y((-x)^{v_0}\Delta(-v, -x + y)u, y).$$

As in [35], we also have

$$\exp(\log(x)v_0)Y(u, y)\exp(-\log(x)v_0) = Y(\exp(\log(x)v_0)u, y).$$

Combined together we get

$$\begin{aligned} & \Delta_{\log}(v, x_2)Y(u, y)\Delta_{\log}(-v, x_2) \\ &= \exp(\log(x)v_0)E^+(-v, -x)Y(u, y)E^+(v, -x)\exp(-\log(x)v_0) \\ &= \exp(\log(x)v_0)Y(x^{-v_0}\Delta(v, x+y)u, y)\exp(-\log(x)v_0) \\ (9.3) \quad &= Y(\exp(\log(x)v_0)x^{-v_0}\Delta(v, x+y)u, y). \end{aligned}$$

Finally, if we take $v \in \bar{V}$, we obtain the wanted formula. \square

Here is the main result of this section

Theorem 9.1. *Let \mathcal{Y} be intertwining operator as above and v as before such that v_0 acts nilpotently on V . Then $\tilde{\mathcal{Y}}(\cdot, x) := \mathcal{Y}(\Delta_{\log}(v, x)\cdot, x)$ is a logarithmic intertwining operator of type*

$$\begin{pmatrix} \tilde{M} \\ V \tilde{M} \end{pmatrix}$$

in the category of \bar{V} -modules.

Proof. The main idea is already in [34]. Because of

$$[L(-1), v_0] = 0, \quad [L(-1), v_n] = -nv_{n-1},$$

we have

$$[L(-1), \Delta_{\log}(u, x)] = -\frac{d}{dx}(\Delta_{\log}(v, x)),$$

which implies

$$[L(-1), \tilde{\mathcal{Y}}(u, x)] = \frac{d}{dx}\tilde{\mathcal{Y}}(u, x),$$

the $L(-1)$ -derivative property.

For the Jacobi identity, we start from the ordinary Jacobi identity for the intertwining operator $\mathcal{Y}(u, x)$, where $u \in V$:

$$\begin{aligned} & x_0^{-1}\delta\left(\frac{x_1 - x_2}{x_0}\right)\tilde{\mathcal{Y}}(u, x_1)\tilde{\mathcal{Y}}(w, x_2) - x_0^{-1}\delta\left(\frac{-x_2 + x_1}{x_0}\right)\tilde{\mathcal{Y}}(w, x_2)\tilde{\mathcal{Y}}(u, x_1) \\ &= x_0^{-1}\delta\left(\frac{x_1 - x_2}{x_0}\right)Y(\Delta(v, x_1)u, x_1)\mathcal{Y}(\Delta_{\log}(v, x_2)w, x_2) \end{aligned}$$

$$\begin{aligned}
& -x_0^{-1}\delta\left(\frac{-x_2+x_1}{x_0}\right)\mathcal{Y}(\Delta_{\log}(v,x_2)w,x_2)Y(\Delta(v,x_1)u,x_1) \\
& = x_2^{-1}\delta\left(\frac{x_1-x_0}{x_2}\right)\mathcal{Y}(Y(\Delta(v,x_1)u,x_0)\Delta_{\log}(v,x_2)w,x_2) \\
& = x_2^{-1}\delta\left(\frac{x_1-x_0}{x_2}\right)\mathcal{Y}(Y(\Delta(v,x_2+x_0)u,x_0)\Delta_{\log}(v,x_2)w,x_2).
\end{aligned}$$

Finally, we apply Lemma 9.1 so the right hand-side becomes

$$x_2^{-1}\delta\left(\frac{x_1-x_0}{x_2}\right)\mathcal{Y}(\Delta_{\log}(v,x_2)Y(u,x_0)w,x_2),$$

which is what we needed to prove. □

By using similar proof to that of Theorem 9.1 we have

Theorem 9.2. *Assume that M_1, M_2 and M_3 are V -modules and that $v \in V$ satisfies conditions (2.1) and (2.2) and v_0 acts nilpotently on $M_i, i = 1, 2, 3$. Let \mathcal{Y} be an intertwining operator of type $\begin{pmatrix} M_3 \\ M_1 \ M_2 \end{pmatrix}$ in the category of V -modules. Then $\tilde{\mathcal{Y}}(\cdot, x) := \mathcal{Y}(\Delta_{\log}(v, x)\cdot, x)$ is a logarithmic intertwining operator of type*

$$\begin{pmatrix} \tilde{M}_3 \\ M_1 \ \tilde{M}_2 \end{pmatrix}$$

in the category of \bar{V} -modules.

Acknowledgments

D.A. would like to thank the Erwin Schrödinger Institute in Vienna for hospitality. We also thank anonymous referee for his/her constructive comments. The second author was partially supported by NSF grant DMS-0802962.

REFERENCES

- [1] T. Abe, A \mathbb{Z}_2 -orbifold model of the symplectic fermionic vertex operator superalgebra. *Math. Z.* **255** (2007), 755–792.
- [2] D. Adamović, Some rational vertex algebras, *Glas. Mat. Ser. III* **29(49)** (1994), 25–40.
- [3] D. Adamović, Classification of irreducible modules of certain subalgebras of free boson vertex algebra, *J. Algebra* **270** (2003) 115–132.
- [4] D. Adamović, A construction of admissible $A_1^{(1)}$ -modules of level $-\frac{4}{3}$, *J. Pure Appl. Algebra* **196** (2005) 119–134.
- [5] D. Adamović and A. Milas, Vertex operator algebras associated to the modular invariant representations for $A_1^{(1)}$, *Math. Res. Lett.* **2** (1995) 563–575.

- [6] D. Adamović and A. Milas, Logarithmic intertwining operators and $\mathcal{W}(2, 2p - 1)$ -algebras, *Journal of Math. Physics* **48**, 073503 (2007).
- [7] D. Adamović and A. Milas, On the triplet vertex algebra $\mathcal{W}(p)$, *Advances in Math.* **217** (2008) 2664–2699; arXiv:0707.1857.
- [8] D. Adamović and A. Milas, The $N = 1$ triplet vertex operator superalgebras, *Comm. Math. Phys.* **288** (2009), 225–270; arXiv:0712.0379.
- [9] D. Adamović and A. Milas, The $N = 1$ triplet vertex operator superalgebras: twisted sector, *SIGMA* **4** (2008) 24 pages, arXiv:0806.3560.
- [10] D. Adamović and A. Milas, An analogue of modular BPZ-equations in logarithmic (super)conformal field theory, *Vertex Operator Algebras and Related Areas* Edited by: Maarten Bergvelt, Gaywalee Yamskulna and Wenhua Zhao, Normal, IL 2009; *Contemporary Mathematics* **497**, 17 pages.
- [11] D. Adamović and A. Milas, On W -algebras associated to $(2, p)$ minimal models and their representations, preprint; arXiv:0908.4053.
- [12] T. Arakawa, Representation theory of superconformal algebras and the Kac-Roan-Wakimoto conjecture, *Duke Math. J.* **130** (2005), 435–478.
- [13] N. Carqueville and M. Flohr, Nonmeromorphic operator product expansion and C_2 -cofiniteness for a family of \mathcal{W} -algebras. *J. Phys. A* **39** (2006), 951–966.
- [14] A. De Sole and V. Kac, Finite vs. affine W -algebras, *Japanese Journal of Math.* **1** (2006) 137–261; arxiv:math/05110055.
- [15] C. Dong, H. Li, and G. Mason, Simple currents and extensions of vertex operator algebras. *Comm. Math. Phys.* **180** (1996) 671–707.
- [16] M. Flohr, On modular invariant partition functions of conformal field theories with logarithmic operators, *Internat. J. Modern Phys. A* **11** (1996), 4147–4172.
- [17] J. Fjelstad, J. Fuchs, S. Hwang, A.M. Semikhatov and I. Yu. Tipunin, Logarithmic conformal field theories via logarithmic deformations, *Nuclear Phys. B* **633** (2002), 379–413.
- [18] B.L. Feigin, A.M. Gaĭnutdinov, A. M. Semikhatov, and I. Yu Tipunin, Modular group representations and fusion in logarithmic conformal field theories and in the quantum group center. *Comm. Math. Phys.* **265** (2006), 47–93.
- [19] B.L. Feigin, A.M. Gaĭnutdinov, A. M. Semikhatov, and I. Yu Tipunin, Kazhdan–Lusztig correspondence for the representation category of the triplet W -algebra in logarithmic CFT, *Theor.Math.Phys.* **148** (2006) 1210–1235; *Teor.Mat.Fiz.* **148** (2006) 398–427.
- [20] B.L. Feigin, A.M. Gaĭnutdinov, A. M. Semikhatov, and I. Yu Tipunin, Logarithmic extensions of minimal models: characters and modular transformations, *Nucl. Phys. B* **757** (2006) 303343.
- [21] J. Fuchs, On non-semisimple fusion rules and tensor categories, *Contemporary Mathematics* **442** (2007) 315–337, arXiv:hep-th/0602051.
- [22] J. Fuchs, S. Hwang, A.M. Semikhatov and I. Yu. Tipunin, Nonsemisimple Fusion Algebras and the Verlinde Formula, *Comm. Math. Phys.* **247** (2004), no. 3, 713–742.
- [23] M. Gaberdiel, Fusion rules and logarithmic representations of a WZW model at fractional level, *Nuclear Phys. B* **618** (2001) 407–436.

- [24] M. Gaberdiel, An algebraic approach to logarithmic conformal field theory, Proceedings of the School and Workshop on Logarithmic Conformal Field Theory and its Applications (Tehran, 2001), *Internat. J. Modern Phys. A* **18** (2003), 4593–4638.
- [25] M. Gaberdiel and H. G. Kausch, A rational logarithmic conformal field theory, *Phys. Lett B* **386** (1996), 131-137, hep-th/9606050
- [26] Y.-Z. Huang, Cofiniteness conditions, projective covers and the logarithmic tensor product theory, *J. Pure Appl. Algebra* **213** (2009) 458-475; arXiv:0712.4109.
- [27] Y.-Z. Huang, J. Lepowsky and L. Zhang, Logarithmic tensor product theory for generalized modules for a conformal vertex algebra, arXiv:0710.2687.
- [28] V. Kac, *Vertex algebras for beginners*, University Lectures Series, Vol. 10, Providence, 1998.
- [29] V. Kac and M. Wakimoto, Modular invariant representations of infinite dimensional Lie algebras and superalgebras, *Proc. Natl. Acad. Sci. USA*, Vol. **85** (1988) 4956-4960.
- [30] V. Kac and M. Wakimoto, On rationality of W-algebras, *Transform. Groups* **13**, No. 3-4 (2008) 671-713, arXiv:0711.2296.
- [31] H. G. Kausch, Extended conformal algebras generated by multiplet of primary fields, *Phys. Lett.* **259 B** (1991), 448-455.
- [32] J. Lepowsky and H. Li, *Introduction to Vertex Operator Algebras and Their Representations*, Progress in Mathematics, Vol. **227**, Birkhäuser, Boston, 2003.
- [33] H. Li, Symmetric invariant bilinear forms on vertex operator algebras, *J. Pure Appl. Algebra* **96** (1994), 279297.
- [34] H. Li, The physics superselection principal in vertex operator algebra theory, *J. Algebra* **196** (1997), 436457.
- [35] A. Milas, Weak modules and logarithmic intertwining operators for vertex operator algebras. Recent developments in infinite-dimensional Lie algebras and conformal field theory (Charlottesville, VA, 2000), 201–225, *Contemp. Math.* **297**, Amer. Math. Soc., Providence, RI, 2002.
- [36] A. Milas, Fusion rings for degenerate minimal models, *J. Algebra* **254** (2002), no. 2, 300–335.
- [37] A. Milas, Logarithmic intertwining operators and vertex operators, *Comm. Math. Phys.* **277** (2008), 497-529; math.QA/0609306.
- [38] M. Miyamoto, Modular invariance of vertex operator algebras satisfying C_2 -cofiniteness. *Duke Math. J.* **122** (2004), 51–91.
- [39] O. Perše, Vertex operator algebras associated to type B affine Lie algebras on admissible half-integer levels, *J. Algebra* **307** (2007) 215-238.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ZAGREB, CROATIA

E-mail address: adamovic@math.hr

DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY AT ALBANY (SUNY), ALBANY, NY 12222

E-mail address: amilas@math.albany.edu