The application Breit-Wigner form with radiative corrections to the resonance fitting

K. Yu. Todyshev *

Budker Institute of Nuclear Physics, 630090, Novosibirsk, Russia

Abstract

The nonrelativistic and relativistic Breit-Wigner forms are conventionally used for the resonance fitting. In this note we consider the application of the Breit-Wigner formula with radiative corrections in initial state.

In this paper the problem of fitting a wide resonance is considered. The general expression for the cross section of e^+e^- annihilation into hadrons via an onium resonance is well known:

$$\sigma(W) = \frac{3\pi}{M^2} \int \frac{\Gamma_{ee}\Gamma_h}{(W\sqrt{1-x}-M)^2 + \frac{\Gamma^2}{4}} \mathcal{F}(x,W) dx , \qquad (1)$$

where W is the collision c.m. energy, Γ , Γ_{ee} and Γ_h are total and partial widths of the resonace, M is its mass and $\mathcal{F}(x, W)$ is the probability to lose the fraction of energy x because of the initial state radiation [1]. To obtain parameters of the resonances like J/ψ and $\psi(2S)$ the additional convolution with a Gaussian distribution is usually needed. Sometime it is possible to avoid double integration if simplify $\mathcal{F}(x, W)$ function. The case of a narrow resonace is described in literature (see [2]). In Ref. [3] the calculation was made for nonrelativistic Breit-Wigner in the soft-photon approximation, in which

$$\mathcal{F}(x,W) = \beta x^{\beta-1} (1+\delta) , \qquad (2)$$
$$(\ln \frac{W}{m_e} - \frac{1}{2}), \ \delta = \frac{\alpha}{\pi} \left(\frac{\pi^2}{3} - \frac{1}{2}\right) + \frac{3}{4}\beta - \frac{\beta^2}{24} \left(\frac{2}{3}\ln \frac{W}{m_e} + 2\pi^2 - \frac{37}{4}\right) .$$

That would be usefull to estimate errors of the functions published in Ref. [3], if make the similar calculations including the contribution of hard photons and the terms $\sim \frac{\Gamma}{W}$.

Let us consider $\sigma(W)$ following [3] in the assumption that $\frac{\Gamma}{W} \sim \beta$. We also include the terms $\sim \beta$ in $\mathcal{F}(x, W)$, which represent the contribution of hard photons in the first order by β . With that accuracy, the $\mathcal{F}(x, W)$ is

$$\mathcal{F}(x,W) = \beta x^{\beta-1}(1+\delta) - \beta (1-\frac{1}{2}x) , \qquad (3)$$

where $\beta = \frac{4\alpha}{\pi}$

^{*}e-mail:todyshev@inp.nsk.su

The integration over variable x gives the following expression

$$\sigma(W) = \frac{3\pi\Gamma_{ee}\Gamma_h}{M^2W\Gamma} \left[(1+\delta)\frac{4\pi\beta}{\sin\pi\beta} \operatorname{Im} f - \beta \mathcal{D}(W) \right], \qquad (4)$$

where

$$f = \left(\frac{M}{W} + i\frac{\Gamma}{2W}\right) \times \left(\frac{W}{M + W + i\frac{\Gamma}{2}}\right)^{1-\beta} \left(\frac{W}{M - W - i\frac{\Gamma}{2}}\right)^{1-\beta}$$
(5)

and

$$\mathcal{D}(W) = \frac{2M}{W} \left(1 + \left(\frac{M}{W}\right)^2 \right) \left(\arctan \frac{2M}{\Gamma} - \arctan \frac{2(M-W)}{\Gamma} \right)$$

$$-\frac{\Gamma}{2W} \left(1 + 3\left(\frac{M}{W}\right)^2 \right) \ln \frac{M^2 + \frac{\Gamma^2}{4}}{(M-W)^2 + \frac{\Gamma^2}{4}}$$

$$+\frac{3M\Gamma}{W^2} \left[1 - \frac{\Gamma}{2W} \left(\arctan \frac{2M}{\Gamma} - \arctan \frac{2(M-W)}{\Gamma} \right) \right]$$

$$+\frac{\Gamma}{2W} - \frac{M\Gamma}{W^2} + \frac{\Gamma^3}{8W^3} \ln \frac{M^2 + \frac{\Gamma^2}{4}}{(M-W)^2 + \frac{\Gamma^2}{4}} .$$
(6)

For relativistic Breit-Wigner form given by expression:

$$\sigma(s) = \int \frac{12\pi\Gamma_{ee}\Gamma_h}{(s(1-x) - M^2)^2 + \Gamma^2 M^2} \mathcal{F}(x,s) dx \tag{7}$$

where $s = W^2$; the integration over the variable x gives

$$\sigma(s) = \frac{12\pi\Gamma_{ee}\Gamma_h}{\Gamma M s} \left[(1+\delta)\frac{\pi\beta}{\sin\pi\beta} \operatorname{Im} f_r - \frac{\beta}{2} \left(1 + \frac{M^2}{s} \right) \left(\arctan\frac{M}{\Gamma} - \arctan\frac{M^2 - s}{\Gamma M} \right) + \frac{\beta\Gamma M}{4s} \ln\frac{\left(\frac{M^2}{s}\right)^2 + \left(\frac{\Gamma M}{s}\right)^2}{\left(1 - \frac{M^2}{s}\right)^2 + \left(\frac{\Gamma M}{s}\right)^2} \right],$$
(8)

where

$$f_r = \left(\frac{M^2}{s} - 1 - i\frac{\Gamma M}{s}\right)^{\beta - 1}.$$
(9)

In the case, where both resonant and non-resonant parts are present in the decay amplitude, the expression for cross section reads

$$\sigma(s) = \int \left| \frac{\sqrt{12\pi\Gamma_{ee}\Gamma_h}}{s(1-x) - M^2 + i\Gamma M} + Ce^{i\phi} \right|^2 \mathcal{F}(x,s)dx , \qquad (10)$$

where C is constant, and ϕ is phase angle of the interference. After integrating over x the cross section over can be represented in the form

$$\sigma(s) = \sigma_R(s) + \sigma_{NR}(s) - 2\frac{\sqrt{12\pi\Gamma_{ee}\Gamma_h}}{s} \frac{C(1+\delta)\pi\beta}{\sin\pi\beta} \operatorname{Re}\left(f_r e^{-i\phi}\right), \qquad (11)$$

where $\sigma_R(s), \sigma_{NR}(s)$ are resonant and non-resonant parts of the cross section, f_r is the function introduced in (9), and the third term corresponds to the interference between the resonant and non-resonant hadron production.

Let us consider the interference between two resonances with masses M_1 and M_2 , in that case, the cross section is :

$$\sigma(s) = \int \left| \frac{\sqrt{12\pi\Gamma_{1,ee}\Gamma_{1,h}}}{s(1-x) - M_1^2 + i\Gamma_1 M_1} + \frac{\sqrt{12\pi\Gamma_{2,ee}\Gamma_{2,h}}}{s(1-x) - M_2^2 + i\Gamma_2 M_2} e^{i\phi} \right|^2 \mathcal{F}(x,s) dx , \qquad (12)$$

where $\Gamma_1, \Gamma_{1,h}, \Gamma_{1,ee}, \Gamma_2, \Gamma_{2,h}, \Gamma_{2,ee}$ are total and partial widths of the resonances, and ϕ is phase angle of the interference. The result of calculation is:

$$\sigma(s) = \sigma_{1,R}(s) + \sigma_{2,R}(s) - \frac{24\pi\sqrt{\Gamma_{1,ee}\Gamma_{1,h}\Gamma_{2,ee}\Gamma_{2,h}}}{s} \frac{(1+\delta)\pi\beta}{\sin\pi\beta} \times \operatorname{Re}\left(\frac{f_1e^{-i\phi} - f_2e^{i\phi}}{M_1^2 - M_2^2 - i(\Gamma_1M_1 + \Gamma_2M_2)}\right),$$
(13)

where $\sigma_{1,R}(s)$, $\sigma_{2,R}(s)$ correspond to resonances cross sections, and the last term represents the interference between resonances. The functions f_1 and f_2 are similar to function introduced in (9) :

$$f_1 = \left(\frac{M_1^2}{s} - 1 - i\frac{\Gamma_1 M_1}{s}\right)^{\beta - 1},$$
 (14)

$$f_2 = \left(\frac{M_2^2}{s} - 1 - i\frac{\Gamma_2 M_2}{s}\right)^{\beta - 1}.$$
 (15)

It should be noted that although we have not discussed the energy dependence of total width, our results are also applicable in such case. If the production cross section has threshold behaviour our approximation can be used if $\frac{\Gamma}{M-\sqrt{s_{thr}}} << 1$.

We also stress that equations (6),(8), (11),(13) are applicable if $\frac{\Gamma}{W} \sim \beta$, $\frac{\Gamma}{W} \ll 1$.

To summarize, the cross section given by integration over the probability to emit certain fraction of energy from initial state has been calculated for the relativistic and nonrelativistic Breit-Wigner forms.

The author would likes to thank A.G. Shamov for helpful discussions and S.I. Eidelman for interest to this work.

References

- [1] E.A. Kuraev and V.S. Fadin, Yad. Fiz. 41, 733 (1985)
- [2] J. D. Jackson and D. D. Scharre, Nucl. Instrum. Methods. 128, 13 (1975)
- [3] Ya. I. Asimov, A. I. Vainshtein, L. N. Lipatov et al., JETP Lett. 21, 172 (1975)