ON THE RATE OF CONVERGENCE OF THE TWO-DIMENSIONAL α -MODELS OF TURBULENCE TO THE NAVIER-STOKES EQUATIONS

YANPING CAO AND EDRISS S. TITI

ABSTRACT. Rates of convergence of solutions of various two-dimensional α -regularization models, subject to periodic boundary conditions, toward solutions of the exact Navier-Stokes equations are given in the $L^{\infty}-L^2$ time-space norm, in terms of the regularization parameter α , when α approaches zero. Furthermore, as a paradigm, error estimates for the Galerkin approximation of the exact two-dimensional Leray- α model are also presented in the $L^{\infty}-L^2$ time-space norm. Simply by the triangle inequality, one can reach the error estimates of the solutions of Galerkin approximation of the α -regularization models toward the exact solutions of the Navier-Stokes equations in the two-dimensional periodic boundary conditions case.

MSC Classification: 35A30, 35Q35, 65M12, 65M15, 76D05.

Keywords: Leray- α Regularization, Navier-Stokes- α , Viscous Camassa-Holm, Modified Leray- α , Simplified Bardina Model, Lagrangian-Averaged-Navier-Stokes- α , Navier-Stokes Equations, Convergence Rate, Error Estimates, Galerkin Approximation.

1. INTRODUCTION

The closure problem of averaged quantities in turbulent flows has been, for many years, an outstanding open challenge for turbulence models. In the recent decade, various α -regularization models (Navier-Stokes- α , Leray- α , Modified Leray- α , Clark- α and simplified Bardina model) were introduced as efficient subgrid scale models of the Navier-Stokes equations (NSE) [2, 6, 10, 11, 12, 15, 18, 19, 25, 26, 27] (see also [37] for an analytical study of a mathematical generalization of the Navier-Stokes $-\alpha$ model). In particular, it was shown in some of these papers that these α -models fit remarkably well with empirical experimental data for a large range of huge Reynolds numbers. Moreover, these models were implemented numerically by various groups [3, 4, 12, 13, 20, 21, 22, 24, 25, 28, 31, 32, 33, 36]. Indeed, the authors of [30] have pushed further this numerical analysis point of view, concerning the α -models. in their study of the MHD- α model (see also [34, 35]). In fact, there have been extensive analytical studies on the global regularity of solutions and finite-dimensionality of global attractor of these models. however, there is much less work on the convergence, especially, the rate of convergence of solutions of various α -models toward the solutions of the exact NSE, when the regularization parameter α approaches zero. The authors of [19] study the convergence of the three-dimensional (3D) Navier-Stokes- α (NS- α) model to the 3D NSE. To be more specific, they show that there exists a subsequence of solutions u^{α_j} of the 3D NS- α that converges to one of the Leray-Hopf weak solutions of the 3D NSE, with periodic boundary conditions. Similar results are reported in [30] concerning the MHD- α model. Later the authors of [14, 45] show that the trajectory attractor of the Leray- α and Navier-Stokes- α , respectively, converges to the trajectory attractor of the 3D Navier-Stokes system, as α approaches zero. Since the uniqueness theorem for global weak solutions (or global existence of strong solutions) of the 3D NSE is not yet proved, the studies mentioned above either consider convergence to a weak solution or consider

Date: October 14, 2009.

To appear in: Numerical Functional Analysis and Optimization.

 $\mathbf{2}$

Y.CAO AND E.S. TITI

the convergence to the trajectory attractor of weak solutions. Recently the authors of [9] study the convergence rate of the Navier-Stokes- α model and obtain a mixed L^1-L^2 time-space norm for small initial data in Besov-type function spaces in which global existence and uniqueness of solutions for 3D NSE can be established with "small" initial data and external forcing. Similar results can also be derived by applying the same techniques for the other α -models under the assumption of existence of strong solutions of the 3D NSE, e.g., when the initial value and external forcing are small enough in the appropriate norms.

It is worth mentioning that inspired by the α -regularization models of turbulence similar regularization schemes were introduced and implemented in [7, 8], in the context of the nonlinear Schrödinger equation, and in [4] for the Leray- α regularization of the inviscid Burgers equation.

We mainly investigate in this paper the rates of convergence of four α -models (NS- α model, Leray- α model, Modified Leray- α model and simplified Bardina model) in the two-dimensional (2D) case, subject to periodic boundary conditions on the periodic box $[0, L]^2$. Since unique strong solution is proved to exist globally in time with any smooth enough initial data in the 2D case, we will show upper bounds, in terms of α , for the difference between solutions of the 2D α -models, u^{α} , and solutions of the 2D NSE system, u, in the L^2 -norm for any time interval [0, T]. Specifically, we show that all the four α -models we include in this study have the same order of convergence and error estimates, i.e., the L^2 -norms of the differences, $||u - u^{\alpha}||_2$, are of the order $O\left(\left(\frac{\alpha}{L}\right)\left(\log\left(\frac{L}{\alpha}\right)\right)^{1/2}\right)$, as $\frac{\alpha}{L}$ tends to zero. These results are presented in detail in Section 4. It is worth mentioning that the Brezis-Gallouet inequality plays an essential role in our error estimates in the 2D case; which in turn results in the logarithmic factor. This logarithmic factor, however, is absent in the 3D case which is treated in [9]. In a forthcoming paper, we will consider the rate of convergence of 3D Leray- α and NS- α models toward the adequate strong solution of the 3D exact NSE system, provided the latter exists, and compare the results to that of [9].

In Section 5, we consider the error estimates of the Galerkin approximation solutions in the 2D case, i.e., we estimate the difference between the solutions of the 2D α -model, u^{α} , and solutions of its corresponding finite-dimensional Galerkin approximation system, u_m^{α} , where $m \geq 1$ is the order of the truncation mode (dimension) of the Galerkin system. We will study, as an example, the Leray- α model and present the detailed proof of the error estimates. One can easily apply similar arguments to the other α -regularization models (NS- α model, Modified Leray- α model and simplified Bardina model) and obtain similar error estimates for these models. For the Leray- α model, the L^2 -norm of the difference, $||u^{\alpha} - u_m^{\alpha}||_2$, is of the order $O\left(\frac{1}{\lambda_{m+1}L^2}(\log(\lambda_{m+1}L^2))^{1/2}\right)$, under the assumption that α is small such that $\alpha^2 \leq \frac{1}{\lambda_{m+1}}$, where λ_{m+1} is the (m+1) - th eigenvalue of the Stokes operator in the 2D case. Applying the triangle inequality, we get error estimates concerning solutions of finite-dimensional Galerkin system of the 2D Leray- α model, u_m^{α} , as an approximation of the exact solution of the 2D NSE system, u. Specifically, we show that the error in the L^2 -norm of the difference between the solution u of the 2D NSE and u_m^{α} , the solution of the Leray- α Galerkin system, $||u - u_m^{\alpha}||_2$, is of the order $O\left(\frac{1}{\lambda_{m+1}L^2}(\log(\lambda_{m+1}L^2)^{1/2})\right)$, provided α is small enough satisfying $\alpha \leq \frac{2\pi}{\lambda_{m+1}L}$.

Before we present the main results of the error estimates, we will first introduce in Section 2 some notations and preliminaries that will be used throughout this paper. In Section 3 we will present all four α -models in functional setting and establish *a priori* estimates for the solutions, by investigating the finite-dimensional Galerkin systems then passing to the limit $m \to \infty$ by Aubin compactness theorem to obtain upper bounds for the exact solutions of the relevant models in certain norms. The main results for the rate of convergence, in terms of the regularization parameter α , of the α -models in the 2D cases will be presented in Section 4. After establishing the rate of convergence of solutions of α -regularization

models toward the solutions of the exact Navier-Stokes system, we will further show in section 5 the error estimates of difference between solutions of the 2D Leray- α model and solutions of its corresponding finite-dimensional Galerkin approximation system.

2. Functional Setting and Preliminaries

In this section, we introduce some preliminary background material following the usual notation used in the context of the mathematical theory of Navier-Stokes equations (NSE) (see, e.g., [16], [40], [41], [42]).

- (i) We denote by L^p and H^m the usual Lebesgue and Sobolev spaces, respectively. And we denote by $|\cdot|$ and (\cdot, \cdot) the L^2 -norm and L^2 -inner product, respectively.
- (ii) Let \mathcal{F} be the set of trigonometric polynomials of two variables with basic periodic domain $\Omega = [0, L]^2$ and spatial average zero, i.e., for every $\phi \in \mathcal{F}$, $\int_{\Omega} \phi(x) dx = 0$. We then set

$$\mathcal{V} = \left\{ \phi \in (\mathcal{F})^2 : \nabla \cdot \phi = 0 \right\}.$$

We denote by H and V the closures of \mathcal{V} in the $(L^2)^2$ and $(H^1)^2$ topologies, respectively. For $u \in H$ and $v \in V$, we denote by

$$|u|^{2} = \int_{\Omega} |u(x)|^{2} dx$$
 and $||v||^{2} = \int_{\Omega} |\nabla v(x)|^{2} dx$

the norms in H and V, respectively. We also note that by Rellich Lemma (see, e.g., [1]), we have that V is compactly embedded in H.

(iii) For any $s \ge 0$, we denote by \dot{H}^s the closure of \mathcal{F} with respect to the $H^s(\Omega)$ topology. Hence, for any $u \in \dot{H}^s$, we can write the Fourier expansion

$$u(x) = \sum_{k \in \mathbb{Z}_0^2} \hat{u}_k e^{i2\pi \frac{k \cdot x}{L}},$$

where the Fourier coefficients \hat{u}_k satisfy the reality condition $\hat{u}_{-k} = \hat{u}_k^*, \forall k \in \mathbb{Z}_0^2 := \mathbb{Z}^2 \setminus \{0, 0\}$. We define the norm on this space as

$$||u||_{\dot{H}^s}^2 = \left(\frac{2\pi}{L}\right)^{2(s-1)} \sum_{k \in \mathbb{Z}_0^2} (1+|k|^2)^s |\hat{u}_k|^2.$$

(iv) We denote by $P_{\sigma}: L^2 \to H$ the Helmholtz-Leray orthogonal projection operator, and by $A = -P_{\sigma}\Delta$ the Stokes operator, subject to periodic boundary conditions, with domain $D(A) = (H^2(\Omega))^2 \cap V$. We note that in the space-periodic case

$$Au = -P_{\sigma}\Delta u = -\Delta u, \quad \text{for all } u \in D(A).$$

The operator A^{-1} is a self-adjoint positive definite compact operator from H into H (see, e.g., [16], [41]). We denote by $0 < (\frac{2\pi}{L})^2 = \lambda_1 \leq \lambda_2 \leq \ldots$ the eigenvalues of A in the 2D case, repeated according to their multiplicities. It is well known that in two dimensions the eigenvalues of the operator A satisfy the Weyl's type formula (see e.g., [16], [42]), namely, there exists a dimensionless constant $c_0 > 0$ such that

$$\frac{j}{c_0} \le \frac{\lambda_j}{\lambda_1} \le c_0 j, \quad \text{for } j = 1, 2, \dots$$
 (1)

We also observe that in the periodic case, $D(A^{n/2}) = (H^n(\Omega))^2 \cap H$, for n > 0. In particular, one can show that $V = D(A^{1/2})$ (see, e.g., [16], [41]).

(v) For every $w \in V$, we have the Poincaré inequality

$$\lambda_1 |w|^2 \le ||w||^2. \tag{2}$$

Y.CAO AND E.S. TITI

Moreover, one can easily show that there is a dimensionless constant c > 0, such that

$$c|Aw| \le ||w||_{H^2} \le c^{-1}|Aw| \qquad \text{for every } w \in D(A), \tag{3}$$

and, by virtue of Poincaré inequality,

$$c|A^{1/2}w| \le ||w||_{H^1} \le c^{-1}|A^{1/2}w|$$
 for every $w \in V$. (4)

Hereafter, c will always denote a generic dimensionless constant.

- Notice that, thanks to (4), the norm of V is equivalent to the usual H^1 norm. (vi) Let $\{w_j\}_{j=1}^{\infty}$ be an orthonormal basis of H consisting of eigenfunctions of the operator A. Denote
- by $H_m = \operatorname{span}\{w_1, w_2, ..., w_m\}$, for $m \ge 1$ and let P_m be the L^2 -orthogonal projection from H onto H_m , then it is easy to see that

$$|(I - P_m)\phi|^2 \le \frac{1}{\lambda_{m+1}} ||\phi||^2, \quad \text{for all } \phi \in V,$$
(5)

$$\|(I - P_m)\phi\|^2 \le \frac{1}{\lambda_{m+1}} |A\phi|^2, \quad \text{for all } \phi \in D(A).$$
(6)

Moreover, one can also easily show that

$$AP_m\phi|^2 \le \lambda_m \|P_m\phi\|^2$$
, for all $\phi \in D(A)$. (7)

(vii) We recall the following 2D interpolation and Gagliardo-Nirenberg-Ladyzhenskaya inequality (see, e.g., [1], [16], [41]) :

$$\|\phi\|_{L^4} \le c \|\phi\|_{L^2}^{1/2} \|\phi\|_{H^1}^{1/2}.$$
(8)

(viii) For every $w_1, w_2 \in \mathcal{V}$, we define the bilinear operators

$$B(w_1, w_2) = P_{\sigma}((w_1 \cdot \nabla)w_2), \tag{9}$$

$$\tilde{B}(w_1, w_2) = -P_{\sigma}(w_1 \times (\nabla \times w_2)).$$
(10)

In the following lemma, we will list certain relevant inequalities and properties of B (see, e.g., [16], [41], [42]) and of \tilde{B} (see [19]).

Lemma 1. The bilinear operator B defined in (9) satisfies the following:

(i) B can be extended as a continuous bilinear map $B: V \times V \to V'$, where V' is the dual space of V.

In particular, the bilinear operator B satisfies the following inequalities:

$$\langle B(u,v), w \rangle_{V'} \leq c |u|^{\frac{1}{2}} ||u||^{\frac{1}{2}} ||v|| ||w||^{\frac{1}{2}} ||w||^{\frac{1}{2}}, \text{ for all } u, v, w \in V,$$

$$(11)$$

$$(B(u,v),w) \le c \|u\|_{\infty} \|v\| \|w|, \quad for \ all \ u \in D(A), \ v \in V, \ w \in H,$$
(12)

$$|(B(u,v),w)| \le c|u| \|\nabla v\|_{\infty} |w|, \quad \text{for all } u \in H, \ v \in D(A^{3/2}), \ w \in H,$$
(13)

$$|\langle B(u,v), w \rangle_{(D(A))'}| \le c|u| \|v\| \|w\|_{\infty}, \text{ for all } u \in H, v \in V, w \in D(A).$$
(14)

Moreover, for every $w_1, w_2, w_3 \in V$, we have

$$\langle B(w_1, w_2), w_3 \rangle_{V'} = -\langle B(w_1, w_3), w_2 \rangle_{V'},$$
(15)

and in particular,

$$\langle B(w_1, w_2), w_2 \rangle_{V'} = 0.$$
 (16)

(ii) In the 2D periodic boundary condition case, we have

$$(B(\varphi,\varphi),A\varphi) = 0, \tag{17}$$

for every $\varphi \in D(A)$.

(iii) \tilde{B} can also be extended as a continuous bilinear map $\tilde{B}: V \times V \to V'$. Furthermore, for every $w_1, w_2, w_3 \in V$, the bilinear operator \tilde{B} satisfies the following inequality:

$$|\langle \tilde{B}(w_1, w_2), w_3 \rangle_{V'}| \le c |w_1|^{1/2} ||w_1||^{1/2} ||w_2|| |w_3|^{1/2} ||w_3||^{1/2}$$

Moreover, for every $w_1, w_2, w_3 \in V$, we have

$$\langle \tilde{B}(w_1, w_2), w_3 \rangle_{V'} = \langle B(w_1, w_2), w_3 \rangle_{V'} - \langle B(w_3, w_2), w_1 \rangle_{V'},$$
(18)

and consequently,

$$\langle \dot{B}(w_1, w_2), w_1 \rangle_{V'} = 0.$$
 (19)

Lemma 2. For every $u \in D(A)$ and $w \in V$, we have

 $|(B(w, u), Au)| \le c ||w|| ||u|| |Au|.$

Proof. Let $w \in \mathcal{V}$ and $u \in D(A)$. Then

$$(B(w,u),Au) = -\sum_{l,j,m=1}^{2} \int_{\Omega} w_{j} (\frac{\partial}{\partial x_{j}} u_{l}) (\frac{\partial^{2}}{\partial x_{m}^{2}} u_{l}) dx$$

$$= \sum_{l,j,m=1}^{2} \left[\int_{\Omega} (\frac{\partial}{\partial x_{m}} w_{j}) (\frac{\partial}{\partial x_{j}} u_{l}) (\frac{\partial}{\partial x_{m}} u_{l}) dx + \int_{\Omega} w_{j} (\frac{\partial}{\partial x_{j}} \frac{\partial}{\partial x_{m}} u_{l}) \frac{\partial}{\partial x_{m}} u_{l}) dx \right].$$

By relation (16) the second term on the right-hand side above is zero. Therefore, by the Cauchy-Schwarz inequality

$$|(B(w,u),Au)| \le c ||w|| ||\nabla u||_{L^4}^2,$$

and by (8), (3) and (4) we have

$$(B(w, u), Au)| \le c ||w|| ||u|| |Au|.$$

From the above and the density of \mathcal{V} in V we conclude our lemma.

Next, we state a two-dimensional periodic boundary condition version of the well-known Brezis-Gallouet inequality [5]. For the sake of completeness, we will present the proof of this version in the Appendix.

Proposition 3. There exists a scale invariant constant c > 0 such that for every $\varphi \in D(A)$,

$$\|\varphi\|_{L^{\infty}} \le c \|\varphi\| \left(1 + \log\left(\frac{L}{2\pi} \frac{|A\varphi|}{\|\varphi\|}\right)\right)^{1/2}.$$

Applying the Brezis-Gallouet inequality above to (12) and (14), we have the following corollary (see also [43] for similar logarithmic inequalities concerning the bilinear term).

Corollary 4. In the two-dimensional case, the bilinear operator B satisfies the following inequalities:

$$|(B(u,v),w)| \le c ||u|| ||v|| ||w| \left(1 + \log \frac{|Au|^2}{||u||^2 \lambda_1}\right)^{1/2} \text{ for all } u \in D(A), \ v \in V, \ w \in H,$$

$$(20)$$

$$|(B(u,v),w)| \le c|u| ||v|| ||w|| \left(1 + \log \frac{|Aw|^2}{||w||^2 \lambda_1}\right)^{1/2}, \text{ for all } u \in H, v \in V, w \in D(A).$$

$$(21)$$

where $\lambda_1 = (\frac{2\pi}{L})^2$ is the first eigenvalue of the Stokes operator A.

3. A Priori Estimates of the 2D Navier-Stokes Equations and the α -Regularization Models

In this section, we will establish a priori estimates for solutions of the 2D NSE and the 2D α -regularization models. These results will be useful for the error estimates in the next two sections.

3.1. *A priori* estimates for the 2D Navier-Stokes equations. We recall that the two dimensional (2D) NSE is equivalent (see, e.g., [16], [41]) to the functional evolution equation in the Hilbert space H

$$\frac{du}{dt} + \nu Au + B(u, u) = f,$$

$$u(0) = u_0.$$
(22)

The corresponding Galerkin system of the 2D NSE is given below as a system of ordinary differential equations in the space H_m , defined in Section 2:

$$\frac{du_m}{dt} + \nu A u_m + P_m B(u_m, u_m) = f_m,$$
(23)
$$u_m(0) = u_{0m}.$$

where $u_{0m} = P_m u_0$, $f_m = P_m f$.

The proof of existence and uniqueness of solutions of the above 2D NSE system (22), subject to periodic boundary conditions, can be established by applying Galerkin approximation procedure [16, 41, 42]. The idea is to establish *a priori* estimates for the solutions of the finite-dimensional system (23) and then by applying Aubin compactness theorem (see also [16, 41, 42]) one can extract a subsequence that converges to the unique solution of the NSE system (22). The details of the proof are textbook material and will not be presented here. In the following proposition we will establish *a priori* estimates for solutions of the Galerkin system (23) that we will use later.

Proposition 5. Let $u_0 \in H, T > 0$ and $f \in L^2((0,T); H)$. Let u_m be the solution of the Galerkin approximation of the 2D NSE, system (23), over the interval [0,T] with initial data u_{0m} for a given $m \ge 1$, then

$$\sup_{0 \le t \le T} \left(|u_m(t)|^2 + \nu \int_0^t ||u_m(s)||^2 \, ds \right) \le K_0^2(\nu, u_0, f, L, T), \tag{24}$$

where $K_0^2 := |u_0|^2 + \frac{1}{\nu\lambda_1} \int_0^T |f(s)|^2 ds$ and $\lambda_1 = (\frac{2\pi}{L})^2$.

Proof. We take the inner product of the first equation in (23) with u_m to obtain

$$\frac{1}{2}\frac{d}{dt}|u_m|^2 + \nu ||u_m||^2 = (P_m f, u_m) \le |f||u_m|.$$

By Poincaré inequality (2) we have

$$\frac{1}{2}\frac{d}{dt}|u_m|^2 + \nu ||u_m||^2 \le \frac{|f|||u_m||}{\lambda_1^{1/2}} \le \frac{|f|^2}{2\nu\lambda_1} + \frac{\nu}{2}||u_m||^2.$$

Hence,

$$\frac{d}{dt}|u_m|^2 + \nu ||u_m||^2 \le \frac{|f|^2}{\nu\lambda_1}$$

and by integration we have

$$|u_m(t)|^2 + \nu \int_0^t ||u_m(s)||^2 \, ds \le |u_0|^2 + \frac{1}{\nu\lambda_1} \int_0^t |f(s)|^2 \, ds \quad \text{for all } t \in [0,T],$$

which concludes the proof.

Remark 1. One needs to further establish estimates on $\frac{du_m}{dt}$ and then, thanks to Aubin compactness theorem [16, 41], one can extract a subsequence $u_{m'}$, that converges to the unique solution u of system (22), as $m' \to \infty$. Following the standard procedure as in [16, 41], one can show that the solution of the

NSE (22) satisfies the same *a priori* estimates, namely, let *u* be the solution of the system (22) over the interval [0, T] with initial data $u_0 \in H$, then

$$\sup_{0 \le t \le T} \left(|u(t)|^2 + \nu \int_0^t ||u(s)||^2 \, ds \right) \le K_0^2(\nu, u_0, f, L, T).$$
(25)

3.2. A priori estimates for the 2D Leray- α model. In this subsection, we will establish a priori estimates for solutions of the 2D Leray- α regularization model. The Leray- α model was introduced and analyzed in [15] and implemented computationally [15, 20, 21, 22, 31] in the context of subgrid scale models of 3D turbulence (see also [4, 24] and [31] for 2D computations with the Leray- α model). The Leray- α model was inspired by the Navier-Stokes- α model (also known as viscous Camassa-Holm or Lagrangian-averaged-Navier-Stokes- α model) [10, 11, 12, 18, 19], and it happened to fit as a member of the general family of regularizations introduced in the seminal work of Leray [29] in the context of establishing the existence of solutions for the 2D and 3D NSE.

The Leray- α regularization model of the NSE is given by the following functional evolution system in the space H:

$$\frac{dv^{\alpha}}{dt} + \nu A v^{\alpha} + B(u^{\alpha}, v^{\alpha}) = f, \qquad (26)$$
$$u^{\alpha} + \alpha^{2} A u^{\alpha} = v^{\alpha},$$
$$u^{\alpha}(0) = u_{0}.$$

Observe that when the regularization parameter $\alpha = 0$ one recovers the exact NSE system (22).

The corresponding Galerkin system of the 2D Leray- α model is given below as a system of ordinary differential equations in the space H_m :

$$\frac{dv_m^{\alpha}}{dt} + \nu A v_m^{\alpha} + P_m B(u_m^{\alpha}, v_m^{\alpha}) = f_m,$$

$$u_m^{\alpha} + \alpha^2 A u_m^{\alpha} = v_m^{\alpha},$$

$$u_m^{\alpha}(0) = u_{0m},$$
(27)

where $u_{0m} = P_m u_0, f_m = P_m f.$

The proof of existence and uniqueness of solution of the above Leray- α system (26) and the other three α -models we cover in this section, subject to periodic boundary conditions, can be established by applying Galerkin approximation procedure. One can follow similar steps as those for the NSE [16, 41, 42] to show the proof. The proof is not the heart of this paper and we will omit the details and only establish *a priori* estimates for solutions of the Galerkin system (27). To be concise and focus on the essential matter of this paper, in this and the following subsections of *a priori* estimates for the 2D α -models, we will simply skip the details of the proof and will not restate these comments. Interested readers can refer to the relevant literature in [16, 41, 42] and references therein to fill in the gap.

Proposition 6. Let $u_0 \in D(A), T > 0$ and $f \in L^2((0,T); H)$. Let u_m^{α} be the solution of the system (27) over the interval [0,T] with initial data u_{0m} for a given $m \ge 1$, then

$$\sup_{0 \le t \le T} \left[\|u_m^{\alpha}(t)\|^2 + \alpha^2 |Au_m^{\alpha}(t)|^2 + \nu \int_0^t (|Au_m^{\alpha}(s)|^2 + \alpha^2 |A^{3/2}u_m^{\alpha}(s)|^2) ds \right] \le \tilde{K}_0^2$$

where $\tilde{K}_0^2 := \|u_0\|^2 + \alpha^2 |Au_0|^2 + \frac{1}{\nu} \int_0^T |f(s)|^2 ds.$

Y.CAO AND E.S. TITI

Proof. We take the inner product of the first equation in system (27) with Au_m^{α} to obtain

$$\frac{1}{2}\frac{d}{dt}(\|u_m^{\alpha}\|^2 + \alpha^2 |Au_m^{\alpha}|^2) + \nu(|Au_m^{\alpha}|^2 + \alpha^2 |A^{3/2}u_m^{\alpha}|^2) + (B(u_m^{\alpha}, v_m^{\alpha}), Au_m^{\alpha}) = (f, Au_m^{\alpha}).$$

By (16) and (17) one has

$$(B(u_m^{\alpha}, v_m^{\alpha}), Au_m^{\alpha}) = (B(u_m^{\alpha}, u_m^{\alpha}), Au_m^{\alpha}) + \alpha^2 (B(u_m^{\alpha}, Au_m^{\alpha}), Au_m^{\alpha}) = 0.$$

Consequently, from the above we have

$$\frac{1}{2}\frac{d}{dt}(\|u_m^{\alpha}\|^2 + \alpha^2 |Au_m^{\alpha}|^2) + \nu(|Au_m^{\alpha}|^2 + \alpha^2 |A^{3/2}u_m^{\alpha}|^2) = (f, Au_m^{\alpha}) \le |f||Au_m^{\alpha}| \le \frac{|f|^2}{2\nu} + \frac{\nu}{2} |Au_m^{\alpha}|^2.$$

Hence

$$\frac{d}{dt}(\|u_m^{\alpha}\|^2 + \alpha^2 |Au_m^{\alpha}|^2) + \nu(|Au_m^{\alpha}|^2 + \alpha^2 |A^{3/2}u_m^{\alpha}|^2) \le \frac{|f|^2}{\nu}$$

and by integration over the interval (0, t), for $t \in [0, T]$ we have

$$\begin{aligned} \|u_m^{\alpha}(t)\|^2 + \alpha^2 |Au_m^{\alpha}(t)|^2 &+ \nu \int_0^t (|Au_m^{\alpha}(s)|^2 + \alpha^2 |A^{3/2}u_m^{\alpha}(s)|^2) \, ds \\ &\leq \frac{1}{\nu} \int_0^T |f(s)|^2 \, ds + \|u_{0m}\|^2 + \alpha^2 |Au_{0m}| \\ &\leq \frac{1}{\nu} \int_0^T |f(s)|^2 \, ds + \|u_0\|^2 + \alpha^2 |Au_0|, \end{aligned}$$

which concludes our proof.

Remark 2. Similar to the 2D NSE, one needs to further establish estimates on $\frac{du_m^{\alpha}}{dt}$ and then, thanks to Aubin compactness theorem [16, 41], one can extract a subsequence $u_{m'}^{\alpha}$ that converges to the unique solution u^{α} of system (26), as $m' \to \infty$. As a result, one can also prove that the solution of the Leray- α system (26) satisfies the same *a priori* estimates as in Proposition 6, namely, let u^{α} be the solution of system (26) over the interval [0, T] with initial data $u_0 \in D(A)$, then

$$\sup_{0 \le t \le T} \left[\|u^{\alpha}(t)\|^2 + \alpha^2 |Au^{\alpha}(t)|^2 + \nu \int_0^t (|Au^{\alpha}(s)|^2 + \alpha^2 |A^{3/2}u^{\alpha}(s)|^2) ds \right] \le \tilde{K}_0^2.$$
(28)

We emphasize here again that the details of the proof of the above results are omitted. In the following subsections of *a priori* estimates of the NS- α , Modified Leray- α and simplified Bardina models, we will also skip the details and will not restate the remark.

3.3. A priori estimates for the 2D NS- α model. In this subsection, we will establish a priori estimates for the 2D NS- α regularization model of the 2D NSE. The NS- α model (also known as the viscous Camassa-Holm or Lagragian-averaged-Navier-Stokes- α model) was introduced and analyzed in [10, 11, 12, 18, 19, 25], which was also the first of the family of the α -models. In addition to the remarkable match of explicit solutions of the 3D NS- α model to experimental data, in the channels and pipes, for a wide range of huge Reynold numbers [10, 11, 12], the validity of NS- α model as a subgrid scale turbulence model was tested numerically in [12, 13, 22, 24, 32, 33, 36].

The NS- α regularization model of the NSE is given by the following functional evolution system in

the space H:

$$\frac{dv^{\alpha}}{dt} + \nu Av^{\alpha} + \tilde{B}(u^{\alpha}, v^{\alpha}) = f,$$

$$u^{\alpha} + \alpha^{2}Au^{\alpha} = v^{\alpha},$$

$$u^{\alpha}(0) = u_{0},$$
(29)

where \tilde{B} is given by (10) and Lemma 1.

Observe that when the regularization parameter $\alpha = 0$ one recovers the exact NSE system (22).

The corresponding Galerkin approximation system of the 2D NS- α model is given below as a system of ordinary differential equations in the space H_m :

$$\frac{dv_m^{\alpha}}{dt} + \nu A v_m^{\alpha} + P_m \tilde{B}(u_m^{\alpha}, v_m^{\alpha}) = f_m,$$

$$u_m^{\alpha} + \alpha^2 A u_m^{\alpha} = v_m^{\alpha},$$

$$u_m^{\alpha}(0) = u_{0m},$$
(30)

where $u_{0m} = P_m u_0, f_m = P_m f.$

The proof of existence and uniqueness of solutions of the NS- α system (29), subject to periodic boundary conditions, is similar to that of the 3D case as it is presented in [19]. In the next proposition we will omit the details and only show the *a priori* estimates for solutions of system (30).

Proposition 7. Let $u_0 \in D(A), T > 0$ and $f \in L^2((0,T); H)$. Let u_m^{α} be the solution of the Galerkin system of 2D NS- α (30) over the interval [0,T] with initial data u_{0m} for a given $m \ge 1$, then

$$\sup_{0 \le t \le T} \left[|u_m^{\alpha}(t)|^2 + \alpha^2 ||u_m^{\alpha}(t)||^2 + \nu \int_0^t (||u_m^{\alpha}(s)||^2 + \alpha^2 |Au_m^{\alpha}(s)|^2) \, ds \right] \le \tilde{\tilde{K}}_{01}^2, \tag{31}$$

where $\tilde{\tilde{K}}_{01}^2 := |u_0|^2 + \alpha^2 ||u_0||^2 + \frac{1}{\nu\lambda_1} \int_0^T |f(s)|^2 ds$, which is independent of m, and $\lambda_1 = (\frac{2\pi}{L})^2$. Moreover,

$$\sup_{0 \le t \le T} \left[\|u_m^{\alpha}(t)\|^2 + \alpha^2 |Au_m^{\alpha}(t)|^2 + \nu \int_0^t (|Au_m^{\alpha}(s)|^2 + \alpha^2 |A^{3/2}u_m^{\alpha}(s)|^2) \, ds \right] \le \tilde{\tilde{K}}_{02}^2, \tag{32}$$

where $\tilde{\tilde{K}}_{02}^2 := \|u_0\|^2 + \alpha^2 |Au_0|^2 + \frac{1}{\nu} \int_0^T |f(s)|^2 ds + \frac{c}{\nu^2} \tilde{\tilde{K}}_{00}^2 \tilde{\tilde{K}}_{01}^2$, and $\tilde{\tilde{K}}_{00}$ is a constant depending on $|Au_0|, \nu, T, f$ and $\tilde{\tilde{K}}_{01}$, which is give explicitly in (36).

Proof. By taking the inner product of the first equation in (30) with u_m^{α} and using (19) we obtain

$$\frac{1}{2}\frac{d}{dt}(|u_m^{\alpha}|^2 + \alpha^2 ||u_m^{\alpha}||^2) + \nu(||u_m^{\alpha}||^2 + \alpha^2 |Au_m^{\alpha}|^2) = (P_m f, u_m^{\alpha}) \le |f| ||u_m^{\alpha}|| \le \frac{|f| ||u_m^{\alpha}||}{\lambda_1^{1/2}} \le \frac{|f|^2}{2\nu\lambda_1} + \frac{\nu ||u_m^{\alpha}||^2}{2},$$

where in the last two steps we used Poincaré inequality (2) and Young's inequality. As a result we have

$$\frac{d}{dt}(|u_m^{\alpha}|^2 + \alpha^2 ||u_m^{\alpha}||^2) + \nu(||u_m^{\alpha}||^2 + \alpha^2 |Au_m^{\alpha}|^2) \le \frac{|f|^2}{\nu\lambda_1}$$

Integrating the above in time we obtain (31).

Next, we prove (32). We take the inner product of the equation in (30) with Au_m^{α} and follow similar steps as above to obtain

$$\frac{1}{2}\frac{d}{dt}(\|u_m^{\alpha}\|^2 + \alpha^2 |Au_m^{\alpha}|^2) + \nu(|Au_m^{\alpha}|^2 + \alpha^2 |A^{3/2}u_m^{\alpha}|^2) + (\tilde{B}(u_m^{\alpha}, v_m^{\alpha}), Au_m^{\alpha}) \\
= (P_m f, Au_m^{\alpha}) \le \frac{|f|^2}{2\nu} + \frac{\nu}{2} |Au_m^{\alpha}|^2.$$
(33)

Observe that

$$(\tilde{B}(u_m^{\alpha}, v_m^{\alpha}), Au_m^{\alpha}) = (B(u_m^{\alpha}, v_m^{\alpha}), Au_m^{\alpha}) - (B(Au_m^{\alpha}, v_m^{\alpha}), u_m^{\alpha}) = \alpha^2 (B(Au_m^{\alpha}, u_m^{\alpha}), Au_m^{\alpha}),$$
(34)

where we use in the above (18), (15), (16), (17) and the relation $v_m^{\alpha} = u_m^{\alpha} + \alpha^2 A u_m^{\alpha}$. From (34) and Lemma 2 we have

$$|(\tilde{B}(u_m^{\alpha}, v_m^{\alpha}), Au_m^{\alpha})| \le c\alpha^2 |A^{3/2}u_m^{\alpha}| \|u_m^{\alpha}\| |Au_m^{\alpha}| \le \frac{\nu}{2}\alpha^2 |A^{3/2}u_m^{\alpha}|^2 + \frac{c\alpha^2}{\nu} \|u_m^{\alpha}\|^2 |Au_m^{\alpha}|^2.$$

From the above and (33) we obtain

$$\frac{d}{dt}(\|u_{m}^{\alpha}\|^{2} + \alpha^{2}|Au_{m}^{\alpha}|^{2}) + \nu(|Au_{m}^{\alpha}|^{2} + \alpha^{2}|A^{3/2}u_{m}^{\alpha}|^{2}) \\
\leq \frac{|f|^{2}}{\nu} + \frac{c\alpha^{2}}{\nu}\|u_{m}^{\alpha}\|^{2}|Au_{m}^{\alpha}|^{2} \\
\leq \frac{|f|^{2}}{\nu} + \frac{c}{\nu}(\|u_{m}^{\alpha}\|^{2} + \alpha^{2}|Au_{m}^{\alpha}|^{2})^{2}.$$
(35)

Therefore, by Gronwall's inequality we have

$$\begin{aligned} \|u_{m}^{\alpha}(t)\|^{2} + \alpha^{2} |Au_{m}^{\alpha}(t)|^{2} &\leq e^{\frac{c}{\nu} \int_{0}^{t} (\|u_{m}^{\alpha}(s)\|^{2} + \alpha^{2} |Au_{m}^{\alpha}(s)|^{2}) ds} (\|u_{0}\|^{2} + \alpha^{2} |Au_{0}|^{2}) \\ &+ \frac{1}{\nu} \int_{0}^{t} e^{\frac{c}{\nu} \int_{\sigma}^{t} (\|u_{m}^{\alpha}(s)\|^{2} + \alpha^{2} |Au_{m}^{\alpha}(s)|^{2}) ds} |f(\sigma)|^{2} d\sigma, \end{aligned}$$

and by (31) we obtain

$$\|u_m^{\alpha}(t)\|^2 + \alpha^2 |A_m^{\alpha}u(t)|^2 \le e^{\frac{c}{\nu^2}\tilde{\tilde{K}}_{01}^2} (\|u_0\|^2 + \alpha^2 |Au_0|^2) + \frac{1}{\nu} e^{\frac{c}{\nu^2}\tilde{\tilde{K}}_{01}^2} \int_0^T |f(\sigma)|^2 \, d\sigma =: \tilde{\tilde{K}}_{00}^2. \tag{36}$$

Integrate (35) and using (31) and (36), we obtain

$$\begin{aligned} \|u_m^{\alpha}(t)\|^2 + \alpha^2 |Au_m^{\alpha}(t)|^2 &+ \nu \int_0^t (|Au_m^{\alpha}(s)|^2 + \alpha^2 |A^{3/2}u_m^{\alpha}(s)|^2) \, ds \\ &\leq \|u_0\|^2 + \alpha^2 |Au_0|^2 + \frac{1}{\nu} \int_0^t |f|^2 \, dt + \frac{c}{\nu^2} \tilde{K}_{00}^2 \tilde{K}_{01}^2, \end{aligned}$$

ludes the proof.

which concludes the proof.

Remark 3. By following similar steps as those of NSE [16, 41, 42], one can show that the exact solutions of the 2D NS- α system have the same *a priori* estimates as those of the solutions of the Galerkin system, namely, let u^{α} be the solution of the system (29) over the interval [0,T] with initial data $u_0 \in D(A)$, then

$$\sup_{0 \le t \le T} \left[|u^{\alpha}(t)|^2 + \alpha^2 ||u^{\alpha}(t)||^2 + \nu \int_0^t (||u^{\alpha}(s)||^2 + \alpha^2 |Au^{\alpha}(s)|^2) \, ds \right] \le \tilde{\tilde{K}}_{01}^2, \tag{37}$$

$$\sup_{0 \le t \le T} \left[\|u^{\alpha}(t)\|^2 + \alpha^2 |Au^{\alpha}(t)|^2 + \nu \int_0^t (|Au^{\alpha}(s)|^2 + \alpha^2 |A^{3/2}u^{\alpha}(s)|^2) \, ds \right] \le \tilde{\tilde{K}}_{02}^2. \tag{38}$$

3.4. A priori estimates for the 2D Modified Leray- α model. In this subsection, we will establish a priori estimates for the 2D Modified Leray- α (ML- α) regularization model of the 2D NSE. Inspired by the NS- α and Leray- α models, the ML- α was introduced and analyzed in [26] and is tested numerically in [22] in the context of 3D sub-grid α -models of turbulence.

The ML- α regularization model of the 2D NSE is given by the following functional evolution system in the space H:

$$\frac{dv^{\alpha}}{dt} + \nu A v^{\alpha} + B(v^{\alpha}, u^{\alpha}) = f, \qquad (39)$$
$$u^{\alpha} + \alpha^{2} A u^{\alpha} = v^{\alpha},$$
$$u^{\alpha}(0) = u_{0}.$$

Observe that when the regularization parameter $\alpha = 0$ one again recovers the exact 2D NSE system (22).

The corresponding Galerkin approximation system of the 2D ML- α model is given below as a system of ordinary differential equations in the space H_m :

$$\frac{dv_m^{\alpha}}{dt} + \nu A v_m^{\alpha} + P_m B(v_m^{\alpha}, u_m^{\alpha}) = f_m,$$

$$u_m^{\alpha} + \alpha^2 A u_m^{\alpha} = v_m^{\alpha},$$

$$u_m^{\alpha}(0) = u_{0m},$$
(40)

where $u_{0m} = P_m u_0, f_m = P_m f$.

Following similar steps as in the 3D case in [26] one can establish the global existence and uniqueness for the system (39) in the case of 2D periodic boundary conditions. In the next proposition we will present *a priori* estimates for the solutions of the Galerkin system (40).

Proposition 8. Let $u_0 \in D(A), T > 0$ and $f \in L^2((0,T); H)$. Let u_m^{α} be the solution of the Galerkin system of 2D ML- α (40) over the interval [0,T] with initial data u_{0m} for a given $m \ge 1$, then

$$\sup_{0 \le t \le T} \left[|u_m^{\alpha}(t)|^2 + \alpha^2 ||u_m^{\alpha}(t)||^2 + \nu \int_0^t (||u_m^{\alpha}(s)||^2 + \alpha^2 |Au_m^{\alpha}(s)|^2) \, ds \right] \le \tilde{\tilde{K}}_{01}^2. \tag{41}$$

Moreover,

$$\sup_{\substack{0 \le t \le T\\z}} \left[\|u_m^{\alpha}(t)\|^2 + \alpha^2 |Au_m^{\alpha}(t)|^2 + \nu \int_0^t (|Au_m^{\alpha}(s)|^2 + \alpha^2 |A^{3/2}u_m^{\alpha}(s)|^2) \, ds \right] \le \tilde{\tilde{K}}_{02}^2, \tag{42}$$

where K_{01} and K_{02} are the same as in Proposition 7.

Proof. Let us take the inner product of the equation in (40) with u_m^{α} to obtain

$$\begin{aligned} \frac{1}{2} \frac{d}{dt} (|u_m^{\alpha}|^2 + \alpha^2 ||u_m^{\alpha}||^2) + \nu (||u_m^{\alpha}||^2 + \alpha^2 |Au_m^{\alpha}|^2) &= (P_m f, u_m^{\alpha}) \le |f| |u_m^{\alpha}| \le |f| \frac{||u_m^{\alpha}||}{\lambda_1^{1/2}} \\ &\le \frac{|f|^2}{2\nu\lambda_1} + \frac{\nu}{2} ||u_m^{\alpha}||^2, \end{aligned}$$

thus we get

$$\frac{d}{dt}(|u_m^{\alpha}|^2 + \alpha^2 ||u_m^{\alpha}||^2) + \nu(||u_m^{\alpha}||^2 + \alpha^2 |Au_m^{\alpha}|^2) \le \frac{|f|^2}{\nu\lambda_1}.$$
(43)

Integrating the above over the interval (0, t), we obtain

$$|u_m^{\alpha}(t)|^2 + \alpha^2 ||u_m^{\alpha}(t)||^2 + \nu \int_0^t (||u_m^{\alpha}(s)||^2 + \alpha^2 |Au_m^{\alpha}(s)|^2) \, ds \, \le \int_0^t \frac{|f(s)|^2}{\nu \lambda_1} \, ds + |u_0|^2 + \alpha^2 ||u_0||^2,$$

which implies (41).

1 .1

Next, we take the inner product of the equation in (40) with Au_m^{α} to obtain

$$\frac{1}{2} \frac{u}{dt} (\|u_m^{\alpha}\|^2 + \alpha^2 |Au_m^{\alpha}|^2) + \nu (|Au_m^{\alpha}|^2 + \alpha^2 |A^{3/2}u_m^{\alpha}|^2) \\
= -(B(v_m^{\alpha}, u_m^{\alpha}), Au_m^{\alpha}) + (P_m f, Au_m^{\alpha}) \\
\leq |(B(v_m^{\alpha}, u_m^{\alpha}), Au_m^{\alpha})| + \frac{|f|^2}{2\nu} + \frac{\nu}{2} |Au_m^{\alpha}|^2.$$
(44)

Next we estimate

 $|B(v_m^{\alpha}, u_m^{\alpha}), Au_m^{\alpha})| = |(B(u_m^{\alpha}, u_m^{\alpha}), Au_m^{\alpha}) + \alpha^2 (B(Au_m^{\alpha}, u_m^{\alpha}), Au_m^{\alpha})|.$ By (17) and Proposition 7 we have

$$\begin{aligned} |(B(v_m^{\alpha}, u_m^{\alpha}), Au_m^{\alpha})| &= \alpha^2 |(B(Au_m^{\alpha}, u_m^{\alpha}), Au_m^{\alpha})| \le c\alpha^2 |A^{3/2} u_m^{\alpha}| ||u_m^{\alpha}|| |Au_m^{\alpha}|| \\ &\le \frac{\nu}{2} \alpha^2 |A^{3/2} u_m^{\alpha}|^2 + \frac{c}{2\nu} ||u_m^{\alpha}||^2 (\alpha^2 |Au_m^{\alpha}|^2) \\ &\le \frac{\nu}{2} \alpha^2 |A^{3/2} u_m^{\alpha}|^2 + \frac{c}{2\nu} (||u_m^{\alpha}||^2 + \alpha^2 |Au_m^{\alpha}|^2)^2. \end{aligned}$$
(45)

From above we have

$$\frac{d}{dt}(\|u_m^{\alpha}\|^2 + \alpha^2 |Au_m^{\alpha}|^2) + \nu(|Au_m^{\alpha}|^2 + \alpha^2 |A^{3/2}u_m^{\alpha}|^2) \le \frac{|f|^2}{\nu} + \frac{c}{\nu}(\|u_m^{\alpha}\|^2 + \alpha^2 |Au_m^{\alpha}|^2)^2,$$

which is exactly the same inequality as (35) (as in the proof of Proposition 7). Following the same steps we obtain (42). \Box

Remark 4. Similarly, the solutions of the exact ML- α system have the same *a priori* estimates as those of the solutions of the Galerkin system, namely, let u^{α} be the solution of the system (39) over the interval [0, T] with initial data $u_0 \in D(A)$, then

$$\sup_{0 \le t \le T} \left[|u^{\alpha}(t)|^2 + \alpha^2 ||u^{\alpha}(t)||^2 + \nu \int_0^t (||u^{\alpha}(s)||^2 + \alpha^2 |Au^{\alpha}(s)|^2) \, ds \right] \le \tilde{\tilde{K}}_{01}^2. \tag{46}$$

Moreover,

$$\sup_{\substack{0 \le t \le T\\ \tilde{\kappa}}} \left[\|u^{\alpha}(t)\|^{2} + \alpha^{2} |Au^{\alpha}(t)|^{2} + \nu \int_{0}^{t} (|Au^{\alpha}(s)|^{2} + \alpha^{2} |A^{3/2}u^{\alpha}(s)|^{2}) \, ds \right] \le \tilde{\tilde{K}}_{02}^{2},\tag{47}$$

where K_{01} and K_{02} are the same as in Proposition 7.

3.5. A priori estimates for the 2D simplified Bardina model. In this subsection, we will establish a priori estimates for the 2D simplified Bardina model for the 2D NSE. The Bardina closure model of turbulence was introduced first by Bardina et al [2] and later simplified and studied further in [27] and in [6]. In particular, global well-posedness for the 3D simplified Bardina model was established in [27]. In [6] these results were slightly improved and the long-time behavior was investigated. Further, it was established in [6] the global regularity of the 3D inviscid version of the simplified Bardina, a property which is still out of reach for the other α -models of turbulence.

The simplified Bardina regularization model of 2D NSE is given by the following functional evolution system in the space H:

$$\frac{dv^{\alpha}}{dt} + \nu A v^{\alpha} + B(u^{\alpha}, u^{\alpha}) = f,$$

$$u^{\alpha} + \alpha^{2} A u^{\alpha} = v^{\alpha},$$

$$u^{\alpha}(0) = u_{0}.$$
(48)

Observe again that when the regularization parameter $\alpha = 0$ one recovers the exact 2D NSE system (22).

The corresponding Galerkin approximation system of the 2D simplified Bardina model is given below as a system of ordinary differential equations in space H_m :

$$\frac{dv_m^{\alpha}}{dt} + \nu A v_m^{\alpha} + P_m B(u_m^{\alpha}, u_m^{\alpha}) = f_m,$$

$$u_m^{\alpha} + \alpha^2 A u_m^{\alpha} = v_m^{\alpha},$$

$$u_m^{\alpha}(0) = u_{0m},$$
(49)

where $u_{0m} = P_m u_0, f_m = P_m f$.

In the following proposition we will establish a prior estimates for the solutions of the Galerkin system (49).

Proposition 9. Let $u_0 \in D(A), T > 0$ and $f \in L^2((0,T); H)$. Let u_m^{α} be the solution of the system (49) over the interval [0,T] with initial data u_{0m} for a given $m \ge 1$, then

$$\sup_{0 \le t \le T} \left[\|u_m^{\alpha}(t)\|^2 + \alpha^2 |Au_m^{\alpha}(t)|^2 + \nu \int_0^t (|Au_m^{\alpha}(s)|^2 + \alpha^2 |A^{3/2}u_m^{\alpha}(s)|^2) \, ds \right] \le \tilde{K}_0^2,$$

where \tilde{K}_0 is given in Proposition 6.

Proof. We take the inner product of the first equation in system (49) with Au_m^{α} and using (17) to obtain

$$\frac{1}{2}\frac{d}{dt}(\|u_m^{\alpha}\|^2 + \alpha^2 |Au_m^{\alpha}|^2) + \nu(|Au_m^{\alpha}|^2 + \alpha^2 |A^{3/2}u_m^{\alpha}|^2) = (P_m f, Au_m^{\alpha}) \le \frac{\nu}{2} |Au_m^{\alpha}|^2 + \frac{|f|^2}{2\nu}.$$

Hence

$$\frac{d}{dt}(\|u_m^{\alpha}\|^2 + \alpha^2 |Au_m^{\alpha}|^2) + \nu(|Au_m^{\alpha}|^2 + \alpha^2 |A^{3/2}u_m^{\alpha}|^2) \le \frac{|f|^2}{\nu}.$$

Integrating the above from 0 to t, we have

$$\|u_m^{\alpha}(t)\|^2 + \alpha^2 |Au_m^{\alpha}(t)|^2 + \nu \int_0^t (|Au_m^{\alpha}(s)|^2 + \alpha^2 |A^{3/2}u_m^{\alpha}(s)|^2) \, ds \le \frac{1}{\nu} \int_0^T |f(s)|^2 \, ds + \|u_0\|^2 + \alpha^2 |Au_0|^2,$$
which concludes our proof.

Remark 5. Similar to the cases of the other α -regularization models, one can show that the solutions of the simplified Bardina model have the same *a priori* estimates as those of the Galerking system, namely, let u^{α} be the solution of the system (48) over the interval [0,T] with initial data $u_0 \in D(A)$, then

$$\sup_{0 \le t \le T} \left[\|u^{\alpha}(t)\|^2 + \alpha^2 |Au^{\alpha}(t)|^2 + \nu \int_0^t (|Au^{\alpha}(s)|^2 + \alpha^2 |A^{3/2}u^{\alpha}(s)|^2) \, ds \right] \le \tilde{K}_0^2, \tag{50}$$

where \tilde{K}_0 is given in Proposition 6.

4. Rates of Convergence of α -Regularization Models to the Navier-Stokes Equations

We aim here to show the convergence rates of solutions of the various α -models toward the corresponding solution of the exact NSE equation when the regularization parameter α approaches zero. We will focus on the L^2 -norm of the difference between u^{α} , solution of the underlying α -model, and u, solution of the exact NSE. From the results concerning all four α -models included in this study, we observe that all the errors $|u - u^{\alpha}|$ are of the same order of $O(\frac{\alpha}{L}(\log \frac{L}{\alpha})^{1/2})$, as $\frac{\alpha}{L}$ goes to zero. Though the four α -models we investigate in this paper share the same order of error estimates with the only difference in the constant, the treatment of each nonlinearity is slightly different and we simply present the details for the readers' benefit. The similar error estimates of these four α -models show that, essentially, these four regularization models are consistent in their convergence to the exact 2D NSE system.

Before we proceed in showing the main results of convergence rates of the α -models, we state an important lemma and a proposition which will play a crucial role in estimating the convergence rates of the various approximation models.

Lemma 10. Let $\alpha > 0$ be a given fixed parameter, and let $\varphi \in H$ and $\delta \in V$. Then

$$|(\varphi - (I + \alpha^2 A)^{-1}\varphi, \delta)| \le \frac{\alpha}{2} |\varphi| ||\delta||.$$

Proof. First we observe that

$$\varphi - (I + \alpha^2 A)^{-1} \varphi = \alpha^2 A (I + \alpha^2 A)^{-1} \varphi$$

Therefore,

$$\begin{aligned} |(\varphi - (I + \alpha^2 A)^{-1} \varphi, \delta)| &= |(\alpha^2 A (I + \alpha^2 A)^{-1} \varphi, \delta)| \\ &= |\alpha \left((\alpha^2 A)^{1/2} (I + \alpha^2 A)^{-1} \varphi, A^{1/2} \delta \right)| \\ &\leq \alpha |(\alpha^2 A)^{1/2} (I + \alpha^2 A)^{-1} \varphi| \|\delta\| \\ &\leq \alpha \|(\alpha^2 A)^{1/2} (I + \alpha^2 A)^{-1}\|_{\mathcal{L}(H)} |\varphi| \|\delta\| \\ &\leq \frac{\alpha}{2} |\varphi| \|\delta\|, \end{aligned}$$

where the last inequality follows from the fact that the operator norm

$$\begin{aligned} \|(\alpha^2 A)^{1/2} (I + \alpha^2 A)^{-1}\|_{\mathcal{L}(H)} &= \sup_{k=1,2,\dots} \left(\frac{(\alpha^2 \lambda_k)^{1/2}}{1 + \alpha^2 \lambda_k}\right) \\ &\leq \sup_{0 \le y < \infty} \frac{y}{1 + y^2} = \frac{1}{2}, \end{aligned}$$
our proof.

which concludes our proof.

Proposition 11. Let $u_0 \in D(A)$ and T > 0. Assume that α is small enough such that $\frac{L}{2\pi\alpha} \ge 1$, and suppose that $u^{\alpha}(t)$ is the solution of system (26) or (48), then

$$\sup_{0 \le t \le T} \|u^{\alpha}(t)\|_{L^{\infty}}^2 \le c \tilde{K}_0^2 \left(1 + \log\left(\frac{L}{2\pi\alpha}\right)\right).$$
(51)

Proof. By the Brezis-Gallouet inequality in Proposition 3, we have

$$\begin{aligned} \|u^{\alpha}(t)\|_{L^{\infty}}^{2} &\leq c \|u^{\alpha}(t)\|^{2} \left(1 + \log\left(\frac{L}{2\pi} \frac{|Au^{\alpha}(t)|}{\|u^{\alpha}(t)\|}\right)\right) \\ &\leq c \left(\|u^{\alpha}(t)\|^{2} + \|u^{\alpha}(t)\|^{2} \log\left(\frac{L}{2\pi\alpha}\right) + \|u^{\alpha}(t)\|^{2} \log\left(\frac{\alpha|Au^{\alpha}(t)|}{\|u^{\alpha}(t)\|}\right)\right). \end{aligned}$$

Therefore, by Proposition 6 for system (26) or Proposition 9 for system (48), we have $\alpha |Au^{\alpha}(t)| \leq \tilde{K}_0$ and $||u^{\alpha}(t)|| \leq \tilde{K}_0$, for all $t \in [0, T]$. Since $\frac{L}{2\pi\alpha} \geq 1$, we obtain

$$\|u^{\alpha}(t)\|_{L^{\infty}}^{2} \leq c \left(\tilde{K}_{0}^{2} + \tilde{K}_{0}^{2} \log\left(\frac{L}{2\pi\alpha}\right) + \tilde{K}_{0}^{2}\left(\frac{\|u^{\alpha}(t)\|}{\tilde{K}_{0}}\right) \log\left(\frac{\tilde{K}_{0}}{\|u^{\alpha}(t)\|}\right)\right).$$

Now, since by Proposition 6 or Proposition 9, $||u^{\alpha}(t)|| \leq \tilde{K}_0$, for all $t \in [0, T]$, then we have

$$\left(\frac{\|u^{\alpha}(t)\|}{\tilde{K}_{0}}\right)\log\left(\frac{\tilde{K}_{0}}{\|u^{\alpha}(t)\|}\right) \leq \frac{1}{e}, \quad \text{for all } t \in [0,T].$$

Consequently, from all the above we have

$$\sup_{0 \le t \le T} \|u^{\alpha}(t)\|_{L^{\infty}}^2 \le c \tilde{K}_0^2 \left(1 + \log\left(\frac{L}{2\pi\alpha}\right)\right).$$

Remark 6. The result above also holds for the solutions u^{α} of systems (29) and (39) with \tilde{K}_0 replaced by \tilde{K}_{02} in the cases of the NS- α and the ML- α models.

Now we are ready to present the convergence rates of various α turbulence models.

4.1. The rate of convergence of the Leray- α model. In this subsection, we will establish error estimates concerning the rate of convergence of the solutions of the 2D Leray- α model to solutions of the 2D NSE, as the regularization parameter α goes to zero. We will proceed by estimating the L^2 -norm of the difference $\delta^{\alpha} = u - u^{\alpha}$, where u is the solution of 2D NSE system (22) and u^{α} is the solution of 2D Leray- α model (26).

From (22) and (26) we observe that $\delta^{\alpha} = u - u^{\alpha}$ satisfies the following equation:

$$\frac{a}{dt}\delta^{\alpha} + \nu A\delta^{\alpha} + B(u,\delta^{\alpha}) + B(\delta^{\alpha},u) - B(\delta^{\alpha},\delta^{\alpha}) - (I+\alpha^2 A)^{-1}B(u^{\alpha},v^{\alpha}) + B(u^{\alpha},u^{\alpha}) = f - (I+\alpha^2 A)^{-1}f.$$
(52)

Theorem 12. Let u^{α} be a solution of the 2D Leray- α system (26) with initial data $u_0 \in D(A)$, and let u be the solution of the 2D NSE system (22) with the same initial data u_0 over the interval [0,T]. Assume that α is small enough such that $\frac{L}{2\pi\alpha} \geq 1$ and let $\delta^{\alpha} = u - u^{\alpha}$, then

$$\sup_{0 \le t \le T} |\delta^{\alpha}(t)|^2 \le \epsilon^2$$

where $\epsilon^2 := \frac{c\alpha^2}{\nu} e^{\frac{cK_0^2}{\nu^2}} \left(T \tilde{K}_0^4 \left(1 + \log \left(\frac{L}{2\pi\alpha} \right) \right) + \|f\|_{L^2((0,T);H)}^2 \right)$, \tilde{K}_0 and K_0 are given in Proposition 6 and 5 respectively, and here c is a dimensionless constant that is independent of ν , α , f or T.

Proof. We take the inner product of the above equation (52) with δ^{α} and use the identity (16) to obtain

$$\frac{1}{2} \frac{d}{dt} |\delta^{\alpha}|^{2} + \nu \|\delta^{\alpha}\|^{2} + (B(\delta^{\alpha}, u), \delta^{\alpha}) + ((I + \alpha^{2}A)^{-1}(B(u^{\alpha}, u^{\alpha}) - B(u^{\alpha}, v^{\alpha})), \delta^{\alpha}) \\ + ((I - (I + \alpha^{2}A)^{-1})B(u^{\alpha}, u^{\alpha}), \delta^{\alpha}) = ((I - (I + \alpha^{2}A)^{-1})f, \delta^{\alpha}).$$

Consequently, one has

$$\frac{1}{2}\frac{d}{dt}|\delta^{\alpha}|^{2} + \nu\|\delta^{\alpha}\|^{2} \le |J_{1}| + |J_{2}| + |J_{3}| + |J_{4}|,$$
(53)

where

1

$$J_1 = (B(\delta^{\alpha}, u), \delta^{\alpha}), \tag{54}$$

$$J_2 = \left((I + \alpha^2 A)^{-1} (B(u^\alpha, u^\alpha) - B(u^\alpha, v^\alpha)), \delta^\alpha \right), \tag{55}$$

$$J_{3} = \left((I - (I + \alpha^{2} A)^{-1}) B(u^{\alpha}, u^{\alpha}), \delta^{\alpha} \right),$$
(56)

$$J_4 = \left((I - (I + \alpha^2 A)^{-1}) f, \delta^{\alpha} \right).$$
(57)

Next, we estimate $|J_1|$, $|J_2|$, $|J_3|$ and $|J_4|$. Thanks to estimate (11) and Young's inequality we have

$$|J_1| = |(B(\delta^{\alpha}, u), \delta^{\alpha})| \le c |\delta^{\alpha}| ||\delta^{\alpha}|| ||u|| \le \frac{\nu}{8} ||\delta^{\alpha}||^2 + \frac{c}{\nu} |\delta^{\alpha}|^2 ||u||^2.$$
(58)

Since $u^{\alpha} + \alpha^2 A u^{\alpha} = v^{\alpha}$, we obtain from (55)

$$|J_2| = |((I + \alpha^2 A)^{-1} B(u^{\alpha}, \alpha^2 A u^{\alpha}), \delta^{\alpha})|$$

= |(B(u^{\alpha}, \alpha^2 A u^{\alpha}), (I + \alpha^2 A)^{-1} \delta^{\alpha})|,

and by applying (15), (12) and Young's inequality we have

$$J_{2}| = |(B(u^{\alpha}, (I + \alpha^{2}A)^{-1}\delta^{\alpha}), \alpha^{2}Au^{\alpha})|$$

$$\leq c\alpha^{2}||u^{\alpha}||_{L^{\infty}}||(I + \alpha^{2}A)^{-1}\delta^{\alpha}|||Au^{\alpha}||$$

$$\leq c\alpha^{2}||u^{\alpha}||_{L^{\infty}}|Au^{\alpha}||\delta^{\alpha}||$$

$$\leq \frac{\nu}{8}||\delta^{\alpha}||^{2} + \frac{c\alpha^{4}}{\nu}|Au^{\alpha}|^{2}||u^{\alpha}||_{L^{\infty}}^{2}.$$

From Proposition 6, we have $\sup_{0 \le t \le T} \alpha^2 |Au^{\alpha}|^2 \le \tilde{K}_0^2$, then by further applying Proposition 11 we have

$$|J_2| \le \frac{\nu}{8} \|\delta^{\alpha}\|^2 + \frac{c\alpha^2}{\nu} \tilde{K}_0^4 \left(1 + \log\left(\frac{L}{2\pi\alpha}\right)\right).$$
(59)

Now, we are ready to estimate $|J_3|$.

By Lemma 10, (56) and Young's inequality we have

$$|J_{3}| \leq \frac{\alpha}{2} |B(u^{\alpha}, u^{\alpha})| \|\delta^{\alpha}\| \leq \frac{\nu}{8} \|\delta^{\alpha}\|^{2} + \frac{\alpha^{2}}{2\nu} |B(u^{\alpha}, u^{\alpha})|^{2}$$
$$\leq \frac{\nu}{8} \|\delta^{\alpha}\|^{2} + \frac{c\alpha^{2}}{\nu} \|u^{\alpha}\|_{L^{\infty}}^{2} \|u^{\alpha}\|^{2}.$$

Since $\sup_{0\leq t\leq T}\|u^\alpha(t)\|^2\leq \tilde{K}_0^2$ by Proposition 6, then by Proposition 11 we reach

$$|J_3| \le \frac{\nu}{8} \|\delta^{\alpha}\|^2 + \frac{c\alpha^2}{\nu} \tilde{K}_0^4 \left(1 + \log\left(\frac{L}{2\pi\alpha}\right)\right).$$

$$(60)$$

Finally, we estimate $|J_4|$. By Lemma 10, (57) and Young's inequality we obtain

$$|J_4| \le \frac{\alpha}{2} |f| \|\delta^{\alpha}\| \le \frac{\nu}{8} \|\delta^{\alpha}\|^2 + \frac{\alpha^2}{2\nu} |f|^2.$$
(61)

Now from (53), (58), (59), (60) and (61) we have

$$\frac{d}{dt}|\delta^{\alpha}|^{2} + \nu\|\delta^{\alpha}\|^{2} \leq \frac{c}{\nu}\|u\|^{2}|\delta^{\alpha}|^{2} + \frac{c\alpha^{2}}{\nu}\left(|f|^{2} + \tilde{K}_{0}^{4}\left(1 + \log\left(\frac{L}{2\pi\alpha}\right)\right)\right).$$

Dropping the $\nu \|\delta^{\alpha}\|^2$ term from the left-hand side and applying Gronwall's inequality we obtain

$$|\delta^{\alpha}(t)|^{2} \leq e^{\frac{c}{\nu}\int_{0}^{t}\|u(\sigma)\|^{2}d\sigma}|\delta^{\alpha}(0)|^{2} + \frac{c\alpha^{2}}{\nu}\int_{0}^{t}e^{\frac{c}{\nu}\int_{\sigma}^{t}\|u(s)\|^{2}ds}\left[|f(\sigma)|^{2} + \tilde{K}_{0}^{4}\left(1 + \log\left(\frac{L}{2\pi\alpha}\right)\right)\right] d\sigma$$

Since $\delta^{\alpha}(0) = 0$, then by Propositon 5 we obtain

$$|\delta^{\alpha}(t)|^{2} \leq \frac{c\alpha^{2}}{\nu} e^{\frac{cK_{0}^{2}}{\nu^{2}}} \left(T\tilde{K}_{0}^{4} \left(1 + \log\left(\frac{L}{2\pi\alpha}\right) \right) + \|f\|_{L^{2}((0,T);H)}^{2} \right) =: \epsilon^{2},$$

which concludes our proof.

4.2. The rate of convergence of the Navier-Stokes- α model. In this subsection, we will establish error estimates concerning the rate of convergence of solutions of the 2D NS- α regularization model to solutions of the 2D NSE, as the regularization parameter α goes to zero. We will proceed by estimating the L^2 -norm of the difference $\delta^{\alpha} = u - u^{\alpha}$, where u is the solution of 2D NSE system (22) and u^{α} is the solution of 2D NS- α model (29).

From (22) and (29) we observe that $\delta^{\alpha} = u - u^{\alpha}$ satisfies the following equation:

$$\frac{d}{dt}\delta^{\alpha} + \nu A\delta^{\alpha} + B(\delta^{\alpha}, u) + B(u, \delta^{\alpha}) - B(\delta^{\alpha}, \delta^{\alpha}) + (I + \alpha^{2}A)^{-1}(B(u^{\alpha}, u^{\alpha}) - B(u^{\alpha}, v^{\alpha})) + (I - (I + \alpha^{2}A)^{-1})B(u^{\alpha}, u^{\alpha}) + (I + \alpha^{2}A)^{-1}B^{*}(u^{\alpha}, v^{\alpha}) = (I - (I + \alpha^{2}A)^{-1})f,$$
(62)

where for every $v \in V$ fixed, the operator $B^*(\cdot, v)$ is the adjoint operator of the operator $B(\cdot, v)$, which is defined by $(B^*(\psi, v), \varphi) = (B(\varphi, v), \psi)$ for every $\varphi, \psi \in \mathcal{V}$ (see relation (18)).

Theorem 13. Let u^{α} be the solution of the 2D NS- α system (29) with initial data $u_0 \in D(A)$, and let u be the solution of the 2D NSE system (22) with the same initial data u_0 over the interval [0,T]. Assume that α is small enough such that $\frac{L}{2\pi\alpha} \geq 1$ and let $\delta^{\alpha} = u - u^{\alpha}$, then

$$\sup_{0 \le t \le T} |\delta^{\alpha}(t)|^2 \le \tilde{\epsilon}^2,$$

where $\tilde{\epsilon}^2 := \frac{c\alpha^2}{\nu} e^{\frac{cK_0^2}{\nu^2}} \left(T \tilde{\tilde{K}}_{02}^4 \left(1 + \log\left(\frac{L}{2\pi\alpha}\right) \right) + \|f\|_{L^2((0,T);H)}^2 \right)$, and K_0 , $\tilde{\tilde{K}}_{02}$ are given in (24) and (32), respectively.

Proof. By taking inner product of (62) with δ^{α} and using (16) and (17) we have

$$\frac{1}{2}\frac{d}{dt}|\delta^{\alpha}|^{2} + \nu \|\delta^{\alpha}\|^{2} + J_{1} + J_{2} + J_{3} + J_{5} = J_{4},$$

where J_1, J_2, J_3 and J_4 are given in (54)-(57) and $J_5 = ((I + \alpha^2 A)^{-1} B^*(u^{\alpha}, v^{\alpha}), \delta^{\alpha})$. Therefore,

$$\frac{1}{2}\frac{d}{dt}|\delta^{\alpha}|^{2} + \nu\|\delta^{\alpha}\|^{2} \le |J_{1}| + |J_{2}| + |J_{3}| + |J_{4}| + |J_{5}|.$$
(63)

The estimates for $|J_1|, |J_2|, |J_3|$ and $|J_4|$ follow exactly the same steps as in the proof of Theorem 12, where \tilde{K}_0 is replaced by \tilde{K}_{02} from Proposition 7 (see also Remark 6).

Next, we estimate $|J_5|$. To this end we first observe that $B^*(u, u) \equiv 0$, because, thanks to (16),

$$\langle B^*(u,u), w \rangle_{V'} = \langle B(w,u), u \rangle_{V'} = 0$$
 for every $w \in V$.

Thus,

$$\begin{aligned} |J_5| &= |((I + \alpha^2 A)^{-1} B^*(u^{\alpha}, v^{\alpha}), \delta^{\alpha})| \\ &= |((I + \alpha^2 A)^{-1} (B^*(u^{\alpha}, v^{\alpha}) - B^*(u^{\alpha}, u^{\alpha})), \delta^{\alpha})| \\ &= \alpha^2 |((I + \alpha^2 A)^{-1} B^*(u^{\alpha}, Au^{\alpha}), \delta^{\alpha})|, \end{aligned}$$

where we used the relation $u^{\alpha} + \alpha^2 A u^{\alpha} = v^{\alpha}$ and the bilinearity of $B^*(\cdot, \cdot)$. Consequently, by the definition of B^* and (15), we have

$$|J_5| = \alpha^2 |(B((I + \alpha^2 A)^{-1} \delta^{\alpha}, Au^{\alpha}), u^{\alpha})| = \alpha^2 |(B((I + \alpha^2 A)^{-1} \delta^{\alpha}, u^{\alpha}), Au^{\alpha})|.$$

Therefore, by Lemma 2 and Young's inequality we have

$$\begin{aligned} |J_5| &\leq c\alpha^2 \| (I + \alpha^2 A)^{-1} \delta^{\alpha} \| \| u^{\alpha} \| |Au^{\alpha}| \\ &\leq c\alpha^2 \| \delta^{\alpha} \| \| u^{\alpha} \| |Au^{\alpha}| \\ &\leq \frac{\nu \| \delta^{\alpha} \|^2}{8} + \frac{c\alpha^4}{\nu} \| u^{\alpha} \|^2 |Au^{\alpha}|^2. \end{aligned}$$

By the above and Proposition 7 we conclude

$$|J_5| \le \frac{\nu \|\delta^{\alpha}\|^2}{8} + \frac{c\alpha^2}{\nu} \tilde{K}_{02}^4.$$
(64)

From (58)-(61) (where \tilde{K}_0 is replaced by $\tilde{\tilde{K}}_{02}$, see also Remark 6), (63) and (64) we obtain

$$\frac{d}{dt} |\delta^{\alpha}|^{2} + \nu \|\delta^{\alpha}\|^{2} \leq \frac{c}{\nu} \|u\|^{2} |\delta^{\alpha}|^{2} + \frac{c}{\nu} \alpha^{2} \left(|f|^{2} + \tilde{\tilde{K}}_{02}^{4} \left(1 + \log\left(\frac{L}{2\pi\alpha}\right) \right) \right).$$

Dropping the $\nu \|\delta^{\alpha}\|^2$ term from the left-hand side and applying Gronwall's inequality we obtain

$$|\delta^{\alpha}(t)|^{2} \leq e^{\frac{c}{\nu}\int_{0}^{t}\|u(\sigma)\|^{2}d\sigma}|\delta^{\alpha}(0)|^{2} + \frac{c\alpha^{2}}{\nu}\int_{0}^{t}e^{\frac{c}{\nu}\int_{\sigma}^{t}\|u(s)\|^{2}ds}\left(|f(\sigma)|^{2} + \tilde{\tilde{K}}_{02}^{4}\left(1 + \log\left(\frac{L}{2\pi\alpha}\right)\right)\right) \, d\sigma.$$

Since $\delta^{\alpha}(0) = 0$, then by Proposition 5 and the above we obtain

$$|\delta^{\alpha}(t)|^{2} \leq \frac{c\alpha^{2}}{\nu} e^{\frac{cK_{0}^{2}}{\nu^{2}}} \left(T\tilde{\tilde{K}}_{02}^{4} \left(1 + \log\left(\frac{L}{2\pi\alpha}\right) \right) + \|f\|_{L^{2}((0,T);H)}^{2} \right) =: \tilde{\epsilon}^{2}.$$

4.3. The rate of convergence of the Modified Leray- α model. In this subsection, we will establish error estimates concerning the rate of convergence of solutions of the 2D ML- α regularization model to solutions of the 2D NSE, as the regularization parameter α goes to zero. We will proceed by estimating the L^2 -norm of the difference $\delta^{\alpha} = u - u^{\alpha}$, where u is the solution of 2D NSE system (22) and u^{α} is the solution of the 2D ML- α model (39).

From (22) and (39) we observe that $\delta^{\alpha} = u - u^{\alpha}$ satisfies the following equation:

$$\frac{d}{dt}\delta^{\alpha} + B(\delta^{\alpha}, u) + B(u, \delta^{\alpha}) - B(\delta^{\alpha}, \delta^{\alpha}) + (I + \alpha^{2}A)^{-1}(B(u^{\alpha}, u^{\alpha}) - B(v^{\alpha}, u^{\alpha}))
+ (I - (I + \alpha^{2}A)^{-1})B(u^{\alpha}, u^{\alpha}) = (I - (I + \alpha^{2}A)^{-1})f.$$
(65)

Theorem 14. Let u^{α} be the solution of the 2D Modified Leray- α system (39) with initial data $u_0 \in D(A)$, and let u be the solution of the 2D NSE system (22) with the same initial data u_0 over the interval [0,T]. Assume that α is small enough such that $\frac{L}{2\pi\alpha} \geq 1$ and let $\delta^{\alpha} = u - u^{\alpha}$, then

$$\sup_{0 \le t \le T} |\delta^{\alpha}(t)|^2 \le \tilde{\epsilon}^2$$

where $\tilde{\epsilon}$ is given in Theorem 13.

Proof. Taking the inner product of the above equation (65) with δ^{α} we obtain

$$\frac{1}{2}\frac{d}{dt}|\delta^{\alpha}|^{2} + \nu\|\delta^{\alpha}\|^{2} \le |J_{1}| + |J_{3}| + |J_{4}| + |J_{6}|, \tag{66}$$

where J_1, J_3 and J_4 are as in (54), (56) and (57) respectively and

$$J_6 = ((I + \alpha^2 A)^{-1} (B(u^{\alpha}, u^{\alpha}) - B(v^{\alpha}, u^{\alpha})), \delta^{\alpha}).$$

Here again the estimates for J_1, J_3 and J_4 are as in the proof of Theorem 12, with \tilde{K}_0 replaced by \tilde{K}_{02} (see also Remark 6). Next, we estimate J_6 . Thanks to (14) we have

$$|J_{6}| \leq |\alpha^{2}((I + \alpha^{2}A)^{-1}B(Au^{\alpha}, u^{\alpha}), \delta^{\alpha})| \\ = |\alpha^{2}(B(Au^{\alpha}, (I + \alpha^{2}A)^{-1}\delta^{\alpha}), u^{\alpha})| \\ \leq c\alpha^{2} ||u^{\alpha}||_{L^{\infty}} ||(I + \alpha^{2}A)^{-1}\delta^{\alpha}|||Au^{\alpha}|| \\ \leq c\alpha^{2} ||u^{\alpha}||_{L^{\infty}} |Au^{\alpha}|||\delta^{\alpha}|| \\ \leq \frac{\nu}{8} ||\delta^{\alpha}||^{2} + \frac{c\alpha^{4}}{\nu} |Au^{\alpha}|^{2} ||u^{\alpha}||_{L^{\infty}}^{2}.$$

Now since $\alpha^2 |Au^{\alpha}|^2 \leq \tilde{\tilde{K}}_{02}^2$ by Proposition 8, then by Proposition 11 (see also Remark 6) we have

$$|J_6| \le \frac{\nu}{8} \|\delta^{\alpha}\|^2 + \frac{c\alpha^4}{\nu} \tilde{K}_{02}^4 \left(1 + \log\left(\frac{L}{2\pi\alpha}\right)\right).$$
(67)

From (58), (60), (61) (where \tilde{K}_0 is replaced by $\tilde{\tilde{K}}_{02}$), (67) and (66) we have

$$\frac{d}{dt}|\delta^{\alpha}|^{2} + \nu \|\delta^{\alpha}\|^{2} \leq \frac{c}{\nu}\|u^{\alpha}\|^{2}|\delta^{\alpha}|^{2} + \frac{c}{\nu}\alpha^{2}\left(|f|^{2} + \tilde{\tilde{K}}_{02}^{4}\left(1 + \log\left(\frac{L}{2\pi\alpha}\right)\right)\right),$$

which concludes the proof of the theorem by Gronwall's inequality.

4.4. The rate of convergence of the simplified Bardina model. Now we present the error estimates for the simplified Bardina regularization model. As before, we will proceed by estimating the L^2 -norm of the difference $\delta^{\alpha} = u - u^{\alpha}$, where u is the solution of 2D NSE system (22) and u^{α} is the solution of the simplified Bardina system (48).

From (22) and (48) we observe that $\delta^{\alpha} = u - u^{\alpha}$ satisfies the following equation:

$$\frac{d}{dt}\delta^{\alpha} + \nu A\delta^{\alpha} + B(u,\delta^{\alpha}) + B(\delta^{\alpha},u) - B(\delta^{\alpha},\delta^{\alpha}) + (I - (I + \alpha^2 A)^{-1})B(u^{\alpha},u^{\alpha})$$
$$= (I - (I + \alpha^2 A)^{-1})f.$$
(68)

Theorem 15. Let u^{α} be the solution of the 2D simplified Bardina system (48) with the initial date $u_0 \in D(A)$, and let u be the solution of the 2D NSE system (22) with the same initial data u_0 over the interval [0,T]. Assume that α is small enough such that $\frac{L}{2\pi\alpha} \geq 1$ and let $\delta^{\alpha} = u - u^{\alpha}$, then

$$\sup_{0 \le t \le T} |\delta^{\alpha}(t)|^2 \le \epsilon^2,$$

where ϵ is given in Theorem 12.

Proof. Taking the inner product of the equation (68) with δ^{α} we obtain

$$\frac{1}{2}\frac{d}{dt}|\delta^{\alpha}|^{2} + \nu\|\delta^{\alpha}\|^{2} \le |J_{1}| + |J_{3}| + |J_{4}|, \tag{69}$$

where J_1, J_3 and J_4 are as in (54), (56) and (57) respectively. Here again the estimates for J_1, J_3 and J_4 are as in the proof of Theorem 12, with exactly the same \tilde{K}_0 . From (58), (60), (61) and (69) we have

$$\frac{d}{dt}|\delta^{\alpha}|^{2}+\nu\|\delta^{\alpha}\|^{2} \leq \frac{c}{\nu}\|u\|^{2}|\delta^{\alpha}|^{2}+\frac{c}{\nu}\alpha^{2}\left(|f|^{2}+\tilde{K}_{0}^{4}\left(1+\log\left(\frac{L}{2\pi\alpha}\right)\right)\right).$$

Y.CAO AND E.S. TITI

Dropping the $\nu \|\delta^{\alpha}\|^2$ term from the left-hand side and applying Gronwall's inequality we obtain

$$\begin{aligned} |\delta^{\alpha}(t)|^{2} &\leq e^{\frac{c}{\nu}\int_{0}^{t}\|u(\sigma)\|^{2}d\sigma}|\delta^{\alpha}(0)|^{2} \\ &+ \frac{c\alpha^{2}}{\nu}\int_{0}^{t}e^{\frac{c}{\nu}\int_{\sigma}^{t}\|u(s)\|^{2}ds}\left(|f(\sigma)|^{2} + \tilde{K}_{0}^{4}\left(1 + \log\left(\frac{L}{2\pi\alpha}\right)\right)\right) \, d\sigma. \end{aligned}$$

Since $\delta^{\alpha}(0) = 0$, then by Proposition 5 we obtain

$$|\delta^{\alpha}(t)|^{2} \leq \frac{c\alpha^{2}}{\nu} e^{\frac{cK_{0}^{2}}{\nu^{2}}} \left(T\tilde{K}_{0}^{4} \left(1 + \log\left(\frac{L}{2\pi\alpha}\right) \right) + \|f\|_{L^{2}((0,T);H)}^{2} \right) =: \epsilon^{2}.$$

5. Error Estimates of The Galerkin Approximation of the α -Regularization Models

In numerical simulation one needs to approximate the exact solutions of the underlying equations, that lie in the infinite dimensional function spaces, by functions that lie in finite dimensional spaces. In this section we estimate the errors between the approximation solutions u_m^{α} , of the finite dimensional ordinary differential equation system (the finite dimensional Galerkin system of order m in this context) and the exact solutions u^{α} of the α -models in L^2 -norm. The errors are given in terms of m and the regularization parameter α . Combining this with the results we establish in the previous section and by applying the triangle inequality, we obtain error estimates of numerical approximation solutions u_m^{α} of the underlying α -model and of the exact solution u of the 2D NSE system. We take the Leray- α model as an example in this section and show the error estimate results. By similar arguments, one can show the error estimates for the other α -regularization models introduced in this paper.

Galerkin approximation for the 2D Leray- α model

Now we present the error estimates for the Galerkin approximation of the Leray- α model. This estimate for the rate of convergence of the Leray- α model is along the same lines of [17, 23, 38, 39] and in the spirit of the work for the 2D NSE.

For u^{α} an exact solution of the Leray- α system (26), we can decompose u^{α} as follows: $u^{\alpha} = p_m + q_m$, where $p_m = P_m u^{\alpha}$ and $q_m = (I - P_m)u^{\alpha}$, P_m is the orthogonal project from H onto H_m , which is defined in Section 2. Rewriting equation (26) as

$$\frac{du^{\alpha}}{dt} + \nu A u^{\alpha} + (I + \alpha^2 A)^{-1} B(u^{\alpha}, (I + \alpha^2 A) u^{\alpha}) = (I + \alpha^2 A)^{-1} f,$$
(70)

and since $u^{\alpha} = p_m + q_m$, we can decompose the above equation (70) into the following coupled system of equations:

$$\frac{dp_m}{dt} + \nu A p_m + (I + \alpha^2 A)^{-1} P_m B(u^\alpha, (I + \alpha^2 A)u^\alpha) = (I + \alpha^2 A)^{-1} P_m f,$$
(71)

$$\frac{dq_m}{dt} + \nu Aq_m + (I + \alpha^2 A)^{-1} (I - P_m) B(u^{\alpha}, (I + \alpha^2 A)u^{\alpha}) = (I + \alpha^2 A)^{-1} (I - P_m) f.$$
(72)

For the Galerkin approximation system of the Leray- α system, we rewrite the equation (27) in the following equivalent form

$$\frac{du_m^{\alpha}}{dt} + \nu A u_m^{\alpha} + (I + \alpha^2 A)^{-1} P_m B(u_m^{\alpha}, (I + \alpha^2 A) u_m^{\alpha}) = (I + \alpha^2 A)^{-1} P_m f.$$
(73)

We will proceed by first estimating the L^2 -norm of q_m and then the L^2 -norm of the difference $\delta_m = p_m - u_m^{\alpha}$, where u_m^{α} is the solution of the Galerkin system of the Leray- α system (73). Then by the triangle inequality and orthogonality of spaces projected by P_m and $(I - P_m)$, we obtain the error estimates

of the L²-norm: $|u^{\alpha} - u_m^{\alpha}|^2 = |q_m|^2 + |p_m - u_m^{\alpha}|^2$.

From (71) and (73) we observe that $\delta_m = p_m - u_m^{\alpha}$ satisfies the following equation:

$$\frac{d\delta_m}{dt} + \nu A\delta_m + (I + \alpha^2 A)^{-1} P_m B(u^{\alpha}, (I + \alpha^2 A)u^{\alpha}) - (I + \alpha^2 A)^{-1} P_m B(u_m^{\alpha}, (I + \alpha^2 A)u_m^{\alpha}) = 0.$$

Since $u^{\alpha} = p_m + q_m$ and $\delta_m = p_m - u_m^{\alpha}$, we can rewrite the above equation as

$$\frac{d\delta_m}{dt} + \nu A\delta_m + (I + \alpha^2 A)^{-1} P_m B(\delta_m + q_m, (I + \alpha^2 A)u^\alpha) - (I + \alpha^2 A)^{-1} P_m B(u_m^\alpha, (I + \alpha^2 A)(\delta_m + q_m)) = 0.$$
(74)

Theorem 16. Let T > 0 and let u^{α} be a solution of the Leray- α system (70) with initial data $u_0 \in D(A)$, and let u_m^{α} be the solution of (73) with initial data $u_{0m} = P_m u_0$ over the interval [0, T]. For a given $m \ge 1$, assume that α is small such that $\alpha^2 \le \frac{1}{\lambda_{m+1}}$, then

$$\sup_{0 \le t \le T} |u^{\alpha}(t) - u^{\alpha}_m(t)|^2 \le e^2,$$

where $e^2 := \frac{1}{\lambda_{m+1}^2} (Q + R + L_m \tilde{U} \tilde{V})$, $L_m = 1 + \log(\frac{\lambda_m}{\lambda_1})$ and $Q, R, \tilde{U}, \tilde{V}$ depend on $\nu, u_0, \log(\frac{L}{2\pi\alpha}), f, T$, and are given explicitly in (80), (81), (92) and (93), respectively.

Remark 7. Here we require $f \in L^{\infty}((0,T); H)$, which is stronger assumption than the condition $f \in L^{2}((0,T); H)$ in the estimate of rates of convergence in section 4.

Proof. First, we estimate the L^2 norm of q_m .

We take the inner product of equation (72) with q_m and get

$$\frac{1}{2} \frac{a}{dt} |q_m|^2 + \nu ||q_m||^2 \leq |((I + \alpha^2 A)^{-1} B(u^\alpha, (I + \alpha^2 A) u^\alpha), q_m)| + |((I + \alpha^2 A)^{-1} f, q_m)| \leq M_1 + M_2.$$
(75)

Next, we estimat M_1 , M_2 . By virtue of (12) we have

$$M_{1} = |((I + \alpha^{2}A)^{-1}B(u^{\alpha}, (I + \alpha^{2}A)u^{\alpha}), q_{m})|$$

$$= |(B(u^{\alpha}, (I + \alpha^{2}A)u^{\alpha}), (I + \alpha^{2}A)^{-1}q_{m})|$$

$$\leq c \|u^{\alpha}\|_{L^{\infty}} \|(I + \alpha^{2}A)u^{\alpha}\||(I + \alpha^{2}A)^{-1}q_{m}|$$

$$\leq c\tilde{K}_{0}(1 + \log(\frac{L}{2\pi\alpha}))^{1/2}(\|u^{\alpha}\| + \alpha^{2}|A^{3/2}u^{\alpha}|)\frac{\|q_{m}\|}{\lambda_{m+1}^{1/2}},$$

where in the last inequality we apply (5) and Proposition 11 to $||u^{\alpha}||_{L^{\infty}}$. Notice that by the assumption that $\alpha^2 \leq \frac{1}{\lambda_{m+1}} \leq \frac{1}{\lambda_1} = (\frac{L}{2\pi})^2$, we consequently have $\frac{L}{2\pi\alpha} \geq 1$, and as a result, it is valid to apply Proposition 11.

Now, by Young's inequality and the a priori estimates obtained in (28), we have

$$M_{1} \leq \frac{\nu}{4} \|q_{m}\|^{2} + \frac{1}{\lambda_{m+1}} \frac{c}{\nu} \tilde{K}_{0}^{2} (1 + \log(\frac{L}{2\pi\alpha})) (\|u^{\alpha}\|^{2} + \alpha^{4} |A^{3/2}u^{\alpha}|^{2})$$

$$\leq \frac{\nu}{4} \|q_{m}\|^{2} + \frac{1}{\lambda_{m+1}} \frac{c}{\nu} \tilde{K}_{0}^{4} (1 + \log(\frac{L}{2\pi\alpha}))$$
(76)

$$4 \qquad \lambda_{m+1} \nu \qquad 2\pi\alpha \\ + \frac{\alpha^2}{\lambda_{m+1}} \frac{c}{\nu} \tilde{K}_0^2 (1 + \log(\frac{L}{2\pi\alpha})) (\alpha^2 |A^{3/2} u^{\alpha}|^2).$$
(77)

Y.CAO AND E.S. TITI

Now, for estimating M_2 , we apply Cauchy-Schwarz inequality, (5) and Young's inequality to obtain

$$M_{2} = |((I + \alpha^{2}A)^{-1}(I - P_{m})f, q_{m})| = |((I - P_{m})f, (I + \alpha^{2}A)^{-1}q_{m})|$$

$$\leq |f||(I + \alpha^{2}A)^{-1}q_{m}| \leq |f|\frac{\|q_{m}\|}{\lambda_{m+1}^{1/2}} \leq \frac{\nu}{4}\|q_{m}\|^{2} + \frac{1}{\lambda_{m+1}}\frac{c}{\nu}|f|^{2}.$$
(78)

Let us substitute the bounds for M_1 , M_2 into (75) to obtain

$$\frac{d}{dt}|q_{m}|^{2} + \nu ||q_{m}||^{2} \leq \frac{1}{\lambda_{m+1}} \frac{c}{\nu} \left(|f|^{2} + \tilde{K}_{0}^{4}(1 + \log(\frac{L}{2\pi\alpha})) \right) + \frac{\alpha^{2}}{\lambda_{m+1}} \frac{c}{\nu} \tilde{K}_{0}^{2}(1 + \log(\frac{L}{2\pi\alpha}))(\alpha^{2}|A^{3/2}u^{\alpha}|^{2}).$$
(79)

By Poincaré inequality and (5), $\lambda_{m+1}|q_m|^2 \leq ||q_m||^2$, we obtain

$$\frac{d}{dt}|q_{m}|^{2} + \nu\lambda_{m+1}|q_{m}|^{2} \leq \frac{1}{\lambda_{m+1}}\frac{c}{\nu}\left(|f|^{2} + \tilde{K}_{0}^{4}(1 + \log(\frac{L}{2\pi\alpha}))\right) + \frac{\alpha^{2}}{\lambda_{m+1}}\frac{c}{\nu}\tilde{K}_{0}^{2}(1 + \log(\frac{L}{2\pi\alpha}))(\alpha^{2}|A^{3/2}u^{\alpha}|^{2})$$

By Gronwall inequality, we get

$$\begin{split} |q_{m}(t)|^{2} &\leq e^{-\nu\lambda_{m+1}t} |q_{m}(0)|^{2} + \int_{0}^{t} e^{-\nu\lambda_{m+1}(t-s)} \frac{1}{\lambda_{m+1}} \frac{c}{\nu} \left(|f|^{2} + \tilde{K}_{0}^{4}(1+\log(\frac{L}{2\pi\alpha})) \right) \, ds \\ &+ \int_{0}^{t} e^{-\nu\lambda_{m+1}(t-s)} \frac{\alpha^{2}}{\lambda_{m+1}} \frac{c}{\nu} \tilde{K}_{0}^{2}(1+\log(\frac{L}{2\pi\alpha})) (\alpha^{2}|A^{3/2}u^{\alpha}|^{2}) \, ds \\ &\leq e^{-\nu\lambda_{m+1}t} \frac{|Aq_{m}(0)|^{2}}{\lambda_{m+1}^{2}} + \frac{1}{\lambda_{m+1}^{2}} (1-e^{-\nu\lambda_{m+1}t}) \frac{c}{\nu^{2}} \left(|f|^{2}_{L^{\infty}((0,T);H)} + \tilde{K}_{0}^{4}(1+\log(\frac{L}{2\pi\alpha})) \right) \\ &+ \frac{\alpha^{2}}{\lambda_{m+1}} \frac{c}{\nu} \tilde{K}_{0}^{2}(1+\log(\frac{L}{2\pi\alpha})) \int_{0}^{t} (\alpha^{2}|A^{3/2}u^{\alpha}|^{2}) \, ds \\ &\leq \frac{1}{\lambda_{m+1}^{2}} |Au_{0}|^{2} + \frac{1}{\lambda_{m+1}^{2}} \frac{c}{\nu^{2}} \left(|f|^{2}_{L^{\infty}((0,T):H)} + \tilde{K}_{0}^{4}(1+\log(\frac{L}{2\pi\alpha})) \right) \\ &+ \frac{\alpha^{2}}{\lambda_{m+1}} \frac{c}{\nu^{2}} \tilde{K}_{0}^{4}(1+\log(\frac{L}{2\pi\alpha})). \end{split}$$

where we apply the *a priori* estimates of solutions of Leray- α model given in Proposition 6. Next, denote by

$$Q := |Au_0|^2 + \frac{c}{\nu^2} \left(|f|^2_{L^{\infty}((0,T):H)} + \tilde{K}^4_0(1 + \log(\frac{L}{2\pi\alpha})) \right), \tag{80}$$

and

$$R := \frac{c}{\nu^2} \tilde{K}_0^4 (1 + \log(\frac{L}{2\pi\alpha})).$$
(81)

we obtain

$$|q_m(t)|^2 \le \frac{1}{\lambda_{m+1}^2}Q + \frac{\alpha^2}{\lambda_{m+1}}R.$$

By the assumption that $\alpha^2 \leq \frac{1}{\lambda_{m+1}}$, we have

$$|q_m|^2 \le \frac{1}{\lambda_{m+1}^2} (Q+R).$$
 (82)

Next, we estimate the L^2 -norm of δ_m . Taking the inner product of equation (74) with δ_m , we get

$$\frac{1}{2}\frac{d}{dt}|\delta_m|^2 + \nu \|\delta_m\|^2 \le M_3 + M_4 + M_5 + M_6,\tag{83}$$

where

$$\begin{split} M_{3} &= |B(\delta_{m}, (I + \alpha^{2}A)u^{\alpha}), (I + \alpha^{2}A)^{-1}\delta_{m})|, \\ M_{4} &= |B(q_{m}, (I + \alpha^{2}A)u^{\alpha}), (I + \alpha^{2}A)^{-1}\delta_{m})|, \\ M_{5} &= |(B(u_{m}^{\alpha}, (I + \alpha^{2}A)\delta_{m}), (I + \alpha^{2}A)^{-1}\delta_{m})|, \\ M_{6} &= |(B(u_{m}^{\alpha}, (I + \alpha^{2}A)q_{m}), (I + \alpha^{2}A)^{-1}\delta_{m})|. \end{split}$$

First, let us estimate M_3 . Thanks to (11) we have

$$M_{3} \leq c|\delta_{m}|^{1/2} \|\delta_{m}\|^{1/2} \|(I + \alpha^{2}A)u^{\alpha}\| \|(I + \alpha^{2}A)^{-1}\delta_{m}\|^{1/2} \|(I + \alpha^{2}A)^{-1}\delta_{m}\|^{1/2}$$

$$\leq c|\delta_{m}| \|\delta_{m}\| (\|u^{\alpha}\| + \alpha^{2}|A^{3/2}u^{\alpha}|))$$

$$\leq \frac{\nu}{16} \|\delta_{m}\|^{2} + \frac{c}{\nu} |\delta_{m}|^{2} (\|u^{\alpha}\|^{2} + \alpha^{4}|A^{3/2}u^{\alpha}|^{2})$$

$$\leq \frac{\nu}{16} \|\delta_{m}\|^{2} + \frac{c}{\nu} |\delta_{m}|^{2} \tilde{K}_{0}^{2} + \frac{1}{\lambda_{m+1}} \frac{c}{\nu} |\delta_{m}|^{2} (\alpha^{2}|A^{3/2}u^{\alpha}|^{2}), \qquad (84)$$

where in the last inequality we use the *a priori* estimates of Leray- α obtained in Proposition 6 and the assumption that $\alpha^2 \leq \frac{1}{\lambda_{m+1}}$. Now, we estimate M_4 . By applying (21) and (7) we obtain

$$M_{4} \leq c|q_{m}|\|(I + \alpha^{2}A)u^{\alpha}\|\|(I + \alpha^{2}A)^{-1}\delta_{m}\|_{L^{\infty}}$$

$$\leq c|q_{m}|(\|u^{\alpha}\| + \alpha^{2}|A^{3/2}u^{\alpha}|)\|(I + \alpha^{2}A)^{-1}\delta_{m}\|\left(1 + \log(\frac{\lambda_{m}}{\lambda_{1}})\right)^{1/2}$$

$$\leq c|q_{m}|(\|u^{\alpha}\| + \alpha^{2}|A^{3/2}u^{\alpha}|)\|\delta_{m}\|L_{m}^{1/2}$$

$$\leq \frac{\nu}{16}\|\delta_{m}\|^{2} + \frac{cL_{m}}{\nu}|q_{m}|^{2}(\|u^{\alpha}\|^{2} + \alpha^{2}(\alpha^{2}|A^{3/2}u^{\alpha}|^{2}))$$

$$\leq \frac{\nu}{16}\|\delta_{m}\|^{2} + \frac{cL_{m}}{\nu}|q_{m}|^{2}(\tilde{K}_{0}^{2} + \alpha^{2}(\alpha^{2}|A^{3/2}u^{\alpha}|^{2})), \qquad (85)$$

where $L_m = 1 + \log(\frac{\lambda_m}{\lambda_1})$, and we apply Proposition 3 and (7) in the second inequality above. Let us now estimate M_5 . By virtue of (11) we have

$$M_5 \leq c |u_m^{\alpha}|^{1/2} ||u_m^{\alpha}||^{1/2} ||(I+\alpha^2 A)\delta_m|| (I+\alpha^2 A)^{-1}\delta_m|^{1/2} ||(I+\alpha^2 A)^{-1}\delta_m|^{1/2} \\ \leq c |u_m^{\alpha}|^{1/2} ||u_m^{\alpha}||^{1/2} (1+\alpha^2 \lambda_m) ||\delta_m||\delta_m|^{1/2} ||\delta_m||^{1/2}.$$

Since $\alpha^2 \lambda_m \leq \alpha^2 \lambda_{m+1} \leq 1$, and c denotes general dimensionless constant, we have

$$M_{5} \leq c \|\delta_{m}\|^{3/2} |\delta_{m}|^{1/2} |u_{m}^{\alpha}|^{1/2} \|u_{m}^{\alpha}\|^{1/2} \leq \frac{\nu}{16} \|\delta_{m}\|^{2} + \frac{c}{\nu} |\delta_{m}|^{2} |u_{m}^{\alpha}|^{2} \|u_{m}^{\alpha}\|^{2} \leq \frac{\nu}{16} \|\delta_{m}\|^{2} + \frac{c}{\nu^{3}} |\delta_{m}|^{2} \frac{\tilde{K}_{0}^{4}}{\lambda_{1}},$$
(86)

where in the last inequality we apply Proposition 6.

For the last term M_6 , we will proceed by applying (15) and (12), then we obtain

$$M_{6} = |(B(u_{m}^{\alpha}, (I + \alpha^{2}A)q_{m}), (I + \alpha^{2}A)^{-1}\delta_{m})|$$

$$= |(B(u_{m}^{\alpha}, (I + \alpha^{2}A)^{-1}\delta_{m}), (I + \alpha^{2}A)q_{m})|$$

$$\leq c ||u_{m}^{\alpha}||_{L^{\infty}} ||(I + \alpha^{2}A)^{-1}\delta_{m}|| |(I + \alpha^{2}A)q_{m}|.$$

By applying Proposition 3, inequality (7), (5), and the facts that

$$\begin{split} \|(I+\alpha^2 A)^{-1}\phi\| &\leq \|\phi\|, \quad \text{for all } \phi \in V, \\ |(I+\alpha^2 A)\phi| &\sim |\phi|+\alpha^2 |A\phi| \quad \text{for all } \phi \in D(A), \end{split}$$

we obtain

$$M_{6} \leq c \|u_{m}^{\alpha}\| \left(1 + \log(\frac{\lambda_{m}}{\lambda_{1}})\right)^{1/2} \|(I + \alpha^{2}A)^{-1}\delta_{m}\||(I + \alpha^{2}A)qm|$$

$$\leq c L_{m}^{1/2}\tilde{K}_{0}\|\delta_{m}\|(|q_{m}| + \alpha^{2}|Aq_{m}|)$$

$$\leq c L_{m}^{1/2}\tilde{K}_{0}\|\delta_{m}\|(|q_{m}| + \alpha^{2}\frac{|A^{3/2}q_{m}|}{\lambda_{m+1}^{1/2}})$$

$$\leq \frac{\nu}{16}\|\delta_{m}\|^{2} + \frac{cL_{m}}{\nu}\tilde{K}_{0}^{2}(|q_{m}|^{2} + \frac{\alpha^{2}}{\lambda_{m+1}}(\alpha^{2}|A^{3/2}u^{\alpha}|^{2}))$$

$$\leq \frac{\nu}{16}\|\delta_{m}\|^{2} + \frac{cL_{m}}{\nu}\tilde{K}_{0}^{2}|q_{m}|^{2} + \frac{1}{\lambda_{m+1}^{2}}\frac{cL_{m}}{\nu}\tilde{K}_{0}^{2}(\alpha^{2}|A^{3/2}u^{\alpha}|^{2}), \qquad (87)$$

where $L_m = 1 + \log(\frac{\lambda_m}{\lambda_1})$, and in the second inequality above we use the fact that $||u_m^{\alpha}|| \leq \tilde{K}_0$ from Proposition 6.

Plugging (84), (85), (86) and (87) into the inequality (83) we obtain

$$\begin{aligned} \frac{d}{dt} |\delta_m|^2 + \nu \|\delta_m\|^2 &\leq \frac{c}{\nu} |\delta_m|^2 (\tilde{K}_0^2 + \frac{1}{\lambda_{m+1}} (\alpha^2 |A^{3/2} u^\alpha|^2) + \frac{K_0^4}{\nu^2 \lambda_1}) \\ &+ \frac{cL_m}{\nu} \tilde{K}_0^2 |q_m|^2 + \alpha^2 \frac{cL_m}{\nu} |q_m|^2 (\alpha^2 |A^{3/2} u^\alpha|^2) \\ &+ \frac{1}{\lambda_{m+1}^2} \frac{cL_m}{\nu} \tilde{K}_0^2 (\alpha^2 |A^{3/2} u^\alpha|^2) \\ &\leq |\delta_m|^2 U + L_m V, \end{aligned}$$

where

$$U(t) := \frac{c}{\nu} (\tilde{K}_0^2 + \frac{1}{\lambda_{m+1}} (\alpha^2 |A^{3/2} u^{\alpha}|^2) + \frac{K_0^4}{\nu^2 \lambda_1}),$$
(88)

and

$$V(t) := \frac{c}{\nu} \tilde{K}_0^2 |q_m|^2 + \alpha^2 \frac{c}{\nu} |q_m|^2 (\alpha^2 |A^{3/2} u^\alpha|^2) + \frac{1}{\lambda_{m+1}^2} \frac{c}{\nu} \tilde{K}_0^2 (\alpha^2 |A^{3/2} u^\alpha|^2).$$
(89)

Applying Gronwall inequality and recalling that $|\delta_m(0)| = 0$, we obtain

$$|\delta_m(t)|^2 \le L_m \int_0^T e^{\int_\tau^t U(s)ds} V(\tau) \, d\tau, \, t \in [0,T].$$
(90)

By the *a priori* estimates in Proposition 6, we know that

$$\nu \int_0^T (\alpha^2 |A^{3/2} u^{\alpha}|^2) \, dt \le \tilde{K}_0^2, \tag{91}$$

which implies that $U \in L^1(0,T)$ and consequently we have

$$\int_{0}^{T} U(t) dt \leq \frac{c}{\nu} (\tilde{K}_{0}^{2}T + \frac{1}{\nu\lambda_{m+1}}\tilde{K}_{0}^{2} + \frac{\tilde{K}_{0}^{4}}{\nu^{2}\lambda_{1}}T) =: \tilde{U}.$$
(92)

Now for estimating V(t), we recall from (82) that $|q_m|^2 \leq \frac{1}{\lambda_{m+1}^2}(Q+R)$. Applying (91), we obtain

$$\int_{0}^{T} V(t) dt \leq \frac{1}{\lambda_{m+1}^{2}} \left(\frac{c}{\nu} \tilde{K}_{0}^{2} (Q+R) T + \frac{1}{\lambda_{m+1}} \frac{c}{\nu^{2}} \tilde{K}_{0}^{2} (Q+R) + \frac{c}{\nu^{2}} \tilde{K}_{0}^{4} \right) =: \frac{1}{\lambda_{m+1}^{2}} \tilde{V}.$$
(93)

Plugging back into the inequality (90), we get

$$\delta_m(t)|^2 \le \frac{L_m}{\lambda_{m+1}^2} \tilde{U}\tilde{V}.$$
(94)

Now by Pythagorean Theorem and orthogonality of the projection spaces P_m and $(I - P_m)$, we have

$$\sup_{0 \le t \le T} |u^{\alpha}(t) - u^{\alpha}_{m}(t)|^{2} \le \sup_{0 \le t \le T} |\delta_{m}|^{2} + \sup_{0 \le t \le T} |q_{m}|^{2} \le \frac{1}{\lambda_{m+1}^{2}} (Q + R + L_{m} \tilde{U} \tilde{V}),$$
(95)

where Q, R and \tilde{U}, \tilde{V} are given in (80), (81), (92) and (93), respectively.

Remark 8. The above result $|u^{\alpha}(t) - u_m^{\alpha}(t)| = O\left(\frac{1}{\lambda_m L^2} (\log(\lambda_m L^2))^{1/2}\right)$ is of the same order as that of the error estimates for the usual Galerkin approximation of NSE. Indeed, in [44] the author points out that for the 2D NSE, the error estimate for the usual Galerkin, $|u - u_m|$ is of order $O\left(\frac{1}{\lambda_m L^2} (\log(\lambda_m L^2))^{1/2}\right)$, where u is the solution of 2D NSE (22) and u_m is the solution of the corresponding Galerkin system (23). Furthermore, this estimate is optimal up to the logarithmic terms provided $f \in L^2$.

Now, simply by applying the triangle inequality, we achieve the error estimates of the solution u_m^{α} for the finite-dimensional Galerkin system of the 2D Leray- α model as approximation of the exact solution u of the 2D NSE system.

Theorem 17. Let T > 0 and let u be the solution of the 2D NSE system (22) with initial data $u_0 \in D(A)$ and u_m^{α} be the solution of finite-dimensional Galerkin approximation of the 2D Leray- α system (73) with initial data $u_{0m} = P_m u_0$ over the interval [0, T]. For a given $m \ge 1$, assume that α is small enough such that $\alpha \le \frac{2\pi}{\lambda_{m+1}L}$, where λ_{m+1} is the (m+1)-th eigenvalue of the Stokes operator A, then

$$\sup_{1 \le t \le T} |u(t) - u_m^{\alpha}(t)|^2 \le C\left(\left(\frac{(2\pi)^2}{\lambda_{m+1}L^2}\right)^2 \log\left(\frac{\lambda_{m+1}L^2}{(2\pi)^2}\right) \right),\tag{96}$$

where C is a constant, which depends on ν , u_0 , f, L and T only.

Proof. By the triangle inequality we have

$$|u(t) - u_m^{\alpha}(t)|^2 \le 2(|u(t) - u^{\alpha}(t)|^2 + |u^{\alpha}(t) - u_m^{\alpha}(t)|^2).$$
(97)

From Theorem 12 we obtain, under the assumption that $\alpha \leq \frac{L}{2\pi}$,

$$\sup_{0 \le t \le T} |u(t) - u^{\alpha}(t)|^2 \le C' \left(\left(\frac{2\pi\alpha}{L}\right)^2 \log\left(\frac{L}{2\pi\alpha}\right) \right), \tag{98}$$

where C' is a constant that depends only on ν, u_0, f, L and T. By virtue of Theorem 16 above, we have, under the assumption that $\alpha^2 \leq \frac{1}{\lambda_{m+1}^2}$,

$$\sup_{0 \le t \le T} |u^{\alpha}(t) - u^{\alpha}_m(t)|^2 \le \tilde{C}\left(\left(\frac{(2\pi)^2}{\lambda_{m+1}L^2}\right)^2 \log\left(\frac{\lambda_{m+1}L^2}{(2\pi)^2}\right)\right),\tag{99}$$

where \tilde{C} is a constant that depends only on ν, u_0, f, L and T. Now we assume that $\alpha \leq \frac{2\pi}{\lambda_{m+1}L}$. Noticing that $\lambda_1 = (\frac{2\pi}{L})^2$, we have

$$\alpha^{2} \leq \left(\frac{2\pi}{\lambda_{m+1}L}\right)\left(\frac{2\pi}{\lambda_{m+1}L}\right) = \frac{\lambda_{1}}{\lambda_{m+1}}\frac{1}{\lambda_{m+1}} \leq \frac{1}{\lambda_{m+1}},\tag{100}$$

where in the last step we use the fact that $\lambda_1 \leq \lambda_{m+1}$.

Since both assumptions for (98) and (99) are satisfied, we combine (97), (98) and (99) to obtain

$$\sup_{0 \le t \le T} |u(t) - u_m^{\alpha}(t)|^2 \le C\left(\left(\frac{(2\pi)^2}{\lambda_{m+1}L^2}\right)^2 \log\left(\frac{\lambda_{m+1}L^2}{(2\pi)^2}\right)\right),$$

which concludes our proof.

Recall that, according to (1), the (m + 1)-th eigenvalue of the 2D Stokes operator satisfies

$$\frac{m+1}{c_0} \le \frac{\lambda_{m+1}}{\lambda_1} \le c_0(m+1),\tag{101}$$

where c_0 is a constant that depends only on L. Applying the above asymptotic estimate and the fact $\lambda_1 = (\frac{2\pi}{L})^2$, one can rewrite the conditions and results of Theorem 17 in the form of the following corollary:

Corollary 18. Let T > 0 and let u be the solution of the 2D NSE system (22) with initial data $u_0 \in D(A)$ and u_m^{α} be the solution of finite-dimensional Galerkin approximation of the 2D Leray- α system (73) with initial data $u_{0m} = P_m u_0$ over the interval [0, T]. For a given $m \ge 1$, assume that α is small enough such that $\alpha \le \frac{L}{2\pi} \frac{c_0}{m+1}$, where c_0 is a constant that depends only on L as stated in (1), then

$$\sup_{0 \le t \le T} |u(t) - u_m^{\alpha}(t)|^2 \le C_0(\frac{1}{m+1})^2 \log(m+1)$$
(102)

where C_0 is a constant, which depends on ν, u_0, f, L and T only.

The error estimates for the other α -models are not presented in this paper in detail. The results are the same as that of the Leray- α model we present above. One can follow the same idea and check the calculation. In fact, the proof of the estimates for the Modified Leray- α model and simplified Bardina model will follow readily from the proof of the Leray- α model since the former two have milder nonlinearity and the estimates on the nonlinear terms will be easier compared to that of the Leray- α model. For the 2D Navier-Stokes- α equation, combining (12)-(14) and (18), we can get similar inequalities for \tilde{B} as those of B and by following same steps obtain similar error estimates for the Navier-Stokes- α model. In a subsequent paper we will present error estimates regarding the three-dimensional case in the spirit of the results reported in [9].

APPENDIX

For the sake of completeness, we present in this section the version of the Brezis-Gallouet inequality presented in Proposition 3.

Proof. Let $\varphi \in D(A)$ such that

$$\varphi = \sum_{k \in \mathbb{Z}^2 \setminus (0,0)} \hat{\varphi}_k e^{i2\pi \frac{x \cdot k}{L}}$$

therefore,

$$\begin{aligned} \|\varphi\|_{L^{\infty}} &\leq \sum_{k \in \mathbb{Z}^{2} \setminus (0,0)} |\hat{\varphi}_{k}| = \sum_{0 < |k| \leq M} |\hat{\varphi}_{k}| + \sum_{|k| > M} |\hat{\varphi}_{k}| \\ &= \sum_{0 < |k| \leq M} \frac{(1+|k|^{2})^{1/2}}{(1+|k|^{2})^{1/2}} |\hat{\varphi}_{k}| + \sum_{|k| > M} \frac{(1+|k|^{2})}{(1+|k|^{2})} |\hat{\varphi}_{k}|, \end{aligned}$$

26

where $M \ge 1$ is a real number to be determined later. Then by the Cauchy-Schwarz inequality we have

$$\begin{split} \|\varphi\|_{L^{\infty}} &\leq \left(\sum_{0<|k|\leq M} (1+|k|^2) |\hat{\varphi}_k|^2\right)^{1/2} \left(\sum_{0<|k|\leq M} \frac{1}{(1+|k|^2)}\right)^{1/2} \\ &+ \left(\sum_{|k|>M} (1+|k|^2)^2 |\hat{\varphi}_k|^2\right)^{1/2} \left(\sum_{|k|>M} \frac{1}{(1+|k|^2)^2}\right)^{1/2} \\ &\leq c \|\varphi\| \left(\int_{|y|\leq M} \frac{dy}{1+|y|^2}\right)^{1/2} + c \frac{L}{2\pi} |A\varphi| \left(\int_{|y|\geq M} \frac{dy}{(1+|y|^2)^2}\right)^{1/2}. \end{split}$$

We observe that in two dimensions

$$\int_{|y| \le M} \frac{dy}{1+|y|^2} = \int_0^{2\pi} d\theta \int_0^M \frac{r \, dr}{1+r^2} = \pi \log(1+M^2),$$

and

$$\int_{|y|\ge M} \frac{dy}{(1+|y|^2)^2} = \int_0^{2\pi} d\theta \int_M^\infty \frac{r \, dr}{(1+r^2)^2} = \pi \frac{1}{1+M^2}.$$

Therefore,

$$\|\varphi\|_{L^{\infty}} \le c \left(\|\varphi\| \left(\log(1+M^2) \right)^{1/2} + \frac{L}{2\pi} \frac{|A\varphi|}{(1+M^2)^{1/2}} \right)$$

for every $M \ge 1$. Notice that by Poincaré inequality we have $\frac{L}{2\pi} \frac{|A\varphi|}{\|\varphi\|} \ge 1$, therefore, one can choose $M = \sqrt{\left(\frac{L}{2\pi} \frac{|A\varphi|}{\|\varphi\|}\right)^2 - 1} + 1$ to conclude the proof.

Acknowledgements

This work was supported in part by the NSF grants no. DMS-0504619 and no. DMS-0708832, and by the ISF grant no. 120/06.

References

- [1] R. A. Adams, Sobolev Spaces, Academic Press, New York (1975).
- [2] J. Bardina, J. H. Ferziger and W. C. Reynolds, Improved subgrid scale models for large eddy simulation, American Institute of Aeronautics and Astronautics, paper 80 (1980) AIAA, 80-1357.
- [3] A.-C. Bennis, R. Lewandowski and E. S. Titi, Simulations de l'écoulement turbulent marin ave un modéle de déconvolution, C. R. A. S., Paris, Série I 347 (2009), 445-450.
- [4] H. S. Bhat and R. C. Fetecau, A Hamiltonian regularization of the Burgers equation, J. Nonlinear Sci. 16 (2006), 615-638.
- [5] H. Brezis and T. Gallouet, Nonlinear Schrödinger evolution equations, Nonlinear Analysis, Theory, Methods & Applications 4 (1980), 677-681.
- [6] Y. Cao, E. M. Lunasin and E. S. Titi, Global well-posedness of three-dimensional viscous and inviscid simplified Bardina turbulence models, Commun. Math. Sci. 4 (2006), 823-848.
- [7] Y. Cao, Z. H. Musslimani and E. S. Titi, Nonlinear Schrödinger-Helmholtz equation as numerical regularization of the nonlinear Schrödinger equation, Nonlinearity 21 (2008), 879-898.
- Y. Cao, Z. H. Musslimani and E. S. Titi, Modulation theory for self-focusing in the nonlinear Schrödinger-Helmholtz equation, Numerical Functional Analysis and Optimization 30 (2009), 46–69.
- [9] L. Chen, R. B. Guenther, S. Kim, E. A. Thomann and E. C. Waymire, A rate of convergence for the LANSα regularization of Navier-Stokes equations, Journal of Mathematical Analysis and Applications 348 (2008), 637-649.
- [10] S. Chen, C. Foias, D. D. Holm, E. Olson, E. S. Titi and S. Wynne, Camassa-Holm equations as a closure model for turbulent channel and pipe flow, Phys. Rev. Lett. 81 (1998), 5338-5341.
- [11] S. Chen, C. Foias, D. D. Holm, E. Olson, E. S. Titi and S. Wynne, A connection between the Camassa-Holm equations and turbulent flows in channels and pipes, Phys. Fluids 11 (1999), 2343-2353.

Y.CAO AND E.S. TITI

- [12] S. Chen, C. Foias, D. D. Holm, E. Olson, E. S. Titi and S. Wynne, The Camassa-Holm equations and turbulence, Phys. D 133 (1999), 49-65.
- [13] S. Chen, D. D. Holm, L. G. Margolin and R. Zhang, Direct numerical simulcations of the Navier-Stokes alpha model, Phys. D 133 (1999), 66-83.
- [14] V. V. Chepyzhov, E. S. Titi and M. I. Vishik, On the convergence of solutions of the Leray-α model to the trajectory attractor of the 3D Navier-Stokes system. Journal of Discrete and Continuous Dynamical Systems - Series A 17 (2007), 33-52.
- [15] A. Cheskidov, D. D. Holm, E. Olson and E. S. Titi, On a Leray-alpha model of turbulence, Roy. Soc. A 461 (2005), 629-649.
- [16] P. Constantin and C. Foias, Navier-Stokes Equations, The University of Chicago Press, 1988.
- [17] C. Devulder, M. Marion and E. S. Titi, On the rate of convergence of the nonlinear Galerkin methods, Mathematics of Computation 60 (1993), 495-514.
- [18] C. Foias, D. D. Holm and E. S. Titi, The Navier-Stokes-α model of fluid turbulence, Phys. D 152 (2001), 505-519.
- [19] C. Foias, D. D. Holm and E. S. Titi, The three dimensional viscous Camassa-Holm equations and their relation to the Navier-Stokes equations and turbulence theory, J. of Dynamics and Differential Equations 14 (2002), 1-35.
- [20] B. Geurts and D. D. Holm, Leray and LANS- α modeling of turbulent mixing, Journal of Turbulence 7 (2006), 1-33.
- [21] B. Geurts and D. D. Holm, Regularization modeling for large-eddy simulation Phys. Fluids 15 (2003), L13-L16.
- [22] B. Geurts, A. Kuczaj and E. S. Titi, Regularization modeling for large-eddy simulation of homogeneous isotropic decaying turbulence, Journal of Physics A 41 (2008), 344008.
- [23] J. G. Heywood, An error estimate uniform in time for spectral Galerkin approximations of the Navier-Stokes problem, Pacific J. Math. 98 (1982), 333-345.
- [24] D. D. Holm and B. Nagida, Modeling mesoscale turbulence in the barotropic double-gyre circulation, J. Phys. Oceanogr. 33 (2003), 2355-2365.
- [25] D. D. Holm and E. S. Titi, Computational models of turbulence: the LANS- α model and the role of global analysis, Feature Article: SIAM News **38(7)** (2006).
- [26] A. Ilyin, E. M. Lunasin and E. S. Titi, A modified-Leray-α subgrid scale model of turbulence, Nonlinearity 19 (2006), 879-897.
- [27] W. Layton and R. Lewandowski, On a well-posed turbulence model, Discrete and Continuous Dyn. Sys. B 6 (2006), 111-128.
- [28] W. Layton and R. Lewandowski, A high accuracy Leray-deconvolution model of turbulence and its limiting behavior, Analysis and Applications 6 (2008), 23-49.
- [29] J. Leray, Eassi sur le mouvement d'un fluide viqueux emplissant l'espace, Acta Math. 63 (1934), 193-248.
- [30] J. S. Linshiz and E. S. Titi, Analytical study of certain magnetophydrodynamic-α models, J. Math. Phys. 48 (2007), 065504.
- [31] E. M. Lunasin, S. Kurien and E. S. Titi, Spectral scaling of the Leray-α model for two-dimensional turbulence, Journal of Physics A: Mathematical and Theoretical 41 (2008), 344014.
- [32] E. M. Lunasin, S. Kurien, M. Taylor and E. S. Titi, A study of the Navier-Stokes- α model for two-dimensional turbulence, Journal of Turbulence 8 (2007), 1-21.
- [33] K. Mohseni, B. Kosović, S. Schkoller and J. E. Marsden, Numerical simulations of the Lagrangian averaged Navier-Stokes equations for homogeneous isotropic turbulence, Phys. of Fluids 15 (2003), 524-544.
- [34] P. D. Mininni, D. C. Montgomery and A. G. Pouquet, Numerical solutions of the three-dimensional magnetohydrodynamic alpha-model, Phys. Rev. E 71 (2005), 046304.
- [35] P. D. Mininni, D. C. Montgomery and A. G. Pouquet, A numerical study of the alpha model for two-dimensional magnetohydrodynamic turbulent flows, Phys. Fluids 17 (2005), 035112-035112-17.
- [36] B. Nadiga and S. Skoller, Enhancement of the inverse-cascade of energy in the two-dimensional Lagrangian-averaged Navier-Stokes equations, Phys. of Fluids 13 (2001), 1528-1531.
- [37] E. Olson and E. S. Titi, Viscosity versus vorticity stretching: global well-posedness for a family of Navier-Stokesalpha-like models, Nonlinear Analysis: Theory, Methods and Applications 66 (2007), 2427–2458.
- [38] R. Rautmann, Eine Fehlerschranke für Galerkin approximationen lokaler Navier-Stokes-Lösungen, Constructive Methods for Nonlinear Boundary Value Problems and Nonlinear Oscillations, ISNM 48 (1979), 110-125, Birkhäuser, Basel.
- [39] R. Rautmann, On the convergence rate of nonstationary Navier-Stokes approximations, Proc. IUTAM Symp, Paderborn (W.-Germany) (1979), Lecture Notes in Math. 771 (1980), 425-449, Springer-Verlag, Berlin and New York.
- [40] G. Sell and Y. You, Dynamics of Evolutionary Equations, Springer-Verlag, New York, 2002.
- [41] R. Temam, Navier-Stokes Equations, Theory and Numerical Analysis, Third Edition, North-Holland, 2001.
- [42] R. Temam, Infinite-Dimensional Dynamical Systems in Mechanics and Physics, Applied Mathematical Sciences 68, Springer-Verlag, New York, 1988.
- [43] E.S. Titi, On a criterion for locating stable stationary solutions to the Navier-Stokes equations, Nonlinear Analysis, Theory, Methods and Applications 11 (1987), 1085–1102.

[44] E. S. Titi, On approximate inertial manifolds to the Navier-Stokes equations, J. Math. Anal. App. 149 (1990), 540-557.

[45] M. I. Vishik, E. S. Titi and V. V. Chepyzhov, On convergence of trajectory attractors of the 3D Navier-Stokes- α model as α approaches 0, Sb. Math. **198** (2007), 1703-1736.

(Y.Cao) DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA, IRVINE, CA 92697-3875, USA *E-mail address:* ycao@math.uci.edu

(E.S. Titi) DEPARTMENT OF MATHEMATICS, AND DEPARTMENT OF MECHANICAL AND AEROSPACE ENGINEERING, UNI-VERSITY OF CALIFORNIA, IRVINE, CA 92697-3875, USA, **ALSO**, DEPARTMENT OF COMPUTER SCIENCE AND APPLIED MATHEMATICS, WEIZMANN INSTITUTE OF SCIENCE, REHOVOT 76100, ISRAEL

E-mail address: etiti@math.uci.edu and edriss.titi@weizmann.ac.il