

THERMODYNAMIC LIMIT FOR THE MALLOWS MODEL ON S_n

SHANNON STARR

ABSTRACT. The Mallows model on S_n is a probability distribution on permutations, $q^{d(\pi,e)}/P_n(q)$, where $d(\pi,e)$ is the distance between π and the identity element, relative to the Coxeter generators. Equivalently, it is the number of inversions: pairs (i,j) where $1 \leq i < j \leq n$, but $\pi_i > \pi_j$. Analyzing the normalization $P_n(q)$, Diaconis and Ram calculated the mean and variance of $d(\pi,e)$ in the Mallows model, which suggests the appropriate $n \rightarrow \infty$ limit has q_n scaling as $1 - \beta/n$. We calculate the distribution of the empirical measure in this limit, $u(x,y) dx dy = \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=1}^n \delta_{(i,\pi_i)}$. Treating it as a mean-field problem, analogous to the Curie-Weiss model, the self-consistent mean-field equations are $\frac{\partial^2}{\partial x \partial y} \ln u(x,y) = 2\beta u(x,y)$, which is an integrable PDE, known as the hyperbolic Liouville equation. The explicit solution also gives a new proof of formulas for the blocking measures in the weakly asymmetric exclusion process, and the ground state of the $\mathcal{U}_q(\mathfrak{sl}_2)$ -symmetric XXZ ferromagnet.

Keywords: Mallows model, random permutation, Liouville equation, ASEP, XXZ model.

MCS numbers: 82B05, 82B10, 60B15

1. INTRODUCTION AND MAIN RESULTS

The Coxeter generators of the symmetric group S_n are the transpositions $(1,2), (2,3), \dots, (n-1,n)$. The height of a permutation is defined distance to the identity element e ,

$$d(\pi,e) = \min\{k \geq 0 : \exists \tau_1, \dots, \tau_k \in \{(1,2), \dots, (n-1,n)\} \text{ such that } \pi = \tau_1 \cdots \tau_k\}.$$

More generally, $d(\pi_1, \pi_2) = d(\pi_2^{-1}\pi_1, e)$. It is easy to see that $d(\pi,e) = d(\pi^{-1}, e)$. In fact, another formula is

$$d(\pi,e) = |\{(i,j) \in \mathbb{Z}^2 : 1 \leq i < j \leq n \text{ and } \pi_i > \pi_j\}|.$$

The pairs (i,j) are called inversions of π . In [8], Diaconis and Ram studied the Mallows measure, which is a probability measure on S_n given by

$$\mathbb{P}_n^q(\pi) = \frac{q^{d(\pi,e)}}{P_n(q)},$$

with $P_n(q)$ being a normalization constant. Actually, Diaconis and Ram studied a Markov chain on S_n for which the Mallows model gives the limiting distribution. This was followed up by another paper on a related topic by Benjamini, Berger, Hoffman and Mossel (BBHM) [3] who related the biased shuffle and the Mallows model to the asymmetric exclusion process and the “blocking” measures (of Liggett, see [17], Chapter VIII, especially Example 2.8 and the end of Section 3). They did this using Wilson’s height functions [20]. We will discuss this more in Section 8. For now, let it suffice that Diaconis and Ram identified the explicit formula for the normalization which they remarked is the “Poincaré polynomial”:

$$P_n(q) = \prod_{i=1}^n \left(\frac{q^i - 1}{q - 1} \right) = [n]_q! = [n]_q \cdots [1]_q, \quad \text{where } [n]_q = \frac{q^n - 1}{q - 1}.$$

Further references for the statistical applications¹ of the Mallows model can be found in their paper.

Date: March 2, 2009.

¹Independently, a similar q -deformed combinatorial formula was explained for a problem in quantum statistical mechanics, the ground state of the ferromagnetic $\mathcal{U}_q(\mathfrak{sl}_2)$ -symmetric XXZ quantum spin chain, by Bolina, Contucci and Nachtergaele [4]. We will comment more on this in Section 8.

Note that, physically speaking, one would define a Hamiltonian energy function $H_n : S_n \rightarrow \mathbb{R}$ as

$$H_n(\pi) = \frac{1}{n-1} \sum_{1 \leq i < j \leq n} \mathbb{1}_{(0,\infty)}(\pi_i - \pi_j).$$

In this case, one thinks of $\pi = (\pi_1, \dots, \pi_n)$ as some type of constrained spin system, where each of the components π_1, \dots, π_n are spins in $\{1, \dots, n\}$ as in a Potts model. The choice of the normalization of the Hamiltonian is then standard for mean-field models. One would be most interested in the free energy

$$f_n(\beta) = -\frac{1}{\beta n} \ln \sum_{\pi \in S_n} e^{-\beta H_n(\pi)}.$$

For our purposes, we prefer to consider the mathematically simpler ‘‘pressure’’

$$p_n(\beta) = \frac{1}{n} \ln \left(\frac{1}{n!} \sum_{\pi \in S_n} e^{-\beta H_n(\pi)} \right).$$

(Note that, contrary to the usual conventions of statistical physics, we have divided the partition function, which is $Z_n(\beta) = \sum_{\pi \in S_n} e^{-\beta H_n(\pi)}$, by the infinite-temperature partition function $Z_n(0) = n!$, which is equivalent to starting with a normalized *a priori* measure rather than counting measure on S_n .) It is trivial to see that this is given precisely by the Poincaré polynomial described by Diaconis and Ram:

$$p_n(\beta) = \frac{1}{n} \ln \frac{P_n(e^{-\beta/(n-1)})}{n!} = \frac{1}{n} \ln \frac{[n]_{e^{-\beta/(n-1)}}!}{n!}.$$

With this scaling, it is also easy to calculate the limit:

$$p(\beta) = \lim_{n \rightarrow \infty} p_n(\beta) = \int_0^1 \ln \left(\frac{1 - e^{-\beta x}}{\beta x} \right) dx,$$

(which can be solved explicitly using the polylogarithm function). From this one can calculate the mean and variance in the limiting Gibbs measure. For instance, one can calculate $\mathbb{E}(d(\pi, e) - n(n-1)/4)^2 \sim n^3/72$ in the uniform measure on S_n .

To go beyond the statistics of $d(\pi, e)$ it seems worthwhile to study the empirical measure of π :

$$\frac{1}{n} \sum_{i=1}^n \delta_{(i, \pi_i)},$$

which is a normalized measure on $\{1, \dots, n\} \times \{1, \dots, n\}$. More specifically, this is a random measure. Rescaling the discrete cube $\{1, \dots, n\} \times \{1, \dots, n\}$ to $[0, 1] \times [0, 1]$, it is easy to see that the random empirical measure converges, in probability, to the non-random Lebesgue measure, when $\beta = 0$ (the uniform case). Our main theorem generalizes this result.

Theorem 1.1. *For any $\beta \in \mathbb{R}$,*

$$\lim_{\epsilon \downarrow 0} \lim_{n \rightarrow \infty} \mathbb{P}_n^{1-\beta/n} \left\{ \left| \frac{1}{n} \sum_{i=1}^n f(i/n, \pi_i/n) - \int_{[0,1] \times [0,1]} f(x, y) u(x, y) dx dy \right| > \epsilon \right\} = 0,$$

for every continuous function $f : [0, 1] \times [0, 1] \rightarrow \mathbb{R}$, where

$$u(x, y) = \frac{(\beta/2) \sinh(\beta/2)}{(e^{\beta/4} \cosh(\beta[x-y]/2) - e^{-\beta/4} \cosh(\beta[x+y-1]/2))^2}.$$

Note that (one can show) the limit $\beta \rightarrow 0$ gives 1. The proof of Theorem 1.1 uses a rigorous version of mean-field theory, as in the solution of the Curie-Weiss model. An interesting feature is that the self-consistent

mean-field equation leads us to the characterization of u as the solution of an integrable PDE

$$\frac{\partial^2}{\partial x \partial y} \ln u(x, y) = 2\beta u(x, y).$$

It is not unusual for mean-field problems to lead to integrable PDE's. We demonstrate this briefly in the next section with the ubiquitous toy model, the Curie-Weiss ferromagnet.

2. TOY MODEL: THE CURIE-WEISS FERROMAGNET

We include this section merely to point out that mean-field problems often do lead to integrable PDE. However the issue is serious: in fact there is a recent paper by Genovese and Barra which we recommend for more details [13]. Our approach merely summarizes their results (in our own words) as well as the earlier paper by Barra, himself [2]. The configuration space of the CW model is $\Omega_N = \{+1, -1\}^N = \{\sigma = (\sigma_1, \dots, \sigma_n) : \sigma_1, \dots, \sigma_n = \pm 1\}$. For technical reasons, we choose the Hamiltonian as

$$H_N(\sigma, t, x) = -\frac{t}{2N} \sum_{i,j=1}^N \sigma_i \sigma_j - x \sum_{i=1}^N \sigma_i,$$

we assume $t \geq 0$ and $x \in \mathbb{R}$. Defining $m_N(\sigma) = N^{-1} \sum_{i=1}^N \sigma_i$, which takes values in $[-1, 1]$, we see that

$$H_N = -N \left(\frac{tm_N^2}{2} + xm_N \right).$$

Therefore, defining

$$p_N(t, x) = \frac{1}{N} \ln \sum_{\sigma \in \Omega_N} e^{-H_N(\sigma, t, x)},$$

we easily see that

$$\frac{\partial}{\partial t} p_N(t, x) = \frac{1}{2} \langle m_N^2 \rangle_{N, t, x},$$

and

$$\frac{\partial^2}{\partial x^2} p_N(t, x) = N (\langle m_N^2 \rangle - \langle m_N \rangle^2),$$

where

$$\langle f \rangle = \langle f \rangle_{N, t, x} = \frac{\sum_{\sigma \in \Omega_N} f(\sigma) e^{-H_N(\sigma, t, x)}}{\sum_{\sigma \in \Omega_N} e^{-H_N(\sigma, t, x)}}.$$

Actually it is easier to consider the ‘‘order parameter,’’

$$u_N(t, x) = \langle m_N \rangle_{N, t, x} = \frac{\partial}{\partial x} p_N(t, x),$$

from which $p_N(t, x)$ can be calculated by solving the ODE:

$$\begin{cases} \frac{\partial}{\partial x} p_N(t, x) = u_N(t, x) & \text{for } x \in \mathbb{R}, \\ p_N(t, x) - |x| \rightarrow \frac{1}{2}t^2 & \text{as } x \rightarrow \pm\infty. \end{cases}$$

Then we see that $u_N(t, x)$ satisfies the viscous Burgers equation (with velocity equal to the negative amplitude):

$$\begin{cases} \frac{\partial}{\partial t} u_N(t, x) = u_N(t, x) \frac{\partial}{\partial x} u_N(t, x) + \frac{1}{2N} \cdot \frac{\partial^2}{\partial x^2} u_N(t, x) & \text{for } t > 0 \text{ and } x \in \mathbb{R}, \\ u_N(0, x) = \tanh(x) & \text{for } x \in \mathbb{R}. \end{cases}$$

This is an integrable PDE, using the Cole-Hopf transform. See, for instance, Chapter 4 of Whitham, [19]. Actually, this leads to a solution in terms of Gaussian integrals. The analogous transform in spin-configuration notation is the Hubbard-Stratonovich transform:

$$e^{Ntm^2/2} = \int_{-\infty}^{\infty} \frac{e^{Nt(mx-x^2/2)}}{\sqrt{2\pi/t}} dx,$$

which “linearizes” the dependence of the Hamiltonian on m_N , in the exponential. This trick is used to solve the Curie-Weiss model. See, for example, Thompson [18].

Note that in the $N \rightarrow \infty$ limit, one obtains $u(t, x) = \lim_{N \rightarrow \infty} u_N(t, x)$ being the vanishing-viscosity solution of the inviscid Burgers equation. Shocks correspond to phase transitions. The Lax-Oleinik variational formula for solutions of hyperbolic conservation laws applies. See for example, Section 3.4.2 of Evans [11]. In this context we claim that this is equivalent to the Gibbs variational formula, in the mean-field limit. We review this next.

3. GIBBS VARIATIONAL FORMULA

Let us begin by considering a general problem in classical statistical mechanics. Suppose that \mathcal{X} is a compact metric space, and suppose that there is a two-body interaction

$$h : \mathcal{X} \times \mathcal{X} \rightarrow \mathbb{R} \cup \{+\infty\}.$$

We assume that h is bounded below. Then for each $N \geq 2$, one can consider the mean-field Hamiltonian $H_N : \mathcal{X}^N \rightarrow \mathbb{R} \cup \{+\infty\}$

$$H_N(x_1, \dots, x_N) = \frac{1}{N-1} \sum_{1 \leq i < j \leq N} h(x_i, x_j).$$

Suppose that there is also an *a priori* measure μ_0 on \mathcal{X} , which we assume is normalized so that

$$\int_{\mathcal{X}} d\mu_0(x) = 1.$$

Then the thermodynamic quantities are the partition function,

$$Z_N(\beta) = \int_{\mathcal{X}^N} e^{-\beta H_N(x_1, \dots, x_N)} d\mu_0(x_1) \cdots d\mu_0(x_N),$$

the pressure,

$$p_N(\beta) = \frac{1}{N} \ln Z_N(\beta),$$

and the Boltzmann-Gibbs measure

$$d\mu_N^\beta(x_1, \dots, x_N) = \frac{e^{-\beta H_N(x_1, \dots, x_N)}}{Z_N(\beta)} d\mu_0(x_1) \cdots d\mu_0(x_N).$$

Physically, it is more correct to consider the free energy rather than the pressure, $f_N(\beta) = -\frac{1}{\beta} p_N(\beta)$. But we will consider $p_N(\beta)$, which seems slightly easier to handle, mathematically.

We will write μ_0^N for the measure $d\mu_0^N(x_1, \dots, x_N) = d\mu_0(x_1) \cdots d\mu_0(x_N)$. Also, if f is a function, then we use the short-hand $\mu(f)$ for $\int f d\mu$. Then, according to the Gibbs variational principle, we have

$$(1) \quad p_N(\beta) = \max_{\mu_N \in \mathcal{M}_{+,1}(\mathcal{X}^N)} \frac{1}{N} [S_N(\mu_N, \mu_0^{\otimes N}) - \beta \mu_N(H_N)],$$

where $S_N(\mu_N, \mu_0^{\otimes N})$ is the relative entropy (and $\mathcal{M}_{+,1}(\mathcal{X}^N)$ denotes all Borel probability measures on \mathcal{X}^N)

$$S_N(\mu_N, \mu_0^{\otimes N}) = \begin{cases} \mu_0^N(\phi(d\mu_N/d\mu_0^N)) & \text{if } \mu_N \text{ is absolutely continuous with respect to } \mu_0^N, \\ -\infty & \text{otherwise,} \end{cases}$$

and $\phi(x) = -x \ln(x)$, which is 0 if $x = 0$. Also, the unique μ_N maximizing the Gibbs variational formula (the “arg-max”) is the Boltzmann-Gibbs measure μ_N^β .

A natural ansatz for the optimizing measure is $\mu_N = \mu^N$, for some measure $\mu \in \mathcal{M}_{+,1}(\mathcal{X})$. Probabilistically, this means that all the x_1, \dots, x_N are independent and identically distributed. Technically, this cannot usually be exact for finite N . But it leads to a simpler formula because

$$S_N(\mu^N, \mu_0^N) = N S_1(\mu, \mu_0) \quad \text{and} \quad \mu^N(H_N) = \frac{N}{2} \mu^2(h),$$

and one hopes that the formula may become exact in the thermodynamic limit. Mark Fannes, Herbert Spohn and Andre Verbeure proved that this approach is rigorous in the $N \rightarrow \infty$ limit [12]:

Proposition 3.1 (Fannes, Spohn, Verbeure 1978). *The limiting pressure exists, $p(\beta) = \lim_{N \rightarrow \infty} p_N(\beta)$, and solves the variational problem*

$$p(\beta) = \max_{\mu \in \mathcal{M}_{+,1}(\mathcal{X})} [S_1(\mu, \mu_0) - \frac{\beta}{2} \mu^2(h)].$$

Moreover, any subsequential limit of the sequence (μ_N^β) is a mixture of infinite product measures μ^∞ , for μ^∞ 's maximizing the right-hand-side of the formula above.

Remark 3.2. *Note that the Gibbs variational principle (1) is true in general for all Hamiltonians whether they are mean-field or not. (See, for instance, Lemma II.3.1 from Israel's monograph [15], or any other textbook on mathematical statistical mechanics, for a rigorous proof which also applies directly in the thermodynamic limit.) But the product ansatz which seems to yield the formula from the proposition is not generally valid, since there are nontrivial correlations in the true Boltzmann-Gibbs state. Nevertheless Fannes, Spohn and Verbeure proved the mean-field limit in the $N \rightarrow \infty$ limit, using de Finetti's theorem (which states that all infinitely exchangeable measures are mixtures of product states) and properties of the relative entropy.*

Because one has $\mu^2 = \mu \times \mu$, one replaces the linear form $\mu_N(H_N)$ by the nonlinear one $\mu^2(h)$. Fannes, Spohn and Verbeure actually proved their theorem more generally for quantum statistical mechanics models, such as the Dicke maser, but it also applies to classical models. For the quantum models, one replaces de Finetti's theorem by the non-commutative analogue, Störmer's theorem. (See [1] and references therein for a detailed survey of de Finetti's theorem, and refer to Fannes, Spohn and Verbeure's paper and references therein for the noncommutative analogue, which we will not need.) With Eugene Kritchevski, we tried to find a simpler proof of the specialization of Proposition 3.1 to the classical case. But there were several errors in our proof, which have been brought to my attention by Alex Opaku, to whom I am grateful. Fortunately, Fannes, Spohn and Verbeure's original paper definitely does also apply to classical models.

4. APPLICATION TO THE MALLOWS MODEL

We take for \mathcal{X} the unit square $[0, 1] \times [0, 1]$. Suppose that $f, g : [0, 1] \rightarrow \mathbb{R}$ are probability densities: $f, g \geq 0$ and $\int_0^1 f(x) dx = \int_0^1 g(y) dy = 1$. For simplicity, later on, we also assume that there are constants $0 < c < C < \infty$ such that $c \leq f, g \leq C$. Then we take the *a priori* measure to be

$$d\mu_0(x, y) = f(x)g(y) dx dy.$$

We take the interaction to be

$$h((x_1, y_1), (x_2, y_2)) = \theta(x_1 - x_2)\theta(y_2 - y_1) + \theta(x_2 - x_1)\theta(y_1 - y_2),$$

where $\theta : \mathbb{R} \rightarrow \mathbb{R}$ is the Heaviside function,

$$\theta(x) = \begin{cases} 1 & \text{if } x > 0, \\ 0 & \text{if } x < 0. \end{cases}$$

Since μ_0 is absolutely continuous with respect to Lebesgue measure, all x_1, \dots, x_N and y_1, \dots, y_N are distinct, with probability 1. (This is why we do not bother to specify θ at the discontinuity point 0.)

Let $X_1 < \dots < X_N$ and $Y_1 < \dots < Y_N$ be any points. Then for any $\sigma, \tau \in S_N$, the symmetric group, we have

$$\frac{d\mu_N^\beta}{d\mu_0^\beta}((X_{\sigma_1}, Y_{\tau_1}), \dots, (X_{\sigma_N}, Y_{\tau_N})) = \mathbb{P}_N^{\exp(-\beta/(N-1))}(\sigma^{-1}\tau),$$

where \mathbb{P}_N^β is the Mallows measure on S_N . So studying the limit of the μ_N^β 's gives us direct information on the limit of $\mathbb{P}_N^{1-\beta/N}$. For any fixed $\sigma \in S_N$, the permutation $\sigma^{-1}\tau$ is uniform on S_N , if τ is. Because of this, we have the following result for the marginal of μ_N^β on (x_1, \dots, x_N) ,

$$\int_{\mathcal{X}^N} U(x_1, \dots, x_N) d\mu_N^\beta((x_1, y_1), \dots, (x_N, y_N)) = \int_{[0,1]^N} U(x_1, \dots, x_N) f(x_1) \cdots f(x_N) dx_1 \cdots dx_N,$$

and the marginal on (y_1, \dots, y_N) ,

$$\int_{\mathcal{X}^N} U(y_1, \dots, y_N) d\mu_N^\beta((x_1, y_1), \dots, (x_N, y_N)) = \int_{[0,1]^N} U(y_1, \dots, y_N) g(y_1) \cdots g(y_N) dy_1 \cdots dy_N,$$

for all bounded, continuous functions $U : (0, 1)^N \rightarrow \mathbb{R}$. Enforcing these conditions on the marginals, Proposition 3.1 yields the following:

$$(2) \quad p(\beta) = \max_{\mu \in \mathcal{M}_{+,1}(f,g)} [S_1(\mu, \mu_0) - \beta \mu^2(h)],$$

where $\mathcal{M}_{+,1}(f, g)$ is the set of all probability measures $\mu \in \mathcal{M}_{+,1}(\mathcal{X})$ such that $d\mu(x, y)$ has marginals $f(x)dx$ and $g(y)dy$.

Suppose that μ is any arg-max of the right-hand-side of (2). Since we have chosen μ_0 to be absolutely continuous with respect to Lebesgue measure on \mathcal{X} , the same must be true of μ . Otherwise the relative entropy would be $-\infty$. So we can write

$$d\mu(x, y) = u(x, y) dx dy.$$

Then it is easy to see that the Euler-Lagrange equations for (2) are

$$(3) \quad \ln u(x, y) = \ln f(x) + \ln g(y) + C - \beta \int_{\mathcal{X}} u(x', y') [\theta(x - x')\theta(y' - y) + \theta(x' - x)\theta(y - y')] dx' dy',$$

for some constant, $C < \infty$. Therefore, u solves the equation

$$(4) \quad \begin{cases} u(x, y) = \frac{1}{Z} f(x)g(y)e^{-\beta \int_{\mathcal{X}} h((x,y),(x',y'))u(x',y') dx' dy'} & \text{for } (x, y) \in \mathcal{X}, \\ \int_0^1 u(x, y) dy = f(x) & \text{for } x \in [0, 1], \\ \int_0^1 u(x, y) dx = g(y) & \text{for } y \in [0, 1], \end{cases}$$

where Z is a normalization constant.

Since $u(x, y)$ solves an integral equation it can be differentiated both with respect to x and y . Doing so yields the partial differential equation

$$(5) \quad \frac{\partial^2}{\partial x \partial y} \ln u(x, y) = 2\beta u(x, y),$$

known as the hyperbolic Liouville equation. This equation arises naturally in differential geometry, related to the problem of choosing a metric on a given manifold. I am very grateful to S.G. Rajeev for important information regarding this PDE. One of the facts he imparted is the symmetry of the differential equation under the following general transformation:

$$(6) \quad \begin{aligned} v(x, y) &= F'(x)G'(y)u(F(x), G(y)) \\ \Rightarrow \frac{\partial^2}{\partial x \partial y} \ln v(x, y) &= \frac{\partial^2}{\partial x \partial y} u(F(x), G(y)) + \frac{\partial}{\partial y} \left(\frac{F''(x)}{F'(x)} \right) + \frac{\partial}{\partial x} \left(\frac{G''(y)}{G'(y)} \right) \\ &= \frac{\partial^2}{\partial x \partial y} u(F(x), G(y)) \\ &= 2\beta F'(x)G'(y)u(F(x), G(y)) \\ &= 2\beta v(x, y). \end{aligned}$$

So, if $\frac{\partial^2}{\partial x \partial y} \ln u = \beta u$ then the same is true for $v(x, y) = F'(x)G'(y)u(F(x), G(y))$.

Our real goal is to solve the Euler-Lagrange equation (4). But as a first step, we want to consider the Cauchy problem for (5). In other words, we want to consider the problem

$$(7) \quad \begin{cases} \frac{\partial^2}{\partial x \partial y} \ln u(x, y) = 2\beta u(x, y) & \text{for } (x, y) \in [0, L_1] \times [0, L_2], \\ u(x, 0) = \phi(x) & \text{for } x \in [0, L_1], \\ u(0, y) = \psi(y) & \text{for } y \in [0, L_2], \end{cases}$$

for some $L_1, L_2 > 0$ and $\phi : [0, L_1] \rightarrow \mathbb{R}$, $\psi : [0, L_2] \rightarrow \mathbb{R}$ both positive and continuous.

Note that $\frac{\partial^2}{\partial x \partial y}$ is a wave operator, with characteristics directed along x and y . Specifically, defining $\xi = (x+y)/\sqrt{2}$ and $\zeta = (x-y)/\sqrt{2}$, we have $\frac{\partial^2}{\partial x \partial y} = \frac{1}{2}(\frac{\partial^2}{\partial \xi^2} - \frac{\partial^2}{\partial \zeta^2})$, the usual wave operator. Therefore, D'Alembert's formula for solutions of the wave equation allow us to reformulate (7) as an integral equation,

$$(8) \quad \ln u(x, y) = \ln \phi(x) + \ln \psi(y) - \ln \alpha + 2\beta \int_{[0,x] \times [0,y]} u(x', y') dx' dy',$$

which we prefer. This equation is supposed to be solved for all $(x, y) \in [0, L_1] \times [0, L_2]$. We have introduced the number $\alpha = \phi(0)$, which we also assumed equals $\psi(0)$, for consistency since both are supposed to give $u(0, 0)$. (Note that the initial surface, $([0, L_1] \times \{0\}) \cup (\{0\} \times [0, L_2])$, is *not* a non-characteristic surface. This is the reason that our Cauchy problem does not require initial data for the tangential derivative of u even though the wave equation is second order.) We refer to Evans textbook for PDE's, (especially Section 2.4 on the wave equation and Section 4.6 on the Cauchy-Kovalevskaya theorem).

As we will see, the symmetry (6) is the key to solving both the Euler-Lagrange equation (4) and the Cauchy problem (8).

5. THE CAUCHY PROBLEM

We start with uniqueness for the Cauchy problem.

Lemma 5.1. *For any $L_1, L_2 > 0$, the Cauchy problem (8) having $\phi = \psi = \alpha = 1$ has at most one solution in the class of nonnegative integrable functions.*

Proof. Since $\phi = \psi = \alpha = 1$, equation (8) simplifies to

$$\ln u(x, y) = 2\beta \int_{[0,x] \times [0,y]} u(x', y') dx' dy'.$$

Assuming that u is nonnegative and integrable, this implies that $\ln u$ is bounded and continuous. Then, using these properties in the right-hand-side of the equation again (similarly as one does to prove elliptic regularity) we deduce that $\ln u$ is continuously differentiable and globally Lipschitz. In particular, it is continuous up to the boundary.

Now suppose that there are two solutions u and v . Letting $z = \ln u - \ln v$, we have

$$z(x, y) = 2\beta \int_0^x \int_0^y [1 - e^{-z(x', y')}] u(x', y') dx' dy'.$$

Since both $\ln u$ and $\ln v$ are bounded, we see that z is as well. Therefore, there exists a constant $K < \infty$ such that $|1 - e^{-z}| \leq K|z|$ for all values of z in the range. So we have

$$|z(x, y)| \leq \beta K \|u\|_\infty \int_0^x \int_0^y |z(x', y')| dx' dy'.$$

A version of Gronwall's lemma then implies that $z \equiv 0$. We outline this now, although our argument can probably be improved.

Let $Z(t) = \sup\{|z(x, y)| : (x, y) \in (0, L_1) \times (0, L_2), xy \leq t\}$. Then we obtain, after making the change of variables $(x, y) \mapsto (x, t)$ where $t = xy$, and using Fubini-Tonelli to integrate over x first,

$$Z(t) \leq \beta K \|u\|_\infty \int_0^t \ln(t/t') Z(t') dt'.$$

We rewrite this as

$$Z(t) \leq \beta K \|u\|_\infty \int_0^t [\ln(t/L_1 L_2) - \ln(t'/L_1 L_2)] Z(t') dt'.$$

Since $\ln(t/L_1 L_2) \leq 0$ for $t \leq L_1 L_2$, and since $Z \geq 0$, we can drop the term $\ln(t/L_1 L_2) Z(t')$ in the integrand to obtain

$$Z(t) \leq \beta K \|u\|_\infty \int_0^t |\ln(t'/L_1 L_2)| Z(t') dt'.$$

Finally, setting $\zeta(t) = \int_0^t |\ln(t'/L_1 L_2)| Z(t') dt'$, this leads to

$$\zeta'(t) \leq \beta K \|u\|_\infty |\ln(t/L_1 L_2)| \zeta(t).$$

By Gronwall's inequality (see for example Appendix B of Evans [11]), we obtain

$$\zeta(t) = e^{\beta K \|u\|_\infty \int_0^t |\ln(t'/L_1 L_2)| dt'} \zeta(0) = e^{\beta K \|u\|_\infty (1 + |\ln(t/L_1 L_2)|) t / L_1 L_2} \zeta(0).$$

But $\zeta(0) = 0$. Hence $\zeta(t) = 0$ for all t . This implies $Z(t) = 0$ for all t which implies $z(x, y) = 0$ for all x, y . \square

Next we derive the explicit solution of (8), for the case $\phi = \psi = \alpha = 1$.

Corollary 5.2. *Suppose that $L_1, L_2 > 0$ and either $\beta \leq 0$ or $L_1 L_2 < 1/\beta$. Then the unique solution of the Cauchy problem (8) with $\phi = \psi = \alpha = 1$ is*

$$u(x, y) = (1 - \beta xy)^{-2}.$$

Proof. Uniqueness was proved in Lemma 5.1, and it is trivial to check that this solves the PDE (5). Therefore, assuming that u is integrable on $[0, L_1] \times [0, L_2]$, we may derive D'Alembert's formula by standard calculus:

$$\begin{aligned} \ln u(x, y) &= \int_0^x \frac{\partial}{\partial x} \ln u(x', y) dx' + \ln \psi(y) \\ &= \int_{(0, x) \times (0, y)} \frac{\partial^2}{\partial x \partial y} \ln u(x', y') dx' dy' + \int_0^x \frac{\partial}{\partial x} \ln \phi(x') dx' + \ln \psi(y) \\ &= 2\beta \int_{(0, x) \times (0, y)} u(x', y') dx' dy' + \ln \psi(y) + \ln \phi(x) - \ln \alpha. \end{aligned}$$

The only issue is to check integrability, which amounts to checking $\inf_{(x, y) \in [0, L_1] \times [0, L_2]} 1 - \beta xy > 0$. This holds if and only if $\beta \leq 0$ or $L_1 L_2 < 1/\beta$. \square

Let us briefly explain one approach to deriving this formula. For nonlinear PDE's one always first guesses a scaling solution, in hopes of finding an explicit formula. Because of the hyperbolic nature it makes sense to look for a solution $u(x, y) = U(xy)$ for some $U(z)$. This leads to the ODE

$$\frac{d}{dz} \ln U(z) + z \frac{d^2}{dz^2} \ln U(z) = 2\beta U(z),$$

which can also be expressed as

$$\frac{d}{dz} \left(z \frac{d}{dz} \ln U(z) \right) = 2\beta U(z).$$

The idea of using a power law solution is natural because the derivative of the logarithm results in a power law, itself. Trying $U(z) = (1 + cz)^p$ leads to

$$\ln U(z) = p \ln(1 + cz) \quad \Rightarrow \quad z \frac{d}{dz} \ln \phi(z) = \frac{cpz}{1 + cz} = p - \frac{p}{1 + cz} \quad \Rightarrow \quad \frac{d}{dz} \left(z \frac{d}{dz} \ln U(z) \right) = \frac{cp}{(1 + cz)^2}.$$

So, taking $p = -2$ and $c = -\beta$, this solves the equation, and gives $U(z) = (1 - \beta z)^{-2} \Rightarrow u(x, y) = (1 - \beta xy)^{-2}$. Finally, we are led to the solution of the general Cauchy problem.

Corollary 5.3. *Suppose that $\phi, \psi : [0, 1] \rightarrow \mathbb{R}$ are continuous and satisfy $c \leq \phi, \psi \leq C$, for some constants $0 < c < C < 1$. Also suppose that $\phi(0) = \psi(0) = \alpha$ for some α . Then the Cauchy problem (8) has a solution if and only if $\beta \leq 0$ or $\int_0^1 \phi(x) dx \int_0^1 \psi(y) dy < \alpha/\beta$. In case a solution exists, it is unique and equals*

$$(9) \quad u(x, y) = \frac{\alpha \phi(x) \psi(y)}{(\alpha - \beta \Phi(x) \Psi(y))^2},$$

where $\Phi(x) = \int_0^x \phi(x') dx'$ and $\Psi(y) = \int_0^y \psi(y') dy'$.

Proof. Suppose that u is any solution of (8). Let v be given by

$$v(x, y) = \frac{u(\Phi^{-1}(\alpha^{1/2}x), \Psi^{-1}(\alpha^{1/2}y))}{\alpha \Phi'(\Phi^{-1}(\alpha^{1/2}x)) \Psi'(\Psi^{-1}(\alpha^{1/2}y))}.$$

Then, by the symmetry (6), v is a solution of Liouville's PDE (5) on the domain $[0, \alpha^{-1/2}\Phi(1)] \times [0, \alpha^{-1/2}\Psi(1)]$. But $v(x, 0) = v(0, y) = 1$ because $\Phi' = \phi$ and $\Psi' = \psi$. So uniqueness, the conditions for existence, and the formula for the solution all follow from Lemma 5.1 and Corollary 5.2. \square

6. SOLVING THE EULER-LAGRANGE EQUATION

By general principles, we know that a solution of (4) always exists: specifically, the optimizer in Proposition 3.1. Next we calculate it, and prove uniqueness.

Lemma 6.1. *If $f = g = 1$, then the unique solution of (4) is given by (9) for*

$$\phi(z) = \psi(z) = \frac{\beta e^{-\beta z}}{1 - e^{-\beta}}, \quad \Phi(z) = \Psi(z) = \frac{1 - e^{-\beta z}}{1 - e^{-\beta}} \quad \text{and} \quad \alpha = \frac{\beta}{1 - e^{-\beta}}.$$

Proof. Suppose u solves (4). Note that $\lim_{x \rightarrow 0} h((x, y), (x', y')) = \theta(y - y')$ for all $(x', y') \in \mathcal{X}$. By the dominated convergence theorem, this implies

$$\lim_{x \rightarrow 0} \int_{\mathcal{X}} h((x, y), (x', y')) u(x', y') dx' dy' = \int_0^1 \int_0^1 \theta(y - y') u(x', y') dx' dy' = \int_0^1 \theta(y - y') dy' = y,$$

where we used the fact that $\int_0^1 u(x', y') dx' = 1$ for all y' . So

$$\psi(y) = \lim_{x \rightarrow 0} u(x, y) = \frac{1}{\mathcal{Z}} e^{-\beta y}.$$

Similar arguments lead to $\phi(x) = \lim_{y \rightarrow 0} u(x, y) = \frac{1}{\mathcal{Z}} e^{-\beta x}$. Since $\int_0^1 u(x, y) dy = 1$ for all $x \in [0, 1]$, it again follows from the dominated convergence theorem, taking the limit $x \rightarrow 0$, that $\int_0^1 \psi(y) dy$ must also be 1. So $\mathcal{Z} = (1 - e^{-\beta})/\beta$. Checking, the reader will easily see that this gives the stated value for ϕ, ψ and α . Integrating, it also leads to Φ and Ψ .

Uniqueness follows from uniqueness of the Cauchy problem, Corollary 5.3. Since this is the only possible solution, and since a solution exists, this must be it. \square

Substituting in, and simplifying leads to the formula

$$(10) \quad u(x, y) = \frac{(\beta/2) \sinh(\beta/2)}{(e^{\beta/4} \cosh(\beta[x - y]/2) - e^{-\beta/4} \cosh(\beta[x + y - 1]/2))^2}.$$

Therefore, we arrive at the final formula.

Corollary 6.2. *As long as $c \leq f, g \leq C$ for some $0 < c < C < 1$, and $\int_0^1 f(x) dx = \int_0^1 g(y) dy = 1$, the unique solution of (4) is*

$$u(x, y) = \frac{(\beta/2) \sinh(\beta/2) f(x) g(y)}{(e^{\beta/4} \cosh(\beta[F(x) - G(y)]/2) - e^{-\beta/4} \cosh(\beta[F(x) + G(y) - 1]/2))^2},$$

where $F(x) = \int_0^x f(x') dx'$ and $G(y) = \int_0^y g(y') dy'$.

Proof. Suppose that u is a solution of (4) under the conditions stated. Define

$$v(x, y) = \frac{u(F^{-1}(x), G^{-1}(y))}{f(F^{-1}(x))g(G^{-1}(y))},$$

analogously to the proof of Corollary 5.3. Note that F and G are continuously, strictly increasing bijections of $[0, 1]$. Using (4), we see that

$$\begin{aligned} \ln v(x, y) &= \ln u(F^{-1}(x), G^{-1}(y)) - \ln f(F^{-1}(x)) - \ln g(G^{-1}(y)) \\ &= -\ln \mathcal{Z} - \beta \int_{\mathcal{X}} h((F^{-1}(x), G^{-1}(y)), (x', y')) u(x', y') dx' dy'. \end{aligned}$$

Making the change-of-variables $x'' = F(x')$ and $y'' = G(y')$, we see that $dx' = dx''/f(F^{-1}(x''))$ and $dy' = dy''/g(G^{-1}(y''))$. So we have

$$\ln v(x, y) = -\ln \mathcal{Z} - \beta \int_{\mathcal{X}} h((F^{-1}(x), G^{-1}(y)), (F^{-1}(x''), G^{-1}(y''))) v(x'', y'') dx'' dy''.$$

But the Heaviside function satisfies $\theta(F(x) - F(x')) = \theta(x - x')$ for any continuous, strictly increasing function F . For this reason,

$$h((F^{-1}(x), G^{-1}(y)), (F^{-1}(x''), G^{-1}(y''))) = h((x, y), (x'', y'')).$$

In other words, v also solves (4), except that

$$\int_0^1 v(x, y) dy = \int_0^1 v(x, y) dx = 1,$$

using the change-of-variables formula, again. So uniqueness and the formula follows from Lemma 6.1. \square

7. PROOF OF MAIN RESULT

We now explain the minor details needed to go from Proposition 3.1 to a proof of Theorem 1.1. According to Fannes, Spohn and Verbeure's result, μ_N^β must converge weakly to a mixture of i.i.d., product measures, each of whose 1-particle marginal optimizes $S_1(\mu, \mu_0) - \frac{\beta}{2} \mu^2(h)$. But μ_N^β has marginals on (x_1, \dots, x_N) and (y_1, \dots, y_N) equal to the product measures of $f(x) dx$ and $g(y) dy$, respectively. Therefore, according to the weak law of large numbers (WLLN), we know that all the μ 's in the support of the directing measure for the limit of μ_N^β , must have x marginal equal to $f(x) dx$ and y marginal $g(y) dy$. Hence, this constraint can be imposed when looking for an optimizer. This is actually a relevant comment because all optimizers, for all choices of *a priori* measure μ_0 , have the same value/pressure: that due to the Mallows measure on S_n . For concreteness, we will now take $f = g = 1$.

Now suppose that μ optimizes the Gibbs formula. It must be absolutely continuous with respect to μ_0 in order to not have the relative entropy equal to $-\infty$. So we can write

$$d\mu(x, y) = u(x, y) dx dy,$$

where $u(x, y)$ is absolutely continuous. Choosing any continuous function $\phi : [0, 1] \times [0, 1] \rightarrow \mathbb{R}$, with

$$\int_{\mathcal{X}} u(x, y) \phi(x, y) dx dy = 0,$$

we can take

$$u_\epsilon(x, y) = (1 + \epsilon \phi(x, y))u(x, y).$$

For $|\epsilon| < 1/\|\phi\|_\infty$, we have that u_ϵ is a probability measure. It is easy to see that

$$S_1(\mu_\epsilon, \mu_0) = S_1(\mu, \mu_0) - \epsilon \int \phi u \ln u - \int [1 + \epsilon \phi] \ln [1 + \epsilon \phi] u.$$

Since $1 + \epsilon \phi$ is bounded away from 0 (and infinity) for the ϵ we are considering, it is clear that both integrals above are well-defined. Moreover, it is clear that

$$\int [1 + \epsilon \phi] \ln [1 + \epsilon \phi] u = o(\epsilon),$$

because $\ln[1 + \epsilon \phi] = \epsilon \phi + o(\epsilon)$ and $\int \phi u = 0$. A similar calculation also shows that

$$\mu_\epsilon^2(h) = \mu^2(h) + \epsilon \mu^2([\phi(x, y) + \phi(x', y')]h) + O(\epsilon^2).$$

Since μ is supposed to be the optimizer, the terms linear in ϵ must vanish:

$$\int \phi u \ln u = \frac{\beta}{2} \mu^2([\phi(x, y) + \phi(x', y')]h).$$

Since h is symmetric, by varying over all ϕ orthogonal to u , we deduce that

$$u(x, y) \ln u(x, y) = \beta u(x, y) \int_{\mathcal{X}} h((x, y), (x', y')) u(x, y) dx' dy' + C u(x, y),$$

for some constant C . (The reason we cannot assume $C = 0$ is because we left out one direction for ϕ , namely the direction parallel to u , so that there is an indeterminacy in this direction, as seen using the Riesz representation theorem.) In other words, we have just deduced equation (3). On the other hand, we have also proved that this equation has a unique solution given by (10). Therefore, μ_N^β does converge weakly to the i.i.d., product measure of μ , where $d\mu(x, y) = u(x, y) dx dy$.

Because of all this, if we take the empirical measure with respect to μ_N^β ,

$$\frac{1}{N} \sum_{i=1}^N \delta_{(x_i, y_i)},$$

then this does satisfy just the type of convergence claimed in Theorem 1.1. But, taking the order statistics $X_1 < \dots < X_N$ and $Y_1 < \dots < Y_N$, we do have $(x_i, y_i) = (X_{\sigma_i}, Y_{\tau_i})$ for some permutations $\sigma, \tau \in S_N$. Moreover (by commutativity of addition)

$$\frac{1}{N} \sum_{i=1}^N f(x_i, y_i) = \frac{1}{N} \sum_{i=1}^N f(X_i, Y_{\pi_i}),$$

where $\pi = \tau \sigma^{-1}$. As noted before, (X_1, \dots, X_N) and (Y_1, \dots, Y_N) are distributed as the order statistics coming from Lebesgue measure, the effect of the Hamiltonian is only present in the Mallow model $\mathbb{P}_N^{\exp(-\beta/(N-1))}$ -measure of π . By the WLLN for the order statistics, we see that, defining

$$g_N(x, y) = \sum_{i, j=1}^N f(X_i, Y_j) \mathbb{1}_{((i-1)/N, i/N]}(x) \mathbb{1}_{((j-1)/N, j/N]}(y),$$

we have that the random function g_N converges in probability to f , everywhere in $(0, 1] \times (0, 1]$. Therefore, since

$$\frac{1}{N} \sum_{i=1}^N f(x_i, y_i) = \frac{1}{N} \sum_{i=1}^N g_N(i/N, \pi_i/N),$$

we do deduce the theorem from the corresponding result for μ_N^β . Finally note that taking $\exp(-\beta/(n-1))$ versus $1 - \beta/n$ in the theorem does not matter, since the probability measures are continuous with respect to β , and $\exp(-\beta/(n-1)) = 1 - \beta(1 + o(1))/n$.

8. APPLICATIONS

The ground state of the $\mathcal{U}_q(\mathfrak{sl}_2)$ -symmetric XXZ quantum spin system, and the invariant measures of the asymmetric exclusion process on an interval can be obtained from \mathbb{P}_N^q . See Koma and Nachtergaele's paper [16] and Gottstein and Werner's paper [14] for information about the XXZ model. For information about the blocking measures and the asymmetric exclusion process, we find it convenient to refer to Benjamini, Berger, Hoffman and Mossel (BBHM), [3]. The reader can easily deduce information for the XXZ model, since there is a perfect dictionary between these two. An excellent reference for this is Caputo's review [6].

An interesting perspective on the ground state of the quantum XXZ ferromagnet was discovered by Bolina, Contucci and Nachtergaele in [4]. They viewed the ground state of the quantum spin system as a thermal Boltzmann-Gibbs state for a classical model at inverse temperature $\beta = \ln(q^{-2})$. The state space they considered was the set of all up-right paths from $(0, 0)$ to $(m, n) \in \mathbb{Z}^2$ (with $m, n \geq 0$). The Hamiltonian energy function for such a path is the energy under the path, and above the x -axis. Note that the Hamiltonian for the Mallows model also has a graphical representations as the number of "crossings" of the permutation. Using their representation, they explained some symmetries of the ground state of the XXZ model, and obtained estimates which were later useful in their follow-up paper, [5]. The two models are related, but only the Mallows model is manifestly a mean-field model.

We consider the (nearest neighbor) asymmetric exclusion process on $\{1, \dots, N\}$, with hopping rate to the left p and hopping rate to the right $1 - p$, and $q = (1 - p)/p$. We no longer use p or p_N for the pressure, instead we use it for the hopping rate as expressed above. As BBHM explain, the invariant measure of the ASEP is a push-forward of \mathbb{P}_N^q . Given a permutation $\pi \in S_N$ and a particle configuration $\eta = (\eta_1, \dots, \eta_N) \in \{0, 1\}^N$, let $\pi\eta = (\eta_{\pi_1}, \dots, \eta_{\pi_N})$. Let $(1^k, 0^{N-k}) = (1, \dots, 1, 0, \dots, 0)$ with k 1's and $N - k$ 0's. Then, taking a random permutation π , distributed according to \mathbb{P}_N^q , and letting

$$\eta^{(k, N-k)} = \pi(1^k, 0^{N-k}),$$

the law of $\eta^{(k, N-k)}$ is the invariant measure for the ASEP, with k particles and $N - k$ holes. As BBHM explain, this is an instance of Wilson's general height function approach to tiling and shuffling [20]².

The question we can answer is the non-random limiting density of $\eta^{(k, N-k)}$ in the scaling limit, $N \rightarrow \infty$, $p_N = \frac{1}{2} + \beta/4N$, $k_N = \lfloor yN \rfloor$. (Note that this corresponds to $q_N = 1 - \beta(1 + o(1))/N$.) Namely, for a continuous function $f : [0, 1] \rightarrow \mathbb{R}$, we have

$$\lim_{\epsilon \downarrow 0} \lim_{N \rightarrow \infty} \mathbb{P}_N^{1-\beta/N} \left\{ \left| \frac{1}{N} \sum_{i=1}^N f(i/N) \eta_i^{(\lfloor yN \rfloor, \lceil (1-y)N \rceil)} - \int_0^1 f(x) \rho(x; y) dx \right| > \epsilon \right\} = 0,$$

for all $y \in [0, 1]$, where

$$\rho(x; y) = \int_0^y u(x, y') dy'.$$

²Because of this, let us note that the ground state of the XXZ model is also a projection, or marginal, of the Mallows model for permutations (using the correspondence between the ASEP and the XXZ model [6]). This raises an interesting point for further consideration: are other integrable models projections of mean-field models?

The scaling $p_N = \frac{1}{2} + \beta/4N$ is the regime typically called “weakly asymmetric.” See, for example, Enaud and Derrida’s paper [10], following the matrix method used, for example by Derrida, Lebowitz and Speer [7]. Note that while they considered the nonequilibrium case, we consider the particle conserving, equilibrium case. On the other hand, we are sure that the formula above is known.

The integral for $\rho(x; y)$ is readily evaluated. Setting ϕ, ψ, Φ, Ψ and α as in Lemma 6.1,

$$\begin{aligned} \rho(x; y) &= \int_0^y \frac{\alpha\phi(x)\psi(y')}{(\alpha - \beta\Phi(x)\Psi(y'))^2} dy' \\ &= \frac{\alpha\phi(x)}{\beta\Phi(x)(\alpha - \beta\Phi(x)\Psi(y'))} \Big|_0^y \\ &= \frac{\phi(x)\Psi(y)}{\alpha - \beta\Phi(x)\Psi(y)}. \end{aligned}$$

Substituting in, and doing minor algebraic simplifications, we obtain

$$\rho(x; y) = \frac{(1 - e^{-\beta y})e^{-\beta x}}{(1 - e^{-\beta}) - (1 - e^{-\beta x})(1 - e^{-\beta y})}.$$

From this formula it is obvious that the $\beta \rightarrow 0$ limit recovers $\rho(x; y) \equiv y$, as it should (for the symmetric case). Also, after further “simplifications,” we obtain

$$\rho(x; y) = \frac{e^{\beta(\frac{1}{2}-x)/2} \sinh(\beta y/2)}{e^{\beta/4} \cosh(\beta[x-y]/2) - e^{-\beta/4} \cosh(\beta[x+y-1]/2)}.$$

In particular, one can observe that the particle-hole/reflection symmetry is manifest in this formula due to the invariance under the transformation $(\beta, x) \mapsto (-\beta, 1-x)$.

Finally, we note that we can partially undo the scaling limit by taking $\beta \rightarrow \infty$ with $x = y + t/\beta$ (assuming $0 < y < 1$). Approximating $\sinh(\beta y/2) \approx \frac{1}{2}e^{\beta y/2}$ and noting that $e^{-\beta/2} \cosh(\beta[x+y-1]/2) \rightarrow 0$ since $|x+y-1| < 1$, we obtain

$$\rho(x; y) \rightarrow \frac{1}{1 + e^t}.$$

This is not correctly normalized due to the fact that $dx = dt/\beta$, and $\beta \rightarrow \infty$. On the other hand, this does recover the actual lattice scaling limit for the density (modulo a reflection), as has been previously calculated for the XXZ model by Dijkgraaf, Orlando and Reffert in Appendix A of [9].

ACKNOWLEDGEMENTS

This research was supported in part by a U.S. National Science Foundation grant, DMS-0706927. I am very grateful to the following people for useful discussions and suggestions: S. G. Rajeev, Alex Opaku, Carl Mueller, Bruno Nachtergaele, Wolfgang Spitzer and Pierluigi Contucci. I also thank the anonymous referees for their useful suggestions for improvement.

REFERENCES

- [1] D. J. Aldous. Exchangeability and related topics. In P.L. Hennequin (ed.), *École d’été de probabilités de Saint-Flour, XII-1983, Lecture Notes in Math. v. 1117*. 1985 Springer, Berlin, pp. 1–198.
- [2] A. Barra. The mean field Ising model through interpolating techniques. *J. Statist. Phys.* **123** (2008), n. 5, pp. 787–809. <http://arxiv.org/abs/0712.1344>.
- [3] I. Benjamini, N. Berger, C. Hoffman and E. Mossel. Mixing times of the biased card shuffling and the asymmetric exclusion process. *Trans. Amer. Math. Soc.* **357** (2005), no. 8, 3013–3029. <http://arxiv.org/abs/math.PR/0207199>.
- [4] O. Bolina, P. Contucci and B. Nachtergaele. Path Integral Representation for Interface States of the Anisotropic Heisenberg Model. *Rev. Math. Phys.* **12** (2000), pp. 1325–1344. <http://arxiv.org/abs/math-ph/9908004>.
- [5] O. Bolina, P. Contucci and B. Nachtergaele. Path Integral Representations for the Spin-Pinned quantum XXZ Chain. *Markov Process. Related Fields* **11** (2005), pp. 211–221. <http://arxiv.org/abs/math-ph/0306057>.

- [6] P. Caputo. Energy gap estimates in XXZ ferromagnets and stochastic particle systems. *Markov Process. Related Fields* **11** (2005), no. 2, 189–210.
- [7] B. Derrida, J. L. Lebowitz, and E. R. Speer. Free energy functional for nonequilibrium systems: an exactly solvable case. *Phys. Rev. Lett.* **87** (2001), no. 15, 150601, 4 pp. <http://arxiv.org/abs/cond-mat/0105110>.
- [8] P. Diaconis and A. Ram. Analysis of systematic scan Metropolis algorithms using Iwahori-Hecke algebra techniques. Dedicated to William Fulton on the occasion of his 60th birthday. *Michigan Math. J.* **48** (2000), 157–190.
- [9] R. Dijkgraaf, D. Orlando and S. Reffert. Quantum crystals and spin chains. *Nuclear Physics B* **811** (2009), no. 3, 463–490. <http://arxiv.org/abs/0803.1927>.
- [10] C. Enaud and B. Derrida Large deviation functional of the weakly asymmetric exclusion process. *J. Statist. Phys.* **114** (2004), no. 3-4, 537–562. <http://arxiv.org/abs/cond-mat/0307023>.
- [11] L. C. Evans. *Partial Differential Equations. Graduate Studies in Mathematics, Volume 19*. 1998 American Mathematical Society, Providence, RI.
- [12] M. Fannes, H. Spohn and A. Verbeure. Equilibrium states for mean field models. *J. Math. Phys.* **21** (1980), no. 2, 355–358.
- [13] G. Genovese and A. Barra. A mechanical approach to mean field spin models. *Preprint*, 2008. <http://arxiv.org/abs/0812.1978>.
- [14] C.-T. Gottstein, R. F. Werner. Ground states of the infinite q-deformed Heisenberg ferromagnet. *Preprint*, (1995). <http://arxiv.org/abs/cond-mat/9501123>.
- [15] R. B. Israel. *Convexity in the Theory of Lattice Gases*. 1979 Princeton University Press, Princeton, New Jersey.
- [16] T. Koma and B. Nachtergaele. The spectral gap of the ferromagnetic XXZ chain. *Lett. Math. Phys.* **40** (1997), no. 1, 1–16. <http://arxiv.org/abs/cond-mat/9512120>.
- [17] T. M. Liggett. *Interacting Particle Systems. Grundlehren der mathematischen Wissenschaften 276*. 1985 Springer-Verlag, New York.
- [18] C. J. Thompson. *Mathematical Statistical Mechanics*. 1972 The Macmillan Company, New York.
- [19] G. B. Whitham. *Linear and Nonlinear Waves*. 1974 John Wiley & Sons, Inc. New York.
- [20] D. B. Wilson. Mixing times of Lozenge tiling and card shuffling Markov chains. *Ann. Appl. Probab.* **14** (2004), no. 1, 274–325. <http://arxiv.org/abs/math.PR/0102193>

DEPARTMENT OF MATHEMATICS, HYLAN BUILDING, UNIVERSITY OF ROCHESTER, ROCHESTER, NY 14627

E-mail address: <http://www.math.rochester.edu/people/faculty/sstarr/>