

# COMPLETE MONOTONICITY OF A FUNCTION INVOLVING A RATIO OF GAMMA FUNCTIONS AND APPLICATIONS

FENG QI AND BAI-NI GUO

ABSTRACT. In the paper, necessary and sufficient conditions are presented for a function involving a ratio of gamma functions to be logarithmically completely monotonic. This extends and generalizes the main result in [*Inequalities and monotonicity for the ratio of gamma functions*, Taiwanese J. Math. **7** (2003), no. 2, 239–247.] and others. As applications, several inequalities involving the volume of the unit ball in  $\mathbb{R}^n$  are derived, which refine, generalize and extend some known inequalities.

## 1. INTRODUCTION

Recall from [4, 21] that a positive real-valued function  $f(x)$  is said to be logarithmically completely monotonic on an interval  $I \subseteq \mathbb{R}$  if it has derivatives of all orders on  $I$  and its logarithm  $\ln f$  satisfies

$$0 \leq (-1)^k [\ln f(x)]^{(k)} < \infty$$

for  $k \in \mathbb{N}$  on  $I$ . For more properties of this class of functions, please refer to [6].

It is general knowledge that the classical Euler gamma function  $\Gamma(x)$  may be defined for  $x > 0$  by

$$\Gamma(x) = \int_0^\infty t^{x-1} e^{-t} dt. \quad (1)$$

The logarithmic derivative of  $\Gamma(x)$ , denoted by  $\psi(x) = \frac{\Gamma'(x)}{\Gamma(x)}$ , is called the psi or digamma function, and  $\psi^{(k)}(x)$  for  $k \in \mathbb{N}$  are called the polygamma functions. It is well known that these functions are fundamental and that they have much extensive applications in mathematical sciences.

In [13, Theorem 2], the following monotonicity was established: The function

$$\frac{[\Gamma(x+y+1)/\Gamma(y+1)]^{1/x}}{x+y+1} \quad (2)$$

is decreasing with respect to  $x \geq 1$  for fixed  $y \geq 0$ . Consequently, for positive real numbers  $x \geq 1$  and  $y \geq 0$ , we have

$$\frac{x+y+1}{x+y+2} \leq \frac{[\Gamma(x+y+1)/\Gamma(y+1)]^{1/x}}{[\Gamma(x+y+2)/\Gamma(y+1)]^{1/(x+1)}}. \quad (3)$$

---

2010 *Mathematics Subject Classification*. Primary 26A48, 33B15; Secondary 26A51, 26D07.

*Key words and phrases*. Necessary and sufficient condition; logarithmically completely monotonic function; gamma function; psi function; polygamma function; volume; unit ball; inequality; application.

The first author was partially supported by the China Scholarship Council and the Science Foundation of Tianjin Polytechnic University.

This paper was typeset using  $\mathcal{A}\mathcal{M}\mathcal{S}\text{-}\mathcal{L}\mathcal{A}\mathcal{T}\mathcal{E}\mathcal{X}$ .

In [22], the function (2) was proved to be logarithmically completely monotonic with respect to  $x \in (0, \infty)$  for  $y \geq 0$ , and so is its reciprocal for  $-1 < y \leq -\frac{1}{2}$ . Consequently, the inequality (3) is valid for  $(x, y) \in (0, \infty) \times [0, \infty)$  and reversed for  $(x, y) \in (0, \infty) \times (-1, -\frac{1}{2}]$ .

For  $(x, y) \in (0, \infty) \times [0, \infty)$  and  $\alpha \in [0, \infty)$ , the function

$$\frac{[\Gamma(x+y+1)/\Gamma(y+1)]^{1/x}}{(x+y+1)^\alpha} \quad (4)$$

was proved in [38] to be strictly increasing (or decreasing, respectively) with respect to the single variable  $x \in (0, \infty)$  if and only if  $0 \leq \alpha \leq \frac{1}{2}$  (or  $\alpha \geq 1$ , respectively), to be strictly increasing with respect to  $y$  on  $[0, \infty)$  if and only if  $0 \leq \alpha \leq 1$ , and to be logarithmically concave with respect to the 2-variable  $(x, y) \in (0, \infty) \times (0, \infty)$  if  $0 \leq \alpha \leq \frac{1}{4}$ .

For given  $y \in (-1, \infty)$  and  $\alpha \in (-\infty, \infty)$ , let

$$h_{\alpha, y}(x) = \begin{cases} \frac{1}{(x+y+1)^\alpha} \left[ \frac{\Gamma(x+y+1)}{\Gamma(y+1)} \right]^{1/x}, & x \in (-y-1, \infty) \setminus \{0\}; \\ \frac{1}{(y+1)^\alpha} \exp[\psi(y+1)], & x = 0. \end{cases} \quad (5)$$

It is clear that the ranges of  $x$ ,  $y$  and  $\alpha$  in the function  $h_{\alpha, y}(x)$  extend the corresponding ones in the functions (2) and (4) which were ever discussed in [13, 22, 38].

The aim of this paper is to present necessary and sufficient conditions such that the function (5) or its reciprocal are logarithmically completely monotonic.

Our main results may be stated as follows.

**Theorem 1.** *For  $y > -1$ , we have the following statements:*

- (1) *the function (5) is logarithmically completely monotonic with respect to  $x \in (-y-1, \infty)$  if and only if  $\alpha \geq \max\{1, \frac{1}{y+1}\}$ ;*
- (2) *if  $\alpha \leq \min\{1, \frac{1}{2(y+1)}\}$ , the reciprocal of the function (5) is logarithmically completely monotonic with respect to  $x \in (-y-1, \infty)$ ;*
- (3) *the necessary condition for the reciprocal of the function (5) to be logarithmically completely monotonic with respect to  $x \in (-y-1, \infty)$  is  $\alpha \leq 1$ .*

As a ready consequence of monotonic results in Theorem 1, the following double inequality may be derived.

**Theorem 2.** *For  $t > 0$ ,  $y+1 > 0$  and  $x+y+1 > 0$ , the double inequality*

$$\left( \frac{x+y+1}{x+y+t+1} \right)^a < \frac{[\Gamma(x+y+1)/\Gamma(y+1)]^{1/x}}{[\Gamma(x+y+t+1)/\Gamma(y+1)]^{1/(x+t)}} < \left( \frac{x+y+1}{x+y+t+1} \right)^b \quad (6)$$

*holds if  $a \geq \max\{1, \frac{1}{y+1}\}$  and  $b \leq \min\{1, \frac{1}{2(y+1)}\}$ .*

In order to show the applicability of Theorem 2, we derive the following double inequalities involving the  $n$ -dimensional volume

$$\Omega_n = \frac{\pi^{n/2}}{\Gamma(1+n/2)} \quad (7)$$

of the unit ball  $\mathbb{B}^n$  in  $\mathbb{R}^n$ .

**Theorem 3.** For  $n \in \mathbb{N}$ , we have

$$\sqrt{\frac{n+2}{n+4}} < \frac{\Omega_{n+2}^{1/(n+2)}}{\Omega_n^{1/n}} < \sqrt[4]{\frac{n+2}{n+4}}, \quad (8)$$

$$\frac{1}{\pi^{2/(n-2)n}} \sqrt{\frac{n+2}{n+4}} < \frac{\Omega_{n+2}^{1/n}}{\Omega_n^{1/(n-2)}} < \frac{1}{\pi^{2/(n-2)n}} \sqrt[8]{\frac{n+2}{n+4}}, \quad (9)$$

$$\sqrt{\frac{n+2}{n+3}} < \frac{\Omega_{n+1}^{1/(n+1)}}{\Omega_n^{1/n}} < \sqrt[4]{\frac{n+2}{n+3}}. \quad (10)$$

In the final section, we will give several remarks about these three theorems.

## 2. LEMMAS

In order to prove our main results, the following lemmas are needed.

**Lemma 1** ([15, p. 107, Lemma 3]). For  $x \in (0, \infty)$  and  $k \in \mathbb{N}$ , we have

$$\ln x - \frac{1}{x} < \psi(x) < \ln x - \frac{1}{2x} \quad (11)$$

and

$$\frac{(k-1)!}{x^k} + \frac{k!}{2x^{k+1}} < (-1)^{k+1} \psi^{(k)}(x) < \frac{(k-1)!}{x^k} + \frac{k!}{x^{k+1}}. \quad (12)$$

**Lemma 2.** For  $x \in (0, \infty)$  and  $k \in \mathbb{N}$ , we have

$$\ln\left(x + \frac{1}{2}\right) - \frac{1}{x} < \psi(x) < \ln(x+1) - \frac{1}{x} \quad (13)$$

and

$$\frac{(k-1)!}{(x+1)^k} + \frac{k!}{x^{k+1}} < (-1)^{k+1} \psi^{(k)}(x) < \frac{(k-1)!}{(x+1/2)^k} + \frac{k!}{x^{k+1}}. \quad (14)$$

*Proof.* In [16, Theorem 1], the following necessary and sufficient conditions are obtained: For real numbers  $\alpha \neq 0$  and  $\beta$ , the function

$$g_{\alpha, \beta}(x) = \left[ \frac{e^x \Gamma(x+1)}{(x+\beta)^{x+\beta}} \right]^\alpha, \quad x \in (\max\{0, -\beta\}, \infty)$$

is logarithmically completely monotonic if and only if either  $\alpha > 0$  and  $\beta \geq 1$  or  $\alpha < 0$  and  $\beta \leq \frac{1}{2}$ . Further considering the fact in [10, p. 98] that a completely monotonic function which is non-identically zero cannot vanish at any point on  $(0, \infty)$  gives

$$(-1)^k [\ln g_{\alpha, \beta}(x)]^{(k)} = (-1)^k \alpha [x + \ln \Gamma(x) + \ln x - (x+\beta) \ln(x+\beta)]^{(k)} > 0$$

for  $k \in \mathbb{N}$  and  $x \in (0, \infty)$  if and only if either  $\alpha > 0$  and  $\beta \geq 1$  or  $\alpha < 0$  and  $\beta \leq \frac{1}{2}$ . As a result, from straightforward calculation and standard arrangement, inequalities (13) and (14) follow. Lemma 2 is thus proved.  $\square$

## 3. PROOFS OF THEOREMS

Now we are in a position to prove our theorems.

*Proof of Theorem 1.* For  $x \neq 0$ , taking the logarithm of  $h_{\alpha,y}(x)$  gives

$$\ln h_{\alpha,y}(x) = \frac{\ln \Gamma(x+y+1) - \ln \Gamma(y+1)}{x} - \alpha \ln(x+y+1).$$

A direct differentiation yields

$$\begin{aligned} [\ln h_{\alpha,y}(x)]^{(k)} &= \frac{k!}{x^{k+1}} \sum_{i=0}^k \frac{(-1)^{k-i} x^i \psi^{(i-1)}(x+y+1)}{i!} \\ &\quad - \frac{(-1)^k k! \ln \Gamma(y+1)}{x^{k+1}} - \frac{(-1)^{k-1} (k-1)! \alpha}{(x+y+1)^k} \end{aligned} \quad (15)$$

for  $k \in \mathbb{N}$ , where  $\psi^{(-1)}(x+y+1)$  and  $\psi^{(0)}(x+y+1)$  stand for  $\ln \Gamma(x+y+1)$  and  $\psi(x+y+1)$  respectively. Furthermore, a simple calculation gives

$$\begin{aligned} \{x^{k+1} [\ln h_{\alpha,y}(x)]^{(k)}\}' &= (-1)^{k-1} x^k \left[ (-1)^{k-1} \psi^{(k)}(x+y+1) \right. \\ &\quad \left. - \frac{(k-1)! \alpha}{(x+y+1)^k} - \frac{k!(y+1)\alpha}{(x+y+1)^{k+1}} \right]. \end{aligned}$$

Utilizing (12) in the above equation leads to

$$\begin{aligned} \frac{(k-1)!(1-\alpha)}{(x+y+1)^k} + \frac{k![1/2 - (y+1)\alpha]}{(x+y+1)^{k+1}} &\leq \frac{(-1)^{k-1}}{x^k} \{x^{k+1} [\ln h_{\alpha,y}(x)]^{(k)}\}' \\ &\leq \frac{(k-1)!(1-\alpha)}{(x+y+1)^k} + \frac{k![1 - (y+1)\alpha]}{(x+y+1)^{k+1}} \end{aligned} \quad (16)$$

for  $k \in \mathbb{N}$ ,  $x \neq 0$ ,  $y \in (-1, \infty)$  and  $\alpha \in (-\infty, \infty)$ . Therefore,

$$\frac{(-1)^{k-1}}{x^k} \{x^{k+1} [\ln h_{\alpha,y}(x)]^{(k)}\}' \begin{cases} \leq 0, & \text{if } \alpha \geq 1 \text{ and } \alpha \geq \frac{1}{y+1} \\ \geq 0, & \text{if } \alpha \leq 1 \text{ and } \alpha \leq \frac{1}{2(y+1)} \end{cases} \quad (17)$$

for  $k \in \mathbb{N}$ ,  $y > -1$  and  $x \neq 0$ . For  $x > 0$ , the equation (17) means

$$\{x^{2k} [\ln h_{\alpha,y}(x)]^{(2k-1)}\}' \begin{cases} \leq 0, & \text{if } \alpha \geq 1 \text{ and } \alpha \geq \frac{1}{y+1} \\ \geq 0, & \text{if } \alpha \leq 1 \text{ and } \alpha \leq \frac{1}{2(y+1)} \end{cases}$$

and

$$\{x^{2k+1} [\ln h_{\alpha,y}(x)]^{(2k)}\}' \begin{cases} \geq 0, & \text{if } \alpha \geq 1 \text{ and } \alpha \geq \frac{1}{y+1} \\ \leq 0, & \text{if } \alpha \leq 1 \text{ and } \alpha \leq \frac{1}{2(y+1)} \end{cases}$$

for  $k \in \mathbb{N}$ . From (15), it is easy to see that

$$\lim_{x \rightarrow 0} \{x^{k+1} [\ln h_{\alpha,y}(x)]^{(k)}\} = 0 \quad (18)$$

for  $k \in \mathbb{N}$  and any given  $y > -1$ . As a result,

$$[\ln h_{\alpha,y}(x)]^{(2k-1)} \begin{cases} < 0, & \text{if } \alpha \geq 1 \text{ and } \alpha \geq \frac{1}{y+1} \\ > 0, & \text{if } \alpha \leq 1 \text{ and } \alpha \leq \frac{1}{2(y+1)} \end{cases} \quad (19)$$

and

$$[\ln h_{\alpha,y}(x)]^{(2k)} \begin{cases} > 0, & \text{if } \alpha \geq 1 \text{ and } \alpha \geq \frac{1}{y+1} \\ < 0, & \text{if } \alpha \leq 1 \text{ and } \alpha \leq \frac{1}{2(y+1)} \end{cases} \quad (20)$$

for  $k \in \mathbb{N}$  and  $x \in (0, \infty)$ , that is,

$$(-1)^k [\ln h_{\alpha,y}(x)]^{(k)} \begin{cases} > 0, & \text{if } \alpha \geq 1 \text{ and } \alpha \geq \frac{1}{y+1} \\ < 0, & \text{if } \alpha \leq 1 \text{ and } \alpha \leq \frac{1}{2(y+1)} \end{cases} \quad (21)$$

for  $k \in \mathbb{N}$  and  $x \in (0, \infty)$ . Hence, the function (5) is logarithmically completely monotonic with respect to  $x$  on  $(0, \infty)$  if  $\alpha \geq 1$  and  $\alpha \geq \frac{1}{y+1}$ , and so is the reciprocal of the function (5) if either  $0 < \alpha \leq 1$  and  $\alpha \leq \frac{1}{2(y+1)}$  or  $\alpha \leq 0$  and  $y > -1$ .

If  $x \in (-y-1, 0)$ , the equation (17) means

$$\{x^{k+1} [\ln h_{\alpha,y}(x)]^{(k)}\}' \begin{cases} \geq 0, & \text{if } \alpha \geq 1 \text{ and } \alpha \geq \frac{1}{y+1} \\ \leq 0, & \text{if } \alpha \leq 1 \text{ and } \alpha \leq \frac{1}{2(y+1)} \end{cases}$$

for  $k \in \mathbb{N}$ . By virtue of (18), it follows that

$$x^{k+1} [\ln h_{\alpha,y}(x)]^{(k)} \begin{cases} \leq 0, & \text{if } \alpha \geq 1 \text{ and } \alpha \geq \frac{1}{y+1} \\ \geq 0, & \text{if } \alpha \leq 1 \text{ and } \alpha \leq \frac{1}{2(y+1)} \end{cases}$$

for  $k \in \mathbb{N}$ , which is equivalent to that the equations (19) and (20) hold for  $x \in (-y-1, 0)$ . As a result, the equation (21) is valid for  $k \in \mathbb{N}$  and  $x \in (-y-1, 0)$ . Therefore, the function  $h_{\alpha,y}(x)$  has the same logarithmically complete monotonicity properties on  $(-y-1, 0)$  as on  $(0, \infty)$ .

Conversely, if  $h_{\alpha,y}(x)$  is logarithmically completely monotonic on  $(-y-1, \infty)$ , then  $[\ln h_{\alpha,y}(x)]' < 0$  on  $(-y-1, \infty)$ , which can be simplified as

$$\alpha \geq (x+y+1) \left[ \frac{1}{x^2} \sum_{i=0}^1 \frac{(-1)^{1-i} x^i \psi^{(i-1)}(x+y+1)}{i!} + \frac{\ln \Gamma(y+1)}{x^2} \right] \quad (22)$$

$$= \frac{1}{x^2} [(x+y+1) \ln \Gamma(y+1) - (y+1)(x+y+1) \psi(x+y+1) + (x+y+1)^2 \psi(x+y+1) - (x+y+1) \ln \Gamma(x+y+1)] \quad (23)$$

$$= \frac{x+y+1}{x} \left[ \frac{x\psi(x+y+1) - \ln \Gamma(x+y+1)}{x} + \frac{\ln \Gamma(y+1)}{x} \right]. \quad (24)$$

From (11), it is easy to see that

$$\lim_{x \rightarrow 0^+} [x^2 \psi(x)] = 0. \quad (25)$$

It is common knowledge that

$$\Gamma(x+1) = x\Gamma(x) \quad (26)$$

for  $x > 0$ . Taking the logarithm on both sides of (26), rearranging, and taking limit lead to

$$\lim_{x \rightarrow 0^+} [x \ln \Gamma(x)] = \lim_{x \rightarrow 0^+} [x \ln \Gamma(x+1)] - \lim_{x \rightarrow 0^+} [x \ln x] = 0. \quad (27)$$

Taking logarithmic derivatives on both sides of (26) yields

$$\psi(x+1) = \frac{1}{x} + \psi(x) \quad (28)$$

for  $x > 0$ , and so

$$\lim_{x \rightarrow 0^+} [x\psi(x)] = 1 + \lim_{x \rightarrow 0^+} [x\psi(x+1)] = 1. \quad (29)$$

Thus, by utilizing (25), (27) and (29), it is revealed that the limit of the function (23) as  $x \rightarrow (-y-1)^+$ , that is, as  $x+y+1 \rightarrow 0^+$ , equals  $\frac{1}{y+1}$ . By L'Hôpital's rule and the double inequality (12) for  $k=1$ , we have

$$\lim_{x \rightarrow \infty} \frac{x\psi(x+y+1) - \ln \Gamma(x+y+1)}{x} = \lim_{x \rightarrow \infty} [x\psi'(x+y+1)] = 1.$$

Hence, the limit of the function (24) as  $x \rightarrow \infty$  equals 1. In a word, the necessary condition for  $h_{\alpha,y}(x)$  to be logarithmically completely monotonic is  $\alpha \geq 1$  and  $\alpha \geq \frac{1}{y+1}$ .

If the reciprocal of  $h_{\alpha,y}(x)$  is logarithmically completely monotonic, then the inequality (22) is reversed. Since the limit of the function (24) equals 1 as  $x \rightarrow \infty$ , as showed above, then the necessary condition  $\alpha \leq 1$  is obtained. The proof of Theorem 1 is complete.  $\square$

*Proof of Theorem 2.* This follows from the monotonicity properties established in Theorem 1.  $\square$

*Proof of Theorem 3.* Letting  $t=1$ ,  $y=0$  and  $x = \frac{n}{2}$  for  $n \in \mathbb{N}$  in (6) reveals that

$$\frac{n+2}{n+4} < \frac{[\Gamma(n/2+1)]^{2/n}}{[\Gamma((n+2)/2+1)]^{2/(n+2)}} < \sqrt{\frac{n+2}{n+4}}$$

which is equivalent to the inequality (8).

If taking  $y=1$ ,  $t=1$  and  $x = \frac{n}{2} - 1$  for  $n \in \mathbb{N}$  in (6), then the inequality (9) follows.

Replacing  $t$  by  $\frac{1}{2}$ ,  $y$  by 0, and  $x$  by  $\frac{n}{2}$  in (6) and simplifying result in (10).  $\square$

#### 4. REMARKS

After proving our theorems, we give several remarks about them.

*Remark 1.* Theorem 1 extends and generalizes the logarithmically complete monotonicity of the function (2) established in [22] and a part of the results in [38].

*Remark 2.* The inequality (6) generalizes and extends the inequality (3) and the main results in [23, 37]: For  $x+y > 0$  and  $y+1 > 0$  the inequality

$$\frac{[\Gamma(x+y+1)/\Gamma(y+1)]^{1/x}}{[\Gamma(x+y+2)/\Gamma(y+1)]^{1/(x+1)}} < \left( \frac{x+y}{x+y+1} \right)^{1/2} \quad (30)$$

is valid if  $x > 1$  and reversed if  $x < 1$  and that the power  $\frac{1}{2}$  is the best possible.

*Remark 3.* When  $n > 2$ , the inequality (10) refines the following double inequality in [2, Theorem 1]:

$$\frac{2}{\sqrt{\pi}} \Omega_{n+1}^{n/(n+1)} \leq \Omega_n < \sqrt{e} \Omega_{n+1}^{n/(n+1)}, \quad n \in \mathbb{N}. \quad (31)$$

For more information on inequalities for the volume of the unit ball  $\mathbb{B}^n$  in  $\mathbb{R}^n$ , please refer to [3, 7, 8, 25], Section 7.5 in [19, pp. 72–73] and related references therein.

*Remark 4.* Theorem 1 may be restated as follows: For  $y \in (0, \infty)$  and  $\alpha \in (-\infty, \infty)$ , the function

$$H_y(x) = \begin{cases} \frac{[\Gamma(x+y)/\Gamma(y)]^{1/x}}{(x+y)^\alpha}, & x \in (-y, \infty) \setminus \{0\} \\ \frac{e^{\psi(y)}}{y^\alpha}, & x = 0 \end{cases} \quad (32)$$

is logarithmically completely monotonic with respect to  $x \in (-y, \infty)$  if and only if  $\alpha \geq \max\{1, \frac{1}{y}\}$ , and so is its reciprocal if  $\alpha \leq \min\{1, \frac{1}{2y}\}$  and only if  $\alpha \leq 1$ .

*Remark 5.* We conjecture that when  $y > -\frac{1}{2}$  the condition  $\alpha \leq \frac{1}{2(y+1)}$  is also necessary for the reciprocal of the function (5) to be logarithmically completely monotonic with respect to  $x \in (-y-1, \infty)$ .

*Remark 6.* For more information on the history, backgrounds, motivations and recent developments of the topic in this paper, please refer to [1, 5, 9, 13, 14, 20, 22, 26, 31, 33, 34] and related references therein.

*Remark 7.* In passing, we survey the history of the notion “logarithmically completely monotonic function”. By searching for the term “logarithmically completely monotonic function” in the database MathSciNet, it is found that this phrase was probably first used in [4], but without an explicit definition. Thereafter, it seems to have not been used by the mathematical community. In early 2004, this terminology was again used in [24] (the preprint of [21, 28]) and it was immediately referenced in [11] and [30] (the preprint of [29]). In [24, Theorem 4], it was proved that a logarithmically completely monotonic function  $f(x)$  on  $I$  must be completely monotonic (i.e., the inequality  $0 \leq (-1)^k f^{(k)}(x) < \infty$  holds for all  $k \geq 0$  on  $I$ ), but not conversely. This result was announced while revising [21]. This conclusion and its proofs were presented once and again in [6] and [32] (the preprint of [12]). More importantly, in the paper [6], the logarithmically completely monotonic functions on  $(0, \infty)$  were characterized as the infinitely divisible completely monotonic functions studied in [17] and all Stieltjes transforms were proved to be logarithmically completely monotonic on  $(0, \infty)$ . For information on the completely monotonic functions, please refer to [18, Chapter XIII] and [36, Chapter IV], especially to the recently published monograph [35].

*Remark 8.* This paper is a part of the preprint [27].

#### REFERENCES

- [1] S. Abramovich, J. Barić, M. Matic and J. Pečarić, *On van de Lune-Alzer’s inequality*, J. Math. Inequal. **1** (2007), no. 4, 563–587. 7
- [2] H. Alzer, *Inequalities for the volume of the unit ball in  $\mathbb{R}^n$* , J. Math. Anal. Appl. **252** (2000), 353–363. 6
- [3] H. Alzer, *Inequalities for the volume of the unit ball in  $\mathbb{R}^n$ , II*, Mediterr. J. Math. **5** (2008), 395–413. 6
- [4] R. D. Atanassov and U. V. Tsoukrovski, *Some properties of a class of logarithmically completely monotonic functions*, C. R. Acad. Bulgare Sci. **41** (1988), no. 2, 21–23. 1, 7
- [5] G. Bennett, *Meaningful inequalities*, J. Math. Inequal. **1** (2007), no. 4, 449–471. 7
- [6] C. Berg, *Integral representation of some functions related to the gamma function*, Mediterr. J. Math. **1** (2004), no. 4, 433–439. 1, 7
- [7] C. Berg and H. L. Pedersen, *A one-parameter family of Pick functions defined by the Gamma function and related to the volume of the unit ball in  $n$ -space*, Proc. Amer. Math. Soc. **139** (2011), in press. 6

- [8] C. Berg and H. L. Pedersen, *A Pick function related to the sequence of volumes of the unit ball in  $n$ -space*, Available online at <http://arxiv.org/abs/0912.2185>. 6
- [9] Ch.-P. Chen, F. Qi, P. Cerone and S. S. Dragomir, *Monotonicity of sequences involving convex and concave functions*, Math. Inequal. Appl. **6** (2003), no. 2, 229–239. 7
- [10] J. Dubourdieu, *Sur un théorème de M. S. Bernstein relatif à la transformation de Laplace-Stieltjes*, Compositio Math. **7** (1939-40), 96–111. 3
- [11] A. Z. Grinshpan and M. E. H. Ismail, *Completely monotonic functions involving the gamma and  $q$ -gamma functions*, Proc. Amer. Math. Soc. **134** (2006), 1153–1160. 7
- [12] B.-N. Guo and F. Qi, *A property of logarithmically absolutely monotonic functions and the logarithmically complete monotonicity of a power-exponential function*, Politehn. Univ. Bucharest Sci. Bull. Ser. A Appl. Math. Phys. **72** (2010), no. 2, 21–30. 7
- [13] B.-N. Guo and F. Qi, *Inequalities and monotonicity for the ratio of gamma functions*, Taiwanese J. Math. **7** (2003), no. 2, 239–247. 1, 2, 7
- [14] B.-N. Guo and F. Qi, *Monotonicity of sequences involving geometric means of positive sequences with monotonicity and logarithmic convexity*, Math. Inequal. Appl. **9** (2006), no. 1, 1–9. 7
- [15] B.-N. Guo and F. Qi, *Two new proofs of the complete monotonicity of a function involving the psi function*, Bull. Korean Math. Soc. **47** (2010), no. 1, 103–111; Available online at <http://dx.doi.org/10.4134/BKMS.2010.47.1.103>. 3
- [16] S. Guo, F. Qi and H. M. Srivastava, *Necessary and sufficient conditions for two classes of functions to be logarithmically completely monotonic*, Integral Transforms Spec. Funct. **18** (2007), no. 11, 819–826. 3
- [17] R. A. Horn, *On infinitely divisible matrices, kernels and functions*, Z. Wahrscheinlichkeitstheorie und Verw. Geb. **8** (1967), 219–230. 7
- [18] D. S. Mitrinović, J. E. Pečarić and A. M. Fink, *Classical and New Inequalities in Analysis*, Kluwer Academic Publishers, 1993. 7
- [19] F. Qi, *Bounds for the ratio of two gamma functions*, J. Inequal. Appl. **2010** (2010), Article ID 493058, 84 pages; Available online at <http://dx.doi.org/10.1155/2010/493058>. 6
- [20] F. Qi, *Inequalities and monotonicity of sequences involving  $\sqrt[n]{(n+k)!/k!}$* , Soochow J. Math. **29** (2003), no. 4, 353–361. 7
- [21] F. Qi and Ch.-P. Chen, *A complete monotonicity property of the gamma function*, J. Math. Anal. Appl. **296** (2004), no. 2, 603–607. 1, 7
- [22] F. Qi and B.-N. Guo, *A logarithmically completely monotonic function involving the gamma function*, Taiwanese J. Math. **14** (2010), no. 4, 1623–1628. 2, 6, 7
- [23] F. Qi and B.-N. Guo, *An extension of an inequality for ratios of gamma functions*, J. Approx. Theory (2011), in press. 6
- [24] F. Qi and B.-N. Guo, *Complete monotonicities of functions involving the gamma and digamma functions*, RGMIA Res. Rep. Coll. **7** (2004), no. 1, Art. 8, 63–72; Available online at <http://rgmia.org/v7n1.php>. 7
- [25] F. Qi and B.-N. Guo, *Monotonicity and logarithmic convexity relating to the volume of the unit ball*, Available online at <http://arxiv.org/abs/0902.2509>. 6
- [26] F. Qi and B.-N. Guo, *Monotonicity of sequences involving convex function and sequence*, Math. Inequal. Appl. **9** (2006), no. 2, 247–254. 7
- [27] F. Qi and B.-N. Guo, *Necessary and sufficient conditions for a function involving a ratio of gamma functions to be logarithmically completely monotonic*, Available online at <http://arxiv.org/abs/0904.1101>. 7
- [28] F. Qi and B.-N. Guo, *Some logarithmically completely monotonic functions related to the gamma function*, J. Korean Math. Soc. **47** (2010), no. 6, 1283–1297; Available online at <http://dx.doi.org/10.4134/JKMS.2010.47.6.1283>. 7
- [29] F. Qi, B.-N. Guo and Ch.-P. Chen, *Some completely monotonic functions involving the gamma and polygamma functions*, J. Aust. Math. Soc. **80** (2006), 81–88. 7
- [30] F. Qi, B.-N. Guo and Ch.-P. Chen, *Some completely monotonic functions involving the gamma and polygamma functions*, RGMIA Res. Rep. Coll. **7** (2004), no. 1, Art. 5, 31–36; Available online at <http://rgmia.org/v7n1.php>. 7
- [31] F. Qi and S. Guo, *On a new generalization of Martins' inequality*, J. Math. Inequal. **1** (2007), no. 4, 503–514. 7
- [32] F. Qi, W. Li and B.-N. Guo, *Generalizations of a theorem of I. Schur*, RGMIA Res. Rep. Coll. **9** (2006), no. 3, Art. 15; Available online at <http://rgmia.org/v9n3.php>. 7



- [33] F. Qi and Q.-M. Luo, *Generalization of H. Minc and L. Sathre's inequality*, Tamkang J. Math. **31** (2000), no. 2, 145–148. 7
- [34] F. Qi and J.-Sh. Sun, *A monotonicity result of a function involving the gamma function*, Anal. Math. **32** (2006), no. 4, 279–282. 7
- [35] R. L. Schilling, R. Song and Z. Vondraček, *Bernstein Functions*, de Gruyter Studies in Mathematics **37**, De Gruyter, Berlin, Germany, 2010. 7
- [36] D. V. Widder, *The Laplace Transform*, Princeton University Press, Princeton, 1946. 7
- [37] Y. Yu, *An inequality for ratios of gamma functions*, J. Math. Anal. Appl. **352** (2009), no. 2, 967–970. 6
- [38] T.-H. Zhao, Y.-M. Chu and Y.-P. Jiang, *Monotonic and logarithmically convex properties of a function involving gamma functions*, J. Inequal. Appl. **2009** (2009), Article ID 728612, 13 pages; Available online at <http://dx.doi.org/10.1155/2009/728612>. 2, 6

(F. Qi) DEPARTMENT OF MATHEMATICS, COLLEGE OF SCIENCE, TIANJIN POLYTECHNIC UNIVERSITY, TIANJIN CITY, 300160, CHINA; SCHOOL OF MATHEMATICS AND INFORMATICS, HENAN POLYTECHNIC UNIVERSITY, JIAOZUO CITY, HENAN PROVINCE, 454010, CHINA

*E-mail address:* [qifeng618@gmail.com](mailto:qifeng618@gmail.com), [qifeng618@hotmail.com](mailto:qifeng618@hotmail.com), [qifeng618@qq.com](mailto:qifeng618@qq.com)  
*URL:* <http://qifeng618.wordpress.com>

(B.-N. Guo) SCHOOL OF MATHEMATICS AND INFORMATICS, HENAN POLYTECHNIC UNIVERSITY, JIAOZUO CITY, HENAN PROVINCE, 454010, CHINA

*E-mail address:* [bai.ni.guo@gmail.com](mailto:bai.ni.guo@gmail.com), [bai.ni.guo@hotmail.com](mailto:bai.ni.guo@hotmail.com)