

IDEALS IN LEFT ALMOST SEMIGROUPS

KAISER MUSHTAQ AND MADAD KHAN

ABSTRACT. A left almost semigroup (LA-semigroup) or an Abel-Grassmann's groupoid (AG-groupoid) is investigated in several papers. In this paper we have discussed ideals in LA-semigroups. Specifically, we have shown that every ideal in an LA-semigroup S with left identity e is prime if and only if it is idempotent and the set of ideals of S is totally ordered under inclusion. We have shown that an ideal of S is prime if and only if it is semiprime and strongly irreducible. We have proved also that every ideal in a regular LA-semigroup S is prime if and only if the set of ideals of S is totally ordered under inclusion. We have proved in the end that every ideal in S is prime if and only if it is strongly irreducible and the set of ideals of S form a semilattice.

1. INTRODUCTION

A left almost-semigroup (LA-semigroup) [6] or Abel-Grassmann's groupoid (AG-groupoid) [10] is a groupoid S with left invertive law:

$$(1) \quad (ab)c = (cb)a, \text{ for all } a, b, c \in S.$$

Every LA-semigroup S satisfy the medial law [5] :

$$(2) \quad (ab)(cd) = (ac)(bd), \text{ for all } a, b, c, d \in S.$$

It has been shown in [6] that if an LA-semigroup contains a left identity then it is unique. It has been proved also that an LA-semigroup with right identity is a commutative monoid, that is, a commutative semigroup with identity element.

2. IDEALS IN AN LA-SEMIGROUP

An LA-semigroup S is called regular if for each a in S there exists x in S such that $a = (ax)a$. An LA-semigroup S is called inverse LA-semigroup if for each a in S there exists x in S such that, $a = (ax)a$ and $x = (xa)x$. If S is a regular LA-semigroup then it is easy to see that $S = S^2$.

Lemma 1. *If S is an LA-semigroup with left identity e then $SS = S$ and $S = eS = Se$.*

Proof. If S is an LA-semigroup with left identity e . Then $x \in S$ implies that

$$x = ex \in SS \text{ and so } S \subseteq SS. \text{ That is}$$

$S = SS$. Now using the facts, $eS = S$, $SS = S$ and (1), we obtain

$$Se = (SS)e = (eS)S = SS = S$$

2000 *Mathematics Subject Classification.* 20M10 and 20N99.

Key words and phrases. LA-semigroup, Regular LA-semigroup, Medial law, ideals, minimal and prime ideals.

Hence $S = eS = Se$. □

Remark 1. *If S is an LA-semigroup with left identity e then $S = S^2$, but the converse is not necessarily true, for example regular LA-semigroup.*

Example 1. *Let $S = \{1, 2, 3, 4, 5, 6, 7\}$ the binary operation “ \cdot ” be defined on S as follows:*

Then (S, \cdot) is an LA-semigroup without left identity, but $S = S^2$.

Proposition 1. *If S is an LA-semigroup with left identity e then $(xy)^2 = (x^2y^2) = (y^2x^2)$, for all x, y in S .*

The proof is easy. Consequently a^2S is an ideal of S .

A subset I of an LA-semigroup S is called a right (left) ideal if $IS \subseteq I$ ($SI \subseteq I$), and is called an ideal if it is a two sided ideal. If S is an LA-semigroup with left identity e , then $S(Sa) \subseteq Sa$ in [10].

Also $(aS)S \subseteq aS$, if a is an idempotent in an LA-semigroup S with left identity.

Proposition 2. *If S is an LA-semigroup with left identity e then every right ideal is an ideal.*

Proof. Let I be a right ideal of an LA-semigroup S and $x \in S, i \in I$. Then by (1), we get $si = (es)i = (is)e \in I$. Hence I is a left ideal. □

Remark 2. (1) *If $S = S^2$ then every right ideal is also a left ideal and $SI \subseteq IS$.*
 (2) *If I is a right ideal of S , then SI is a left and IS is a right ideal of S .*

It is obvious that the intersection of two or more ideals is an ideal. Similarly the union of two or more ideals is also an ideal.

Lemma 2. *If I is a left ideal of an LA-semigroup S with left identity e , then aI is a left ideal of S .*

Proof. If I is a left ideal of S , then by (1) and (2), and the fact that $ai \in aI$, we get

$$s(ai) = (es)(ai) = (ea)(si) = a(si) \in aI.$$

□

Lemma 3. *If I is a right ideal of an LA-semigroup S with left identity e then I^2 is an ideal of S .*

Proof. Let $x \in I^2$ then $x = ij$ where $i, j \in I$. Now using (1), we get $xs = (ij)s = (sj)i \in II = I^2$. This implies that I^2 is a right ideal and by proposition 1, it becomes a left ideal. \square

Remark 3. *If I is a left ideal of S then I^2 is an ideal of S .*

Proposition 3. *A proper ideal M of an LA-semigroup S with left identity e , is minimal if and only if $M = a^2M$, for all $a \in S$.*

Proof. Let M be the minimal ideal of S , as M^2 is an ideal so $M = M^2$. Now using (1) we see that a^2M is an ideal of S , contain in M and as M is minimal so $M = a^2M$.

Conversely, assume that $M = a^2M$, for all $a \in S$. Let A be the minimal ideal properly contain in M containing a , then $M = a^2M \subseteq A$, which is a contradiction. Hence M is a minimal ideal. \square

An ideal I of an LA-semigroup S is called minimal if and only if it does not contain any ideal of S other than itself.

Theorem 1. *If I is a minimal left ideal of an LA-semigroup S with left identity e , then a^2I^2 is a minimal ideal of S .*

Proof. Let $ai \in aI$ where I is a minimal left ideal of an LA-semigroup S with left identity e . Then using (1) and Proposition 1, we get, $(a^2I^2)S = (SI^2)a^2 \subseteq I^2a^2 = a^2I^2$, which shows that a^2I^2 is a right ideal of S and by proposition 2, it becomes left. Let H be a non-empty ideal of S properly contained in a^2I^2 . Define $H' = \{r \in I : ar \in H\}$. Then $a(sy) = s(ay) \in SH \subseteq H$ imply that H' is a left ideal of S properly contained in I . But this is a contradiction to the minimality of I . Hence a^2I^2 is a minimal ideal of S . \square

An ideal P of an LA-semigroup S is called prime if $AB \subseteq P$ implies that either $A \subseteq P$ or $B \subseteq P$, for all ideals A and B in S . An ideal P of an LA-semigroup S is said to be semiprime if $I^2 \subseteq P$ implies that $I \subseteq P$, for any ideal I of S . An LA-semigroup S is said to be fully semiprime if every ideal of S is semiprime. An LA-semigroup S is called fully prime if every ideal of S is prime. The set of ideals of an LA-semigroup S is called totally ordered under inclusion if for all ideals A, B of S , either $A \subseteq B$ or $B \subseteq A$ and is denoted by $\text{ideal}(S)$.

Theorem 2. *An LA-semigroup S with left identity e is fully prime if and only if every ideal is idempotent and $\text{ideal}(S)$ is totally ordered under inclusion.*

Proof. Assume that an LA-semigroup S is fully prime. Let I be the ideal of S . Then by Lemma 3, I^2 becomes an ideal of S and obviously $I^2 \subseteq I$. Now

$$II \subseteq I^2 \text{ yields } I \subseteq I^2 \text{ and hence}$$

$I = I^2$. Let P, Q be ideals of S and $PQ \subseteq P, PQ \subseteq Q$ imply that $PQ \subseteq P \cap Q$. Since $P \cap Q$ is prime, so $P \subseteq P \cap Q$ or $Q \subseteq P \cap Q$ which further imply that $P \subseteq Q$ or $Q \subseteq P$. Hence $\text{ideal}(S)$ is totally ordered under inclusion.

Conversely, assume that every ideal of S is idempotent and $\text{ideal}(S)$ is totally ordered under inclusion. Let I, J and P be the ideals of S with $IJ \subseteq P$ such that $I \subseteq J$. Since I is idempotent, $I = I^2 = II \subseteq IJ \subseteq P$ implies that $I \subseteq P$ and so S is fully prime. \square

If S is an LA-semigroup with left identity e then the principal left ideal generated by a is defined by $\langle a \rangle = Sa = \{sa : s \in S\}$, where a is any element of S . Let P be a left ideal of an LA-semigroup S , P is called quasi-prime if for left ideals A, B of S such that $AB \subseteq P$, we have $A \subseteq P$ or $B \subseteq P$. P is called quasi-semiprime if for any left ideal I of S such that $I^2 \subseteq P$, we have $I \subseteq P$.

Theorem 3. *If S is an LA-semigroup with left identity e , then a left ideal P of S is quasi-prime if and only if $a(Sb) \subseteq P$ implies that either $a \in P$ or $b \in P$, where $a, b \in S$.*

Proof. Let P be a left ideal of an LA semigroup S with identity e . Now suppose that $a(Sb) \subseteq P$. Then by (2), we get

$$\begin{aligned} S(a(Sb)) &\subseteq SP \subseteq P, \text{ that is} \\ S(a(Sb)) &= (SS)(a(Sb)) = (Sa)(S(Sb)) = (Sa)((SS)(Sb)) \\ &= (Sa)((bS)(SS)) = (Sa)(Sb). \end{aligned}$$

Hence, either $a \in P$ or $b \in P$.

Conversely, assume that $AB \subseteq P$ where A and B are left ideals of S such that $A \not\subseteq P$. Then there exists $x \in A$ such that $x \notin P$. Now $x(Sy) \subseteq A(SB) \subseteq AB \subseteq P$, for all $y \in B$. Hence by hypothesis, $y \in P$ for all $y \in B$. This shows that P is quasi-prime. \square

Corollary 1. *If S is an LA-semigroup with left identity e , then a left ideal P of S is quasi-semiprime if and only if $a(Sa) \subseteq P$ implies $a \in P$, for all $a \in S$.*

Lemma 4. *If I is a proper right (left) ideal of an LA-semigroup S with left identity e then $e \notin I$.*

Proof. Suppose that $e \in I$. Then $S = eS \subseteq IS \subseteq I$ imply that $S \subseteq I$ and so $I = S$. This is a contradiction to the fact that I is proper. Hence $e \notin I$. \square

It is easy to see from the above results that if I is a proper ideal of S with left identity e then $H(a) = \{x \in S : ax = e\} \not\subseteq I$.

A left inverse \acute{a} of a in an LA-semigroup with left identity e , becomes the right inverse of a . If all elements of S are right invertible then S is called a right invertible LA-semigroup.

Theorem 4. *If I is a ideal of a right invertible LA-semigroup S with left identity e , then I is a proper ideal if and only if for all a in I there does not exist x in S such that $ax = e$.*

Proof. Let I be a proper right ideal of S and $a \in I$ be a right invertible element of S . Then there exists $x \in S$ such that $e = ax \in IS \subseteq I$ which gives $e \in I$ and so by Lemma 4, $I = S$. But this is a contradiction. Hence for all a in I there does not exist x in S such that $ax = e$.

The converse is straight forward. \square

An ideal I of an LA-semigroup S is said to be strongly irreducible if and only if for ideals H and K of S , $H \cap K \subseteq I$ implies that $H \subseteq I$ or $K \subseteq I$.

Proposition 4. *An ideal I of an LA-semigroup S is prime if and only if it is semiprime and strongly irreducible.*

The proof is obvious.

Theorem 5. *Let S be an LA semigroup and $\{P_i : i \in N\}$ be a family of prime ideals totally ordered under inclusion in S . Then $\cap P_i$ is a prime ideal.*

The proof is same as in [9].

Theorem 6. *For each ideal I there exists a minimal prime ideal of I in an LA-semigroup S .*

The proof is same as in [9].

3. IDEALS IN A REGULAR LA-SEMIGROUP

Lemma 5. *Every right ideal of a regular LA-semigroup is an ideal.*

Proof. Let I be a right ideal of S . Then for each s in S there exists x such that $s = (sx)s$. If $i \in I$ then, $si = ((sx)s)i = (is)(sx) \subseteq I$ imply that I is a left ideal of S . \square

It is easy to see that $SI \subseteq IS$.

Lemma 6. *If S is a regular LA-semigroup then $PQ = P \cap Q$, where P is a right and Q is a left ideal.*

Proof. Let P and Q be right and left ideals of S with $PQ \subseteq P \cap Q$. If $a \in P \cap Q$, then there exists b in S such that $a = (ab)a \in PQ$. Hence $PQ = P \cap Q$. \square

Note that if I is a left ideal of an LA-semigroup then by Lemma 6, $I = I^2$. Also it is worth mentioning that ideals of a regular LA-semigroup are semiprime.

The following Theorem is an easy consequence of Lemma 6.

Theorem 7. *The set of ideals of a regular LA-semigroup S form a semilattice, (L_S, \wedge) , where $A \wedge B = AB$, A and B are ideals of S .*

Theorem 8. *A regular LA-semigroup S is fully prime if and only if $\text{ideal}(S)$ is totally ordered under inclusion.*

The proof follows from Lemma 6 and Theorem 2.

Theorem 9. *Every ideal in a regular LA-semigroup S is prime if and only if it is strongly irreducible.*

Proof. Assume that P is a prime ideal of S . Then there exist ideals A and B in S such that $AB \subseteq P$. Then by Lemma 6, $AB = A \cap B$ implies that either $A \subseteq P$ or $B \subseteq P$. Hence P is strongly irreducible.

Conversely, assume that every ideal of a regular LA-semigroup S is strongly irreducible. Let P be a strongly irreducible ideal of S . Then $A \cap B \subseteq P$ (where A and B are any ideals of S) implies that either $A \subseteq P$ or $B \subseteq P$. But by Lemma 6, $AB = A \cap B$. Hence P is prime. \square

REFERENCES

- [1] Birkhoff, G, *Lattice Theory*, AMS Colloq; 1954.
- [2] Clifford, A. H and G. B. Preston, *The Algebraic Theory of semigroups*, Vols, 1, 2. AMS Surveys, 7, (1961, 1967).
- [3] Golan, J. S, *The theory of semirings with applications in mathematics and theoretical computer science*, Pitman Monographs and surveys in pure and applied Mathematics, 54, England, 1992.
- [4] Howie, J. M, *Introduction to semigroup theory*, Academic Press, London, 1976.

- [5] Kazim, M. A. and M. Naseeruddin, On almost-semigroups, The Alig. Bull. Math., 2 (1972), 1 – 7.
- [6] Mushtaq, Q and S. M. Yusuf, On LA-semigroups, The Alig. Bull; of Math., 8 (1978), 65 – 70.
- [7] Mushtaq, Q, Abelian groups defined by LA-semigroups, Studia Scient. Math. Hungar, 18 (1983), 427 – 428.
- [8] Mushtaq, Q and Q. Iqbal, Decomposition of a locally associative LA-semigroup, Semigroup Forum, 41 (1990), 154 – 164.
- [9] Pilz, G, *Near rings, The theory and its applications*, North Holland Publishing Company Amsterdam. New York. Revised edition, 1983.
- [10] Protic, P. V and N. Stevanovic, AG-test and some general properties of Abel-Grassmann's groupoids, PU. M. A, 4, 6 (1995), 371 – 383.

DEPARTMENT OF MATHEMATICS, QUAID-I-AZAM UNIVERSITY, ISLAMABAD, PAKISTAN.

E-mail address: qmushtaq@isb.apollo.net.pk

E-mail address: madadmath@yahoo.com