MORE SUPPLEMENTS TO A CLASS OF LOGARITHMICALLY COMPLETELY MONOTONIC FUNCTIONS ASSOCIATED WITH THE GAMMA FUNCTION

SENLIN GUO AND FENG QI

ABSTRACT. In this article, a necessary and sufficient condition and a necessary condition are established for a function involving the gamma function to be logarithmically completely monotonic on $(0, \infty)$. As applications of the necessary and sufficient condition, some inequalities for bounding the psi and polygamma functions and the ratio of two gamma functions are derived.

This is a continuator of the paper [12].

1. Introduction

Recall [3, 23] that a positive function f is said to be logarithmically completely monotonic on an interval I if f has derivatives of all orders on I and

$$(-1)^n [\ln f(x)]^{(n)} \ge 0. \tag{1}$$

This kind of functions has very closer relationships with the Laplace transforms, Stieltjes transforms and infinitely divisible completely monotonic functions. For more detailed information, please refer to [4, 8, 11, 12, 20, 21, 24] and related references therein.

It is well-known that the classical Euler gamma function is defined for x > 0 by

$$\Gamma(z) = \int_0^\infty t^{x-1} e^{-t} \, \mathrm{d}t. \tag{2}$$

The logarithmic derivative of $\Gamma(z)$, denoted by $\psi(z) = \frac{\Gamma'(z)}{\Gamma(z)}$, is called the psi function, and $\psi^{(k)}$ for $k \in \mathbb{N}$ are called the polygamma functions.

For $\alpha \in \mathbb{R}$ and $\beta \geq 0$, define

$$f_{\alpha,\beta,\pm 1}(x) = \left[\frac{e^x \Gamma(x+\beta)}{x^{x+\beta-\alpha}}\right]^{\pm 1}, \quad x \in (0,\infty).$$
 (3)

The investigation of the function $f_{\alpha,\beta,\pm 1}(x)$ has a long history. In what follows, we would like to give a short survey in the literature.

In [15, Theorem 1], for showing

$$\frac{b^{b-1}}{a^{a-1}}e^{a-b} < \frac{\Gamma(b)}{\Gamma(a)} < \frac{b^{b-1/2}}{a^{a-1/2}}e^{a-b}$$
(4)

for b > a > 1, monotonic properties of the functions $\ln f_{\alpha,0,+1}(x)$ and $\ln f_{\alpha,0,+1}(x)$ on $(1,\infty)$ were obtained.

In [17, Theorem 2.1], the function $f_{\alpha,0,+1}(x)$ for $\alpha \leq 1$ was proved to be logarithmically completely monotonic on $(0,\infty)$. In [13, Theorem 2.1], the functions

The second author was supported partially by the China Scholarship Council.

This paper was typeset using A_MS -IAT_EX.

²⁰⁰⁰ Mathematics Subject Classification. Primary 33B15, 26A48, 26A51; Secondary 26D20, 65R10.

Key words and phrases. Necessary and sufficient condition, logarithmically completely monotonic function, inequality, ratio, gamma function, psi function, polygamma function, application.

 $f_{\alpha,0,+1}(x)$ and $f_{\alpha,0,-1}(x)$ were proved to be logarithmically completely monotonic on $(0,\infty)$ if and only if $\alpha \leq \frac{1}{2}$ and $\alpha \geq 1$ respectively. These results were mentioned in [14, Theorem 2.1] later. However, we do not think the proof in [13] for the necessity is convincible.

In [2, Theorem 3.2], it was recovered that the function $f_{1/2,0,+1}(x)$ is decreasing and logarithmically convex from $(0,\infty)$ onto $(\sqrt{2\pi},\infty)$ and that the function $f_{1,0,+1}(x)$ is increasing and logarithmically concave from $(0,\infty)$ onto $(1,\infty)$.

In [1, p. 376, Theorem 2], it was presented that the function $f_{\alpha,0,+1}(x)$ is decreasing on (c,∞) for $c\geq 0$ if and only if $\alpha\leq \frac{1}{2}$ and increasing on (c,∞) if and only if

$$\alpha \ge \begin{cases} c[\ln c - \psi(c)] & \text{if } c > 0, \\ 1 & \text{if } c = 0. \end{cases}$$
 (5)

The necessary and sufficient conditions for the functions $f_{\alpha,0,+1}(x)$ and $f_{\alpha,0,-1}(x)$ to be logarithmically completely monotonic on $(0,\infty)$, stated in [13, Theorem 2.1] and [14, Theorem 2.1] and mentioned above, were recovered in [5, Theorem 2]. Moreover, the function $f_{\alpha,\beta,+1}(x)$ was proved in [5, Theorem 1] to be logarithmically completely monotonic on $(0,\infty)$ if $2\alpha \le 1 \le \beta$. Using monotonic properties of the functions $f_{1/2,0,+1}(x)$ and $f_{1,0,-1}(x)$, the inequality (4) was extended in [5, Remark 1] from b > a > 1 to b > a > 0.

In [25], after proving once again the logarithmically completely monotonic property of the functions $f_{1/2,0,+1}(x)$ and $f_{1,0,-1}(x)$, in virtue of Jensen's inequality for convex functions, the upper and lower bounds were established: For positive numbers x and y, the inequality

$$\frac{x^{x-1/2}y^{y-1/2}}{[(x+y)/2]^{x+y-1}} \le \frac{\Gamma(x)\Gamma(y)}{[\Gamma((x+y)/2)]^2} \le \frac{x^{x-1}y^{y-1}}{[(x+y)/2]^{x+y-2}} \tag{6}$$

holds, where the middle term in (6) is called Gurland's ratio [16]. The left-hand side inequality in (6) is same as the corresponding one in [16, Theorem 1], but their upper bounds do not include each other.

Recently, some new conclusions on logarithmically completely monotonic properties of the function $f_{\alpha,\beta,+1}(x)$ were procured in [12, Theorem 1]:

- (1) If $\beta \in (0, \infty)$ and $\alpha \leq 0$, then $f_{\alpha,\beta,+1}(x)$ is logarithmically completely monotonic on $(0,\infty)$;
- (2) If $\beta \in (0, \infty)$ and $f_{\alpha, \beta, +1}(x)$ is a logarithmically completely monotonic function on $(0, \infty)$, then $\alpha \leq \min\{\beta, \frac{1}{2}\}$;
- (3) If $\beta \geq 1$, then $f_{\alpha,\beta,+1}(x)$ is logarithmically completely monotonic on $(0,\infty)$ if and only if $\alpha \leq \frac{1}{2}$.

As direct consequences of these monotonic properties, it is deduced immediately that if x and y are positive numbers with $x \neq y$, then

(1) the inequality

$$\frac{\Gamma(x+\beta)}{\Gamma(y+\beta)} < \frac{x^{x+\beta-\alpha}}{y^{y+\beta-\alpha}} e^{1/(y-x)} \tag{7}$$

for $\beta \geq 1$ and x > y > 0 holds true if and only if $\alpha \leq \frac{1}{2}$;

(2) the inequality (16) for $\beta \in (0, \infty)$ holds true also if $\alpha \leq 0$.

In this paper, for continuing the work in the paper [12], we consider logarithmically completely monotonic properties of the function $f_{\alpha,\beta,-1}(x)$ on $(0,\infty)$.

The main results of this paper are as follows.

Theorem 1. If the function $f_{\alpha,\beta,-1}(x)$ is logarithmically completely monotonic on $(0,\infty)$, then either $\beta > 0$ and $\alpha \ge \max\{\beta,\frac{1}{2}\}$ or $\beta = 0$ and $\alpha \ge 1$.

Theorem 2. If $\beta \geq \frac{1}{2}$, the necessary and sufficient condition for the function $f_{\alpha,\beta,-1}(x)$ to be logarithmically completely monotonic on $(0,\infty)$ is $\alpha \geq \beta$.

As the first application of Theorem 2, the following inequalities are derived by using logarithmically completely monotonic properties of the function $f_{\alpha,\beta,\pm 1}(x)$ on $(0,\infty)$.

Theorem 3. Let β be a positive number.

(1) For $k \in \mathbb{N}$, double inequalities

$$\ln x - \frac{1}{x} \le \psi(x) \le \ln x - \frac{1}{2x} \tag{8}$$

and

$$\frac{(k-1)!}{x^k} + \frac{k!}{2x^{k+1}} \le (-1)^{k+1} \psi^{(k)}(x) \le \frac{(k-1)!}{x^k} + \frac{k!}{x^{k+1}}$$
(9)

hold in $(0, \infty)$.

(2) When $\beta > 0$, inequalities

$$\psi(x+\beta) \le \ln x + \frac{\beta}{x} \tag{10}$$

and

$$(-1)^k \psi^{(k-1)}(x+\beta) \ge \frac{(k-2)!}{x^{k-1}} - \frac{\beta(k-1)!}{x^k} \tag{11}$$

hold on $(0, \infty)$ for $k \geq 2$.

(3) When $\beta \geq \frac{1}{2}$, inequalities

$$\psi(x+\beta) \ge \ln x \quad and \quad (-1)^k \psi^{(k-1)}(x+\beta) \le \frac{(k-2)!}{x^{k-1}}$$
 (12)

hold on $(0, \infty)$ for $k \geq 2$.

(4) When $\beta \geq 1$, inequalities

$$\psi(x+\beta) \le \ln x + \frac{\beta - 1/2}{x} \tag{13}$$

and

$$(-1)^k \psi^{(k-1)}(x+\beta) \ge \frac{(k-2)!}{x^{k-1}} - \frac{(\beta - 1/2)(k-1)!}{x^k}$$
(14)

holds on $(0, \infty)$ for $k \geq 2$.

As the second application of Theorem 2, the following inequalities are derived by using logarithmically convex properties of the function $f_{\alpha,\beta,\pm 1}(x)$ on $(0,\infty)$.

Theorem 4. Let $n \in \mathbb{N}$, $x_k > 0$ for $1 \le k \le n$, $p_k \ge 0$ satisfying $\sum_{k=1}^n p_k = 1$. If either $\beta > 0$ and $\alpha \le 0$ or $\beta \ge 1$ and $\alpha \le \frac{1}{2}$, then

$$\frac{\prod_{k=1}^{n} [\Gamma(x_k + \beta)]^{p_k}}{\Gamma(\sum_{k=1}^{n} p_k x_k + \beta)} \ge \frac{\prod_{k=1}^{n} x_k^{p_k(x_k + \beta - \alpha)}}{(\sum_{k=1}^{n} p_k x_k)^{\sum_{k=1}^{n} p_k x_k + \beta - \alpha}}.$$
 (15)

If $\alpha \geq \beta \geq \frac{1}{2}$, then the inequality (15) reverses.

As the final application of Theorem 2, the following inequality may be derived by using the decreasingly monotonic property of the function $f_{\alpha,\beta,-1}(x)$ on $(0,\infty)$.

Theorem 5. If $\alpha \geq \beta \geq \frac{1}{2}$, then

$$I(x,y) < \left[\left(\frac{x}{y} \right)^{\alpha-\beta} \frac{\Gamma(x+\beta)}{\Gamma(y+\beta)} \right]^{1/(x-y)}$$
(16)

holds true for $x, y \in (0, \infty)$ with $x \neq y$, where

$$I(a,b) = \frac{1}{e} \left(\frac{b^b}{a^a}\right)^{1/(b-a)} \tag{17}$$

for a > 0 and b > 0 with $a \neq b$ is the identric or exponential mean.

Remark 1. For $\beta \in \mathbb{R}$, let

$$h_{\beta,\pm 1}(x) = \left[\frac{e^x \Gamma(x+1)}{(x+\beta)^{x+\beta}}\right]^{\pm 1} \tag{18}$$

on $(\max\{0, -\beta\}, \infty)$. In [10, 11], it was showed that the functions $h_{\beta,+1}(x)$ and $h_{\beta,-1}(x)$ are logarithmically completely monotonic if and only if $\beta \geq 1$ and $\beta \leq \frac{1}{2}$ respectively. As consequences of monotonic results of the function $h_{\beta,\pm 1}(x)$, the following two-sided inequality was derived in [11]:

$$\frac{(x+1)^{x+1}}{(y+1)^{y+1}} e^{y-x} < \frac{\Gamma(x+1)}{\Gamma(y+1)} < \frac{(x+1/2)^{x+1/2}}{(y+1/2)^{y+1/2}} e^{y-x}$$
(19)

for y > x > 0, where the constants 1 and $\frac{1}{2}$ in the very left and the very right sides of the two-sided inequality (19) cannot be replaced, respectively, by smaller and larger numbers.

Remark 2. In [9], it was showed that the function

$$h(x) = \frac{e^x \Gamma(x)}{x^{x[1-\ln x + \psi(x)]}} \tag{20}$$

on $(0, \infty)$ has a unique maximum e at x = 1, with the following two limits

$$\lim_{x \to 0^+} h(x) = 1 \quad \text{and} \quad \lim_{x \to \infty} h(x) = \sqrt{2\pi}. \tag{21}$$

As consequences of the monotonicity of h(x), it was concluded in [9] that the following inequality

$$\frac{x^{x[\ln x - \psi(x) - 1]}}{y^{y[\ln y - \psi(y) - 1]}} e^{y - x} < \frac{\Gamma(y)}{\Gamma(x)}$$

$$(22)$$

holds true for $y > x \ge 1$. If $0 < x < y \le 1$, the inequality (22) is reversed.

Remark 3. It is worthwhile to point out that [20, Thorem 1.3] is equivalent to necessary and sufficient conditions in [5, Theorem 2] and [13, Theorem 2.1] for the functions $f_{1/2,0,+1}(x)$ and $f_{1,0,-1}(x)$ to be logarithmically completely monotonic on $(0,\infty)$. See also [11, 12] and related references therein.

2. Proofs of main results

Now we are in a position to prove our theorems.

Proof of Theorem 1. Suppose that $f_{\alpha,\beta,-1}(x)$ is logarithmically completely monotonic on $(0,\infty)$. Then

$$[\ln f_{\alpha,\beta,-1}(x)(x)]' = \ln x - \psi(x+\beta) + \frac{\beta - \alpha}{x} \le 0,$$
 (23)

from which we have

$$\beta - \alpha \le x[\psi(x+\beta) - \ln x], \quad x \in (0, \infty). \tag{24}$$

If $\beta > 0$, then

$$\beta - \alpha \le \lim_{x \to 0^+} [x\psi(x+\beta) - x \ln x] = 0.$$

That is

$$\alpha \ge \beta.$$
 (25)

Using the asymptotic formula

$$\psi(x) = \ln x - \frac{1}{2x} + O\left(\frac{1}{x^2}\right), \quad x \to \infty, \tag{26}$$

see [6, p. 47], in (24) for $\beta > 0$, we obtain

$$\begin{split} \beta - \alpha & \leq \lim_{x \to \infty} x \bigg[\ln(x + \beta) - \frac{1}{2(x + \beta)} + O\bigg(\frac{1}{x^2}\bigg) - \ln x \bigg] \\ & = \lim_{x \to \infty} \bigg[x \ln\bigg(1 + \frac{\beta}{x}\bigg) \bigg] - \frac{1}{2} \\ & = \beta \lim_{x \to \infty} \bigg[\frac{x}{\beta} \ln\bigg(1 + \frac{\beta}{x}\bigg) \bigg] - \frac{1}{2} \\ & = \beta - \frac{1}{2}, \end{split}$$

from which we get

$$\alpha \ge \frac{1}{2}.\tag{27}$$

Combining (25) and (27) yields

$$\alpha \ge \max \left\{ \beta, \frac{1}{2} \right\} \quad \text{if} \quad \beta > 0.$$
 (28)

If $\beta = 0$, considering $f_{\alpha,0,-1}(x) = f_{\alpha,1,-1}(x)$ and (28) yields $\alpha \ge \max\{1, \frac{1}{2}\} = 1$. The proof is complete.

Proof of Theorem 2. By Theorem 1, the necessary condition is obtained readily. Differentiating (23) and making use of

$$\psi^{(n)}(x) = (-1)^{n+1} \int_0^\infty \frac{t^n}{1 - e^{-t}} e^{-xt} \, \mathrm{d}t, \quad x \in (0, \infty)$$

and

$$\frac{1}{x^n} = \frac{1}{\Gamma(n)} \int_0^\infty t^{n-1} e^{-xt} \, \mathrm{d}t, \quad x \in (0, \infty),$$

see [7, p. 884], gives

$$(-1)^{n} [\ln f_{\alpha,\beta,-1}(x)]^{(n)} = \frac{(n-2)!}{x^{n-1}} - (-1)^{n} \psi^{(n-1)}(x+\beta) - \frac{(\beta-\alpha)(n-1)!}{x^{n}}$$

$$= \int_{0}^{\infty} t^{n-2} e^{-xt} dt - \int_{0}^{\infty} \frac{t^{n-1}}{1-e^{-t}} e^{-(x+\beta)t} dt - (\beta-\alpha) \int_{0}^{\infty} t^{n-1} e^{-xt} dt$$

$$= \int_{0}^{\infty} \left[\alpha - \beta - \frac{1}{t} \left(e^{(1-\beta)t} \frac{t}{e^{t} - 1} - 1 \right) \right] t^{n-1} e^{-xt} dt$$
(29)

for $n \geq 2$. The inequality

$$\frac{t}{e^t-1}<\frac{1}{e^{t/2}},\quad t\in(0,\infty) \tag{30}$$

was ever used in [18, 19, 22]. Substituting it into (29) leads to

$$(-1)^n [\ln f_{\alpha,\beta,-1}(x)]^{(n)} \ge \int_0^\infty \left[\alpha - \beta - \frac{e^{(1/2-\beta)t} - 1}{t} \right] t^{n-1} e^{-xt} \, \mathrm{d}t, \quad n \ge 2.$$

Since the function

$$\frac{e^{(1/2-\beta)t}-1}{t}, \quad t \in (0,\infty)$$
(31)

is increasing, if $\alpha \geq \beta \geq \frac{1}{2}$, then

$$(-1)^{n} [\ln f_{\alpha,\beta,-1}(x)]^{(n)} \ge (\alpha - \beta) \int_{0}^{\infty} t^{n-1} e^{-xt} \, \mathrm{d}t \ge 0, \quad n \ge 2.$$
 (32)

By using (26), it follows that

$$[\ln f_{\alpha,\beta,-1}(x)(x)]' = \ln \left(1 + \frac{\beta}{x}\right) - \frac{1}{2(x+\beta)} + \frac{\alpha-\beta}{x} + O\left(\frac{1}{x^2}\right)$$

as $x \to \infty$, thus for all α and β ,

$$\lim_{x \to \infty} [\ln f_{\alpha,\beta,-1}(x)(x)]' = 0.$$
 (33)

From (33) and (32), we have

$$[\ln f_{\alpha,\beta,-1}(x)]' \le 0 \quad \text{if } \alpha \ge \beta \ge \frac{1}{2}.$$

Thus $(-1)^n [\ln f_{\alpha,\beta,-1}(x)]^{(n)} \geq 0$ are valid for all $n \in \mathbb{N}$. The proof is complete. \square

Proof of Theorem 3. If $f_{\alpha,\beta,-1}(x)$ is logarithmically completely monotonic on $(0,\infty)$, then

$$(-1)^k [\ln f_{\alpha,\beta,-1}(x)]^{(k)} \ge 0$$

on $(0,\infty)$ for $k\in\mathbb{N}$, which is equivalent to

$$\psi(x+\beta) \ge \ln x + \frac{\beta - \alpha}{x} \tag{34}$$

and, for $k \geq 2$,

$$(-1)^k \psi^{(k-1)}(x+\beta) \le \frac{(k-2)!}{x^{k-1}} - \frac{(\beta-\alpha)(k-1)!}{x^k}.$$
 (35)

Hence, Theorem 2 implies inequalities in (12).

If $\beta \geq 1$, Theorem 1 in [12] said that the function $f_{\alpha,\beta,+1}(x)$ is logarithmically completely monotonic on $(0,\infty)$ if and only if $\alpha \leq \frac{1}{2}$, this means that inequalities in (34) and (35) are reversed, and so inequalities in (13) and (14) are valid.

If $\beta > 0$ and $\alpha \leq 0$, Theorem 1 in [12] also said that the function $f_{\alpha,\beta,+1}(x)$ is logarithmically completely monotonic on $(0,\infty)$, this means that inequalities in (34) and (35) are also reversed, and so inequalities (10) and (11) are valid.

When $\beta = 0$, several mathematicians have proved that the functions $f_{\alpha,0,+1}(x)$ and $f_{\alpha,0,-1}(x)$ are logarithmically completely monotonic on $(0,\infty)$ if and only if $\alpha \leq \frac{1}{2}$ and $\alpha \geq 1$ respectively, which implies by reasoning as above the double inequalities (8) and (9). The proof of Theorem 3 is complete.

Proof of Theorem 4. The first conclusion in [12, Theorem 1] implies that the function $f_{\alpha,\beta,+1}(x)$ is logarithmically convex for $\beta > 0$ and $\alpha \leq 0$ on $(0,\infty)$. Combining this with Jensen's inequality for convex functions yields

$$\ln \frac{\exp\left(\sum_{k=1}^{n} p_k x_k\right) \Gamma\left(\sum_{k=1}^{n} p_k x_k + \beta\right)}{\left(\sum_{k=1}^{n} p_k x_k\right)^{\sum_{k=1}^{n} p_k x_k + \beta - \alpha}} \le \sum_{k=1}^{n} p_k \ln \frac{\exp(x_k) \Gamma(x_k + \beta)}{x_k^{x_k + \beta - \alpha}}, \tag{36}$$

where $n \in \mathbb{N}$, $x_k > 0$ for $1 \le k \le n$, $p_k \ge 0$ satisfying $\sum_{k=1}^n p_k = 1$, $\beta > 0$ and $\alpha \le 0$. Rearranging it leads to the inequality (15).

The third conclusion in [12, Theorem 1] implies that the function $f_{\alpha,\beta,+1}(x)$ is also logarithmically convex for $\beta \geq 1$ and $\alpha \leq \frac{1}{2}$ on $(0,\infty)$, hence, the inequality (36) is also valid for $\beta \geq 1$ and $\alpha \leq \frac{1}{2}$.

Theorem 2 above implies that the function $f_{\alpha,\beta,+1}(x)$ is logarithmically concave for $\alpha \geq \beta \geq \frac{1}{2}$ on $(0,\infty)$, therefore, the inequality (36) is reversed. Theorem 4 is proved.

Proof of Theorem 5. Theorem 2 implies that the function $f_{\alpha,\beta,-1}(x)$ is decreasing on $(0,\infty)$ if $\alpha \geq \beta \geq \frac{1}{2}$, this is,

$$\frac{e^{y}\Gamma(y+\beta)}{y^{y+\beta-\alpha}} > \frac{e^{x}\Gamma(x+\beta)}{x^{x+\beta-\alpha}}$$

for y > x > 0, which can be rearranged as

$$\frac{\Gamma(y+\beta)}{\Gamma(x+\beta)} > e^{x-y} \frac{y^{y+\beta-\alpha}}{x^{x+\beta-\alpha}},$$

$$\left[\left(\frac{y}{x} \right)^{\alpha-\beta} \frac{\Gamma(y+\beta)}{\Gamma(x+\beta)} \right]^{1/(y-x)} > \frac{1}{e} \left(\frac{y^y}{x^x} \right)^{1/(y-x)}.$$

The proof of Theorem 5 is complete.

References

- H. Alzer, On some inequalities for the gamma and psi functions, Math. Comp. 66 (1997), no. 217, 373–389.
- [2] G. D. Anderson, R. W. Barnard, K. C. Richards, M. K. Vamanamurthy and M. Vuorinen, Inequalities for zero-balanced hypergeometric functions, Trans. Amer. Math. Soc. 347 (1995), no. 5, 1713–1723.
- [3] R. D. Atanassov and U. V. Tsoukrovski, Some properties of a class of logarithmically completely monotonic functions, C. R. Acad. Bulgare Sci. 41 (1988), 21–23.
- [4] C. Berg, Integral representation of some functions related to the gamma function, Mediterr. J. Math. 1 (2004), no. 4, 433–439.
- [5] Ch.-P. Chen and F. Qi, Logarithmically completely monotonic functions relating to the gamma function, J. Math. Anal. Appl. 321 (2006), no. 1, 405–411.
- [6] A. Erdélyi(ed), Higher Transcendental Functions, Vol 1, McGraw-Hill, New York, 1953.
- [7] I. S. Gradshteyn and I. M. Ryzhik, Table of Integrals, Series and Products, 6th Ed., Academic Press, New York, 2000.
- [8] A. Z. Grinshpan and M. E. H. Ismail, Completely monotonic functions involving the Gamma and q-gamma functions, Proc. Amer. Math. Soc. 134 (2006), 1153–1160.
- [9] B.-N. Guo, Y.-J. Zhang and F. Qi, Refinements and sharpenings of some double inequalities for bounding the gamma function, J. Inequal. Pure Appl. Math. 9 (2008), no. 1, Art. 17; Available online at http://jipam.vu.edu.au/article.php?sid=953.
- [10] S. Guo, Some classes of completely monotonic functions involving the gamma function, Internat. J. Pure Appl. Math. 30 (2006), no. 4, 561–566.
- [11] S. Guo, F. Qi, and H. M. Srivastava, Necessary and sufficient conditions for two classes of functions to be logarithmically completely monotonic, Integral Transforms Spec. Funct. 18 (2007), no. 11, 819–826.
- [12] S. Guo, F. Qi, and H. M. Srivastava, Supplements to a class of logarithmically completely monotonic functions associated with the gamma function, Appl. Math. Comput. 197 (2008), no. 2, 768–774.
- [13] M. E. H. Ismail, L. Lorch, and M. E. Muldoon, Completely monotonic functions associated with the gamma function and its q-analogues, J. Math. Anal. Appl. 116 (1986), 1–9.
- [14] M. E. H. Ismail and M. E. Muldoon, Inequalities and monotonicity properties for gamma and q-gamma functions, in: R.V.M. Zahar (Ed.), Approximation and Computation: A Festschrift in Honour of Walter Gautschi, ISNM, Vol. 119, BirkhRauser, Basel, 1994, 309–323.
- [15] J. D. Kečlić and P. M. Vasić, Some inequalities for the gamma function, Publ. Inst. Math. (Beograd) (N. S.) 11 (1971), 107–114.
- [16] M. Merkle, Gurland's ratio for the gamma function, Comput. Math. Appl. 49 (2005), 389–406.
- [17] M. E. Muldoon, Some monotonicity properties and characterizations of the gamma function, Aequationes Math. 18 (1978), 54–63.
- [18] F. Qi, An integral expression and some inequalities of Mathieu type series, Rostock. Math. Kolloq. 58 (2004), 37–46.
- [19] F. Qi, Integral expression and inequalities of Mathieu type series, RGMIA Res. Rep. Coll. 6 (2003), no. 2, Art. 10; Available online at http://www.staff.vu.edu.au/rgmia/v6n2.asp.
- [20] F. Qi, Three classes of logarithmically completely monotonic functions involving gamma and psi functions, Integral Transforms Spec. Funct. 18 (2007), no. 7, 503–509.
- [21] F. Qi and Ch.-P. Chen, A complete monotonicity property of the gamma function, J. Math. Anal. Appl. 296 (2004), no. 2, 603–607.
- [22] F. Qi, Ch.-P. Chen, and B.-N. Guo, Notes on double inequalities of Mathieu's series, Internat. J. Math. Sci. 2005 (2005), no. 16, 2547–2554.
- [23] F. Qi and B.-N. Guo, Complete monotonicities of functions involving the gamma and digamma functions, RGMIA Res. Rep. Coll. 7 (2004), no. 1, Art. 8, 63-72; Available online at http://www.staff.vu.edu.au/rgmia/v7n1.asp.

- [24] F. Qi, B.-N. Guo and Ch.-P. Chen, Some completely monotonic functions involving the gamma and polygamma functions, J. Aust. Math. Soc. 80 (2006), 81–88.
- [25] Y.-J. Wei, S.-L. Zhang, and Ch.-P. Chen, Logarithmically completely monotonic functions and Gurland's ratio for the gamma function, Adv. Stud. Contemp. Math. 15 (2007), no. 2, 253–257.
- (S. Guo) Department of Mathematics, Zhongyuan University of Technology, Zhengzhou City, Henan Province, 450007, China

 $E ext{-}mail\ address: sguo@hotmail.com, senlinguo@gmail.com}$

(F. Qi) RESEARCH INSTITUTE OF MATHEMATICAL INEQUALITY THEORY, HENAN POLYTECHNIC UNIVERSITY, JIAOZUO CITY, HENAN PROVINCE, 454010, CHINA

 $E\text{-}mail\ address:$ qifeng6180gmail.com, qifeng6180hotmail.com, qifeng6180qq.com URL: http://qifeng618.spaces.live.com