POWERS OF COMPONENTWISE LINEAR IDEALS

JÜRGEN HERZOG, TAKAYUKI HIBI AND HIDEFUMI OHSUGI

Dedicated to the memory of Professor Masayoshi Nagata

ABSTRACT. We give criteria for graded ideals to have the property that all their powers are componentwise linear. Typical examples to which our criteria can be applied include the vertex cover ideals of certain finite graphs.

INTRODUCTION

Let K be a field, let $S = K[x_1, \ldots, x_n]$ the polynomial ring in n variables over K, and let $\mathfrak{m} = (x_1, \ldots, x_n)$ be its graded maximal ideal. Let $I \subset S$ be a graded ideal. The ideal I is said to be *componentwise linear*, if for all j the ideal $I_{\langle j \rangle} = (I_j)$, generated by the jth component of I, has a linear resolution. It is known that I is componentwise linear if $(I_{[j]})$ has a linear resolution for all $j \leq \operatorname{reg} I$, see [10]. In particular, ideals with linear resolution are componentwise linear. Typical examples of componentwise linear ideals are stable and squarefree stable ideals.

Naively one would expect that for an ideal with linear resolution, all its powers have a linear resolution as well. But this is not the case. A first counterexample was given by Terai who observed that, if the characteristic of K is zero, then the Stanley–Reisner ideal of the natural triangulation of the real projective plane has a linear resolution, but the square does not. For ideals with linear resolution there exist criteria, among them the socalled x-condition [12], that allow to test whether all of its powers have a linear resolution as well. For example it was shown [12], by using the x-condition, that all powers of a monomial ideal with linear resolution generated in degree 2 have linear resolutions. But this condition fails in many other cases, for example for the ideal of 2-minors of a generic symmetric 3×3 -matrix which as we shall see has the property that all of its powers have a linear resolution. To prove this we shall use the following criterion (Corollary 1.5) which is one of the main results of the paper. Let K be an infinite field and $I \subset S$ a graded ideal. Then all powers of I are componentwise linear if and only if a generic sequence of linear forms generating the K-vector space S_1 is a d-sequence with respect to the Rees ring R(I) of I. Proposition 1.7 provides a Gröbner basis condition on the defining ideal R(I) that guarantees that a given K-basis of S_1 is a d-sequence with respect to R(I). In [19, Proposition 3.1] Römer presents a result related to our main theorem, characterizing standard bigraded K-algebras with x-regularity 0.

Our criterion can be easily checked by any computer algebra system for graded ideals whose number of generators is not too big. Interesting examples are the ideals defining

¹⁹⁹¹ Mathematics Subject Classification. Primary: 13A30 Secondary: 13D45.

This paper was written while the first author was staying at Osaka University.

rational normal scrolls. In all cases we checked it turned out that all powers of these ideals have a linear resolution. Due to these computations and due to other known cases [1] and [7], one may expect that all powers of ideals defining rational normal scrolls have a linear resolution. One even may expect this property holds true for any graded ideal I for which S/I is Cohen-Macaulay with minimal multiplicity. However Conca [4] gave an example of a graded ideal I whose residue class ring is Cohen-Macaulay with minimal multiplicity, but for which I^2 does not have a linear resolution. The best one could hope is that any reduced graded ideal $I \subset S$ for which S/I is Cohen-Macaulay with minimal multiplicity has the property that all its powers have a linear resolution.

In the second section of this paper we consider vertex cover ideals of chordal graphs. The vertex cover ideal of a graph G is the Alexander dual of the edge ideal of G. In [13] it was shown that the edge ideal of a chordal graph is Cohen-Macaulay if and only if it is unmixed. This result indicated that edge ideals of chordal graphs might be sequentially Cohen-Macaulay, and this was indeed shown by Francisco and Van Tuyl [9, Theorem 3.2]. According to [10, Theorem 2.1], this result is equivalent to the statement that the vertex cover ideal of a chordal graph is componentwise linear. In this paper we show that all powers of the vertex cover ideal of a star graph are componentwise linear (Theorem 2.3), and that those of a Cohen-Macaulay chordal graphs. By these results and computational evidence we are lead to conjecture that all powers of the vertex cover ideal of a chordal graphs.

1. The criterion

The criterion we are going to prove is based on results on approximation complexes in [15] and on a result of Römer [18] and Yanagawa [20, Proposition 4.9], which they proved independently. This result appeared first in Römer's dissertation. A new proof of their result in a more general frame is given in [16, Theorem 5.6].

Let M be a finitely generated graded S-module with graded minimal free resolution (G, φ) . Replacing all entries which are not linear by zero in the matrices describing the differentials φ_i in G, one obtains a complex, which is called the *linear part* of G. The linear part of G can be described more naturally as follows: we define a filtration \mathcal{F} on G by setting $\mathcal{F}_j G_i = \mathfrak{m}^{j-i} G_i$ for all i and j, and denote by $\operatorname{gr}_{\mathfrak{m}}(G)$ the associated graded complex. This complex is isomorphic to the linear part of G.

The module M is called *componentwise linear*, if for all j the submodules generated by M_j have a linear resolution.

Theorem 1.1 (Römer, Yanagawa). Let M be a finitely generated graded S-module with minimal graded free resolution G. The following conditions are equivalent:

- (a) M is componentwise linear.
- (b) $\operatorname{gr}_{\mathfrak{m}}(G)$ is acyclic.

If the equivalent conditions hold, $H_0(\operatorname{gr}_{\mathfrak{m}}(G)) \cong \operatorname{gr}_{\mathfrak{m}}(M)$.

The link to approximation complexes is given by the next theorem. As usual, the \mathcal{M} -complex of M with respect to \mathfrak{m} is denoted by $\mathcal{M}(\mathfrak{m}; M)$. It is a linear complex of

free S-modules. In other words, all entries of the matrices describing the differentials of this complex are linear forms. We refer the reader to [14] for a detailed description of approximation complexes. The following result ([15, Theorem 5.1]) provides another interpretation of the linear part of a resolution.

Theorem 1.2 (Herzog, Simis, Vasconcelos). Let M be a finitely generated graded S-module with minimal graded free resolution G. Then

$$\operatorname{gr}_{\mathfrak{m}}(G) \cong \mathcal{M}(\mathfrak{m}; M).$$

In view of Theorem 1.1 and Theorem 1.2 we would like to know when the \mathcal{M} -complex is acyclic. An answer to this question is given by the next result [15, Theorem 4.1].

Theorem 1.3 (Herzog, Simis, Vasconcelos). Let M be a finitely generated graded S-module and I a proper, graded ideal. Then the following conditions are equivalent:

- (a) $\mathcal{M}(I; M)$ is acyclic.
- (b) I is generated by a d-sequence with respect to M.

The concept of *d*-sequences was introduced by Huneke [17]. Recall that a sequence $\mathbf{z} = z_1, \ldots, z_m$ of elements of *S* with $I = (\mathbf{z})$ is called a *d*-sequence, if

$$(z_1, \ldots, z_{i-1})M : {}_M z_i \cap IM = (z_1, \ldots, z_{i-1})M$$
 for $i = 1, \ldots, m$.

After these preparations we are ready to state and prove our main result.

Theorem 1.4. Let K be an infinite field, and let $I \subset S = K[x_1, \ldots, x_n]$ be a graded ideal. If there exists a K-basis $\mathbf{z} = z_1, \ldots, z_n$ of S_1 such that \mathbf{z} is a d-sequence with respect to the Rees algebra R(I), then all powers of I are componentwise linear.

Conversely, let \mathbf{z} be a generic K-basis of S_1 . If all powers of I are componentwise linear, then \mathbf{z} is a d-sequence with respect to R(I).

Proof. Let $\mathbf{z} = z_1, \ldots, z_n$ be a K-basis of S_1 which is a d-sequence with respect to the Rees algebra R(I). Then

(1)
$$(z_1, \ldots, z_{i-1})R(I): z_i \cap \mathfrak{m}R(I) = (z_1, \ldots, z_{i-1})R(I)$$

The Rees ring R(I) is naturally \mathbb{Z} -graded with $R(I)_k = I^k t^k$ for all k. Since the elements z_i in this grading are of degree 0, equality (4) yields the equalities

$$(z_1,\ldots,z_{i-1})I^k$$
: $z_i \cap \mathfrak{m}I^k = (z_1,\ldots,z_{i-1})I^k$, $k = 0, 1, 2, \ldots$

In other words, \mathbf{z} is a *d*-sequence with respect to I^k for all k. Therefore Theorem 1.3 implies that the approximation complexes $\mathcal{M}(\mathfrak{m}; I^k)$ are acyclic. Let $G^{(k)}$ be the graded minimal free *S*-resolution of I^k . It follows from Theorem 1.2 that the associated graded complex $\operatorname{gr}_{\mathfrak{m}}(G^{(k)})$ is acyclic for all k. According Theorem 1.1 this implies that I^k is componentwise linear.

Conversely assume that all powers of I are componentwise linear. Then Theorem 1.1 and Theorem 1.2 imply that all the approximation complexes $\mathcal{M}(\mathfrak{m}; I^k)$ are acyclic. Hence Theorem 1.3 asserts that for each k there exists a K-basis of S_1 which is a d-sequence d-sequence with respect to I^k . The proof of [15, Theorem 4.1] shows how the d-sequence is constructed: choose any K-basis $\mathbf{z} = z_1, \ldots, z_n$ of S_1 with the property that for all i > 0 and all $j \in \{1, \ldots, n\}$ we have $H_i(z_1, \ldots, z_j; \operatorname{gr}_{\mathfrak{m}}(I^k))_{\ell} = 0$ for all $\ell \gg 0$. Then under the assumption that $\mathcal{M}(\mathfrak{m}; I^k)$ is acyclic, this sequence is a *d*-sequence with respect to I^k .

Suppose we can choose a K-basis \mathbf{z} of S_1 such that

(2)
$$((z_1,\ldots,z_j)\operatorname{gr}_{\mathfrak{m}}(I^k):z_{j+1})/(z_1,\ldots,z_j)\operatorname{gr}_{\mathfrak{m}}(I^k)$$

has finite length for all k and all j, then, since $\operatorname{reg}(\operatorname{gr}_{\mathfrak{m}}(I^k)) = 0$, [8, Proposition 20.20] implies that

(3)
$$(((z_1, \ldots, z_j) \operatorname{gr}_{\mathfrak{m}}(I^k) : z_{j+1})/(z_1, \ldots, z_j) \operatorname{gr}_{\mathfrak{m}}(I^k))_{\ell} = 0 \text{ for all } \ell > 0.$$

Observe that the regularity of $\operatorname{gr}_{\mathfrak{m}}(I^k)$ is indeed zero, because $\operatorname{gr}_{\mathfrak{m}}(I^k)$ is generated in degree 0 and the approximation complex $\mathcal{M}(\mathfrak{m}; I^k)$ provides a linear resolution of $\operatorname{gr}_{\mathfrak{m}}(I^k)$.

For a sequence satisfying (3) for all k it follows that $H_i(z_1, \ldots, z_j; \operatorname{gr}_{\mathfrak{m}}(I^k))_{\ell} = 0$ for all i > 0, all k and $\ell > 0$, and so z is a d-sequence with respect to all I^k , as explained before. This is equivalent to say that z is a d-sequence with respect to R(I). But how can we find a sequence \mathbf{z} such that property (2) (and consequently property (3)) is satisfied for all powers of I?

Suppose we have already chosen z_1, \ldots, z_j satisfying (2). We apply the graded version of [3, Proposition 2.5] to the standard graded $S/(z_1,\ldots,z_j)$ -algebra

$$A = \bigoplus_{k \ge 0} \operatorname{gr}_{\mathfrak{m}}(I^k)(z_1, \dots, z_j) \operatorname{gr}_{\mathfrak{m}}(I^k)$$

to conclude that

$$\mathcal{A} = \bigcup_{k \ge 0} \operatorname{Ass}(\operatorname{gr}_{\mathfrak{m}}(I^k)(z_1, \dots, z_j) \operatorname{gr}_{\mathfrak{m}}(I^k))$$

is a finite set. Hence since K is infinite we may choose a linear form $z_{i+1} \in S/(z_1, \ldots, z_i)$ which is contained in no prime ideal of \mathcal{A} different from the graded maximal ideal of $S/(z_1,\ldots,z_i)$. This is the desired next linear form in our sequence. Hence in each step of the constructing of \mathbf{z} we have an open choice. In other words, generic sequences will be d-sequences with respect to R(I).

Corollary 1.5. Let K be an infinite field, $I \subset S$ a graded ideal and \mathbf{z} a generic K-basis of S_1 . The following conditions are equivalent:

- (a) All powers of I are componentwise linear;
- (b) \mathbf{z} is a d-sequence with respect to R(I).

Let us analyze what it means that a sequence z_1, \ldots, z_n is a *d*-sequence with respect to R(I). After a change of coordinates we may assume that our given sequence is the sequence x_1, \ldots, x_n . Let f_1, \ldots, f_m a homogeneous system of generators of I, and let $T = S[y_1, \ldots, y_m]$ be the polynomial ring over S in the variables y_1, \ldots, y_m . We consider the surjective S-algebra homomorphism $T \to R(I)$ with $y_j \mapsto f_j$ for $j = 1, \ldots, m$. Then $R(I) \cong T/J$ where J is the kernel of the map $T \to R(I)$. Identifying R(I) with T/J, we see that x_1, \ldots, x_n is a *d*-sequence with respect to R(I) if and only if

$$(4)(x_1, \dots, x_{i-1}) + J \colon x_i \cap ((x_1, \dots, x_n) + J) = (x_1, \dots, x_{i-1}) + J \quad \text{for} \quad i = 1, \dots, n.$$

This is a condition which can be easily checked by any computer algebra system. For example, if we let I be the ideal of 2-minors of the generic symmetric matrix

$$\left(\begin{array}{ccc}a&b&c\\b&d&e\\c&e&f\end{array}\right),$$

one easily finds with CoCoA that a, b, e, d + f, c, f is a d-sequence on R(I), and hence all powers of I have a linear resolution.

Passing to the initial ideal of J with respect to some term order we can deduce a sufficient condition for the property that all powers of an ideal are componentwise linear. We denote by J_i the ideal which is the image of J under the canonical epimorphism $T \to T/(x_1, \ldots, x_i)$. Then (4) implies that x_1, \ldots, x_n is a *d*-sequence on R(I), if and only if

(5)
$$J_{i-1}: x_i \cap (x_i, \dots, x_n) + J_{i-1} = J_{i-1}$$
 for $i = 1, \dots, n$

Let < be a monomial order. We denote by in(I) the initial ideal with respect to this order.

Lemma 1.6. Suppose there exists a monomial order < on $T/(x_1, \ldots, x_{i-1})$ such that

(6)
$$\operatorname{in}(J_{i-1}): x_i \cap ((x_i, \dots, x_n) + \operatorname{in}(J_{i-1})) = \operatorname{in}(J_{i-1})$$

Then (5) holds for this integer i.

Proof. Let $f \in J_{i-1}: x_i \cap ((x_i, \ldots, x_n) + J_{i-1})$. Then $x_i f \in J_{i-1}$, and so $x_i \operatorname{in}(f) \in \operatorname{in}(J_{i-1})$. Therefore, $\operatorname{in}(f) \in \operatorname{in}(J_{i-1}): x_i$. On the other hand, since $f \in (x_i, \ldots, x_n) + J_{i-1}$, it follows that $\operatorname{in}(f) \in \operatorname{in}(x_1, \ldots, x_i) + J_{i-1}) \subset (x_1, \ldots, x_i) + \operatorname{in}(J_{i-1})$. Thus our hypothesis implies that $\operatorname{in}(f) \in \operatorname{in}(J_{i-1})$. Hence there exists $g \in J_{i-1}$ such that $\operatorname{in}(g) = \operatorname{in}(f)$. We may assume that the leading coefficients of f and g are 1, and set h = f - g. Then $x_i h = x_i f - x_i g \in J_{j-1}$ and $h \in (x_i, \ldots, x_n) + J_{i-1}$. Since $\operatorname{in}(h) < \operatorname{in}(f)$, we may assume by induction that $h \in J_{i-1}$. This then implies that $f \in J_{i-1}$, as desired. \Box

For a monomial ideal L we denote as usual by G(L) the unique minimal set of monomial generators of L. We also set $y^b = y_1^{b_1} \cdots y_m^{b_m}$ for any integer vector $b = (b_1, \ldots, b_m)$. Now the preceding considerations yield

Proposition 1.7. The sequence x_1, \ldots, x_n is d-sequence on R(I), and hence all powers of I are componentwise linear, if for each i there exists monomial order on $T/(x_1, \ldots, x_{i-1})$ such that whenever x_i divides $u \in G(in(J_{i-1}))$, then u the form $u = x_i y^b$ and $x_j y^b \in in(J_{i-1})$ for all $j \ge i$.

In order to prove this proposition we just have to see that equation (6) holds for all i. This will follow from

Lemma 1.8. Let $I \subset T$ be a monomial ideal. Then the following conditions are equivalent:

- (a) $I: x_1 \cap (\mathfrak{m}T + I) = I.$
- (b) If x_1 divides $u \in G(I)$, then u is of the form $u = x_1y^b$ and $x_jy^b \in I$ for all j = 1, ..., n.

Proof. (a) \Rightarrow (b): Let $u \in G(I)$ such that x_1 divides u. Say, $u = x_1^a u'$ with $a \ge 1$ and x_1 does not divide u'. Suppose a > 1; then $x_1^{a-1}u' \notin I$. But $x_1^{a-1}u' \in I : x_1 \cap (\mathfrak{m}T + I)$, a contradiction. Hence we see that $u = x_1u'$. Suppose some x_i divides u' for some i > 1. Then $u' \in I : x_1 \cap (\mathfrak{m}T + I)$, but $u' \notin I$, again a contradiction. Thus $u = x_1y^b$ for some exponent vector b. The monomial x_iy^b belongs to $I : x_1 \cap (\mathfrak{m}T + I)$ for all i, and hence $x_iy^b \in I$ for all i.

(b) \Rightarrow (a): We only need to prove the inclusion $I : x_1 \cap (\mathfrak{m}T + I) \subset I$. Our assumptions imply that G(I) can be written as $\{x_1y^{b_1}, \ldots, x_1y^{b_k}, u_{k+1}, \ldots, u_m\}$ with some integer vectors b_i , and with monomials $u_j \in G(I)$ which are not divisible by x_1 for $j = k + 1, \ldots, m$. It follows that

$$I: x_1 = (y^{b_1}, \dots, y^{b_k}, u_{k+1}, \dots, u_m).$$

Thus if $u \in I : x_1 \cap \mathfrak{m}T + I$, then u is either a multiple of one of the u_j , in which case $u \in I$, or $u = vy^{b_i}$ for some i, and x_j divides v for some j. Write $v = x_jv'$; then $u = v'x_jy^{b_i} \in I$, according to the assumptions made in (b).

We demonstrate the use of Proposition 1.7 with a simple example.

Example 1.9. Let I be the ideal of 2-minors of

$$\left(\begin{array}{rrr} x_1 & a & b \\ x_2 & b & c \end{array}\right).$$

We have $I = (-x_2a + x_1b, -x_2b + x_1c, -b^2 + ac)$ and $J = (-by_1 + ay_2 - xy_3, cy_1 - by_2 + x_2y_3)$.

We want to show that x_1, x_2, a, c, b is a *d*-sequence. We apply Proposition 1.7 by using the reverse lexicographic order induced by $x_1 > x_2 > a > b > c > y_1 > y_2 > y_3$. Then

$$in(J_0) = (cy_1, by_1, b^2y_2), \quad in(J_1) = (cy_1, by_1, b^2y_2), \quad in(J_2) = (cy_1, by_1, b^2y_2),$$

 $in(J_3) = (cy_1, by_1, b^2y_2) \quad and \quad in(J_4) = (by_1, by_2).$

Thus we see that the conditions of Proposition 1.7 are satisfied, and hence x_1, x_2, a, c, b is indeed a *d*-sequence on R(I).

Example 1.9 naturally leads us to pose the following question. Let I be the ideal of 2-minors of

$$\left(\begin{array}{cccc} x & a_1 & a_2 & \cdots & a_{n-1} & a_n \\ y & a_2 & a_3 & \cdots & a_n & a_{n+1} \end{array}\right)$$

This ideal defines a rational scroll. Is it true that the sequence $x, y, a_1, a_{n+1}, a_n, \ldots, a_2$ is a *d*-sequence on R(I). This is the case for $n \leq 6$.

2. Chordal graphs

Let $[n] = \{1, 2, ..., n\}$ denote the vertex set and G a finite graph on [n] with no loop and no multiple edge. We write E(G) for the set of edges of G. The edge ideal of Gis the ideal I(G) of $S = K[x_1, ..., x_n]$ which is generated by those monomials $x_i x_j$ with $\{i, j\} \in E(G)$. A subset C of [n] is called *vertex cover* of G if, for each edge $\{i, j\}$ of G, one has either $i \in C$ or $j \in C$. A minimal vertex cover of G is a vertex cover C with the property that any proper subset of C cannot be a vertex cover of G. We say that G is unmixed if all minimal vertex covers have the same cardinality. A mixed graph is a graph which is not unmixed. The vertex cover ideal of G is the ideal of S which is generated by those squarefree monomials $\prod_{i \in C} x_i$ such that C is a minimal vertex cover of G.

The vertex cover ideal of G is the Alexander dual of I(G); in other words, the ideal

$$I_G := I(G)^{\vee} = \bigcap_{\{i,j\} \in E(G)} (x_i, x_j)$$

Recall that a finite graph G is called *chordal* if each cycle of G of length > 3 possesses a chord. In the paper [9, Theorem 3.2] by Francisco and Van Tuyl it is shown that every vertex cover ideal of a chordal graph is componentwise linear.

Example 2.1. (a) Let G be the line of length 3. Thus $E(G) = \{\{1,2\},\{2,3\},\{3,4\}\}$. Then G is unmixed and $I_G = (x_1x_3, x_2x_3, x_2x_4)$. Since

 $x_1, x_3, x_2 + x_4, x_2$

is a d-sequence with respect to $R(I_G)$, all powers of I_G are (componentwise) linear.

(b) Let G be the line of length 4. Thus $E(G) = \{\{1,2\},\{2,3\},\{3,4\},\{4,5\}\}$. Then G is mixed and $I_G = (x_1x_3x_4, x_1x_3x_5, x_2x_4, x_2x_3x_5)$. Since

 $x_1, x_3, x_5, x_2 + x_4, x_2$

is a d-sequence with respect to $R(I_G)$, all powers of I_G are componentwise linear.

A complete graph on [n] is a finite graph such that $\{i, j\} \in E(G)$ for all $1 \leq i < j \leq n$. Let G_n denote the complete graph on [n]. Its vertex cover ideal I_{G_n} is generated by all squarefree monomials of S of degree n - 1; in other words, I_{G_n} is an ideal of Veronese type ([11]). In particular, I_{G_n} is a polymatroidal ideal ([5]). Hence all powers of I_{G_n} have linear resolutions. Theorem 1.4 then guarantees that a generic K-basis $\mathbf{z} = z_1, \ldots, z_n$ of S_1 is a d-sequence with respect to $R(I_{G_n})$.

Let $m \ge 1$ and Γ a connected graph on [n+m] such that the induced subgraph of G on [n] is G_n and that $\{i, j\} \notin E(G)$ if $n < i < j \le n+m$. Such a graph is called a *star* graph based on G_n . A star graph based on G_n is a chordal graph.

Example 2.2. Let n = 3 and m = 3. Let G be the star graph based on G_3 with the edges

 $\{1,4\},\{2,4\},\{2,5\},\{3,5\},\{1,6\},\{3,6\}$

together with all edges of G_3 . Since both $\{1, 2, 3\}$ and $\{2, 3, 4, 6\}$ are minimal vertex covers of G, it follows that G is mixed. Let K be a field of characteristic 0. Since the K-basis

$$x_4, x_5, x_6, x_1 + x_2 + x_3, 2x_1 + 3x_2 + 5x_3, 7x_1 + 11x_2 + 13x_3$$

of S_1 is a *d*-sequence with respect to $R(I_G)$, all powers of I_G are componentwise linear.

The simple computational observation done in Example 2.2 now grows the following

Theorem 2.3. All powers of the vertex cover ideal of a star graph based on G_n are componentwise linear.

Proof. Let G be a star graph on the vertex set $\{x_1, \ldots, x_n, t_1, \ldots, t_m\}$ based on G_n with vertex set $\{x_1, \ldots, x_n\}$.

Let $T = S[t_1, \ldots, t_m]$ denote the polynomial ring in n + m variables over K and $I_G \subset T$ the vertex cover ideal of G.

First, we observe that I_G is generated by the following monomials:

$$u_{i} = \prod_{\substack{j=1\\ j \neq i}}^{n} x_{j} \prod_{\substack{k \\ \{x_{i}, t_{k}\} \in E(G)}} t_{k}, \quad i = 1, \dots, n,$$

and

$$u_{n+1} = \prod_{j=1}^{n} x_j$$

It is clear that all these monomials belong to I_G . Conversely, suppose that C is an arbitrary vertex cover of G and u_C its corresponding monomial. If $\{x_1, \ldots, x_n\} \subset C$, then u_{n+1} divides u_C . Thus assume that $x_i \notin C$ for some i. Then $x_j \in C$ for all $j \neq i$, because if $x_j \notin C$ for some $j \neq i$, then the edge $\{x_i, x_j\}$ is not covered by C. It follows that $\{x_1, \ldots, x_{i-1}, x_{i+1}, \ldots, x_n\} \subset C$. Moreover since $x_i \notin C$ we must have $t_k \in C$ for all kwith $\{x_i, t_k\} \in E(G)$. This shows that u_i divides u_C .

Notice that u_{n+1} is not always part of the minimal monomial set $G(I_G)$ of generators of I_G . In fact, $u_{n+1} \in G(I_G)$ if and only if for each i = 1, ..., n there exists k such that $\{x_i, y_k\} \in E(G)$. For the following considerations, however, it is not important whether u_{n+1} belongs to $G(I_G)$ or not.

Second, we write the Rees algebra of I_G as the factor ring $R(I_G) = T[y_1, \ldots, y_{n+1}]/J$, where J is the kernel of the T-algebra homomorphism given by $y_j \mapsto u_j$ for $j = 1, \ldots, n+1$.

We claim that J is generated by the binomials

$$f_i = x_i y_i - (\prod_k t_k) y_{n+1}, \quad i = 1, \dots n$$

where for each *i* the product $\prod_k t_k$ in f_i is taken over all *k* with $\{x_i, t_k\} \in E(G)$.

To see why this is true, we write L for the ideal generated by the binomials f_1, \ldots, f_n , and have to show that L = J. We first observe that f_1, \ldots, f_n is a regular sequence. Indeed, let < be a lexicographic monomial order with $y_i > y_{n+1} > x_j, t_k$ for all i, j, k and $i \neq n+1$. Then $in_{\leq}(f_i) = x_i y_i$ for $i = 1, \ldots, n$. Since $gcd(in_{\leq}(f_i), in_{\leq}(f_j)) = 1$ for all $i \neq j$, it follows that f_1, \ldots, f_n is a Gröbner basis of L, and that f_1, \ldots, f_n is a regular sequence.

Now it follows that dim $T[y_1, \ldots, y_{n+1}]/L = \dim T + 1 = \dim R(I_G)$. Hence, if L is a prime ideal, then one has L = J. To show that L is a prime ideal, we consider the ideal $(L, x_1y_1) = (vy_{n+1}, x_1y_1, f_2, \ldots, f_n)$, where $v = \prod_k t_k$ and where the product is taken over all k with $\{x_1, t_k\} \in E(G)$. The initial terms of the generators of the ideal (L, x_1y_1) are $vy_{n+1}, x_1y_1, \ldots, x_ny_n$. Since these initial terms form a regular sequence of monomials, it follows as above that f_1, \ldots, f_n, x_1y_1 is a regular sequence. In particular the variable x_1 is a nonzero divisor on $T[y_1, \ldots, y_{n+1}]/L$. By the same reason all x_i are nonzero divisors on $T[y_1, \ldots, y_{n+1}]/L$. Let $x = \prod_{i=1}^n x_i$; then the natural homomorphism $T[y_1, \ldots, y_{n+1}]/L \to (T[y_1, \ldots, y_{n+1}]/L)_x$ is a monomorphism. In the localized polynomial ring $T[y_1, \ldots, y_{n+1}]_x$ the ideal L_x is generated by the elements $x_i^{-1}f_i = y_i - x_i^{-1}(\prod_k t_k)y_{n+1}$. Thus we see that $(T[y_1, \ldots, y_{n+1}]/L)_x \cong T[y_{n+1}]_x$, which is a domain. It follows that $T[y_1, \ldots, y_{n+1}]/L$ is a domain. In other words, one has L = J, as desired.

Now we have that $in_{\leq}(J) = (x_1y_1, \ldots, x_ny_n)$. Hence t_1, \ldots, t_m is a regular sequence on $T[y_1, \ldots, y_{n+1}]/in_{\leq}(J)$, and consequently is a regular sequence on the Rees algebra $R(I_G) = T[y_1, \ldots, y_{n+1}]/J$. Since any regular sequence is a *d*-sequence, it remains to be shown that we can find a basis z_1, \ldots, z_n of S_1 which is a *d*-sequence on $\overline{R(I_G)} = R(I_G)/(t_1, \ldots, t_m)$. Note that

$$\overline{R(I_G)} \cong S[y_1, \dots, y_n]/(x_1y_1, \dots, x_ny_n).$$

Thus the theorem follows form the following Lemma 2.4.

Lemma 2.4. Let $R = S[y_1, \ldots, y_n]/(x_1y_1, \ldots, x_ny_n)$, and z_1, \ldots, z_n a generic K-basis of S_1 . Then z_1, \ldots, z_n is a d-sequence on R.

Proof. We will apply Theorem 1.3 and have to show that the approximation complex $\mathcal{M}(\mathfrak{m}; R)$ is acyclic, where $\mathfrak{m} = (x_1, \ldots, x_n)$. Recall that

$$\mathcal{M}(\mathfrak{m};R)_i = H_i(\mathfrak{m};R) \otimes A(-i),$$

where $A = R[e_1, \ldots, e_n]$ is a polynomial ring over R in the variables e_1, \ldots, e_n , and where $H_i(\mathfrak{m}; R)$ is the *i*th Koszul homology of the sequence x_1, \ldots, x_n with values in R. Since R is a complete intersection, it follows from [2, Theorem 2.3.9] that $H(\mathfrak{m}; R)$ is the *i*th exterior algebra of $H_1(\mathfrak{m}; R)$. Since $H_1(\mathfrak{m}; R)$ is a free $K[y_1, \ldots, y_n]$ -modules whose basis is given by the homology classes $[y_i e_i]$ of the cycles $y_i e_i$, we see that $H_i(\mathfrak{m}; R)$ is free $(R/\mathfrak{m})[y_1, \ldots, y_n]$ -module with basis $\{b_J \colon J = \{j_1, j_2, \ldots, j_i\} \ 1 \le j_1 < j_2 < \ldots < j_i \le n\}$, where

$$b_J = [y_{j_1}y_{j_2}\cdots y_{j_i}e_{j_1} \wedge e_{j_2} \wedge \cdots \wedge e_{j_i}].$$

Thus $\mathcal{M}(\mathfrak{m}; R)_i$ can be identified with the free *B*-module $\bigoplus_{J \mid J \mid = i} b_J B$, where

$$B = K[y_1, \ldots, y_n, e_1, \ldots, e_n].$$

After this identification the differential of $\mathcal{M}(\mathfrak{m}; R)$ is given by


$$\partial(b_J) = \sum_{k=1}^{i} (-1)^{k+1} y_{j_k} e_{j_k} b_{J \setminus \{j_k\}} \quad \text{for} \quad J = \{j_1 < j_2 < \dots < j_i\}$$

Hence $\mathcal{M}(\mathfrak{m}; R)$ is isomorphic to the Koszul complex $K(y_1e_1, \ldots, y_ne_n; B)$. Since the sequence y_1e_1, \ldots, y_ne_n is a regular sequence, it follows then that $\mathcal{M}(\mathfrak{m}; R)$ is acyclic, as desired.

Theorem 2.3 and computational evidence, including Example 2.6 leads us to present the following

Conjecture 2.5. All powers of the vertex cover ideal of a chordal graph are componentwise linear.

Example 2.6. Let G be the chordal graph drawn below. The graph G is no longer a star graph.



Then G is mixed and I_G is generated by

 $x_1x_3x_5x_6,\ x_2x_3x_4x_5x_6,\ x_1x_2x_4x_6x_7,\ x_1x_2x_3x_4x_6,$

 $x_2x_3x_4x_5x_7$, $x_1x_2x_3x_5x_7$, $x_1x_2x_5x_6x_7$, $x_2x_4x_5x_6x_7$.

We can easily find a *d*-sequence which is a "generic" *K*-basis of S_1 created by CoCoA with "Randomized." Hence all powers of the vertex cover ideal of *G* are componentwise linear.

In [13] the Cohen-Macaulay chordal graphs are classified. Let G be a finite graph on [n], and $\Delta(G)$ its clique complex, that is to say, the simplicial complex whose faces are the cliques (i.e. complete subgraphs) of G. It is shown in [13] that if G is chordal, then G is Cohen-Macaulay if and only if [n] is the disjoint union of those facets of $\Delta(G)$ which have a free vertex. A vertex of a facet is *free*, if it belongs to no other facet.

In support of Conjecture 2.5 we have the following result.

Theorem 2.7. Let G be a graph on [n], and suppose that [n] is the disjoint union of those facets of the clique complex of G with a free vertex. Then all powers of I_G have a linear resolution.

Proof. Let F_1, \ldots, F_m be the facets of $\Delta(G)$ which have a free vertex. Since $[n] = F_1 \cup F_2 \cup \cdots \cup F_s$ is a disjoint union, we may assume that if $i \in F_p$, $j \in F_q$ and p < q, then i < j. In particular, $1 \in F_1$ and $n \in F_s$. Moreover, we may assume that if $i_1, i_2 \in F_i$ where i_1 is a nonfree vertex and i_2 is a free vertex, then $i_1 < i_2$.

Observe that any minimal vertex cover of G is of the following form:

$$(F_1 \setminus \{a_1\}) \cup (F_2 \setminus \{a_2\}) \cup \cdots \cup (F_s \setminus \{a_s\}), \text{ where } a_j \in F_j.$$

In particular, G is unmixed and all generators of I_G have degree n-s.

Now let $R(I_G)$ be the Rees algebra of vertex cover ideal of G. Suppose u_1, \ldots, u_m is the minimal set of monomial generators of I_G . Then there is a surjective K-algebra homomorphism

$$K[x_1, \ldots, x_n, y_1, \ldots, y_m] \longrightarrow R(I_G) \quad x_i \mapsto x_i \quad \text{and} \quad y_j \mapsto u_j,$$

whose kernel J is a binomial ideal. Let < be the lexicographic order induced by the ordering $x_1 > x_2 > \cdots > x_n > y_1 > \cdots > y_m$. We are going to show that the generators of $i_{n>(J)}$ are at most of degree 1 in the x_i . This is the so-called x-condition and it implies that all powers of I_G have a linear resolution, see [12]. Suppose that $x_{i_1}x_{i_2}\cdots x_{i_p}y_{j_1}y_{j_2}\cdots y_{j_q}$ with $i_1 \leq i_2 \leq \ldots \leq i_p$ is a minimal generator of $i_{n<(J)}$. Then

(7)
$$x_{i_1}x_{i_2}\cdots x_{i_p}y_{j_1}y_{j_2}\cdots y_{j_q} - x_{k_1}x_{k_2}\cdots x_{k_p}y_{\ell_1}y_{\ell_2}\cdots y_{\ell_q} \in J.$$

It follows that $i_1 < \min\{k_1, \ldots, k_p\}$, and there exists an index j_r such that x_{i_1} does not divide u_{j_r} . Say, $i_1 \in F_c$. Then (7) implies that there exists $d \in [p]$ with $k_d \in F_c$. In particular, $i_1 \neq \max\{i: i \in F_d\}$. Let $i_0 = \max\{i: i \in F_d\}$. Since i_0 is a free vertex, it follows that $x_{i_1}(u_{j_r}/x_{i_0})$ is a minimal generator of I_G , say, u_g . Therefore, $f = x_{i_1}y_{j_r} - x_{i_0}y_g \in J$ and $\inf(f) = x_{i_1}y_{j_r}$ divides $x_{i_1}x_{i_2}\cdots x_{i_p}y_{j_1}y_{j_2}\cdots y_{j_q}$, as desired.

References

- Akin, D. Buchsbaum and J. Weyman, Resolution of determinantal ideals: the submaximal minors, Adv. Math. 39 (1980), 1–30.
- [2] W. Bruns and J. Herzog, Cohen-Macaulay rings (Revised), Cambridge University Press (1998).
- [3] M. Brodmann, Asymptotic depth and connectedness of schemes, Proc. Amer. Math. Soc. 108 (1990), 573–581.
- [4] A. Conca, Regularity jumps for powers of ideals, Commutative algebra, Lect. Notes Pure Appl. Math. 244, Chapman & Hall/CRC, Boca Raton, FL, (2006), 21–32.
- [5] A. Conca and J. Herzog, Castelnuovo–Mumford regularity of products of ideals, *Collect. Math.* 54 (2003), 137–152.
- [6] A. Conca, J. Herzog and T. Hibi, Rigid resolutions and big Betti numbers, Comm. Math. Helv. 79 (2004), 826–838.
- [7] A. Conca, J. Herzog and G. Valla, Sagbi bases with applications to blow-up algebras, J. Reine Angew. Math. 474 (1996), 113–138.
- [8] D. Eisenbud, Commutative Algebra; with a view towards algebraic geometry, Springer (1995).
- C. A. Francisco and A. Van Tuyl, Sequentially Cohen-Macaulay edge ideals, Proc. Amer. Math. Soc. 135 (2007), 2327–2337.
- [10] J. Herzog and T. Hibi, Componentwise linear ideals, Nagoya Math. J. 153 (1999), 141–153.
- [11] J. Herzog and T. Hibi, Cohen-Macaulay polymatroidal ideals, European J. Combin. 27 (2006), 513– 517.
- [12] J. Herzog, T. Hibi and X. Zheng, Monomial ideals whose powers have a linear resolution, Math. Scand. 95 (2004), 23 – 32.
- [13] J. Herzog, T. Hibi and X. Zheng, Cohen-Macaulay Chordal Graphs, J. Combin. Theory Ser. A 113 (2006), 911–916.
- [14] J. Herzog, A. Simis and W. Vasconcelos, Approximation complexes and blowing-up rings, J. of Alg. 74 (1982), 466–493.
- [15] J. Herzog, A. Simis and W. Vasconcelos, Approximation complexes and blowing-up rings II, J. of Alg. 82 (1983), 53–83.
- [16] S. Iyengar and T. Römer, Linearity defects of modules over commutative rings, to appear in J. Alg.
- [17] C. Huneke, The theory of d-sequences and powers of ideals, Adv. Math. 46 (1982), 249–279.
- [18] T. Römer, On minimal graded free resolutions, Disseration, Universität Duisburg-Essen, 2001.
- [19] T. Römer, Homological properties of bigraded algebras, Illinois J. Math. 45 (2001), 1361–1376.
- [20] K. Yanagawa, Alexander Duality for Stanley–Reisner Rings and Squarefree Nⁿ-Graded Modules, J. of Alg. 225 (2000), 630–645.

JÜRGEN HERZOG, FACHBEREICH MATHEMATIK, UNIVERSITÄT DUISBURG-ESSEN, CAMPUS ESSEN, 45117 ESSEN, GERMANY

E-mail address: juergen.herzog@uni-essen.de

TAKAYUKI HIBI, DEPARTMENT OF MATHEMATICS, GRADUATE SCHOOL OF SCIENCE, OSAKA UNIVER-SITY, TOYONAKA, OSAKA 560-0043, JAPAN

E-mail address: hibi@math.sci.osaka-u.ac.jp

HIDEFUMI OHSUGI, DEPARTMENT OF MATHEMATICS, COLLEGE OF SCIENCE, RIKKYO UNIVERSITY, TOKYO 171-8501, JAPAN

E-mail address: ohsugi@rkmath.rikkyo.ac.jp