

# NON-NEGATIVE LEGENDRIAN ISOTOPY IN $ST^*M$

VLADIMIR CHERNOV AND STEFAN NEMIROVSKI

ABSTRACT. It is shown that if the universal cover of a manifold  $M$  is an open manifold, then two different fibres of the spherical cotangent bundle  $ST^*M$  cannot be connected by a non-negative Legendrian isotopy. This result is applied to the study of causality in globally hyperbolic spacetimes. It is also used to strengthen a result of Eliashberg, Kim, and Polterovich on the existence of a partial order on  $\widetilde{\text{Cont}}_0(ST^*M)$ .

**1. Introduction.** Let  $M$  be a connected not necessarily orientable manifold of dimension  $m \geq 2$  and let  $\pi_M : ST^*M \rightarrow M$  be its spherical cotangent bundle. It is well-known that  $ST^*M$  carries a canonical co-oriented contact structure. An isotopy  $\{L_t\}_{t \in [0,1]}$  of Legendrian submanifolds in a co-oriented contact manifold is called *non-negative* if it can be parameterised in such a way that the tangent vectors of the trajectories of individual points lie in the non-negative tangent half-spaces defined by the contact structure, see Definition 2.1. For a generic Legendrian isotopy in  $ST^*M$ , this property can be expressed by saying that the co-oriented wave fronts  $\pi_M(L_t) \subset M$  move in the direction of their co-orientation.

**Theorem 1.1.** *Assume that the universal cover of  $M$  is an open manifold. Then there does not exist a non-negative Legendrian isotopy connecting two different (nonoriented) fibres of  $ST^*M$ .*

In the special case when  $M$  can be covered by an open subset of  $\mathbb{R}^m$ , this statement was proved by Colin, Ferrand, and Pushkar [8]. Independently, a slightly stronger result was obtained by the present authors in the course of the proof of the so-called Legendrian Low Conjecture from Lorentz geometry [6]. Theorem 1.1 allows us to extend the results of [6] to a wider class of Lorentz manifolds, see §10.

A closely related notion of non-negative contact isotopy plays a key role in the orderability problem for contactomorphism groups, see [11], [10], and [4]. Theorem 1.1 can be applied to settle a question left open in [10], see §9.

It is easy to see that the assertion of Theorem 1.1 is false if  $M$  carries a Riemann metric turning it into a  $Y_\ell^x$ -manifold, see Example 8.3. In particular, it is false if  $M$  is a metric quotient of the standard sphere. Hence, the hypothesis of Theorem 1.1 cannot be weakened for surfaces (this is obvious) and 3-manifolds (this follows from Perelman's work on the Poincaré conjecture [17, 18, 19]). On the other hand, it seems very likely that the result can be strengthened in all higher dimensions.

The proof of Theorem 1.1 is based on Viterbo's invariants of generating functions [20]. However, it is different from the arguments in [8] and [6] already in the case when  $M = \mathbb{R}^m$ . In particular, no use is made of the identification  $ST^*\mathbb{R}^m \cong \mathcal{J}^1(S^{m-1})$ .

All manifolds, maps etc. are assumed to be smooth unless the opposite is explicitly stated, and the word *smooth* means  $C^\infty$ .

---

The second author was supported by grants from DFG, RFBR, Russian Science Support Foundation, and the programme "Leading Scientific Schools of Russia."

**2. Non-negative Legendrian isotopies.** Let  $(Y, \ker \alpha)$  be a contact manifold with a co-oriented contact structure defined by a contact form  $\alpha$ .

**Definition 2.1.** A Legendrian isotopy  $\{L_t\}_{t \in [0,1]}$  in  $(Y, \ker \alpha)$  is called *non-negative* if it has a parameterisation  $F : L_0 \times [0, 1] \rightarrow Y$  such that  $(F^*\alpha) \left( \frac{\partial}{\partial t} \right) \geq 0$ . If the latter inequality is strict, the isotopy is called *positive*.

Clearly, this definition does not depend on the choice of the parameterisation  $F$  of the Legendrian isotopy and on the choice of the contact form defining the co-oriented contact structure. It is also obvious that (co-orientation preserving) contactomorphisms preserve the property of being non-negative or positive.

**Lemma 2.2.** *Let  $\{L_t\}_{t \in [0,1]}$  be a non-negative Legendrian isotopy of compact submanifolds such that  $L_0 \cap L_1 = \emptyset$ . Then there exists a ( $C^\infty$ -close) positive Legendrian isotopy with the same ends.*

*Sketch of proof.* Let  $L'_t = \psi_{\varepsilon t}(L_t)$ , where  $\psi_t$  is the Reeb flow on  $(Y, \alpha)$  and  $\varepsilon > 0$ . The isotopy  $\{L'_t\}$  is positive. If  $\varepsilon$  is small enough, then  $L'_1$  is Legendrian isotopic to  $L_1$  in  $Y \setminus L_0$  and therefore there exists a contactomorphism  $\varphi$  supported in  $Y \setminus L_0$  such that  $\varphi(L'_1) = L_1$ . Thus,  $L''_t := \varphi(L'_t)$  is a positive Legendrian isotopy connecting  $L_0$  and  $L_1$ .  $\square$

An advantage of positive isotopies is that positivity is an open condition and hence one can make a positive Legendrian isotopy generic by a small perturbation.

**Definition 2.3.** An isotopy  $\{L_t\}_{t \in [0,1]}$  is in general position with respect to a submanifold  $\Lambda \subset Y$  of codimension  $\dim_{\mathbb{R}} L_0$  if it has a parameterisation  $F : L_0 \times [0, 1] \rightarrow Y$  such that

- a)  $F^{-1}(\Lambda)$  is a 1-dimensional submanifold in  $L_0 \times [0, 1]$ ;
- b) the projection  $F^{-1}(\Lambda) \rightarrow [0, 1]$  has isolated critical points;
- c)  $F^{-1}(\Lambda)$  is transverse to  $L_0 \times \{0\}$  and  $L_0 \times \{1\}$ .

Note that a point  $(x, \tau) \in F^{-1}(\Lambda)$  which is not critical for the projection  $F^{-1}(\Lambda) \rightarrow [0, 1]$  lies on the graph of a section of this projection over a non-trivial closed interval  $[t', t''] \ni \tau$ . In other words, there exists a curve  $\gamma : [t', t''] \rightarrow L_0$  such that  $\gamma(\tau) = x$  and  $F(\gamma(t), t) \in \Lambda$  for all  $t \in [t', t'']$ .

**3. Exact pre-Lagrangian submanifolds.** Suppose that  $\Lambda$  is an  $m$ -dimensional submanifold of a  $(2m - 1)$ -dimensional contact manifold  $(Y, \ker \alpha)$  such that

$$df = e^h \alpha|_{\Lambda} \tag{3.1}$$

for some functions  $f, h : \Lambda \rightarrow \mathbb{R}$ . Then  $\Lambda$  is said to be *exact pre-Lagrangian* and  $f$  is called a *contact potential* on  $\Lambda$ . (The terminology will be explained in §4.)

The following lemma shows that the contact potential is non-decreasing along any curve traced on  $\Lambda$  by a non-negative Legendrian isotopy.

**Lemma 3.1.** *Let  $L_t = \varphi_t(L_0)$ ,  $t \in [0, 1]$ , be a non-negative Legendrian isotopy. Suppose that  $\gamma : [0, 1] \rightarrow L_0$  is a curve such that  $\varphi_t(\gamma(t)) \in \Lambda$ , where  $\Lambda$  is an exact pre-Lagrangian submanifold with contact potential  $f$ . Then the function  $t \mapsto f(\varphi_t(\gamma(t)))$  is non-decreasing.*

*Proof.* This follows from the definitions and the chain rule. Indeed,

$$\begin{aligned} \frac{d}{dt} f(\varphi_t(\gamma(t))) &= df \left( \frac{d\varphi_t}{dt}(\gamma(t)) + d\varphi_t(\gamma(t)) \frac{d\gamma}{dt}(t) \right) \\ &= e^h \left[ \alpha \left( \frac{d\varphi_t}{dt}(\gamma(t)) + \alpha(d\varphi_t(\gamma(t)) \frac{d\gamma}{dt}(t)) \right) \right]. \end{aligned}$$

The first summand in square brackets is non-negative by the definition of non-negative Legendrian isotopy and the second one is zero because the isotopy is Legendrian and  $\frac{d\gamma}{dt}$  is tangent to  $L_0$ . Hence, the derivative of our function is non-negative.  $\square$

**4. Symplectisation.** Let  $(Y, \ker \alpha)$  be a contact manifold. Its symplectisation  $Y^{\text{symp}}$  is the (exact) symplectic manifold  $(Y \times \mathbb{R}, d(e^s \alpha))$ .

**Example 4.1.** Let  $Y \subset T^*M$  be the unit sphere bundle with respect to a Riemann metric on  $M$ . Then  $\alpha := \lambda_{\text{can}}|_Y$  is a contact form defining the canonical contact structure on  $Y \cong ST^*M$ . The map

$$Y^{\text{symp}} \ni (\xi, s) \longmapsto e^s \xi \in T^*M$$

is a symplectomorphism onto the complement of the zero section of  $T^*M$  such that the pull-back of the canonical 1-form  $\lambda_{\text{can}}$  is precisely  $e^s \alpha$ .  $\square$

A contactomorphism  $\varphi : Y \rightarrow Y$  lifts to a symplectomorphism  $\tilde{\varphi} : Y^{\text{symp}} \rightarrow Y^{\text{symp}}$  defined by the formula

$$\tilde{\varphi}(x, s) := (\varphi(x), s - \eta(x)),$$

where  $\eta : Y \rightarrow \mathbb{R}$  is the function such that  $\varphi^* \alpha = e^\eta \alpha$ . It follows from this definition that if  $\{\varphi_t\}_{t \in [0,1]}$  is a contact isotopy of  $Y$ , then  $\{\tilde{\varphi}_t\}_{t \in [0,1]}$  is a *Hamiltonian* isotopy of  $Y^{\text{symp}}$ . (It can be defined by the Hamiltonian function  $\tilde{H}(x, s, t) = -\alpha(\frac{d\varphi_t}{dt})$ .)

An exact pre-Lagrangian submanifold  $\Lambda$  with a contact potential  $f$  such that  $df = e^h \alpha|_\Lambda$  lifts to an exact Lagrangian submanifold

$$\tilde{\Lambda} = \{(x, h(x)) \in Y^{\text{symp}} \mid x \in \Lambda\} \subset Y^{\text{symp}}.$$

Indeed, the function  $\tilde{f} : \tilde{\Lambda} \rightarrow \mathbb{R}$ ,  $\tilde{f}(x, h(x)) = f(x)$ , is a primitive for the 1-form  $e^s \alpha|_{\tilde{\Lambda}}$ . Note that for any contactomorphism  $\varphi : Y \rightarrow Y$ , the image  $\varphi(\Lambda)$  with the contact potential  $f \circ \varphi^{-1}$  is exact pre-Lagrangian and  $\widetilde{\varphi(\Lambda)} = \tilde{\varphi}(\tilde{\Lambda})$ .

**5. Generating functions.** Let  $M$  be a manifold, which may be open or have boundary. Consider the product  $M \times \mathbb{R}^N$  for some  $N \geq 0$  and let  $\pi : M \times \mathbb{R}^N \rightarrow M$  be the projection onto  $M$ . For a function  $\mathcal{S} : M \times \mathbb{R}^N \rightarrow \mathbb{R}$ , consider the set of its fibre critical points

$$\text{FCrit}(\mathcal{S}) := \{z \in M \times \mathbb{R}^N \mid d\mathcal{S}(z)|_{\mathbb{R}^N} = 0\}.$$

Note that there is a natural fibrewise map

$$d_M \mathcal{S} : \text{FCrit}(\mathcal{S}) \rightarrow T^*M$$

which associates to a point  $z \in \text{FCrit}(\mathcal{S})$  the linear form  $v \mapsto d\mathcal{S}(z)(\hat{v})$  on  $T_{\pi(z)}M$ , where  $\hat{v} \in T_z(M \times \mathbb{R}^N)$  is any tangent vector such that  $d\pi(\hat{v}) = v \in T_{\pi(z)}M$ .

A function  $\mathcal{S} : M \times \mathbb{R}^N \rightarrow \mathbb{R}$  is called a *generating function* for a Lagrangian submanifold  $L \subset T^*M$  if it satisfies the following two conditions:

- (GF1) its set of fibre critical points is cut out transversely;
- (GF2) the map  $d_M \mathcal{S} : \text{FCrit}(\mathcal{S}) \rightarrow T^*M$  is a diffeomorphism onto  $L$ .

Note that  $\mathcal{S} \circ (d_M \mathcal{S})^{-1} : L \rightarrow \mathbb{R}$  is a primitive for  $\lambda_{\text{can}}|_L$ . Hence, a Lagrangian submanifold of  $T^*M$  admitting a generating function is exact.

A generating function  $\mathcal{S} : M \times \mathbb{R}^N \rightarrow \mathbb{R}$  is called *quadratic at infinity* if furthermore

- (GF3)  $\mathcal{S}(y, \xi) = \sigma(y, \xi) + Q(\xi)$ , where  $Q$  is a non-degenerate quadratic form on  $\mathbb{R}^N$  and the projection  $\pi : \text{supp } \sigma \rightarrow M$  is a proper map.

Note that a Lagrangian submanifold  $L \subset T^*M$  admitting a quadratic at infinity generating function is *properly embedded*, i. e., the projection  $L \rightarrow M$  is a proper map.

**Proposition 5.1.** *Let  $\{L_t\}_{t \in [0,1]}$  be a compactly supported isotopy of properly embedded Lagrangian submanifolds in  $T^*M$ . Suppose that*

- a)  $L_0$  admits a quadratic at infinity generating function;
- b) there exists a family of functions  $f_t : L_t \rightarrow \mathbb{R}$  such that  $df_t = \lambda_{\text{can}}|_{L_t}$ .

*Then there exists a family  $\mathcal{S}_t : M \times \mathbb{R}^N \rightarrow \mathbb{R}$  of quadratic at infinity generating functions for  $L_t$  such that  $\mathcal{S}_t \circ (d_M \mathcal{S}_t)^{-1} = f_t$  for all  $t \in [0, 1]$ .*

*Proof.* This is a minor extension of the Laudenbach–Sikorav theorem [13]. It can be obtained, for instance, by applying the version of Chekanov’s theorem [5] for properly embedded Legendrian submanifolds [9, Sec. 4] to the Legendrian isotopy

$$\widehat{L}_t := \{(x, f_t(x)) \in \mathcal{J}^1(M) \mid x \in L_t\}$$

in the 1-jet bundle of  $M$ . □

**6. Critical values of quadratic at infinity functions.** Let  $S : \mathbb{R}^N \rightarrow \mathbb{R}$  be a function quadratic at infinity in the sense that

$$S(z) = \sigma(z) + Q(z),$$

where  $\sigma$  has compact support and  $Q$  is a non-degenerate quadratic form on  $\mathbb{R}^N$ . (We will eventually take  $S$  to be the restriction of a quadratic at infinity generating function  $\mathcal{S} : M \times \mathbb{R}^N \rightarrow \mathbb{R}$  to the fibre  $\{x\} \times \mathbb{R}^N$  over a point  $x \in M$ .) Following Viterbo [20, §2], let us define an invariant  $c_-(S) \in \mathbb{R}$  of such a function.

Consider the sublevel sets

$$S^c := \{z \in \mathbb{R}^N \mid S(z) \leq c\}$$

and denote by  $S^{-\infty}$  the set  $S^c$  for a sufficiently negative  $c \ll 0$ . Pick a  $Q$ -negative linear subspace  $V \subset \mathbb{R}^N$  of maximal possible dimension  $\varkappa$ . The relative homology class  $[V] \in H_{\varkappa}(\mathbb{R}^N, S^{-\infty})$  does not depend on the choice of  $V$ . Set

$$c_-(S) := \inf\{c \in \mathbb{R} \mid [V] \in \iota_* H_{\varkappa}(S^c, S^{-\infty})\},$$

where  $\iota_* : H_{\varkappa}(S^c, S^{-\infty}) \rightarrow H_{\varkappa}(\mathbb{R}^N, S^{-\infty})$  is the homomorphism of relative homology groups induced by the inclusion  $\iota : S^c \rightarrow \mathbb{R}^N$ .

By Morse theory,  $c_-(S)$  is a critical value of  $S$ . In particular, if  $S$  has a single critical point  $z_0 \in \mathbb{R}^N$ , then  $c_-(S) = S(z_0)$ .

We shall need the following version of Viterbo’s monotonicity lemma [20, Lemma 4.7] adapted to our situation, cf. Lemma 3.1.

**Lemma 6.1.** *Let  $\{S_t\}_{t \in [0,1]}$  be a family of quadratic at infinity functions on  $\mathbb{R}^N$  and let*

$$C := \{(z, \tau) \in \mathbb{R}^N \times [0, 1] \mid dS_{\tau}(z) = 0\} = \bigcup_{\tau \in [0,1]} \text{Crit}(S_{\tau}) \times \{\tau\}.$$

*Suppose that for any  $(z, \tau)$  from a dense subset  $C' \subseteq C$  there exists a non-trivial closed interval  $[t', t''] \ni \tau$  and a curve  $\gamma : [t', t''] \rightarrow \mathbb{R}^N$  such that*

- a)  $\gamma(\tau) = z$ ;
- b)  $\gamma(t) \in \text{Crit}(S_t)$  for all  $t \in [t', t'']$ ;
- c) the function  $t \mapsto S_t(\gamma(t))$  is non-decreasing on  $[t', t'']$ .

*Then  $t \mapsto c_-(S_t)$  is a non-decreasing (continuous) function on  $[0, 1]$ .*

*Proof.* According to [20, Lemma 4.7], the claim will follow if we show that  $\frac{\partial S_\tau}{\partial t}(z) \geq 0$  for all  $(z, \tau) \in C$ . If  $(z, \tau) \in C'$ , we have

$$0 \leq \frac{d}{dt} S_t(\gamma(t))|_{t=\tau} = \frac{\partial S_\tau}{\partial t}(z) + dS_\tau(z) \frac{d\gamma}{dt}(\tau) = \frac{\partial S_\tau}{\partial t}(z),$$

where  $\gamma$  is a curve satisfying conditions (a)–(c) for  $\tau$  and  $z$ . Since  $C'$  is dense in  $C$ , this inequality is valid for all  $(z, \tau) \in C$ .  $\square$

**7. The Importance of Being Open.** Let  $M$  be an open manifold. We identify  $ST^*M$  with the unit sphere bundle in  $T^*M$  for some Riemann metric on  $M$  and view the complement to the zero section of  $T^*M$  as the symplectisation of  $ST^*M$ , see Example 4.1. Let  $\pi_M$  denote the bundle projection  $T^*M \rightarrow M$ .

**Lemma 7.1.** *There exists a function  $\Phi : M \rightarrow \mathbb{R}$  without critical points.*

*Proof.* This is well-known, see [12, Lemma 1.15].  $\square$

**Definition 7.2.** Let  $\Lambda_\Phi := \left\{ \frac{d\Phi(x)}{\|d\Phi(x)\|} \mid x \in M \right\} \subset ST^*M$ .

It is clear from the definition of the canonical 1-form that  $\Lambda_\Phi$  is an exact pre-Lagrangian submanifold of  $ST^*M$  and the function

$$f_\Phi(\zeta) = \Phi(\pi_M(\zeta))$$

is a contact potential on  $\Lambda_\Phi$ . The associated Lagrangian lift

$$\tilde{\Lambda}_\Phi = \{d\Phi(x) \mid x \in M\} \subset T^*M$$

is just the graph of the differential of  $\Phi$ . It has an obvious generating function

$$\mathcal{S}_\Phi : M \times \mathbb{R}^0 \rightarrow \mathbb{R}, \quad \mathcal{S}_\Phi(x \times \{\text{pt}\}) := \Phi(x).$$

**8. Proof of Theorem 1.1.** Let  $M$  be a manifold (universally) covered by an open manifold. Suppose that there exists a non-negative Legendrian isotopy  $\{L_t\}_{t \in [0,1]}$  connecting two different fibres of  $ST^*M$ . Since such an isotopy lifts to the spherical cotangent bundle of the covering manifold, we may assume that  $M$  is itself an open manifold. By Lemma 2.2, we may also assume that the Legendrian isotopy is positive.

Let  $\Lambda_\Phi$  be the exact pre-Lagrangian submanifold with contact potential  $f_\Phi = \Phi \circ \pi_M$  defined in §7. Applying a global contactomorphism induced by a suitable diffeomorphism of  $M$ , we can arrange that  $L_0 = ST_{x_0}^*M$  and  $L_1 = ST_{x_1}^*M$ , where the points  $x_0, x_1 \in M$  are such that  $\Phi(x_0) > \Phi(x_1)$ . Furthermore, we can put the isotopy in general position with respect to  $\Lambda_\Phi$  in the sense of Definition 2.3, leaving  $L_0$  and  $L_1$  fixed (because they are already transversal to  $\Lambda_\Phi$ ).

Let  $\{\varphi_t\}_{t \in [0,1]}$  be a compactly supported contact isotopy of  $ST^*M$  such that  $L_t = \varphi_t(L_0)$  for all  $t \in [0,1]$ . (Such an isotopy exists by the Legendrian isotopy extension theorem.) Consider the Hamiltonian isotopy of exact Lagrangian submanifolds  $(\tilde{\varphi}_t)^{-1}(\tilde{\Lambda}_\Phi) \subset T^*M$  and the functions  $\tilde{f}_\Phi \circ \tilde{\varphi}_t$  on these manifolds, see §4. By Proposition 5.1, there exists a family of quadratic at infinity generating functions

$$\mathcal{S}_t : M \times \mathbb{R}^N \rightarrow \mathbb{R}$$

for  $(\tilde{\varphi}_t)^{-1}(\tilde{\Lambda}_\Phi) \subset T^*M$  such that

$$\mathcal{S}_t \circ (d_M \mathcal{S}_t)^{-1} = \tilde{f}_\Phi \circ \tilde{\varphi}_t.$$

Let

$$S_t := \mathcal{S}_t(x_0, \cdot) : \mathbb{R}^N \rightarrow \mathbb{R}$$

be the restrictions of  $\mathcal{S}_t$  to the fibre  $\{x_0\} \times \mathbb{R}^N$ . By construction, the map

$$\text{Crit}(S_t) \ni z \longmapsto \varphi_t \left( \frac{d_M \mathcal{S}_t(z)}{\|d_M \mathcal{S}_t(z)\|} \right) \in L_t$$

establishes a bijective correspondence between the set of critical points of  $S_t$  and the intersection  $L_t \cap \Lambda_\Phi$ . Furthermore, the value of  $S_t$  at a point  $z \in \text{Crit } S_t$  is equal to the value of  $f_\Phi$  at the corresponding point in  $\Lambda_\Phi$ .

In particular,  $S_0$  and  $S_1$  each have a single critical point corresponding to the intersection of  $\Lambda_\Phi$  with  $L_0 = ST_{x_0}^* M$  and  $L_1 = ST_{x_1}^* M$ , respectively. Since  $f_\Phi = \Phi \circ \pi_M$ , we see that

$$c_-(S_0) = \Phi(x_0) \quad \text{and} \quad c_-(S_1) = \Phi(x_1).$$

Hence,

$$c_-(S_0) > c_-(S_1) \tag{8.1}$$

by our choice of the points  $x_0$  and  $x_1$ .

On the other hand, it follows from Lemma 3.1 and the discussion after Definition 2.3 that the family of functions  $\{S_t\}_{t \in [0,1]}$  satisfies the hypotheses of Lemma 6.1. Thus,  $c_-(S_t)$  is a non-decreasing function of  $t$  and therefore  $c_-(S_0) \leq c_-(S_1)$ , which contradicts (8.1).

This contradiction shows that a non-negative Legendrian isotopy cannot connect two different fibres of  $ST^*M$ .  $\square$

**Corollary 8.1.** *If the universal cover of  $M$  is an open manifold, then there does not exist a positive Legendrian loop in the Legendrian isotopy class of the fibre of  $ST^*M$ .*

*Proof.* Suppose that  $\{L_t\}_{t \in [0,1]}$  is a positive Legendrian isotopy such that  $L_0 = L_1 = ST_x^* M$ . If  $\{\varphi_t\}_{t \in [0,1]}$  is a contact isotopy such that  $\varphi_0 = \text{id}$ ,  $\varphi_1(ST_x^* M) = ST_y^* M$  for some  $y \neq x$ , and  $\|\frac{d\varphi_t}{dt}\|$  is sufficiently small, then the isotopy  $\{\varphi_t(L_t)\}_{t \in [0,1]}$  is positive and connects two different fibres of  $ST^*M$ , which contradicts Theorem 1.1.  $\square$

**Remark 8.2.** With a little more work, it can be shown that any *non-negative* Legendrian loop in the Legendrian isotopy class of the fibre must be constant, cf. [6, Corollaries 5.5 and 6.2].

**Example 8.3.** Suppose that there exists a Riemann metric  $g$  on  $M$  such that  $(M, g)$  is a  $Y_\ell^x$ -manifold for some  $x \in M$  and  $\ell > 0$ , i. e., such that all  $g$ -geodesics starting from  $x$  return to  $x$  in time  $\ell$ , see [3, Definition 7.7(c)]. Then moving the fibre  $ST_x^* M$  along the (co-)geodesic flow on  $ST^*M$  defines a positive Legendrian loop based at  $ST_x^* M$ . Thus, Corollary 8.1 and Theorem 1.1 do not hold for such a manifold  $M$ .

Note that if  $\dim M = 2$  or  $3$ , then either the universal cover of  $M$  is open or  $M$  admits a Riemann metric turning it into a  $Y_\ell^x$ -manifold. For  $\dim M = 2$ , this statement follows immediately from the classification of surfaces. For  $\dim M = 3$ , the Poincaré conjecture proved by Perelman [17, 18, 19] implies that the universal cover of  $M$  is either non-compact or diffeomorphic to  $S^3$ . In the latter case, the elliptisation conjecture also proved by Perelman guarantees that  $M$  is diffeomorphic to a quotient of the standard round  $S^3$  by the action of a finite group of isometries and the quotient metric turns  $M$  into a  $Y_\ell^x$ -manifold. Thus, Theorem 1.1 fails for every surface or 3-manifold such that its universal cover is not open.

The weak form of the Bott–Samelson theorem proved by Bérard-Bergery, see [2] and [3, Theorem 7.37], says that if  $(M, g)$  is a  $Y_\ell^x$ -manifold, then the universal cover of  $M$  is compact and the rational cohomology ring  $H^*(M, \mathbb{Q})$  is generated by one element. In view of the preceding discussion, it seems natural to ask whether the latter property is also shared by all manifolds  $M$  such that there exists a positive Legendrian loop in the Legendrian isotopy class of the fibre of  $ST^*M$ .  $\square$

**Remark 8.4.** Let  $p : Z \rightarrow N$  be a Legendrian fibration of a co-oriented contact manifold. Suppose that there exists a contact covering  $ST^*M \rightarrow Z$  such that  $M$  is open and the pre-image of any fibre of  $p$  is a union of fibres of  $ST^*M$ . (Note that  $p$  does not have to be locally trivial and its fibres do not have to be spheres.) Then Theorem 1.1 and Corollary 8.1 hold for the fibres of  $p$ .

**9. Orderability of  $ST^*M$ .** Let  $(Y, \ker \alpha)$  be a connected contact manifold. Consider the identity component  $\text{Cont}_0(Y)$  of the group of compactly supported contactomorphisms of  $(Y, \ker \alpha)$  and let  $\widetilde{\text{Cont}}_0(Y)$  denote the universal cover of this group corresponding to the base point  $\text{id}_Y \in \text{Cont}_0(Y)$ . For  $f, g \in \widetilde{\text{Cont}}_0(Y)$ , write  $f \preceq g$  if the element  $gf^{-1}$  can be represented by a path  $\varphi_t \in \text{Cont}_0(Y)$  such that the contact Hamiltonian  $H := \alpha(\frac{d\varphi_t}{dt})$  is non-negative. Following Eliashberg and Polterovich [11], we say that the contact manifold  $(Y, \ker \alpha)$  is *orderable* if the relation  $\preceq$  defines a genuine partial order on  $\widetilde{\text{Cont}}_0(Y)$ .

Eliashberg, Kim, and Polterovich [10, Theorem 1.18] used contact homology to prove that  $ST^*M$  is orderable for a closed manifold  $M$  such that its fundamental group  $\pi_1(M)$  is either finite or has infinitely many conjugacy classes. It is an open problem whether an infinite finitely presented group can have finitely many conjugacy classes, see [1, Problem (FP19)]. The following result shows that the orderability of  $ST^*M$  does not depend on the solution of that problem.

**Corollary 9.1.**  *$ST^*M$  is orderable for any closed manifold  $M$ .*

*Proof.* By [11, Criterion 1.2.C], a closed contact manifold  $(Y, \ker \alpha)$  is orderable if and only if there does not exist a contractible loop of contactomorphisms  $\varphi_t \in \text{Cont}_0(Y)$ ,  $t \in [0, 1]$ , such that  $\varphi_0 = \varphi_1 = \text{id}_Y$  and the corresponding contact Hamiltonian is everywhere positive. It is clear that applying a contact isotopy of  $\text{id}_Y$  generated by a positive contact Hamiltonian to any Legendrian submanifold  $L \subset Y$ , we obtain a positive Legendrian isotopy of  $L$ . Thus, if  $Y$  is not orderable, then every Legendrian isotopy class contains a positive (contractible) Legendrian loop.

Suppose that  $ST^*M$  is not orderable. By [10, Theorem 1.18], the fundamental group of  $M$  is infinite and hence the universal cover of  $M$  is open. In that case, however, the Legendrian isotopy class of the fibre of  $ST^*M$  does not contain positive Legendrian loops by Corollary 8.1, a contradiction.  $\square$

**Example 9.2.** The proof of Corollary 9.1 shows that if  $\pi_1(M)$  is infinite, then there are no positive loops in  $\text{Cont}_0(ST^*M)$ , contractible or not. On the other hand, the (co-)geodesic flow of the standard round metric on the  $m$ -sphere  $S^m$  defines a *non-contractible* positive loop in  $\text{Cont}_0(ST^*S^m)$ , cf. Example 8.3.  $\square$

**Corollary 9.3.** *Let  $p : ST^*M \rightarrow Z$  be a contact covering of a closed contact manifold  $Z$ . Then  $Z$  is orderable.*

*Proof.* Suppose that  $Z$  is not orderable and argue by contradiction. By [11, Criterion 1.2.C], there exists a contractible loop of contactomorphisms of  $Z$  based at  $\text{id}_Z$  and generated by a positive contact Hamiltonian. Since this loop is contractible, it lifts to a loop of contactomorphisms of  $ST^*M$  with the same properties. If  $M$  is closed, we conclude that  $ST^*M$  is not orderable by [11, Criterion 1.2.C], which contradicts Corollary 9.1. If  $M$  is open, then the argument from the proof of that corollary shows that there exists a positive Legendrian loop based at a fibre of  $ST^*M$ , which contradicts Corollary 8.1.  $\square$

**10. Legendrian Low Conjecture.** Here we give a very brief exposition of the relevant material from Lorentz geometry. Further details and references may be found in [6].

Let  $(X, g)$  be a globally hyperbolic spacetime with a smooth spacelike Cauchy surface  $M \subset X$ . That is to say,  $(X, g)$  is a connected time-oriented Lorentz manifold and  $M$  is a smooth spacelike hypersurface in  $X$  such that every inextensible future directed curve in  $X$  meets  $M$  exactly once. (The time orientation on  $X$  is a continuous choice of the future and past hemicones  $C_x^\uparrow$  and  $C_x^\downarrow$  in the non-spacelike cone at each  $x \in X$ . A piecewise smooth curve  $\gamma = \gamma(t)$  is called future directed if  $\dot{\gamma}(t) \in C_{\gamma(t)}^\uparrow$  for all  $t$ .)

**Definition 10.1.** Two points  $x, y \in X$  are called *causally related* if they can be connected by a future or past directed curve.

Let  $\mathfrak{N}$  be the set of all future directed non-parameterised null geodesics in  $(X, g)$  or, in other words, the set of all light rays of our spacetime.  $\mathfrak{N}$  has a canonical structure of a contact manifold, see [16, pp. 252–253]. There is a contactomorphism

$$\rho_M : \mathfrak{N} \xrightarrow{\cong} ST^*M$$

that associates to a null geodesic  $\gamma \in \mathfrak{N}$  the equivalence class of the (non-zero) linear form  $v \mapsto g(\dot{\gamma}, v)$  on  $T_{\gamma \cap M}M$ , where  $\dot{\gamma}$  is a future pointing tangent vector to  $\gamma$  at  $\gamma \cap M$ .

The set  $\mathfrak{S}_x$  of all null geodesics passing through a point  $x \in X$  is a Legendrian sphere in  $\mathfrak{N}$  called the *sky* of that point. Note that two skies intersect if and only if the corresponding points lie on the same null geodesic. Note also that  $\rho_M(\mathfrak{S}_x) = ST_x^*M$  for any  $x \in M$ .

Since  $X$  is connected, the skies of any two points are Legendrian isotopic in  $\mathfrak{N}$ . However, Legendrian links formed by unions of disjoint skies may be quite different.

A basic observation is that all Legendrian links  $\mathfrak{S}_x \sqcup \mathfrak{S}_y$  corresponding to causally *unrelated* points  $x, y \in X$  belong to the same Legendrian isotopy class, see [6, Lemma 4.3]. Let us denote this isotopy class of Legendrian links by  $\mathcal{U}$  (as in *unrelated* and *unlinked*). A natural way to represent  $\mathcal{U}$  is to pick the points  $x$  and  $y$  on the Cauchy surface  $M$  so that  $\rho_M$  identifies  $\mathfrak{S}_x \sqcup \mathfrak{S}_y$  with  $ST_x^*M \sqcup ST_y^*M \subset ST^*M$ .

Two skies  $\mathfrak{S}_x, \mathfrak{S}_y \subset \mathfrak{N}$  are said to be *Legendrian linked* if either  $\mathfrak{S}_x \cap \mathfrak{S}_y \neq \emptyset$  or the Legendrian link  $\mathfrak{S}_x \sqcup \mathfrak{S}_y$  does *not* belong to  $\mathcal{U}$ . We have just seen that if  $\mathfrak{S}_x$  and  $\mathfrak{S}_y$  are Legendrian linked, then the points  $x$  and  $y$  are causally related.

**Definition 10.2.** The *Legendrian Low Conjecture* holds for a globally hyperbolic spacetime if two points in it are causally related if and *only if* their skies are Legendrian linked.

**Remark 10.3.** The general problem of describing causal relations in terms of linking in the space of null geodesics originates from the work of Robert Low that was apparently inspired by a question raised by Penrose, see e.g. [14] and [15]. The Legendrian Low Conjecture was explicitly stated by Natário and Tod [16, Conjecture 6.4] in the case when the Cauchy surface  $M$  is diffeomorphic to an open subset of  $\mathbb{R}^3$ .

It was shown in our paper [6] that the Legendrian Low Conjecture holds for any globally hyperbolic spacetime such that its Cauchy surface has a cover diffeomorphic to an open subset of  $\mathbb{R}^m$ ,  $m \geq 2$ . Using Theorem 1.1 instead of [6, Corollary 6.2], we can now extend our result to a wider class of spacetimes.

**Theorem 10.4.** *The Legendrian Low Conjecture holds for any globally hyperbolic spacetime such that the universal cover of its Cauchy surface is not compact.*

*Proof.* Let  $x, y \in X$  be two points such that their skies are disjoint and there exists a future directed curve connecting  $x$  to  $y$ . By [6, Proposition 4.2], there exists a non-negative Legendrian isotopy connecting  $\mathfrak{S}_y$  to  $\mathfrak{S}_x$ . Suppose that the link  $\mathfrak{S}_x \sqcup \mathfrak{S}_y$  belongs to  $\mathcal{U}$ . Then the link  $\rho_M(\mathfrak{S}_x \sqcup \mathfrak{S}_y) \subset ST^*M$  is Legendrian isotopic to a link formed by a pair of fibres of  $ST^*M$ . Since Legendrian isotopic links are ambiently contactomorphic, we



obtain a non-negative Legendrian isotopy connecting two different fibres of  $ST^*M$ , which contradicts Theorem 1.1. Thus,  $\mathfrak{S}_x$  and  $\mathfrak{S}_y$  are Legendrian linked.  $\square$

Combining this theorem with Perelman's proof of the Poincaré conjecture, we see that the Legendrian Low Conjecture holds for any  $(3+1)$ -dimensional globally hyperbolic spacetime such that the universal cover of its Cauchy surface is not diffeomorphic to  $S^3$ . (This result was obtained in [6] by a more involved argument using the full strength of the geometrisation conjecture.) On the other hand, if the Cauchy surface is a quotient of  $S^3$ , then the Legendrian Low Conjecture may fail because of the following general construction, cf. [7, Example 3].

**Example 10.5.** If  $(M, \bar{g})$  is a Riemann  $Y_\ell^x$ -manifold (see Example 8.3), then the Legendrian Low Conjecture does not hold for the globally hyperbolic spacetime  $(M \times \mathbb{R}, \bar{g} \oplus -dt^2)$ . Indeed, null geodesics in this spacetime have the form  $\gamma(s) = (\bar{\gamma}(s), s)$ , where  $\bar{\gamma}$  is a  $\bar{g}$ -geodesic on  $M$  and  $s$  is the natural parameter on  $\bar{\gamma}$ . In particular,  $\rho_{M \times \{0\}}(\mathfrak{S}_{(x,\ell)}) = ST_x^*M$  by the definition of a  $Y_\ell^x$ -manifold. Thus, the skies  $\mathfrak{S}_{(x',0)}$  and  $\mathfrak{S}_{(x,\ell)}$  are *not* Legendrian linked if  $x' \neq x \in M$ . However, the points  $(x', 0)$  and  $(x, \ell)$  in  $M \times \mathbb{R}$  are causally related if  $x'$  is sufficiently close to  $x$  in  $M$ .  $\square$

**Remark 10.6.** One can use Remark 8.2 and the proof of [6, Theorem C] to show that if the universal cover of the Cauchy surface of a globally hyperbolic spacetime  $X$  is non-compact, then the Legendrian links  $\mathfrak{S}_x \sqcup \mathfrak{S}_y$  and  $\mathfrak{S}_y \sqcup \mathfrak{S}_x$  are different for any pair of causally related points  $x, y \in X$  with disjoint skies.

## REFERENCES

- [1] G. Baumslag, A. G. Myasnikov, V. Shpilrain, *Open problems in combinatorial group theory. Second edition*, Combinatorial and geometric group theory (New York, 2000/Hoboken, NJ, 2001), Contemp. Math. **296**, Amer. Math. Soc., Providence, RI (2002), 1–38.
- [2] L. Bérard-Bergery, *Quelques exemples de variétés riemanniennes où toutes les géodésiques issues d'un point sont fermées et de même longueur, suivies de quelques résultats sur leur topologie*, Ann. Inst. Fourier (Grenoble) **27** (1977), 231–249.
- [3] A. L. Besse, *Manifolds all of whose geodesics are closed*, with appendices by D. B. A. Epstein, J.-P. Bourguignon, L. Bérard-Bergery, M. Berger and J. L. Kazdan. Ergebnisse der Mathematik und ihrer Grenzgebiete, 93. Springer-Verlag, Berlin-New York, 1978.
- [4] M. Bhupal, *A partial order on the group of contactomorphisms of  $\mathbb{R}^{2n+1}$  via generating functions*, Turkish J. Math. **25** (2001), 125–135.
- [5] Yu. V. Chekanov, *Critical points of quasifunctions, and generating families of Legendrian manifolds*, Funktsional. Anal. i Prilozhen. **30**:2 (1996), 56–69 (Russian); English transl. in Funct. Anal. Appl. **30**:2 (1996), 118–128.
- [6] V. Chernov, S. Nemirovski, *Legendrian links, causality, and the Low conjecture*, Preprint [arXiv:0810.5091](https://arxiv.org/abs/0810.5091).
- [7] V. Chernov (Tchernov), Yu. Rudyak, *Linking and causality in globally hyperbolic space-times*, Comm. Math. Phys. **279** (2008), 309–354.
- [8] V. Colin, E. Ferrand, P. Pushkar, *Positive isotopies of Legendrian submanifolds*, Preprint available at <http://people.math.jussieu.fr/~ferrand/publi/PIL.pdf>
- [9] Ya. Eliashberg, M. Gromov, *Lagrangian intersection theory: finite-dimensional approach*, Geometry of differential equations, 27–118, Amer. Math. Soc. Transl. Ser. 2, **186**, AMS, Providence, RI, 1998.
- [10] Y. Eliashberg, S. S. Kim, L. Polterovich, *Geometry of contact transformations and domains: orderability versus squeezing*, Geom. Topol. **10** (2006), 1635–1747; *Erratum*, Geom. Topol. **13** (2009), 1175–1176.
- [11] Y. Eliashberg, L. Polterovich, *Partially ordered groups and geometry of contact transformations*, Geom. Funct. Anal. **10** (2000), 1448–1476.
- [12] C. Godbillon, *Feuilletages. Études géométriques*, Progress in Mathematics, 98. Birkhäuser Verlag, Basel, 1991.

- [13] F. Laudenbach, J.-C. Sikorav, *Persistence d'intersection avec la section nulle au cours d'une isotopie hamiltonienne dans un fibré cotangent*, Invent. Math. **82** (1985), 349–357.
- [14] R. J. Low, *Causal relations and spaces of null geodesics*, PhD Thesis, Oxford University (1988).
- [15] R. J. Low, *The space of null geodesics*, Proceedings of the Third World Congress of Nonlinear Analysts, Part 5 (Catania, 2000). Nonlinear Anal. **47** (2001), 3005–3017.
- [16] J. Natário, P. Tod, *Linking, Legendrian linking and causality*, Proc. London Math. Soc. (3) **88** (2004), 251–272.
- [17] G. Perelman, *The entropy formula for the Ricci flow and its geometric applications*, Preprint [math.DG/0211159](#).
- [18] G. Perelman, *Ricci flow with surgery on three-manifolds*, Preprint [math.DG/0303109](#).
- [19] G. Perelman, *Finite extinction time for the solutions to the Ricci flow on certain three-manifolds*, Preprint [math.DG/0307245](#).
- [20] C. Viterbo, *Symplectic topology as the geometry of generating functions*, Math. Ann. **292** (1992), 685–710.

DEPARTMENT OF MATHEMATICS, 6188 KEMENY HALL, DARTMOUTH COLLEGE, HANOVER, NH 03755, USA

*E-mail address:* [Vladimir.Chernov@dartmouth.edu](mailto:Vladimir.Chernov@dartmouth.edu)

STEKLOV MATHEMATICAL INSTITUTE, 119991 MOSCOW, RUSSIA;

MATHEMATISCHES INSTITUT, RUHR-UNIVERSITÄT BOCHUM, 44780 BOCHUM, GERMANY

*E-mail address:* [stefan@mi.ras.ru](mailto:stefan@mi.ras.ru)