

About amenability of subgroups of the group of diffeomorphisms of the interval.

E.T.Shavgulidze

Department of Mechanics and Mathematics, Moscow State University, Moscow, 119991 Russia

Averaging linear functional on the space continuous functions of the group of diffeomorphisms of interval is found. Amenability of several discrete subgroups of the group of diffeomorphisms $\text{Diff}^3([0, 1])$ of interval is prove. In particular, a solution of the problem of amenability of the Thompson's group F is given. See about the Thompson's group F in [1].

Let $\text{Diff}_+^1([0, 1])$ be the group of all diffeomorphisms of class C^1 of interval $[0, 1]$ that preserve the endpoints of interval, and let $\text{Diff}_+^3([0, 1])$ be the subgroup of $\text{Diff}_+^1([0, 1])$ consisting of all diffeomorphisms of class $C^3([0, 1])$, and

$$\text{Diff}_0^3([0, 1]) = \{f \in \text{Diff}_+^3([0, 1]) : f'(0) = f'(1) = 1\}.$$

The group $\text{Diff}_+^1([0, 1])$ is equipped with the topology inherited from the space $C^1([0, 1])$.

For any positive $\delta < 1$, denote by $C_0^{1,\delta}([0, 1])$ the set of all functions $f \in C^1([0, 1])$ such that $f(0) = 0$ and $\exists C > 0 \forall t_1, t_2 \in [0, 1] |f'(t_2) - f'(t_1)| < C|t_2 - t_1|^\delta$. Define a Banach structure on the linear space $C_0^{1,\delta}([0, 1])$ by a norm

$$\|f\|_{1,\delta} = |f'(0)| + \sup_{t_1, t_2 \in [0, 1]} \frac{|f'(t_2) - f'(t_1)|}{|t_2 - t_1|^\delta}$$

for any function $f \in C_0^{1,\delta}([0, 1])$.

Let $\text{Diff}_+^{1,\delta}([0, 1]) = \text{Diff}_+^1([0, 1]) \cap C_0^{1,\delta}([0, 1])$. It is easy to see that $\text{Diff}_+^{1,\delta}([0, 1])$ is a subgroup of the group $\text{Diff}_+^1([0, 1])$. The subgroup $\text{Diff}_+^{1,\delta}([0, 1])$ is equipped with the topology inherited from the space $C_0^{1,\delta}([0, 1])$.

Let $C_b(\text{Diff}_+^{1,\delta}([0, 1]))$ be the linear space of all bounded continuous functions on the space $\text{Diff}_+^{1,\delta}([0, 1])$, and let $C_b(\text{Diff}_+^1([0, 1]))$ be the linear space of all bounded continuous functions on the space $\text{Diff}_+^1([0, 1])$.

Introduce the functions $e_{1,\delta} : \text{Diff}_+^{1,\delta}([0, 1]) \rightarrow \mathbf{R}$, $e_{1,0} : \text{Diff}_+^1([0, 1]) \rightarrow \mathbf{R}$ by setting $e_{1,\delta}(g) = 1$ for any $g \in \text{Diff}_+^{1,\delta}([0, 1])$ and $e_{1,0}(f) = 1$ for any $f \in \text{Diff}_+^1([0, 1])$. Let $F_g(f) = F(g^{-1} \circ f)$ for any $g \in \text{Diff}_0^3([0, 1])$, $f \in \text{Diff}_+^{1,\delta}([0, 1])$ and $F \in C_b(\text{Diff}_+^{1,\delta}([0, 1]))$.

Theorem 1. *For any positive $\delta < \frac{1}{2}$, there exists a linear functional $L_\delta : C_b(\text{Diff}_+^{1,\delta}([0, 1])) \rightarrow \mathbf{R}$ such that $L_\delta(e_{1,\delta}) = 1$, $|L_\delta(F)| \leq \sup_{f \in \text{Diff}_+^{1,\delta}([0, 1])} |F(f)|$,*

$L_\delta(F) \geq 0$ for any nonnegative function $F \in C_b(\text{Diff}_+^{1,\delta}([0, 1]))$, and

$L_\delta(F_g) = L_\delta(F)$ for any $g \in \text{Diff}_0^3([0, 1])$ and $F \in C_b(\text{Diff}_+^{1,\delta}([0, 1]))$.

The restriction of any function of the space $C_b(\text{Diff}_+^1([0, 1]))$ on $\text{Diff}_+^{1,\delta}([0, 1])$ belong to the space $C_b(\text{Diff}_+^{1,\delta}([0, 1]))$. Hence we obtain the following assertion.

Corollary 1.1. *There exists a linear functional $L_0 : C_b(\text{Diff}_+^1([0, 1])) \rightarrow \mathbf{R}$ such that $L_0(e_{1,0}) = 1$, $|L_0(F)| \leq \sup_{f \in \text{Diff}_+^1([0, 1])} |F(f)|$, $L_0(F) \geq 0$ for any nonnegative*

function $F \in C_b(\text{Diff}_+^1([0, 1]))$, and $L_0(F_g) = L_0(F)$ for any $g \in \text{Diff}_0^3([0, 1])$ and $F \in C_b(\text{Diff}_+^1([0, 1]))$.

We say that a discrete subgroup G of $\text{Diff}_0^3([0, 1])$ satisfies condition (a), if there is a such $C > 0$ that

$$\sup_{t \in [0, 1]} |\ln(g_1'(t)) - \ln(g_2'(t))| \geq C \text{ for any } g_1, g_2 \in G, g_1 \neq g_2.$$

Theorem 2. *If a discrete subgroup G of $\text{Diff}_0^3([0, 1])$ satisfies condition (a), then the subgroup G is amenable.*

Proof. Let $B(G)$ be the linear space of all bounded functions on the group G .

Let positive $\delta < \frac{1}{2}$, let

$$p(f) = |\ln(f'(0))| + \sup_{t_1, t_2 \in [0, 1]} \frac{|\ln(f'(t_2)) - \ln(f'(t_1))|}{|t_2 - t_1|^\delta}$$

and $r(f) = \inf_{h \in G} p(h^{-1} \circ f)$ for $f \in \text{Diff}_+^{1, \delta}([0, 1])$, $\theta(t) = 1 - t$ for $t \in [0, 1]$ and $\theta(t) = 0$ for $t > 1$.

For any fixed $f \in \text{Diff}_+^{1, \delta}([0, 1])$, $C > 0$, the set of functions $\{\psi : \psi(t) = \ln(g'(t)), g \in G, p_n(g \circ f) < C\}$ contain in a compact subset of the space $C([0, 1])$, therefore it is finite according to condition (a). Hence, we can define the linear mapping $\pi_\delta : B(G) \rightarrow C_b(\text{Diff}_+^{1, \delta}([0, 1]))$ by setting

$$\pi_\delta F(f) = \frac{\sum_{h \in G} \theta(p_\delta(h^{-1} \circ f) - r(f))F(h)}{\sum_{h \in G} \theta(p_\delta(h^{-1} \circ f) - r(f))}.$$

Assign a linear functional $l : B(G) \rightarrow \mathbf{R}$ by setting $l(F) = L_\delta(\pi_\delta F)$.

It is easy to see that

$$|l(F)| = |L_\delta(\pi_\delta F)| \leq \sup_{f \in \text{Diff}_+^{1, \delta}([0, 1])} |\pi_\delta F(f)| \leq \sup_{g \in G} |F(g)|,$$

$l(F) \geq 0$ for any nonnegative function $F \in B(G)$, and $l(e_G) = 1$, where $e_G(g) = 1$ for all $g \in G$.

Denote by $F_g(h) = F(g^{-1} \circ h)$ for $F \in B(G)$, $g, h \in G$.

We have

$$\begin{aligned} \pi_\delta F_g(f) &= \frac{\sum_{h \in G} \theta(p_\delta(h^{-1} \circ f) - r(f))F(g^{-1} \circ h)}{\sum_{h \in G} \theta(p_\delta(h^{-1} \circ f) - r(f))} = \\ &= \frac{\sum_{h \in G} \theta(p_\delta(h^{-1} \circ g \circ f) - r(g \circ f))F(h)}{\sum_{h \in G} \theta(p_\delta(h^{-1} \circ g \circ f) - r(g \circ f))} = \pi_\delta F(g \circ f), \end{aligned}$$

hence $l(F_g) = L_\delta(\pi_\delta F_g) = L_\delta(\pi_\delta F) = l(F)$,

which implies the assertion of Theorem 2.

In [2] E.Ghys and V.Sergiescu proved that The Thompson's group F is isomorphic to a discrete subgroup G of $\text{Diff}_0^3([0, 1])$ which satisfies condition (a), and therefore we obtain the following assertion.

Corollary 2.1. *The Thompson's group F is amenable.*

For the proof Theorem 1 we need the following auxiliary assertions.

Lemma 1. *There exist positive constants c_1, c_2 such that*

$$\begin{aligned} c_1 2^{n+1} (n+1)! &\leq \int_{-\infty}^{+\infty} \dots \int_{-\infty}^{+\infty} \frac{dx_1 \dots dx_n}{\sqrt{(1+x_1^2)(1+(x_2-x_1)^2) \dots (1+(x_n-x_{n-1})^2)(1+x_n^2)}} \leq \\ &\leq c_2 2^{n+1} (n+1)! \end{aligned}$$

for any natural n .

Proof. The Fourier transform of the function $\frac{1}{\sqrt{1+x^2}}$ is given by the Bessel function of the second kind

$$\frac{2}{\sqrt{2\pi}} K_0(y) = \frac{2}{\sqrt{2\pi}} \int_0^{+\infty} \frac{\cos(xy) dx}{\sqrt{1+x^2}} = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{+\infty} \frac{e^{-ixy} dx}{\sqrt{1+x^2}}$$

(see [6]).

Hence we have

$$\begin{aligned} I &= \int_{-\infty}^{+\infty} \dots \int_{-\infty}^{+\infty} \frac{dx_1 \dots dx_n}{\sqrt{(1+x_1^2)(1+(x_2-x_1)^2) \dots (1+(x_n-x_{n-1})^2)(1+x_n^2)}} = \\ &= \frac{2^n}{\pi} \int_{-\infty}^{+\infty} (K_0(y))^{n+1} dx = \frac{2^{n+1}}{\pi} \int_0^{+\infty} (K_0(y))^{n+1} dx. \end{aligned}$$

There exist positive constants $\varepsilon < 1$, c such that $-\ln \frac{y}{\varepsilon} < K_0(y) < -\ln \frac{y}{2}$ for any $y \in (0, \varepsilon)$, and $0 < K_0(y) < ce^{-y}$ for any $y \geq \varepsilon$.

Thus,

$$\begin{aligned} I &\leq \frac{2^{n+1}}{\pi} \left[\int_0^{\varepsilon} \left(-\ln\left(\frac{y}{2}\right)\right)^{n+1} dy + \int_{\varepsilon}^{+\infty} (ce^{-y})^{n+1} dy \right] \leq \\ &\leq \frac{2^{n+1}}{\pi} \left[\int_0^2 \left(-\ln\left(\frac{y}{2}\right)\right)^{n+1} dy + c^{n+1} \int_{\varepsilon}^{+\infty} e^{-(n+1)y} dy \right] = \\ &= \frac{2^{n+1}}{\pi} \left[2((n+1)!) + \frac{c^{n+1} e^{-(n+1)\varepsilon}}{n+1} \right], \end{aligned}$$

and

$$I \geq \frac{2^{n+1}}{\pi} \int_0^{\varepsilon} \left(-\ln\left(\frac{y}{\varepsilon}\right)\right)^{n+1} dy = \frac{\varepsilon}{\pi} 2^{n+1} (n+1)!,$$

which implies the assertion of lemma 1.

Let

$$v_1(\tau) = \int_{-\infty}^{+\infty} \frac{d\tau_1}{\sqrt{(1+\tau_1^2)(1+(\tau-\tau_1)^2)}}$$

for $\tau \in \mathbf{R}$.

Lemma 2. *The derivation $v_1'(t)$ is negative for any $t > 0$, and $|v_1'(t)| \leq \frac{4}{|t|} v_1(t)$ for any $t \neq 0$.*

Proof. The derivation of the function $v_1(t)$ is equal

$$v_1'(t) = - \int_{-\infty}^{+\infty} \frac{(t - \tau) d\tau}{\sqrt{(1 + \tau^2)(1 + (t - \tau)^2)^3}}.$$

Taking $\tau_1 = t - \tau$ we receive

$$\begin{aligned} v_1'(t) &= - \int_{-\infty}^{+\infty} \frac{\tau_1 d\tau_1}{\sqrt{(1 + \tau_1^2)^3(1 + (\tau_1 - t)^2)}} = \\ &= - \int_0^{+\infty} \left(\frac{1}{\sqrt{1 + (\tau_1 - t)^2}} - \frac{1}{\sqrt{1 + (\tau_1 + t)^2}} \right) \frac{\tau_1 d\tau_1}{\sqrt{(1 + \tau_1^2)^3}} = \\ &= - \int_0^{+\infty} \frac{4 t \tau_1^2 d\tau_1}{(\sqrt{1 + (\tau_1 - t)^2} + \sqrt{1 + (\tau_1 + t)^2}) \sqrt{(1 + \tau_1^2)^3(1 + (\tau_1 - t)^2)(1 + (\tau_1 + t)^2)}}. \end{aligned}$$

Hence, if $t > 0$, then $v_1'(t) < 0$.

If $t \neq 0$, then

$$\begin{aligned} |v_1'(t)| &\leq \int_0^{+\infty} \frac{4 |t| \tau_1^2 d\tau_1}{(1 + \tau_1^2)(1 + (\tau_1 + |t|)^2) \sqrt{(1 + (\tau_1 - t)^2)(1 + \tau_1^2)}} \leq \\ &\leq \frac{4}{|t|} \int_0^{+\infty} \frac{d\tau_1}{\sqrt{(1 + (\tau_1 - t)^2)(1 + \tau_1^2)}} = \frac{4}{|t|} v_1(t), \end{aligned}$$

which implies the assertion of lemma 2.

Let $v(t) = v_1(\ln(t + \sqrt{t^2 - 1}))$ for $t \geq 1$.

Let us denote $D_n = \{(x_1, \dots, x_{n-1}) : 0 < x_1 < \dots < x_{n-1} < 1\}$

and $x_{-1} = x_{n-1} - 1$, $x_0 = 0$, $x_n = 1$.

Let

$$u_{1,n}(x_1, \dots, x_{n-1}) = \prod_{k=1}^n \frac{1}{x_k - x_{k-1}} v\left(\frac{x_k - x_{k-2}}{2\sqrt{(x_k - x_{k-1})(x_{k-1} - x_{k-2})}}\right),$$

$$J_n = \int_0^1 \int_{x_1}^1 \dots \int_{x_{n-2}}^1 u_{1,n}(x_1, x_2, \dots, x_{n-1}) dx_1 dx_2 \dots dx_{n-1},$$

$$u_n(x_1, \dots, x_{n-1}) = \frac{1}{J_n} u_{1,n}(x_1, \dots, x_{n-1}).$$

Lemma 3. *The following equality is fulfilled*

$$J_n = 2^{n-1} \int_{-\infty}^{+\infty} \dots \int_{-\infty}^{+\infty} \frac{dt_1 dt_2 \dots dt_{2n-1}}{\sqrt{(1 + t_1^2)(1 + (t_2 - t_1)^2) \dots (1 + (t_{2n-1} - t_{2n-2})^2)(1 + t_{2n-1}^2)}}$$

for any natural n , and $c_1 2^{3n-1} (2n)! \leq J_n \leq c_2 2^{3n-1} (2n)! .$

Proof. Taking $y_k = \frac{x_k - x_{k-1}}{1 - x_{n-1}}$ we have

$$J_n = \int_0^1 \int_{x_1}^1 \dots \int_{x_{n-2}}^1 u_{1,n}(x_1, x_2, \dots, x_{n-1}) dx_1 dx_2 \dots dx_{n-1} =$$

$$= \int_0^{+\infty} \int_0^{+\infty} \dots \int_0^{+\infty} v\left(\frac{y_1+1}{2\sqrt{y_1}}\right) v\left(\frac{1+y_{n-1}}{2\sqrt{y_{n-1}}}\right) \prod_{k=2}^{n-1} v\left(\frac{y_k+y_{k-1}}{2\sqrt{y_k y_{k-1}}}\right) \frac{dy_1 dy_2 \dots dy_{n-1}}{y_1 y_2 \dots y_{n-1}}.$$

Making substitutions $t_{2k} = \frac{1}{2} \ln(y_k)$ and taking into account

$$\begin{aligned} \ln\left(\frac{y_k+y_{k-1}}{2\sqrt{y_k y_{k-1}}}\right) + \sqrt{\left(\frac{y_k+y_{k-1}}{2\sqrt{y_k y_{k-1}}}\right)^2 - 1} &= |t_{2k} - t_{2k-2}|, \\ v\left(\frac{y_k+y_{k-1}}{2\sqrt{y_k y_{k-1}}}\right) &= v_1(|t_{2k} - t_{2k-2}|) = \\ &= 2 \int_{-\infty}^{+\infty} \frac{dt_{2k-1}}{\sqrt{(1+(t_{2k-1}-t_{2k-2})^2)(1+(t_{2k}-t_{2k-1})^2)}}, \\ v\left(\frac{y_1+1}{2\sqrt{y_1}}\right) &= 2 \int_{-\infty}^{+\infty} \frac{dt_1}{\sqrt{(1+t_1^2)(1+(t_2-t_1)^2)}}, \\ v\left(\frac{1+y_{n-1}}{2\sqrt{y_{n-1}}}\right) &= 2 \int_{-\infty}^{+\infty} \frac{dt_{2n-1}}{\sqrt{(1+t_{2n-1}^2)(1+(t_{2n-1}-t_{2n-2})^2)}}, \end{aligned}$$

we receive

$$J_n = 2^{n-1} \int_{-\infty}^{+\infty} \dots \int_{-\infty}^{+\infty} \frac{dt_1 dt_2 \dots dt_{2n-1}}{\sqrt{(1+t_1^2)(1+(t_2-t_1)^2) \dots (1+(t_{2n-1}-t_{2n-2})^2)(1+t_{2n-1}^2)}}.$$

Thus the assertion of lemma 3 is an immediate consequence of lemma 1.

Lemma 4. For arbitrary positive ε , it exist positive constants c_3 such that

$$v\left(\frac{y_1+a}{2\sqrt{y_1 a}}\right) v\left(\frac{a+y_2}{2\sqrt{a y_2}}\right) \leq c_3 v\left(\frac{y_1+y_2}{2\sqrt{y_1 y_2}}\right)$$

for all a, y_1, y_2 , satisfying $\varepsilon \leq a \leq 1, y_1 > 0, y_2 > 0, y_1 + y_2 \leq 1$.

Proof. Let $t_1 = -\frac{1}{2} \ln(y_1), t_2 = -\frac{1}{2} \ln(y_2), \alpha = -\frac{1}{2} \ln(a)$ then $0 \leq \tau_0 \leq r = -\frac{1}{2} \ln(\varepsilon)$ and

$$v\left(\frac{y_1+a}{2\sqrt{y_1 a}}\right) = v_1(t_1 - \alpha), v\left(\frac{a+y_2}{2\sqrt{a y_2}}\right) = v_1(t_2 - \alpha), v\left(\frac{y_1+y_2}{2\sqrt{y_1 y_2}}\right) = v_1(t_2 - t_1).$$

It follows from the lemma 2 that the derivation $v_1'(t)$ is negative for any $t > 0$ and the function $v_1(t)$ decrease on $[0, +\infty)$.

Thus, $v_1(t) \leq v_1(0) = \int_{-\infty}^{+\infty} \frac{d\tau}{(1+\tau^2)} = \pi$ for any $t \geq 0$.

As $v_1(t) > 0$ for any $t \in \mathbf{R}$, the function $v_1(t)$ is continuous and

$$\lim_{t \rightarrow +\infty} \frac{v_1(t-r)}{v_1(t)} = 1,$$

then it exists positive constant c^* such that $v_1(t-\tau) \leq c^* v_1(t)$ for any $t > 0, \tau \in [0, r]$.

Let $c_3 = \pi(c^*)^2$.

If $t_2 \geq t_1$, then $t_2 \geq t_2 - t_1 \geq 0$ and

$$v_1(t_1 - \alpha) v_1(t_2 - \alpha) \leq (c^*)^2 v_1(t_1) v_1(t_2) \leq \pi(c^*)^2 v_1(t_2) \leq \pi(c^*)^2 v_1(t_2 - t_1) = c_3 v_1(t_2 - t_1).$$

If $t_1 \geq t_2$, then $v_1(t_1 - \alpha)v_1(t_2 - \alpha) \leq c_3v_1(t_1 - t_2) = c_3v_1(t_2 - t_1)$.

Hence, $v(\frac{y_1+a}{2\sqrt{y_1a}})v(\frac{a+y_2}{2\sqrt{ay_2}}) \leq c_3v(\frac{y_1+y_2}{2\sqrt{y_1y_2}})$.

Let us define a function $\vartheta(t) = 1$ for $t \in [0, 1]$ and $\vartheta(t) = 0$ for $t \in (-\infty, 0) \cup (1, +\infty)$.

Lemma 5. *The following equality is valid*

$$\lim_{n \rightarrow \infty} \left(\int_0^1 \int_{x_1}^1 \dots \int_{x_{n-2}}^1 (1 - \vartheta(\frac{1}{\varepsilon} \max_{1 \leq k \leq n} (x_k - x_{k-1}))) u_n(x_1, x_2, \dots, x_{n-1}) dx_1 dx_2 \dots dx_{n-1} \right) = 0$$

for any positive $\varepsilon < 1$.

Proof. We have

$$\begin{aligned} & \int_0^1 \int_{x_1}^1 \dots \int_{x_{n-2}}^1 (1 - \vartheta(\frac{1}{\varepsilon} \max_{1 \leq k \leq n} (x_k - x_{k-1}))) u_n(x_1, x_2, \dots, x_{n-1}) dx_1 dx_2 \dots dx_{n-1} \leq \\ & \leq \sum_{k=1}^n \int_0^1 \int_{x_1}^1 \dots \int_{x_{n-2}}^1 (1 - \vartheta(\frac{1}{\varepsilon} (x_k - x_{k-1}))) u_n(x_1, x_2, \dots, x_{n-1}) dx_1 dx_2 \dots dx_{n-1}. \end{aligned}$$

Let $1 \leq k \leq n$. Let us take $r = x_k - x_{k-1}$ ($\varepsilon \leq r \leq 1$),

$$y'_{-1} = x_{n-1} - 1, y'_0 = 0, y'_1 = x_1, \dots, y'_{k-1} = x_{k-1},$$

$$y'_k = x_{k+1} - r, \dots, y'_{n-2} = x_{n-1} - r, y'_{n-1} = 1 - r$$

and $y_l = \frac{y'_l}{1-r}$ for $l = -1, 0, 1, \dots, n-1$.

It follows from lemma 4 that it exists positive c_3 such that

$$\begin{aligned} & v\left(\frac{x_k - x_{k-2}}{2\sqrt{(x_k - x_{k-1})(x_{k-1} - x_{k-2})}}\right) v\left(\frac{x_{k+1} - x_{k-1}}{2\sqrt{(x_{k+1} - x_k)(x_k - x_{k-1})}}\right) \leq \\ & \leq c_3 v\left(\frac{x_{k+1} - x_k + x_{k-1} - x_{k-2}}{2\sqrt{(x_{k+1} - x_k)(x_{k-1} - x_{k-2})}}\right) = c_3 v\left(\frac{y'_k - y'_{k-2}}{2\sqrt{(y'_k - y'_{k-1})(y'_{k-1} - y'_{k-2})}}\right) \end{aligned}$$

Hence,

$$\begin{aligned} I_k &= \int_0^1 \int_{x_1}^1 \dots \int_{x_{n-2}}^1 (1 - \vartheta(\frac{1}{\varepsilon} (x_k - x_{k-1}))) u_n(x_1, x_2, \dots, x_{n-1}) dx_1 dx_2 \dots dx_{n-1} \leq \\ & \leq \frac{c_3}{J_n} \int_{\varepsilon}^1 \frac{1}{r} \left[\int_0^{1-r} \int_{y'_1}^{1-r-r} \dots \int_{y'_{n-3}}^{1-r} u_{1,n-1}(y'_1, y'_2, \dots, y'_{n-1}) dy'_1 dy'_2 \dots dy'_{n-2} \right] dr = \\ & = \frac{c_3}{J_n} \int_{\varepsilon}^1 \frac{(1-r) dr}{r} \cdot \int_0^1 \int_{y_1}^1 \dots \int_{y_{n-3}}^1 u_{1,n-1}(y_1, y_2, \dots, y_{n-1}) dy_1 dy_2 \dots dy_{n-2} \leq \\ & \leq |\ln \varepsilon| \frac{c_3 J_{n-1}}{J_n} \leq |\ln \varepsilon| \frac{c_3 c_2}{16c_1(2n-1)n}. \end{aligned}$$

Thus,

$$0 \leq \int_0^1 \int_{x_1}^1 \dots \int_{x_{n-2}}^1 (1 - \vartheta(\frac{1}{\varepsilon} \max_{1 \leq k \leq n} (x_k - x_{k-1}))) u_n(x_1, x_2, \dots, x_{n-1}) dx_1 dx_2 \dots dx_{n-1} \leq$$

$$\leq n |\ln \varepsilon| \frac{c_3 c_2}{16c_1(2n-1)n} = |\ln \varepsilon| \frac{c_3 c_2}{16c_1(2n-1)},$$

which implies the assertion of lemma 5.

Lemma 6. *The following equality is fulfilled*

$$\lim_{n \rightarrow \infty} \left(\int_0^1 \int_{x_1}^1 \dots \int_{x_{n-2}}^1 \vartheta \left(\frac{1}{r} \min_{1 \leq k \leq n} \frac{x_k - x_{k-2}}{2\sqrt{(x_k - x_{k-1})(x_{k-1} - x_{k-2})}} \right) u_n(x_1, x_2, \dots, x_{n-1}) dx_1 dx_2 \dots dx_{n-1} \right) = 0$$

for any $r > 1$.

Proof. We have

$$\begin{aligned} & \int_0^1 \int_{x_1}^1 \dots \int_{x_{n-2}}^1 \vartheta \left(\frac{1}{r} \min_{1 \leq k \leq n} \frac{x_k - x_{k-2}}{2\sqrt{(x_k - x_{k-1})(x_{k-1} - x_{k-2})}} \right) \\ & \quad u_n(x_1, x_2, \dots, x_{n-1}) dx_1 dx_2 \dots dx_{n-1} \leq \\ & \leq \sum_{k=1}^n \int_0^1 \int_{x_1}^1 \dots \int_{x_{n-2}}^1 \vartheta \left(\frac{1}{r} \frac{x_k - x_{k-2}}{2\sqrt{(x_k - x_{k-1})(x_{k-1} - x_{k-2})}} \right) \\ & \quad u_n(x_1, x_2, \dots, x_{n-1}) dx_1 dx_2 \dots dx_{n-1}. \end{aligned}$$

Let $1 \leq k \leq n$. Making the same substitutions $y_l = \frac{x_l - x_{l-1}}{1 - x_{n-1}}$ и $t_{2l} = \frac{1}{2} \ln(y_l)$ as at lemma 3 we receive

$$\begin{aligned} I_k &= \int_0^1 \int_{x_1}^1 \dots \int_{x_{n-2}}^1 \vartheta \left(\frac{1}{r} \frac{x_k - x_{k-2}}{2\sqrt{(x_k - x_{k-1})(x_{k-1} - x_{k-2})}} \right) \\ & \quad u_n(x_1, x_2, \dots, x_{n-1}) dx_1 dx_2 \dots dx_{n-1} = \\ & = \frac{1}{J_n} \int_{-\infty}^{+\infty} \dots \int_{-\infty}^{+\infty} \vartheta \left(\frac{1}{r} \frac{e^{2t_{2k}} + e^{2t_{2(k-1)}}}{2e^{t_{2k}} e^{t_{2(k-1)}}} \right) \\ & \quad dt_1 dt_2 \dots dt_{2n-1} \\ & \quad \frac{1}{\sqrt{(1+t_1^2)(1+(t_2-t_1)^2) \dots (1+(t_{2n-1}-t_{2n-2})^2)(1+t_{2n-1}^2)}}. \end{aligned}$$

The inequality $\frac{e^{2t_{2k}} + e^{2t_{2(k-1)}}}{2e^{t_{2k}} e^{t_{2(k-1)}}} \leq r$ is valid if and only if

$$|t_{2k} - t_{2k-2}| \leq a = \ln(r + \sqrt{r^2 - 1}).$$

Hence, $\vartheta \left(\frac{1}{r} \frac{e^{2t_{2k}} + e^{2t_{2(k-1)}}}{2e^{t_{2k}} e^{t_{2(k-1)}}} \right) = \vartheta \left(\frac{1}{a} |t_{2k} - t_{2k-2}| \right)$ и

$$I_k = \frac{2^{n-1}}{J_n} \int_{-\infty}^{+\infty} \dots \int_{-\infty}^{+\infty} \frac{\vartheta \left(\frac{1}{a} |t_{2k} - t_{2k-2}| \right) dt_1 dt_2 \dots dt_{2n-1}}{\sqrt{(1+t_1^2)(1+(t_2-t_1)^2) \dots (1+(t_{2n-1}-t_{2n-2})^2)(1+t_{2n-1}^2)}}.$$

As $|t_{2k} - t_{2k-2}| \leq a$ we have $|t_{2k} - t_{2k-3}| \leq |t_{2k-2} - t_{2k-3}| + a$. Thus

$$1 + (t_{2k} - t_{2k-3})^2 \leq (1 + (t_{2k-2} - t_{2k-3})^2) 4(1+a)^2$$

and

$$\frac{1}{1 + (t_{2k-2} - t_{2k-3})^2} \leq \frac{4(1+a)^2}{1 + (t_{2k} - t_{2k-3})^2}.$$

Taking into account

$$\int_{-\infty}^{+\infty} \frac{dt_{2k-1}}{\sqrt{(1+(t_{2k}-t_{2k-1})^2)(1+(t_{2k-1}-t_{2k-2})^2)}} = v_1(|t_{2k}-t_{2k-2}|) \leq \pi$$

we receive

$$\begin{aligned} I_k &\leq \frac{\pi 2^n (1+a)}{J_n} \int_{-\infty}^{+\infty} \dots \int_{-\infty}^{+\infty} \int_{-\infty}^{+\infty} \dots \int_{-\infty}^{+\infty} \left(\int_{t_{2k-a}}^{t_{2k+a}} dt_{2k-2} \right) \\ &\quad \frac{dt_1 dt_2 \dots dt_{2k-3} dt_{2k} \dots dt_{2n-1}}{\sqrt{(1+t_1^2)(1+(t_2-t_1)^2) \dots (1+(t_{2k-3}-t_{2k})^2) \dots (1+(t_{2n-1}-t_{2n-2})^2)(1+t_{2n-1}^2)}} = \\ &= \frac{8\pi a(1+a)J_{n-1}}{J_n} \leq \frac{c_2 \pi a(1+a)}{c_1(2n-1)n}. \end{aligned}$$

Hence

$$\begin{aligned} 0 &\leq \int_0^1 \int_{x_1}^1 \dots \int_{x_{n-2}}^1 \vartheta \left(\frac{1}{r} \min_{1 \leq k \leq n} \frac{x_k - x_{k-2}}{2\sqrt{(x_k - x_{k-1})(x_{k-1} - x_{k-2})}} \right) \\ &\quad u_n(x_1, x_2, \dots, x_{n-1}) dx_1 dx_2 \dots dx_{n-1} \leq \\ &\leq n \frac{c_2 \pi a(1+a)}{c_1(2n-1)n} = \frac{c_2 \pi a(1+a)}{c_1(2n-1)}, \end{aligned}$$

which implies the assertion of Lemma 6.

Define the mapping $A : \text{Diff}_+^1([0, 1]) \rightarrow C_0([0, 1])$ by setting

$$A(q)(t) = \ln(f'(t)) - \ln(q'(0)) \quad \forall t \in [0, 1].$$

The mapping A is a topological isomorphism between the space $\text{Diff}_+^1([0, 1])$, $C_0([0, 1])$ moreover

$$A^{-1}(\xi)(t) = \frac{\int_0^t e^{\xi(\tau)} d\tau}{\int_0^1 e^{\xi(\tau)} d\tau}.$$

Introduce the Wiener measure w on the space $C_0([0, 1])$. Define a Borel measure ν on $\text{Diff}_+^1([0, 1])$ by setting $\nu(X) = w(A(X))$ for any Borel subset X of topological space $\text{Diff}_+^1([0, 1])$.

Let $\delta \in (0, \frac{1}{2})$. It follows from the properties of Wiener measure w (see [5]) that measure ν is concentrated on the set $E_\delta = \text{Diff}_+^{1,\delta}([0, 1])$, i.d. $\nu(E_\delta) = 1$, moreover the Borel subsets of metric space E_δ is measurable with respect to the measure ν .

As was proved in [3], the measure ν is quasi-invariant with respect to the left action of subgroup $\text{Diff}_+^3([0, 1])$ on the group $\text{Diff}_+^1([0, 1])$, moreover

$$\nu(gX) = \frac{1}{\sqrt{g'(0)g'(1)}} \int_X e^{\frac{g''(0)}{g'(0)} f'(0) - \frac{g''(1)}{g'(1)} f'(1) + \int_0^1 S_g(q(t))(q'(t))^2 dt} \nu(dq),$$

for any Borel subset X of topological space $\text{Diff}_+^1([0, 1])$, and any $g \in \text{Diff}_+^3([0, 1])$, where $gX = \{g \circ q : q \in X\}$ and $S_g(\tau) = \frac{g'''(\tau)}{g'(\tau)} - \frac{3}{2} \left(\frac{g''(\tau)}{g'(\tau)} \right)^2$ (Schwartz derivative of function g).

Lemma 7. *The following equality is valid*

$$\int_{E_\delta} (q'(0))^l \nu(dq) = \int_{E_\delta} (q'(1))^l \nu(dq)$$

for any natural l .

Proof. Let $\xi = A(q)$, i.d. $\xi(t) = \ln(f'(t)) - \ln(q'(0))$. Then

$$q'(0) = \frac{1}{\int_0^1 e^{\xi(\tau)} d\tau}, \quad q'(1) = \frac{e^{\xi(1)}}{\int_0^1 e^{\xi(\tau)} d\tau}.$$

Let us take

$$\begin{aligned} M_l &= \int_{E_\delta} (q'(1))^l \nu(dq) = \int_{\text{Diff}_+^1([0,1])} (q'(1))^l \nu(dq) = \\ &= \int_{C_0([0,1])} \left(\frac{e^{\xi(1)}}{\int_0^1 e^{\xi(\tau)} d\tau} \right)^l \nu(dq) = \int_{C_0([0,1])} \left(\frac{1}{\int_0^1 e^{\xi(1-\tau) - \xi(1)} d\tau} \right)^l \nu(dq) \end{aligned}$$

Let $\zeta(t) = \xi(1-t) - \xi(1)$. The Wiener measure w is invariant with respect to the action $\zeta \mapsto \xi$, thus,

$$\begin{aligned} M_l &= \int_{C_0([0,1])} \left(\frac{1}{\int_0^1 e^{\zeta(\tau)} d\tau} \right)^l \nu(dq) = \\ &= \int_{\text{Diff}_+^1([0,1])} (q'(0))^l \nu(dq) = \int_{E_\delta} (q'(0))^l \nu(dq), \end{aligned}$$

which implies the assertion of Lemma 7.

Introduce the measure $\nu_n = \nu \otimes \dots \otimes \nu$ on the space $E_\delta^n = E_\delta \times \dots \times E_\delta$.

Let $c_4 = 1 + M_1 + M_2 + \int_{E_1} \int_0^1 (q'(t))^2 dt \nu(dq)$.

For any $r > 0$, $g \in \text{Diff}_+^3([0,1])$, $\bar{x} = (x_1, \dots, x_{n-1}) \in D_n$, we write

$$C_g = 1 + \max_{0 \leq t \leq 1} \left(\left| \frac{g''(t)}{g'(t)} \right| + \left(\frac{g''(t)}{g'(t)} \right)^2 + \left| \frac{g'''(t)}{g'(t)} \right| \right) \mathfrak{H}$$

$$X_{r,g,\bar{x}} = \{(q_1, \dots, q_n) : q_1, \dots, q_n \in E_\delta\}$$

$$\begin{aligned} & \left| \sum_{k=1}^n [(x_k - x_{k-1}) \left(\frac{g''(x_{k-1})}{g'(x_{k-1})} q'_k(0) - \frac{g''(x_k)}{g'(x_k)} q'_k(1) \right) + \right. \\ & \left. + (x_k - x_{k-1})^2 \int_0^1 S_g(x_{k-1} + (x_k - x_{k-1})q_k(t)) (q'_k(t))^2 dt] \right| \leq 4c_4 C_g r \}. \end{aligned}$$

Lemma 8. *If $\epsilon \in (0,1)$, then the following inequality is fulfilled*

$\nu_n(E_\delta^n \setminus X_{\sqrt[3]{\epsilon}, g, \bar{x}}) \leq 2\sqrt[3]{\epsilon}$ for any $g \in \text{Diff}_+^3([0,1])$, for any positive integer n and $\bar{x} = (x_1, \dots, x_{n-1}) \in D_n$, satisfying the inequality

$$\max_{1 \leq k \leq n} (x_k - x_{k-1}) < \epsilon.$$

Proof. Let

$$f_1(q_1, \dots, q_n) = \sum_{k=1}^n (x_k - x_{k-1}) \left(\frac{g''(x_{k-1})}{g'(x_{k-1})} q'_k(0) - \frac{g''(x_k)}{g'(x_k)} q'_k(1) \right).$$

Then

$$I_1 = \int_{E_\delta} \dots \int_{E_\delta} f_1(q_1, \dots, q_n) \nu(dq_1) \dots \nu(dq_n) = M_1 \sum_{k=1}^n (x_k - x_{k-1}) \left(\frac{g''(x_{k-1})}{g'(x_{k-1})} - \frac{g''(x_k)}{g'(x_k)} \right).$$

As $\left| \frac{g''(x_{k-1})}{g'(x_{k-1})} - \frac{g''(x_k)}{g'(x_k)} \right| \leq C_g (x_k - x_{k-1})$, we have

$$|I_1| \leq M_1 C_g \sum_{k=1}^n (x_k - x_{k-1})^2 \leq M_1 C_g \epsilon \sum_{k=1}^n (x_k - x_{k-1}) = M_1 C_g \epsilon.$$

If $k \neq l$ then

$$\begin{aligned} & \int_{E_\delta} \int_{E_\delta} \left(\frac{g''(x_{k-1})}{g'(x_{k-1})} (q'_k(0) - M_1) - \frac{g''(x_k)}{g'(x_k)} (q'_k(1) - M_1) \right) \\ & \left(\frac{g''(x_{l-1})}{g'(x_{l-1})} (q'_l(0) - M_1) - \frac{g''(x_l)}{g'(x_l)} (q'_l(1) - M_1) \right) \nu(dq_k) \nu(dq_l) = 0, \end{aligned}$$

therefore

$$\begin{aligned} I_2 &= \int_{E_\delta} \dots \int_{E_\delta} (f_1(q_1, \dots, q_n) - I_1)^2 \nu(dq_1) \dots \nu(dq_n) = \\ & \sum_{k=1}^n (x_k - x_{k-1})^2 \int_{E_\delta} \left[\frac{g''(x_{k-1})}{g'(x_{k-1})} (q'_k(0) - M_1) - \frac{g''(x_k)}{g'(x_k)} (q'_k(1) - M_1) \right]^2 \nu(dq_k) \leq \\ & \leq 2 \sum_{k=1}^n (x_k - x_{k-1})^2 \left[\left(\frac{g''(x_{k-1})}{g'(x_{k-1})} \right)^2 \int_{E_\delta} (q'_k(0) - M_1)^2 \nu(dq_k) + \right. \\ & \quad \left. + \left(\frac{g''(x_k)}{g'(x_k)} \right)^2 \int_{E_\delta} (q'_k(1) - M_1)^2 \nu(dq_k) \right] = \\ & = 2 \sum_{k=1}^n (x_k - x_{k-1})^2 \left[\left(\frac{g''(x_{k-1})}{g'(x_{k-1})} \right)^2 + \left(\frac{g''(x_k)}{g'(x_k)} \right)^2 \right] \left[\int_{E_\delta} (q'_k(0))^2 \nu(dq_k) - (M_1)^2 \right] \leq \\ & \leq 4M_2 C_g \sum_{k=1}^n (x_k - x_{k-1})^2 \leq 4M_2 C_g \epsilon \sum_{k=1}^n (x_k - x_{k-1}) = 4M_2 C_g \epsilon. \end{aligned}$$

Hence,

$$\nu_n(\{(q_1, \dots, q_n) : |f_1(q_1, \dots, q_n) - I_1| \geq 2c_4 C_g \sqrt[3]{\epsilon}\}) \leq \frac{I_2}{(2c_4 C_g \sqrt[3]{\epsilon})^2} \leq \frac{4M_2 C_g \epsilon}{(2c_4 C_g \sqrt[3]{\epsilon})^2} \leq \sqrt[3]{\epsilon}.$$

Thus

$$\nu_n(\{(q_1, \dots, q_n) : |f_1(q_1, \dots, q_n)| \geq 3c_4 C_g \sqrt[3]{\epsilon}\}) \leq \sqrt[3]{\epsilon}.$$

Let

$$f_2(q_1, \dots, q_n) = \sum_{k=1}^n ((x_k - x_{k-1})^2 \int_0^1 S_g(x_{k-1} + (x_k - x_{k-1})q_k(t)) (q'_k(t))^2 dt).$$

Then

$$I_3 = \int_{E_\delta} \dots \int_{E_\delta} |f_2(q_1, \dots, q_n)| \nu(dq_1) \dots \nu(dq_n) \leq$$

$$\begin{aligned}
&\leq 2C_g \sum_{k=1}^n (x_k - x_{k-1})^2 \int_{E_\delta} \left(\int_0^1 (q'_k(t))^2 dt \right) \nu(dq_k) \leq \\
&\leq 2c_4 C_g \epsilon \sum_{k=1}^n (x_k - x_{k-1}) = 2c_4 C_g \epsilon.
\end{aligned}$$

Thus

$$\nu_n(\{(q_1, \dots, q_n) : |f_2(q_1, \dots, q_n)| \geq 2c_4 C_g \sqrt[3]{\epsilon}\}) \leq \frac{I_3}{2c_4 C_g \sqrt[3]{\epsilon}} \leq \frac{2c_4 C_g \epsilon}{2c_4 C_g \sqrt[3]{\epsilon}} = (\sqrt[3]{\epsilon})^2 \leq \sqrt[3]{\epsilon}.$$

Hence,

$$\begin{aligned}
&\nu_n(E_\delta^n \setminus X_{\sqrt[3]{\epsilon}, g, \bar{x}}) = \\
&= \nu_n(\{(q_1, \dots, q_n) : |f_1(q_1, \dots, q_n) + f_2(q_1, \dots, q_n)| \geq 4c_4 C_g \sqrt[3]{\epsilon}\}) \leq \\
&\leq \nu_n(\{(q_1, \dots, q_n) : |f_1(q_1, \dots, q_n)| \geq 2c_4 C_g \sqrt[3]{\epsilon}\}) + \\
&+ \nu_n(\{(q_1, \dots, q_n) : |f_2(q_1, \dots, q_n)| \geq 2c_4 C_g \sqrt[3]{\epsilon}\}) \leq 2\sqrt[3]{\epsilon},
\end{aligned}$$

which implies the assertion of Lemma 8.

Lemma 9. For any $g \in \text{Diff}_0^3([0, 1])$, $\epsilon > 0$, there are $r > 1$ and $\delta_1 \in (0, 1)$ such that the inequality is valid

$$\left| \prod_{k=1}^n \frac{v\left(\frac{g(x_k) - g(x_{k-2})}{2\sqrt{(g(x_k) - g(x_{k-1})) (g(x_{k-1}) - g(x_{k-2})))} }\right)}{v\left(\frac{x_k - x_{k-2}}{2\sqrt{(x_k - x_{k-1})(x_{k-1} - x_{k-2})}}\right)} - 1 \right| \leq \epsilon,$$

$g(x_{-1}) = g(x_{n-1}) - 1$ for any natural n and any $\bar{x} = (x_1, \dots, x_{n-1}) \in D_n$ satisfying the inequality $\max_{1 \leq k \leq n} (x_k - x_{k-1}) < \delta_1$, $\min_{1 \leq k \leq n} \left(\frac{x_k - x_{k-2}}{2\sqrt{(x_k - x_{k-1})(x_{k-1} - x_{k-2})}} \right) > r$, where $x_0 = 0$, $x_n = 1$, $x_{-1} = x_{n-1} - 1$.

Proof. Let $\epsilon \in (0, 1)$. Let $C = \max_{t_1, t_2 \in [0, 1]} |g''(t_1)|$, $\delta_1 = \frac{1}{400(C+1)}$ и $r = e^{\frac{8000(C+1)}{\epsilon}}$.

For any k ($1 \leq k \leq n$), there are x_{k-1}^* , $x_{k-1}^{**} \in (0, 1)$ such that

$$\begin{aligned}
g(x_k) - g(x_{k-1}) &= g'(x_{k-1})(x_k - x_{k-1}) + \frac{1}{2}g''(x_{k-1}^*)(x_k - x_{k-1})^2 = \\
&= g'(x_{k-1})(x_k - x_{k-1}) \left(1 + \frac{g''(x_{k-1}^*)}{2g'(x_{k-1})}(x_k - x_{k-1})\right), \\
g(x_{k-1}) - g(x_{k-2}) &= g'(x_{k-1})(x_{k-1} - x_{k-2}) + \frac{1}{2}g''(x_{k-1}^{**})(x_{k-1} - x_{k-2})^2 \\
&= g'(x_{k-1})(x_{k-1} - x_{k-2}) \left(1 + \frac{g''(x_{k-1}^{**})}{2g'(x_{k-1})}(x_{k-1} - x_{k-2})\right),
\end{aligned}$$

for $k \geq 2$ and for $k = 1$

$$\begin{aligned}
g(x_1) - g(x_0) &= g'(x_0)(x_1 - x_0) + \frac{1}{2}g''(x_0^*)(x_1 - x_0)^2 = \\
&= g'(x_0)(x_1 - x_0) \left(1 + \frac{g''(x_0^*)}{2g'(x_0)}(x_1 - x_0)\right),
\end{aligned}$$

$$\begin{aligned}
g(x_0) - g(x_{-1}) &= g(1) - g(x_{n-1}) = g'(1)(1 - x_{n-1}) + \frac{1}{2}g''(x_0^{**})(1 - x_{n-1})^2 = \\
&= g'(x_0)(x_0 - x_{-1}) \left(1 + \frac{g''(x_0^{**})}{2g'(x_0)}(x_0 - x_{-1})\right).
\end{aligned}$$

Hence

$$\begin{aligned} & \frac{g(x_k) - g(x_{k-2})}{2\sqrt{(g(x_k) - g(x_{k-1}))(g(x_{k-1}) - g(x_{k-2}))}} = \\ & = \frac{x_k - x_{k-2}}{2\sqrt{(x_k - x_{k-1})(x_{k-1} - x_{k-2})}}(1 + \lambda_k(x_k - x_{k-2})), \end{aligned}$$

where

$$\lambda_k = \frac{1}{x_k - x_{k-2}} \left(\frac{1 + \frac{g''(x_{k-1}^*)(x_k - x_{k-1})^2 + g''(x_{k-1}^*)(x_{k-1} - x_{k-2})^2}{2g'(x_{k-1})(x_k - x_{k-2})}}{\sqrt{(1 + \frac{g''(x_{k-1}^*)}{2g'(x_{k-1})}(x_k - x_{k-1})) (1 + \frac{g''(x_{k-1}^*)}{2g'(x_{k-1})}(x_{k-1} - x_{k-2}))}} - 1 \right).$$

As $C(x_k - x_{k-2}) < C\delta_1 = \frac{1}{400}$, then $|\lambda_k| < 4C$.

Let $t = \frac{x_k - x_{k-2}}{2\sqrt{(x_k - x_{k-1})(x_{k-1} - x_{k-2})}}$ and $\alpha = \lambda_k(x_k - x_{k-2})$. Then

$$t(1 + \alpha) = \frac{g(x_k) - g(x_{k-2})}{2\sqrt{(g(x_k) - g(x_{k-1}))(g(x_{k-1}) - g(x_{k-2}))}}$$

and $|\alpha| < \frac{1}{80}$, $t > 4$.

Let us obtain $\tau = \ln(t + \sqrt{t^2 - 1}) > \ln t > 1$ and

$$\beta = \ln\left[1 + \alpha\left(\frac{t}{t + \sqrt{t^2 - 1}}\right)\left(1 + \frac{(2 + \alpha)t}{\sqrt{t^2 - 1} + \sqrt{t^2(1 + \alpha)^2 - 1}}\right)\right].$$

Then $\frac{1}{4}|\alpha| \leq |\beta| \leq 4|\alpha|$, $|\beta| \leq \frac{1}{20}$ and

$$\begin{aligned} & \ln(t(1 + \alpha) + \sqrt{t^2(1 + \alpha)^2 - 1}) = \ln(t + \sqrt{t^2 - 1}) + \\ & \ln\left[1 + \alpha\left(\frac{t}{t + \sqrt{t^2 - 1}}\right)\left(1 + \frac{(2 + \alpha)t}{\sqrt{t^2 - 1} + \sqrt{t^2(1 + \alpha)^2 - 1}}\right)\right] = \tau + \beta. \end{aligned}$$

There exists $\theta \in (0, 1)$ such that $v(t(1 + \alpha)) = v_1(\tau + \beta) = v_1(\tau) + v_1'(\tau + \theta\beta)\beta$.

It follows from the lemma 2 that $|v_1'(\tau + \theta\beta)| \leq \frac{4}{\tau + \theta\beta}v_1(\tau + \theta\beta)$. As $|\theta\beta| \leq \frac{1}{20}$, we have $|v_1'(\tau + \theta\beta)| \leq \frac{10}{\tau}v_1(\tau)$.

Thus it exists $\omega(\alpha, t)$ such that $|\omega(\alpha, t)| < 10$ и $\frac{v_1(\tau + \beta)}{v_1(\tau)} = 1 + \omega(\alpha, t)\frac{\beta}{\tau}$. Taking account $\frac{1}{4}|\lambda_k(x_k - x_{k-2})| = \frac{1}{4}|\alpha| \leq |\beta| \leq 4|\alpha| = 4|\lambda_k(x_k - x_{k-2})|$, we receive

$$\begin{aligned} & \frac{v\left(\frac{g(x_k) - g(x_{k-2})}{2\sqrt{(g(x_k) - g(x_{k-1}))(g(x_{k-1}) - g(x_{k-2}))}}\right)}{v\left(\frac{x_k - x_{k-2}}{2\sqrt{(x_k - x_{k-1})(x_{k-1} - x_{k-2})}}\right)} = \\ & = \frac{v_1(\tau + \beta)}{v_1(\tau)} = 1 + \omega_k \frac{x_k - x_{k-2}}{\ln \frac{x_k - x_{k-2}}{2\sqrt{(x_k - x_{k-1})(x_{k-1} - x_{k-2})}}}, \end{aligned}$$

where $\omega_k = \omega(\alpha, t)\frac{\beta \ln t}{\tau(x_k - x_{k-2})}$ и $|\omega_k| \leq 200C$.

Hence

$$\begin{aligned} \sigma & = \ln\left(\prod_{k=1}^n \frac{v\left(\frac{g(x_k) - g(x_{k-2})}{2\sqrt{(g(x_k) - g(x_{k-1}))(g(x_{k-1}) - g(x_{k-2}))}}\right)}{v\left(\frac{x_k - x_{k-2}}{2\sqrt{(x_k - x_{k-1})(x_{k-1} - x_{k-2})}}\right)}\right) = \\ & = \sum_{k=1}^n \ln\left(1 + \omega_k \frac{x_k - x_{k-2}}{\ln \frac{x_k - x_{k-2}}{2\sqrt{(x_k - x_{k-1})(x_{k-1} - x_{k-2})}}}\right) \end{aligned}$$

and

$$|\sigma| \leq 2 \sum_{k=1}^n |\omega_k| \frac{x_k - x_{k-2}}{\ln \frac{x_k - x_{k-2}}{2\sqrt{(x_k - x_{k-1})(x_{k-1} - x_{k-2})}}} \leq \frac{400C}{\ln r} \sum_{k=1}^n (x_k - x_{k-2}) = \frac{800C}{\ln r} \leq \frac{\epsilon}{10},$$

therefore

$$\begin{aligned} & \left| \prod_{k=1}^n \frac{v\left(\frac{g(x_k) - g(x_{k-2})}{2\sqrt{(g(x_k) - g(x_{k-1}))(g(x_{k-1}) - g(x_{k-2}))}}\right)}{v\left(\frac{x_k - x_{k-2}}{2\sqrt{(x_k - x_{k-1})(x_{k-1} - x_{k-2})}}\right)} - 1 \right| = \\ & = |e^\sigma - 1| \leq e^{\frac{\epsilon}{10}} - e^{-\frac{\epsilon}{10}} \leq \frac{\epsilon}{5} + \frac{\epsilon}{5} < \epsilon, \end{aligned}$$

which implies the assertion of Lemma 9.

Introduce the mapping $Q_n : D_n \times E_\delta^n \rightarrow E_\delta = \text{Diff}_+^{1,\delta}([0, 1])$ by setting $f_n \circ (l_n)^{-1} = Q_n(x_1, \dots, x_{n-1}, \varphi_1, \dots, \varphi_n)$, where

$$\begin{aligned} f_n(t) &= x_{k-1} + (x_k - x_{k-1})\varphi_k\left(n\left(t - \frac{k-1}{n}\right)\right), \\ l_n(t) &= \frac{1}{x_1 - x_0 + \sum_{m=2}^n (x_m - x_{m-1}) \frac{\varphi'_2(0)\varphi'_3(0)\dots\varphi'_m(0)}{\varphi'_1(1)\varphi'_2(1)\dots\varphi'_{m-1}(1)}} \cdot \\ & \cdot \left(x_1 - x_0 + \sum_{m=2}^{k-1} (x_m - x_{m-1}) \frac{\varphi'_2(0)\varphi'_3(0)\dots\varphi'_m(0)}{\varphi'_1(1)\varphi'_2(1)\dots\varphi'_{m-1}(1)} + \right. \\ & \left. + (x_k - x_{k-1}) \frac{\varphi'_2(0)\varphi'_3(0)\dots\varphi'_k(0)}{\varphi'_1(1)\varphi'_2(1)\dots\varphi'_{k-1}(1)} n\left(t - \frac{k-1}{n}\right)\right) \end{aligned}$$

for $t \in [\frac{k-1}{n}, \frac{k}{n}]$, $(x_1, \dots, x_{n-1}) \in D_n$, $(\varphi_1, \dots, \varphi_n) \in E_\delta^n$.

The function $f = f_n \circ (l_n)^{-1}$ belongs to $\text{Diff}_+^{1,\delta}([0, 1])$, because the left derivation

$$\begin{aligned} f'((l_n)^{-1}\left(\frac{k-1}{n} - 0\right)) &= n(x_{k-1} - x_{k-2})\varphi_{k-1}(1) \cdot \\ & \cdot \frac{x_1 - x_0 + \sum_{m=2}^n (x_m - x_{m-1}) \frac{\varphi'_2(0)\varphi'_3(0)\dots\varphi'_m(0)}{\varphi'_1(1)\varphi'_2(1)\dots\varphi'_{m-1}(1)}}{(x_{k-1} - x_{k-2}) \frac{\varphi'_2(0)\varphi'_3(0)\dots\varphi'_{k-1}(0)}{\varphi'_1(1)\varphi'_2(1)\dots\varphi'_{k-2}(1)} n} = \\ & = (x_1 - x_0 + \sum_{m=2}^n (x_m - x_{m-1}) \frac{\varphi'_2(0)\varphi'_3(0)\dots\varphi'_m(0)}{\varphi'_1(1)\varphi'_2(1)\dots\varphi'_{m-1}(1)}) \frac{\varphi'_1(1)\varphi'_2(1)\dots\varphi'_{k-1}(1)}{\varphi'_2(0)\varphi'_3(0)\dots\varphi'_{k-1}(0)}. \end{aligned}$$

is equal to the right derivation

$$\begin{aligned} f'((l_n)^{-1}\left(\frac{k-1}{n} + 0\right)) &= n(x_k - x_{k-1})\varphi_k(0) \cdot \\ & \cdot \frac{x_1 - x_0 + \sum_{m=2}^n (x_m - x_{m-1}) \frac{\varphi'_2(0)\varphi'_3(0)\dots\varphi'_m(0)}{\varphi'_1(1)\varphi'_2(1)\dots\varphi'_{m-1}(1)}}{(x_k - x_{k-1}) \frac{\varphi'_2(0)\varphi'_3(0)\dots\varphi'_k(0)}{\varphi'_1(1)\varphi'_2(1)\dots\varphi'_{k-1}(1)} n} = \\ & = (x_1 - x_0 + \sum_{m=2}^n (x_m - x_{m-1}) \frac{\varphi'_2(0)\varphi'_3(0)\dots\varphi'_m(0)}{\varphi'_1(1)\varphi'_2(1)\dots\varphi'_{m-1}(1)}) \frac{\varphi'_1(1)\varphi'_2(1)\dots\varphi'_{k-1}(1)}{\varphi'_2(0)\varphi'_3(0)\dots\varphi'_{k-1}(0)}. \end{aligned}$$

Write

$$L_{\delta,n}(F) = \int_0^1 \int_{x_1}^1 \dots \int_{x_{n-2}}^1 \int_{E_\delta} \dots \int_{E_\delta} F(Q_n(x_1, x_2, \dots, x_{n-1}, \varphi_1, \dots, \varphi_n)) \cdot u(x_1, x_2, \dots, x_{n-1}) dx_1 dx_2 \dots dx_{n-1} \nu(d\varphi_1) \dots \nu(d\varphi_n)$$

for any function $F \in C_b(E_\delta) = C_b(\text{Diff}_+^{1,\delta}([0, 1]))$.

Theorem 3. *a function F belongs to the space $C_b(\text{Diff}_+^{1,\delta}([0, 1]))$, let a diffeomorphism g belongs to the group $\text{Diff}_0^3([0, 1])$.*

Then $\lim_{n \rightarrow \infty} |L_{\delta,n}(F_g) - L_{\delta,n}(F)| = 0$.

Proof. Let $F \in C_b(\text{Diff}_+^{1,\delta}([0, 1]))$, $C = \sup_{g \in E_\delta} |F(g)|$.

Let $\epsilon \in (0, 1)$ and $g \in \text{Diff}_0^3([0, 1])$. Let us take positive ϵ_1 satisfying the inequalities $\epsilon_1 < \frac{1}{8}\epsilon$, $e^{4c_5 C_g \sqrt[3]{\epsilon_1}} - e^{-4c_4 C_g \sqrt[3]{\epsilon_1}} < \epsilon$.

It follows from the lemma 9 that there are $r > 1$ and $\delta_1 \in (0, 1)$ such that

$$\left| \prod_{k=1}^n \frac{v\left(\frac{g(x_k) - g(x_{k-2})}{2\sqrt{(g(x_k) - g(x_{k-1})) (g(x_{k-1}) - g(x_{k-2})))}\right)}{v\left(\frac{x_k - x_{k-2}}{2\sqrt{(x_k - x_{k-1})(x_{k-1} - x_{k-2})}}\right)} - 1 \right| \leq \epsilon$$

for any positive integer n and for $\bar{x} = (x_1, \dots, x_{n-1}) \in D_n$ satisfying the inequalities

$$\max_{1 \leq k \leq n} (x_k - x_{k-1}) < \delta_1, \quad \min_{1 \leq k \leq n} \left(\frac{x_k - x_{k-2}}{2\sqrt{(x_k - x_{k-1})(x_{k-1} - x_{k-2})}} \right) > r.$$

It follows from the lemma 8 that the inequality is valid $\nu_n(E_\delta^n \setminus X_{\sqrt[3]{\epsilon_1}, g, \bar{x}}) \leq 2\sqrt[3]{\epsilon_1} \leq \epsilon$ for any positive integer n and for any $\bar{x} = (x_1, \dots, x_{n-1}) \in D_n$ satisfying the inequalities $\max_{1 \leq k \leq n} (x_k - x_{k-1}) < \epsilon_1$.

It follows from the lemma 5, 6 that there exists a positive integer N such that

$$\int_0^1 \int_{x_1}^1 \dots \int_{x_{n-2}}^1 (1 - \vartheta\left(\frac{1}{\min(\epsilon_1, \delta)} \max_{1 \leq k \leq n} (x_k - x_{k-1})\right)) u_n(x_1, x_2, \dots, x_{n-1}) dx_1 dx_2 \dots dx_{n-1} < \epsilon,$$

$$\int_0^1 \int_{x_1}^1 \dots \int_{x_{n-2}}^1 \vartheta\left(\frac{1}{r} \min_{1 \leq k \leq n} \frac{x_k - x_{k-2}}{2\sqrt{(x_k - x_{k-1})(x_{k-1} - x_{k-2})}}\right)$$

$$u_n(x_1, x_2, \dots, x_{n-1}) dx_1 dx_2 \dots dx_{n-1} < \epsilon$$

for any integer $n > N$.

Let

$$\phi(x_1, x_2, \dots, x_{n-1}) = \vartheta\left(\frac{1}{\min(\epsilon_1, \delta)} \max_{1 \leq k \leq n} (x_k - x_{k-1})\right)$$

$$\left(1 - \vartheta\left(\frac{1}{r} \min_{1 \leq k \leq n} \frac{x_k - x_{k-2}}{2\sqrt{(x_k - x_{k-1})(x_{k-1} - x_{k-2})}}\right)\right),$$

then

$$\int_0^1 \int_{x_1}^1 \dots \int_{x_{n-2}}^1 \phi(x_1, x_2, \dots, x_{n-1}) u_n(x_1, x_2, \dots, x_{n-1}) dx_1 dx_2 \dots dx_{n-1} \geq 1 -$$

$$\begin{aligned}
& - \int_0^1 \int_{x_1}^1 \dots \int_{x_{n-2}}^1 (1 - \vartheta(\frac{1}{\min(\epsilon_1, \delta)} \max_{1 \leq k \leq n} (x_k - x_{k-1}))) u_n(x_1, x_2, \dots, x_{n-1}) dx_1 dx_2 \dots dx_{n-1} \\
& \quad - \int_0^1 \int_{x_1}^1 \dots \int_{x_{n-2}}^1 \vartheta(\frac{1}{r} \min_{1 \leq k \leq n} \frac{x_k - x_{k-2}}{2\sqrt{(x_k - x_{k-1})(x_{k-1} - x_{k-2})}}) \\
& \quad \quad u_n(x_1, x_2, \dots, x_{n-1}) dx_1 dx_2 \dots dx_{n-1} > 1 - 2\epsilon.
\end{aligned}$$

Let $y_k = g(x_k)$, $\bar{y} = (y_1, \dots, y_{n-1}) \in D_n$,
 $g^{-1}(\bar{y}) = (x_1, \dots, x_{n-1}) \in D_n$, $\phi_g(y_1, \dots, y_{n-1}) = \phi(x_1, \dots, x_{n-1})$ and

$$gX_{\mathfrak{Z}_{\epsilon_1, g, g^{-1}}(\bar{y})} = \{(\varphi_1, \dots, \varphi_n) : (q_1, \dots, q_{n-1}) \in X_{\mathfrak{Z}_{\epsilon_1, g, \bar{x}}}\},$$

$$Q_n(y_1, \dots, y_{n-1}, \varphi_1, \dots, \varphi_n) = g \circ (Q_n(x_1, x_2, \dots, x_{n-1}, q_1, \dots, q_{n-1})).$$

It is easy to see that $\varphi_k(t) = \frac{g(x_{k-1} + (x_k - x_{k-1})q_k(nt - k + 1)) - g(x_{k-1})}{g(x_k) - g(x_{k-1})}$, because

$$\frac{(y_k - y_{k-1})\varphi'_2(0)\varphi'_3(0)\dots\varphi'_k(0)}{(y_1 - y_0)\varphi'_1(1)\varphi'_2(1)\dots\varphi'_{k-1}(1)} = \frac{(x_k - x_{k-1})q'_2(0)q'_3(0)\dots q'_k(0)}{(x_1 - x_0)q'_1(1)q'_2(1)\dots q'_{k-1}(1)}$$

We have

$$\begin{aligned}
& \int_0^1 \int_{y_1}^1 \dots \int_{y_{n-2}}^1 \phi_g(\bar{y}) u_n(\bar{y}) \nu_n(gX_{\mathfrak{Z}_{\epsilon_1, g, g^{-1}}(\bar{y})}) dy_1 dy_2 \dots dy_{n-1} = \\
& = \int_0^1 \int_{x_1}^1 \dots \int_{x_{n-2}}^1 \phi(\bar{x}) u_n(\bar{x}) \left(\int_{X_{\mathfrak{Z}_{\epsilon_1, g, \bar{x}}}} \exp\left(\sum_{k=1}^n [(x_k - x_{k-1}) \left(\frac{g''(x_{k-1})}{g'(x_{k-1})} q'_k(0) - \frac{g''(x_k)}{g'(x_k)} q'_k(1)\right)]\right) \right. \\
& \quad \left. + (x_k - x_{k-1})^2 \int_0^1 S_g(x_{k-1} + (x_k - x_{k-1})q_k(t)) (q'_k(t))^2 dt \right) \nu(dq_1) \dots \nu(dq_n) \\
& \quad \prod_{k=1}^n \frac{v\left(\frac{g(x_k) - g(x_{k-2})}{2\sqrt{(g(x_k) - g(x_{k-1}))(g(x_{k-1}) - g(x_{k-2}))}}\right)}{v\left(\frac{x_k - x_{k-2}}{2\sqrt{(x_k - x_{k-1})(x_{k-1} - x_{k-2})}}\right)} dx_1 dx_2 \dots dx_{n-1} \geq \\
& \geq (1 - \epsilon)^2 \int_0^1 \int_{x_1}^1 \dots \int_{x_{n-2}}^1 \phi(\bar{x}) u_n(\bar{x}) \nu_n(X_{\mathfrak{Z}_{\epsilon_1, g, \bar{x}}}) dx_1 dx_2 \dots dx_{n-1} \geq \\
& \geq (1 - \epsilon)^3 \int_0^1 \int_{x_1}^1 \dots \int_{x_{n-2}}^1 \phi(\bar{x}) u_n(\bar{x}) dx_1 dx_2 \dots dx_{n-1} \geq (1 - \epsilon)^3 (1 - 2\epsilon).
\end{aligned}$$

Hence,

$$\begin{aligned}
& |L_{\delta, n}(F_g) - \int_0^1 \int_{y_1}^1 \dots \int_{y_{n-2}}^1 \phi_g(\bar{y}) u_n(\bar{y}) \left(\int_{gX_{\mathfrak{Z}_{\epsilon_1, g, g^{-1}}(\bar{y})}} F_g(Q_n(\bar{y}, \varphi_1, \dots, \varphi_n)) \right. \\
& \quad \left. \nu(d\varphi_1) \dots \nu(d\varphi_n) \right) dy_1 dy_2 \dots dy_{n-1}| \leq C(1 - (1 - \epsilon)^3(1 - 2\epsilon))
\end{aligned}$$

and

$$|L_{\delta, n}(F) - \int_0^1 \int_{x_1}^1 \dots \int_{x_{n-2}}^1 \phi_g(\bar{x}) u_n(\bar{x}) \left(\int_{X_{\mathfrak{Z}_{\epsilon_1, g, \bar{x}}}} F(Q_n(\bar{x}, q_1, \dots, q_n)) \right.$$

$$\nu(dq_1)\dots\nu(dq_n)dx_1dx_2\dots dx_{n-1}| \leq C(1 - (1 - \epsilon)(1 - 2\epsilon)).$$

We have

$$\begin{aligned} & \left| \int_0^1 \int_{y_1}^1 \dots \int_{y_{n-2}}^1 \phi_g(\bar{y})u_n(\bar{y}) \left(\int_{gX_{\mathfrak{F}\epsilon_1, g, g^{-1}(\bar{y})}} F_g(Q_n(\bar{y}, \varphi_1, \dots, \varphi_n)) \right. \right. \\ & \quad \left. \left. \nu(d\varphi_1)\dots\nu(d\varphi_n)dy_1dy_2\dots dy_{n-1} - \right. \right. \\ & \quad \left. \left. - \int_0^1 \int_{x_1}^1 \dots \int_{x_{n-2}}^1 \phi_g(\bar{x})u_n(\bar{x}) \left(\int_{X_{\mathfrak{F}\epsilon_1, g, \bar{x}}} F(Q_n(\bar{x}, q_1, \dots, q_n)) \right. \right. \right. \\ & \quad \left. \left. \left. \nu(dq_1)\dots\nu(dq_n)dx_1dx_2\dots dx_{n-1} \right| \leq \right. \\ & \leq \int_0^1 \int_{x_1}^1 \dots \int_{x_{n-2}}^1 \phi(\bar{x})u_n(\bar{x}) \left(\int_{X_{\mathfrak{F}\epsilon_1, g, \bar{x}}} \left| \exp\left(\sum_{k=1}^n [(x_k - x_{k-1}) \left(\frac{g''(x_{k-1})}{g'(x_{k-1})}q'_k(0) - \frac{g''(x_k)}{g'(x_k)}q'_k(1)\right) + \right. \right. \right. \\ & \quad \left. \left. \left. + (x_k - x_{k-1})^2 \int_0^1 S_g(x_{k-1} + (x_k - x_{k-1})q_k(t))(q'_k(t))^2 dt \right] \right. \right. \\ & \quad \left. \left. \prod_{k=1}^n \frac{v\left(\frac{g(x_k) - g(x_{k-2})}{2\sqrt{(g(x_k) - g(x_{k-1}))(g(x_{k-1}) - g(x_{k-2}))}\right)}}{v\left(\frac{x_k - x_{k-2}}{2\sqrt{(x_k - x_{k-1})(x_{k-1} - x_{k-2})}\right)} \right) - 1 \right| \\ & \quad \left| F(Q_n(\bar{x}, q_1, \dots, q_n)) \right| \nu(dq_1)\dots\nu(dq_n)u_n(\bar{x})dx_1dx_2\dots dx_{n-1} \leq \\ & \leq C\epsilon(2 + \epsilon) \int_0^1 \int_{x_1}^1 \dots \int_{x_{n-2}}^1 \phi(\bar{x})u_n(\bar{x})\nu_n(X_{\mathfrak{F}\epsilon_1, g, \bar{x}})dx_1dx_2\dots dx_{n-1} \leq C\epsilon(2 + \epsilon), \end{aligned}$$

which implies the assertion of Theorem 3.

Define a ultrafilter \mathfrak{F} on the set positive integers such that \mathfrak{F} contains the sets $\{n, n+1, \dots\}$ for any positive integer n . We set $L_\delta(F) = \lim_{\mathfrak{F}} L_{\delta, n}(F)$ for any function $F \in C_b(E_\delta)$.

Note that the limit always exists because $|L_{\delta, n}(F)| \leq \sup_{f \in E_\delta} |F(f)|$.

It is easy to see that $L(e_{1, \delta}) = 1$, $|L_\delta(F)| \leq \sup_{f \in E_\delta} |F(f)|$, and $L(F) \geq 0$ for any nonnegative function $F \in B(G)$. In turn, Theorem 1 follows from Theorem 3.

Acknowledgements. The author acknowledge I.K. Babenko, R. Grigorchuk, P. de la Harpe, V.Sergiescu, O.G. Smolyanov, A.I. Stern for the discussions and interest about this work.

References

1. Cannon J.W., Floyd W.J., Parry W.R. "Introductory notes on Richard Thompson's groups Enseign Math., vol 42, issue 2 (1996), pages 215–256.
2. Ghys È., Sergiescu V. "Sur un groupe remarquable de difféomorphismes du cercle Comment.Math.Helvetici 62 (1987) 185–239.
3. Shavgulidze E.T. "Some Properties of Quasi-Invariant Measures on Groups of Diffeomorphisms of the Circle Russ. J. Math. Physics, vol 7, issue 4 (2000), pages 464–472.
4. Shavgulidze E.T. "Amenability of discrete subgroups of the group of diffeomorphisms of the circle Russ. J. Math. Physics, vol 16, issue 1 (2009), pages 138–140

5. Hui-Hsiuhg Kuo "Gaussian measures in Banach space Lecture Notes in Mathematics 463, Springer-Verlag 1975.
6. G.H.Watson "A Treatise on the Theory of Bessel Functions Cambridge 1944.