

Investigating the Disk-Corona Relation in a Blue AGN Sample

Jie-Ying Liu^{1,2} and B. F. Liu¹

¹ National Astronomical Observatories/Yunnan Observatory, Chinese Academy of Sciences, PO Box 110, Kunming 650011; ljy0807@ynao.ac.cn

² Graduate School of Chinese Academy of Sciences, Beijing 100049

Received [year] [month] [day]; accepted [year] [month] [day]

Abstract We compile a blue AGN sample from SDSS and investigate the ratio of hard X-ray to bolometric luminosity in dependence on Eddington ratio and black hole mass. Our sample comprises 240 radio-quiet Seyfert 1 galaxies and QSOs. We find that the fraction of hard X-ray luminosity ($\log(L_{2-10\text{kev}}/L_{\text{bol}})$) decreases with the increase of Eddington ratio. We also find that the fraction of hard X-ray luminosity is independent on the black hole mass for the radio-quiet AGNs. The relation of $\log(L_{2-10\text{kev}}/L_{\text{bol}})$ decreasing with increasing Eddington ratio indicates that X-ray bolometric correction is not a constant, from a larger sample supporting the results of Vasudevan & Fabian (2007). We interpret our results by the disk corona evaporation/condensation model (Meyer et al. 2000; Liu et al. 2002a; Liu et al. 2007). In the frame of this model, the Compton cooling becomes efficient in cooling of the corona at high accretion rate (in units of Eddington rate), leading to condensation of corona gas to the disk. Consequently, the relative strength of corona to the disk becomes weaker at higher Eddington ratio. Therefore, the fraction of hard X-ray emission to disk emission and hence to the bolometric emission is smaller at higher Eddington ratio. The independence of the fraction of hard X-ray luminosity on the mass of the black hole can also be explained by the disk corona model since the corona structure and luminosity (in units of Eddington luminosity) are independent on the mass of black holes.

Key words: accretion:accretion disks — galaxies: active — X-rays:galaxies

1 INTRODUCTION

The accretion onto the supermassive black hole is the key process to produce the spectral energy distribution in AGNs, including the Big Blue Bump, soft X-ray excess, Fe $K\alpha$ lines at 6.4 keV and hard X-ray tail. These observed spectral properties indicate that the cold and hot gas coexist near the central supermassive black hole. According to the commonly accepted theory the optical/UV radiations are emitted from a geometrically thin and optically thick disk (i.e. the cold gas) (e.g., Shakura & Sunyaev 1973, Lynden-Bell & Pringle 1974), and the hard X-rays arise from hot optically thin and geometrically thick accretion flows, such as disk corona (Haardt & Maraschi 1991, 1993; Nakamura & Osaki 1993; Svensson & Zdziarski 1994; Kawaguchi et al. 2001, Cao 2009) or advection-dominated accretion flows (ADAF)(e. g., Narayan & Yi 1994, 1995; Narayan et al. 1998).

The disk corona model was proposed by Liang & Price (1977) to explain the observed X-ray emission in Cyg X-1. The accretion in the corona is supplied with material and energy through disk evaporation (e.g., Meyer & Meyer-Hofmeister 1994; Liu et al. 1999) or magnetic process (e.g. Galeev et

al. 1979; Stella et al. 1984; Liu et al. 2002b, 2003). In AGNs a hot corona sandwiched by a disk is thought responsible to emit the X-rays (e.g. Haardt & Maraschi 1991; Nakamura & Osaki 1993; Kawaguchi et al. 2001; Liu et al. 2002b, 2003). In the context of these studies it is often assumed that a fraction (f) of accretion energy is released in the corona in fits to individual AGN spectrum. Questions arise as, is the fraction f dependent on the Eddington ratio and black hole? how does the relative strength of the corona to the disk vary with the Eddington ratio and black hole mass?

In this work, we attempt to investigate these issues from observation. We compile a large blue AGN sample of 240 objects derived from Dong et al. 2008 (hereafter Dong08), in which the objects are selected from the Sloan Digital Sky Survey (SDSS) DR4 spectral data and less affected by dust extinction. We investigate the correlation between the corona component and Eddington ratio. We show how the coronal luminosity fraction varies with the Eddington ratio and black hole mass for radio-quiet (RQ) AGNs. The results are compared with previous results of Wang et al. (2004) for a sample of 56 RQ AGNs observed by ASCA and Yang et al. (2007) for a combined sample from different observational satellites. Our aim is to study the physical relation between the disk and corona.

The outline of the paper is in the following. In Sect.2 the black hole mass, the optical and X-ray luminosities are calculated. In Sect.3 the overall properties of the sample and the correlation analyses are present. In the Sect.4 the statistical results are interpreted by the disk corona model. The discussion and conclusions are given in Sect.5 and Sect.6 respectively.

2 THE SAMPLE AND DATA REDUCTION

Our sample is derived from a blue AGN sample in Dong08. These blue AGNs are selected from the Sloan Digital Sky Survey (SDSS) DR4 spectral data, including Seyfert 1 galaxies and QSOs. They are less affected by dust extinction. In order to investigate the relation between the disk and the corona, whether and how the black hole mass and Eddington ratio affect this relation, we select the objects by the following criteria:

- (1) The objects are detected or covered by Faint Images of Radio Sky at Twenty Centimeters (FIRST) survey.
- (2) The X-ray flux is available.
- (3) Combining the radio with the optical data, we can determine the radio loudness and the RQ AGNs are selected.

According to the above criteria, we analyze the sample in Dong08. Given the upper limit to the radio flux, 404 AGNs are detected or covered by FIRST. After matching to ROSAT bright and faint SRC Catalogue database (ROSAT-SRC) with a distance criterion between the optical and X-ray source of $\Delta \leq 60''$ (Voges et al. 1994), we obtain a sample containing 258 AGNs.

We use the radio-loudness definition (Ivezić et al. 2002) to divide RQ and radio-loud(RL) AGNs, that is,

$$R_i \equiv \log\left(\frac{F_{20\text{cm}}}{F_i}\right) = 0.4(i - t). \quad (1)$$

where F_i and $F_{20\text{cm}}$ are flux densities at I-band and 20cm respectively, i is the I-band magnitude, and $t = -2.5 \log(F_{20\text{cm}}/3631\text{Jy})$. Objects with $R_i > 1$ are RL and objects with $R_i < 1$ are RQ. According to this definition, 240 (93%) objects in the sample are RQ and 18 (7%) objects are RL. We mainly analyze these 240 RQ objects in order to constraint the accretion mechanism.

The black hole mass, M_{BH} , is a fundamental parameter of AGNs. Different methods have been developed to estimate the black hole mass (Woo & Urry 2002 and reference therein; Vestergaard & Peterson 2006), of which the reverberation mapping is the most widely used method (Peterson 1993; Peterson et al. 2004). In the reverberation mapping method, the distance of the broad line region (BLR) from the central black hole can be deduced from the time lag between continuum and emission lines. By combining the distance with the measured emission line width, the black hole mass can be determined. In our sample, we take the $FWHM(H\beta)$ as the circular velocity in BLR. The size/distance of the

BLR is given by an empirical relation with the continuum luminosity at 5100Å (Vestergaard & Peterson 2006). Thus, the mass of black hole for RQ AGNs can be estimated as

$$\log M_{\text{BH}}(H\beta) = \log \left\{ \left[\frac{FWHM(H\beta)}{1000 \text{ km s}^{-1}} \right]^2 \left[\frac{\lambda L_{\lambda}(5100\text{\AA})}{10^{44} \text{ ergs s}^{-1}} \right]^{0.5} \right\} + (6.9 \pm 0.02). \quad (2)$$

The bolometric luminosity of AGNs is approximately calculated from optical continuum luminosity, $L_{\text{bol}} \approx 9\lambda L_{\lambda}(5100\text{\AA})$ (Elvis et al. 1994). The Eddington ratio ($L_{\text{bol}}/L_{\text{Edd}}$) is then calculated from the bolometric luminosity and black hole mass,

$$L_{\text{bol}}/L_{\text{Edd}} = 0.072 \frac{\lambda L_{\lambda}(5100\text{\AA})}{10^{44} \text{ ergs s}^{-1}} \left(\frac{M_{\text{BH}}}{10^8 M_{\odot}} \right)^{-1}, \quad (3)$$

where the Eddington luminosity is $L_{\text{Edd}} = 1.25 \times 10^{38} M_{\text{BH}}/M_{\odot}$ ergs s⁻¹.

We get the X-ray flux densities from the count rates using the energy to counts conversion factor (ECF) for power-law spectra and Galactic absorption, where the power-law photon indices and the corresponding absorption column densities (N_H) are estimated from the two hardness ratios given by the ROSAT-SRC. With the integrated flux densities in 0.1–2.4 keV and power-law indices Γ we deduce the monochromatic flux at 2 keV. After deriving the flux at 2 keV we extrapolate X-ray flux (F_{int}) in the range from 2 keV to 10 keV by assuming a constant hard X-ray photon index, $\Gamma = 1.9$ (Pounds et al. 1990; Reeves & Turner 2000). The X-ray luminosity is then calculated by the standard luminosity-flux relation,

$$L = 4\pi d_L^2 F_{\text{int}} (1+z)^{-(1-\alpha)}, \quad (4)$$

where $\alpha = \Gamma - 1$ and d_L is the luminosity distance related to the redshift and cosmology model.

Throughout this paper, we adopt the cosmology model with $H_0 = 70 \text{ km s}^{-1} \text{ Mpc}^{-1}$, $\Omega_M = 0.3$, $\Omega_\Lambda = 0.7$.

Our sample is listed in Table 1 of Appendix A, where column 1 is the object name, column 2 the redshift, column 3 the derived mass of black hole, column 4 the optical luminosity at 5100Å (in unit of $10^{44} \text{ ergs s}^{-1}$), column 5 the radio flux at 20cm, column 6 the radio-loudness, column 7 and 8 are the derived X-ray data in 0.1–2.4 keV and 2–10 keV respectively.

3 DATA ANALYSIS AND RESULTS

3.1 Properties of the Sample

Fig.1 shows the histogram of black hole mass and the Eddington ratio of our sample. It can be seen that the mass of black holes covers a large range from 10^6 to $10^{10} M_{\odot}$. The center of the distribution of M_{BH} for the sources is greater than $10^7 M_{\odot}$. The histogram of the Eddington ratio of the sample shows that the Eddington ratio ranges from $10^{-2.5}$ to 1.0, most of the objects are at high accretion rate. If the accretion mode in AGNs is similar to that in black hole X-ray binaries, the high accretion rates of our sample indicate that the thin disk extends down to the innermost stable circular orbit, hence the hard X-ray arises from a disk corona rather than an ADAF.

3.2 Correlation Analysis and Results

We plot the ratio of X-ray luminosity to the bolometric luminosity vs. the mass of black holes and vs. the Eddington ratio for RQ sources in Fig.2. In the upper two panels the vertical axis is logarithm of the fraction of hard X-ray luminosity to bolometric luminosity ($\log(L_{2-10\text{keV}}/L_{\text{bol}})$). While in the lower panels, the vertical axis is logarithm of the fraction of soft X-ray luminosity (0.1–2.4 keV) to the bolometric luminosity. Comparing the upper and lower panels one can see that the distribution trends are different for soft X-ray and hard X-ray fraction.

We test the correlations between $\log(L_{2-10\text{keV}}/L_{\text{bol}})$ and $\log(M_{\text{BH}})$ and between $\log(L_{2-10\text{keV}}/L_{\text{bol}})$ and Eddington ratio by the method of Spearman's rank correlation analysis




Fig. 1 The left plot is the histogram of black hole mass for our sample. We can see that the center of the distribution of M_{BH} of the RQ sources is nearly $10^{8.0} M_{\odot}$. The right panel plots the histogram of Eddington ratio for the sources. Most of sources are at high accretion rate.

(William et al. 2000). We calculate the Spearman's rank correlation coefficient(ρ) and the two-sided significance of its deviation from zero (P_{Null}), representing the probability for randomness of the observed correlation of two variables. Significant correlation of two variables is accepted if P_{null} is less than 0.05. For our sample, the coefficient ρ between $\log(L_{2-10\text{keV}}/L_{\text{bol}})$ and $\log M_{\text{BH}}$ is 0.001, and P_{Null} is 0.988. This means that the fraction of hard X-ray is independent of black hole mass. For the relation between the hard X-ray fraction and Eddington ratio, we get $\rho = -0.17$, $P_{\text{Null}} = 0.008$. This indicates that for RQ AGNs the hard X-ray fraction correlates with Eddington ratio, with increase of the Eddington ratio, the hard X-ray fraction decreases. This result is qualitatively consistent with the work of Wang et al. (2004).

In addition, we analyze the correlation of soft X-ray fraction with the black hole mass and Eddington ratio. It is found that the soft X-ray fraction in the bolometric luminosity depends on black hole mass ($\rho = -0.26$, $P_{\text{Null}} = 4.085e-5$), in contrast to the case of hard X-rays. The soft X-ray fraction also positively correlated with the Eddington ratio ($\rho = 0.18$, $P_{\text{Null}} = 0.005$), contrary to the trend of hard X-ray fraction.

4 INTERPRETATION OF THE CORRELATIONS WITH DISK CORONA MODEL

The hard X-ray radiation from RQ AGNs with high Eddington ratio is commonly thought to be produced from a disk corona as a result of Comptonization of the softer photons arising from the accretion disk (e.g. Haardt & Maraschi 1993; Kawaguchi et al. 2001; Liu et al. 2003; Cao 2009). If the bolometric luminosity is contributed dominantly by the disk (L_d) and corona (L_c), the fraction of hard X-ray luminosity in the total luminosity represents the strength of the corona relative to the disk. Thus, the negative correlation between the fraction of hard X-ray and Eddington ratio for RQ AGNs indicates that the corona relative to the disk becomes weaker as the Eddington-scaled accretion rate increases.

Theoretical calculation of radiations from the disk corona (Haardt & Maraschi 1991; 1993) shows that the spectral shape does not change with accretion rate. In other words, ratio (f) of X-ray luminosity and bolometric luminosity is predicted to be independent on the accretion rate. To explain the observational correlation between f and Eddington accretion rate, Wang et al. (2004), Yang et al. (2007) and Cao(2009) consider magnetic fields and different shear stress tensors. Here we introduce the disk corona evaporation and/or condensation model (Liu et al. 2002a). We show that even without magnetic fields the correlation can be simply a consequence of efficient condensation of coronal gas to the disk in the case of high accretion rates.




Fig.2 The fraction of X-ray luminosity in bolometric luminosity vs. the mass of black hole and Eddington ratio for our sample. The top panels show $\log(L_{2-10\text{keV}}/L_{\text{bol}})$ in dependence on $\log M_{\text{BH}}$ and $\log(L_{\text{bol}}/L_{\text{Edd}})$. The bottom panels describe the trend of $\log(L_{0.1-2.4\text{keV}}/L_{\text{bol}})$ varying with $\log M_{\text{BH}}$ and $\log(L_{\text{bol}}/L_{\text{Edd}})$. We can see that $\log(L_{0.1-2.4\text{keV}}/L_{\text{bol}})$ increases with the decrease of black hole mass and the increase of the Eddington ratio.

Assuming that the energy of photons from the disk is amplified by a factor of $A \equiv \Delta\epsilon/\epsilon$ in the process of collisions with electrons when going through the corona, for non-relativistic thermal distributions of electrons A is a function of electron temperature (T_e) and optical depth (τ_{es}) of the corona,

$$A = \frac{4kT_e}{m_e c^2} \tau_{es}. \quad (5)$$

The fraction of hard X-ray luminosity is expressed as,

$$\frac{L_{2-10\text{keV}}}{L_{\text{bol}}} = \frac{AL_d}{AL_d + L_d} = \frac{1}{1 + \frac{1}{A}}. \quad (6)$$

In the case of high accretion rate, disk radiations are strong, which leads to efficient inverse Compton scattering in the corona. As a consequence of over-cooling, part of the corona gas condenses into the disk, thereby the optical depth decreases and hence the amplification factor is low. The higher the accretion rate, the lower density of the corona is. Therefore, from Eq.6 one can see that the fraction of X-ray decreases at high accretion rates. Thus, the negative correlation between the fraction of hard X-ray luminosity and the Eddington ratio can be explained by the corona condensation to the disk as a consequence of pressure and energy balance between the disk and corona.

Detailed calculations of condensation rate and luminosity have been carried out by Liu et al. (2007) and Taam et al. (2008). In this model, when the Compton scattering is the dominant cooling mechanism, the luminosities from the corona and disk are respectively as follows (Taam et al. 2008),

$$\begin{aligned} L_c &= \int_{3R_s}^{R_{out}} 2\pi RH \frac{4kT_e}{m_e c^2} n_e \sigma_T 4\sigma T_{\text{eff}}^4(R) dR \\ &= 4.11 \times 10^{38} \alpha^{-\frac{7}{5}} m \dot{m}_c^{\frac{7}{5}} \dot{m}_d^{\frac{3}{5}} \int_3^{r_{out}} \left(\frac{3}{r}\right)^{\frac{23}{10}} \left[1 - \left(\frac{3}{r}\right)^{\frac{1}{2}}\right]^{\frac{3}{5}} dr. \end{aligned} \quad (7)$$

$$\begin{aligned} L_d &= \int_{3R_s}^{R_{out}} \sigma T_{\text{eff}}^4(R) 4\pi R dR \\ &= \int_{3R_s}^{R_{out}} \frac{3GM\dot{M}_d}{8\pi R^3} \left[1 - \left(\frac{3R_s}{R}\right)^{\frac{1}{2}}\right] 4\pi R dR \\ &= 9.38 \times 10^{38} m \dot{m}_d \int_3^{r_{out}} \frac{1}{r^2} \left[1 - \left(\frac{3}{r}\right)^{\frac{1}{2}}\right] dr. \end{aligned} \quad (8)$$

where $r = R/R_s$ (R_s the Schwarzschild radius), $m = M_{\text{BH}}/M_\odot$, \dot{m}_c is the Eddington-scaled accretion rates in the corona and \dot{m}_d the one in the disk. Note that for a truncated inner disk lying under a corona, \dot{m}_d comes only from the condensation of coronal gas; While in the case of a full disk coexisting with a corona for AGNs discussed here, the accretion rate in the disk is mainly from outer region, $\dot{m}_d = \dot{m} - \dot{m}_c$. We then derive the ratio of corona luminosity and disk luminosity,

$$L_c/L_d \propto \alpha^{-\frac{7}{5}} \dot{m}_c^{\frac{7}{5}} \dot{m}_d^{-\frac{2}{5}} = \alpha^{-\frac{7}{5}} \dot{m}_c^{\frac{7}{5}} (\dot{m} - \dot{m}_c)^{-\frac{2}{5}}. \quad (9)$$

In the disk corona evaporation and/or condensation model, the coronal accretion flow is supplied by disk evaporation in the region around $300R_s$. The coronal flow rate \dot{m}_c can be as high as a few percent of Eddington value if the accretion rate exceeds a few percent of Eddington value. However, the coronal flow can condense partially into the disk in the inner region as a result of Compton cooling, and the condensation rate is larger at higher accretion rate. This means that the coronal flow (\dot{m}_c) is lower at higher accretion rate. From Eq.(9) one can see that the ratio of corona luminosity and disk luminosity decreases with increasing accretion rate \dot{m} . Noting that the accretion rate in unit of Eddington value equals to the Eddington ratio, the disk corona evaporation and condensation model predicts that the ratio of X-ray luminosity and disk luminosity decreases with increasing Eddington ratio, which can explain the correlation obtained from our RQ AGN sample.

If magnetic fields are also taken into account as additional heating (Liu et al. 2003; Cao 2009), the coronal accretion flow can be higher and the corona can be stronger than that without magnetic fields. But the trend that condensation is stronger at higher accretion rates does not change by the magnetic fields. Therefore, the variation of relative strength of the disk and corona with accretion rates keeps the same trend. Thus, with inclusion of magnetic fields the model will still work in interpreting the correlations.

Another important feature of the disk corona evaporation and condensation model is that the corona structure and evaporation/condensation properties are independent on the mass of black hole if scaled properly. In fact, the model can be applied in both X-ray binaries and AGNs. Eq.(9) shows that the luminosity ratio from the corona and disk indeed does not vary with the mass of black hole. Therefore, the model can explain why for our sample the fraction of hard X-ray luminosity is independent on the mass of black hole.

5 DISCUSSION

5.1 Implication to the Hard X-ray Bolometric Correction

The bolometric luminosity is either estimated from the measured luminosity at 5100Å or from hard X-ray luminosity with fixed correction factors. However, our data analysis shows that the hard X-ray fraction in the bolometric luminosity decreases with increasing Eddington ratio for RQ AGNs. This means that bolometric correction factor from hard X-rays, $k_{2-10\text{keV}} = L_{\text{bol}}/L_{2-10\text{keV}}$, should increase with increasing Eddington ratio. This result is in good agreement with that of Vasudevan & Fabian (2007), which is drawn from 54 AGNs with available SEDs. Note that in our study we calculate the bolometric luminosity from $9\lambda L_\lambda(5100\text{\AA})$. If the bolometric correction factor for $L_{5100\text{\AA}}$ also tends to increase with Eddington ratio (see Fig.15 of Vasudevan & Fabian 2007), the hard X-ray fraction in bolometric luminosity would be even smaller at higher Eddington ratio for our sample. Therefore, our statistical results support that the bolometric correction for hard X-rays increases with increasing Eddington ratio.

5.2 The Constant Photon Index

In order to investigate the strength of corona ie. to get the hard X-ray flux, we use the same photon indices for all objects in our sample to extrapolate the hard X-ray emissiong from 2keV flux. From above analysis, we get a smaller absolute coefficient value (ρ) between the fraction of hard X-ray luminosity and Eddington ratio. However, if we note that for some individual objects the spectra become softer when a source is brighter (e.g., Lu & Yu 1999, Bian et al. 2005, Middleton et al. 2008; Saez et al. 2008) and this effect is taken into account, the hard X-ray luminosity fraction should change with the Eddington ratio more steeply than our results since here we assume the photon index is constant in calculating the hard X-ray luminosity.

5.3 The Origin of Soft X-rays in 0.1-2.4 keV

Our study also indicates that for the same RQ AGNs the fraction of soft X-ray luminosity in the ROSAT energy band (0.1 – 2.4keV) anticorrelates on the mass of black hole, as shown in Fig.2. If the soft X-ray is also from the corona, the correlation should be similar to that for hard X-rays, that is, the fraction of soft X-ray is independent on the balck hole mass. On the other hand, if the disk emits dominantly in the soft X-ray, the fraction of the soft X-ray to the total luminosity should also be independent on the mass of black hole since both the total luminosity and disk luminosity are proportional to the mass of black hole. We note that the effective temperature of the disk decreases with the mass of black hole (e.g., Thorne K. S. 1974, Bonning et al. 2007), $\sigma T_{\text{eff}}^4 = \frac{3GM}{8\pi R^3} [1 - (3R_s/R)^{1/2}]$, reaching a maximum at $R = (49/36)(3R_S)$ (see Liu et al. 2007),

$$T_{\text{eff,max}} = 1.3 \times 10^5 (M_{\text{BH}}/10^8 M_\odot)^{-1/4} \dot{m}^{1/4} K. \quad (10)$$

For AGN with a high mass of black hole, the disk does not emit in the soft X-rays but in the optical/UV band. With mass decreases, the high energy part of the disk spectrum could extend to soft X-ray. Consequently, the fraction of disk contribution to the soft X-ray increases with decreasing of M_{BH} . This could lead to a negative correlation between the fraction of soft X-ray and the mass of black hole, qualitatively consistent with the observational results shown in Section 3. Therefore, the different correlations with black hole mass of the soft X-ray and hard X-ray may indicate that the soft X-rays are contributed from not only the corona but also the disk. Zhou et al. (1997) separated two components in the soft X-ray region through statistic analyzing the Big Blue Bump.

6 CONCLUSION

We compile a blue AGNs sample containing Seyfert 1 galaxies and QSOs with 240 AGNs being RQ. We mainly investigate the dependence of the fraction of hard X-ray luminosity in bolometric luminosity

$(\log(L_{2-10\text{keV}}/L_{\text{bol}}))$ on the black hole mass and Eddington ratio in RQ AGNs by using the Spearman's rank correlation. We find that the fraction of hard X-ray luminosity in bolometric luminosity decreases with the increase of Eddington ratio for RQ AGNs. While the fraction of hard X-ray luminosity in the bolometric luminosity is independent on black hole mass for a wide range of black hole mass. The correlation between the hard X-ray and bolometric luminosity suggests that the hard X-ray bolometric correction increase at high Eddington ratio, confirming with a large sample Vasudevan & Fabian (2007)'s results. These observational features support the disk corona model developed by Meyer et al. (2000) and Liu et al. (2002a). Combining the theoretical model and the wide range of black hole mass in our sample, we suggest that the intrinsic accretion process in AGNs and black hole X-ray binaries be similar.

Acknowledgements We are grateful to the anonymous referee for constructive and enlightening comments. Liu J. Y. greatly thanks Yuan W. for the instruction in X-ray data deducing and the statistical analysis. We also acknowledge Dong X. B. for the SDSS data analysis. This work is supported by the National Natural Science Foundation of China (grants 10533050 and 10773028) and National Basic Research Program of China-973 Program 2009CB824800.

References

- Bian W. H., 2005, ChJAS, 5, 289
 Bonning E.W., Cheng L. et al., 2007, ApJ, 659, 211
 Cao Xinwu, 2009, MNRAS, 394, 207
 Dong X.B., Wang T.G. et al., 2008, MNRAS, 383, 581(Dong08)
 Elvis M., et al., 1994, ApJS, 95, 1
 Galeev A. A., Rosner R., Vaiana G. S., 1979, ApJ, 229, 318
 Haardt, F., & Maraschi, L., 1991, ApJ, 380, L51
 Haardt, F., & Maraschi, L., 1993, ApJ, 413, 507
 Ivezić Ž., et al., 2002, AJ, 124, 2364
 Kawaguchi, T., Shimura, T., & Mineshige, S., 2001, ApJ, 546, 966
 Liang E. P. T., Price R. H., 1977, ApJ, 218, 247
 Liu B. F., Yuan W., Meyer F., et al., 1999, ApJ, 527, L17
 Liu B. F., Mineshige S., Meyer F., et al. 2002, ApJ, 575, 117 (2002a)
 Liu B. F., Mineshige S., Shibata K., 2002, ApJ, 572, L173 (2002b)
 Liu B. F., Mineshige S., Ohsuga K., 2003, ApJ, 587, 571
 Liu B. F., Taam Ronald E., Meyer-Hofmeister E., Meyer F., 2007, ApJ, 671, 695
 Lu Y. & Yu Q., 1999, ApJ, 570, L47
 Lynden-Bell D. & Pringle J.E., 1974, MNRAS, 168, 603
 Meyer F. & Meyer-Hofmeister E., 1994, A&A, 288, 175
 Meyer F., Liu B. F., et al., 2000, A&A, 361, 175
 Middleton M., Done C. and Schurch N., 2008, MNRAS, 383, 1501
 Nakamura, K & Osaki, Y., 1993, PASJ, 45, 775
 Narayan R. & Yi I., 1994, ApJ, 428, L13
 Narayan R., Mahadevan R., & Quataert E., 1998, in The Theory of Black Hole Accretion Discs, eds. M. A. Abramowicz, G. Bjornsson, & J. E. Pringle (Cambridge University, Cambridge)
 Narayan R., & Yi, I. 1995, ApJ, 452, 710
 Peterson B. M., 1 Narayan et al. 1993, PASP, 100, 18
 Peterson B. M., Ferrarese L., et al., 2004, ApJ, 613, 682
 Pounds K. A., Nandran K., Stewart G. C. et al., 1990, Nature, 344, 132

- Reeves J. N., Turner M. J. L., 2000, MNRAS, 316, 234
Saez C., Chartas G., et al. 2008, ApJ, 135, 1505
Shakura N.I., Sunyaev R.A., 1973, A&A, 24, 337
Stella L., Rosner R., 1984, ApJ, 227, 312
Svensson R., Zdziarski A. A., 1994, ApJ, 436, 599
Taam, R.E, Liu, B. F, et al., 2008, ApJ, 688, 527
Thorne K. S., 1974, ApJ, 191, 507
Vasudevan R .V., Fibian A. C., 2007, MNRAS,381, 1235
Vestergaard M., & Peterson B. M., 2006, ApJ, 641, 689
Voges W., Gruber R., Haberl F., et al., 1994, ROSAT NEWS No.32
Wang J. M., Watarai K. Y., Mineshige, S., 2004, ApJ, 607, L107
William H.P., Saul A.T. et al., 2000, Numerical Recipes in Fortran (2nd, Cambridge University Press, Cambridge)
Woo J. H. & Urry C. M., 2002, ApJ, 578, 530
Yang F., Hu C. et al., 2007, ChJAA, 7, 353
Zhou Y. Y., Yu K. N., et al., 1997, ApJ, L9

Appendix A: TABLE

Table1.The Blue AGNs Sample

Object SDSS (1)	Z (2)	$\log(M_{BH})$ (M_{\odot}) (3)	$L_{5100\text{\AA}}$ (10^{44} ergs s $^{-1}$) (4)	$F_{20\text{cm}}$ (mJy) (5)	R_i (6)	$\log(L_{0.1-2.4\text{keV}})$ (ergs s $^{-1}$) (7)	$\log(L_{2-10\text{keV}})$ (ergs s $^{-1}$) (8)
000710.01+005329.0	0.31620	9.13	3.85	1.44	0.55	44.949	44.238
000834.72+003156.1	0.26303	7.71	2.71	<1.00	0.43	44.665	42.915
000943.14−090839.1	0.20958	8.52	2.09	<1.00	0.21	44.057	43.225
001247.93−084700.4	0.22006	8.01	1.95	<1.00	0.21	44.629	43.965
004319.74+005115.3	0.30807	9.36	3.08	1.60	0.80	44.425	44.099
010226.31−033904.6	0.29421	7.92	9.63	<1.00	−0.10	45.012	43.656
011254.92+000313.0	0.23855	8.14	1.98	<1.00	0.42	44.398	43.816
014017.07−005003.0	0.33424	8.92	13.40	<1.00	0.06	45.232	44.927
014234.40−011417.3	0.224453	7.14	0.68	<1.00	0.69	44.078	44.060
015530.01−085704.0	0.16465	8.48	1.78	<1.00	−0.04	44.212	43.739
015910.04+010514.6	0.21716	8.08	1.99	<1.00	0.22	44.164	43.731
022347.48−083655.5	0.26077	7.52	0.97	1.04	0.81	44.330	43.631
022417.16−092549.3	0.31148	8.79	1.72	<1.00	0.67	44.751	43.763
072937.04+375435.0	0.20364	7.23	1.39	<1.00	0.40	44.558	43.179
073503.49+431153.5	0.26246	8.11	1.86	<1.00	0.59	44.302	43.979
074645.05+314149.2	0.32696	7.55	3.31	<1.00	0.54	44.972	43.504
074820.97+340752.6	0.34304	8.09	3.08	<1.00	0.68	44.482	43.986
074948.26+345444.0	0.13181	7.71	1.16	0.82	−0.01	44.129	43.526
075217.84+193542.2	0.11723	9.43	2.59	22.95	0.84	43.173	42.932
075245.60+261735.7	0.08216	6.92	0.46	1.27	0.13	43.924	42.985
075819.68+421935.1	0.21125	7.97	2.84	1.81	0.36	44.958	43.865
075949.54+320023.9	0.18805	7.60	1.92	<1.00	0.35	44.488	43.353
080559.94+260602.3	0.13594	8.31	1.34	<1.00	−0.09	43.371	42.525
080644.65+384318.3	0.34539	7.98	3.28	<1.00	0.71	44.954	44.363
081054.73+501319.5	0.32184	9.13	5.47	<1.00	0.37	44.258	43.940
081317.91+435620.7	0.25455	8.30	1.81	<1.00	0.60	44.351	43.472
081427.69+437051.2	0.22418	7.61	1.27	<1.00	0.47	44.205	43.334
081738.33+242330.0	0.28259	8.20	4.23	4.45	0.86	44.599	44.382
082633.51+074248.4	0.31064	8.14	7.54	<1.00	0.12	44.876	43.978
082640.73+063041.5	0.17092	7.82	0.80	<1.00	0.50	44.165	43.175
083120.99+483154.4	0.33466	7.56	3.30	<1.00	0.62	45.087	44.444
083225.34+370736.1	0.09191	9.22	1.20	11.73	0.75	44.291	43.776
083417.91+491439.2	0.17315	7.40	0.93	<1.00	0.26	44.082	43.225
083443.80+382632.7	0.28822	8.59	2.94	<1.00	0.48	45.016	43.780
083553.46+055317.1	0.20438	7.49	2.24	<1.00	0.15	44.031	43.165
083658.91+442602.4	0.25443	8.74	13.58	10.25	0.64	45.314	44.562
084230.51+495802.3	0.30508	8.53	2.88	<1.00	0.54	44.447	43.552
084504.20+542612.0	0.30317	7.97	1.53	<1.00	0.88	44.427	43.895
084853.09+282411.8	0.19820	8.29	0.81	<1.00	0.60	44.265	43.715
085259.22+031320.6	0.29708	8.31	8.80	<1.00	−0.02	44.334	43.979
085632.39+504114.0	0.23471	8.20	5.11	<1.00	−0.10	44.005	43.129
085828.69+342343.8	0.25666	8.70	6.43	<1.00	−0.08	45.039	43.989
085900.48+383211.6	0.34551	7.73	2.94	<1.00	0.65	45.459	43.663
085915.65+011800.5	0.28206	8.40	3.25	<1.00	0.46	44.582	43.603
090137.99+532051.1	0.16165	7.11	1.05	<1.00	0.25	44.464	43.359
090151.14+103020.4	0.20093	7.92	4.13	<1.00	−0.14	44.572	43.903
090455.00+511444.6	0.22463	8.20	2.55	<1.00	0.10	44.364	43.564
090519.67+440139.1	0.34246	7.50	1.64	<1.00	0.80	44.898	43.344
090654.47+391455.4	0.24067	7.39	1.85	<1.00	0.51	44.593	43.695
090851.25+444611.2	0.32021	8.34	3.66	<1.00	0.42	44.130	43.885

Object SDSS	Z	$\log(M_{\text{BH}})$ (M_{\odot})	$L_{\text{5100}\text{\AA}}$ (10^{44} ergs s $^{-1}$)	$F_{20\text{cm}}$ (mJy)	R_i	$\log(L_{0.1-2.4\text{keV}})$ (ergs s $^{-1}$)	$\log(L_{2-10\text{keV}})$ (ergs s $^{-1}$)
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)
090932.04+503019.6	0.26728	7.56	1.14	<1.00	0.79	43.933	43.046
091010.00+481341.7	0.11700	8.14	0.86	<1.00	0.00	44.297	43.446
091702.38-004417.5	0.32231	8.71	8.21	<1.00	0.15	45.337	44.643
091755.02+053749.7	0.34898	8.43	1.86	<1.00	0.74	44.573	43.461
091955.35+552137.1	0.12295	8.46	2.33	<1.00	-0.38	44.338	43.412
092309.87+453046.4	0.29238	7.63	2.16	<1.00	0.54	44.672	43.779
092554.44+453544.2	0.32948	7.97	3.10	<1.00	0.47	44.534	43.631
092909.79+464424.0	0.23996	8.64	6.15	<1.00	-0.12	43.833	43.103
092933.58+095617.0	0.23194	7.53	0.95	<1.00	0.70	44.199	43.811
093701.04+010543.7	0.05054	6.92	0.30	<1.00	-0.04	43.885	43.308
093939.69+375705.8	0.23127	7.79	2.24	<1.00	0.34	43.739	42.863
094621.27+471131.3	0.23049	7.67	1.37	<1.00	0.50	44.233	43.890
095048.38+392650.5	0.20564	8.41	3.45	<1.00	-0.16	44.807	44.329
095302.64+380145.2	0.27291	8.45	2.60	<1.00	0.45	44.740	43.853
095823.45+065506.5	0.34581	8.30	5.20	<1.00	0.53	44.715	44.356
095833.95+560224.4	0.21639	6.99	0.74	0.83	0.61	44.113	42.926
095915.65+050355.1	0.16230	7.88	1.42	<1.00	0.10	44.040	42.836
095931.67+504449.0	0.14323	7.54	1.12	<1.00	0.02	43.771	42.984
100033.88+104723.7	0.22648	7.99	2.25	2.17	0.29	44.776	43.656
100201.76+620816.3	0.13379	6.98	0.35	<1.00	0.49	44.093	43.173
100420.13+051300.4	0.16049	7.61	2.61	<1.00	-0.12	43.401	42.547
100541.86+433240.4	0.17843	7.79	2.67	2.81	0.47	44.844	43.917
100627.94+603043.6	0.21025	7.15	1.00	<1.00	0.56	44.164	43.081
100744.54+500746.6	0.21204	7.67	2.51	<1.00	0.18	44.594	43.861
101044.51+004331.3	0.17757	8.83	3.15	0.77	-0.36	44.556	43.211
101401.86+461953.7	0.32112	8.38	5.56	<1.00	0.22	44.707	43.696
101415.14+091839.3	0.25224	8.62	1.18	<1.00	0.58	44.500	43.999
101437.46+440639.1	0.20014	7.62	1.38	<1.00	0.31	44.251	43.458
101730.97+470225.0	0.33499	7.91	3.62	<1.00	0.48	44.629	43.732
101852.45+495800.3	0.15481	7.00	0.41	<1.00	0.63	43.931	43.436
102309.48+082602.1	0.34411	8.02	2.69	<1.00	0.60	44.492	43.585
102512.85+480853.2	0.33156	8.50	2.35	<1.00	0.66	44.289	43.386
102531.28+514034.8	0.04488	7.05	0.29	0.60	-0.43	44.038	42.994
102745.84+051558.9	0.31480	7.56	3.02	<1.00	0.45	44.648	43.411
103421.70+605318.1	0.22775	8.07	1.97	<1.00	0.29	44.158	43.292
103457.29-010209.0	0.32801	7.64	3.20	<1.00	0.51	44.791	43.891
104041.50+600239.3	0.29711	7.63	2.53	<1.00	0.54	44.318	43.656
104541.76+520235.5	0.28393	7.87	5.73	<1.00	0.16	44.855	43.895
105007.75+113228.6	0.13344	7.65	2.50	1.50	-0.10	44.264	43.445
105055.14+552723.2	0.33196	7.86	5.20	<1.00	0.33	45.165	44.059
105118.23+605008.2	0.27603	7.98	1.97	<1.00	0.61	44.031	42.963
105444.70+483139.0	0.28651	8.73	10.08	1.56	-0.00	45.219	44.523
105752.69+105037.9	0.22033	7.23	1.20	<1.00	0.48	44.458	43.585
105830.13+601600.3	0.14868	7.56	1.40	<1.00	-0.11	44.300	43.041
110016.19+393524.5	0.31265	7.71	1.60	<1.00	0.88	44.998	43.979
110540.09+064225.7	0.23046	8.36	1.20	<1.00	0.36	44.178	44.159
111006.95+612521.4	0.26234	7.60	1.46	<1.00	0.51	44.403	43.527
111706.39+441333.3	0.14382	8.73	4.02	<1.00	-0.45	43.323	42.475
111740.48+530151.3	0.15851	7.56	0.83	<1.00	0.36	43.783	43.081
111830.28+402554.0	0.15457	7.67	3.16	2.32	0.06	44.670	43.826
112114.21+032546.8	0.15203	7.62	1.12	2.20	0.59	43.858	43.871
112417.79+602026.7	0.20470	7.27	1.06	<1.00	0.49	44.288	43.619

Object SDSS	Z (1)	$\log(M_{\text{BH}})$ (M_{\odot}) (2)	$L_{\text{5100}\text{\AA}}$ (10^{44} ergs s $^{-1}$) (3)	$F_{20\text{cm}}$ (mJy) (4)	R_i (5)	$\log(L_{0.1-2.4\text{keV}})$ (ergs s $^{-1}$) (6)	$\log(L_{2-10\text{keV}})$ (ergs s $^{-1}$) (7)	$\log(L_{2-10\text{keV}})$ (ergs s $^{-1}$) (8)
112439.18+420145.0	0.22503	8.18	7.01	<1.00	-0.30	44.623	43.072	
112646.43-013417.9	0.34095	8.32	2.30	<1.00	0.80	44.001	43.866	
112941.93+512050.6	0.23385	7.47	2.50	1.26	0.36	44.719	43.577	
113105.04+610405.0	0.33802	7.95	3.64	<1.00	0.46	44.206	43.415	
113422.48+041127.6	0.10800	8.19	1.04	<1.00	-0.26	44.366	43.591	
113706.84+013948.0	0.19262	8.73	2.64	<1.00	0.07	44.714	44.322	
113738.04+103930.1	0.17454	8.06	2.55	1.65	0.20	44.405	43.707	
113908.97+591154.8	0.06127	7.86	0.76	<1.00	-0.48	43.543	43.303	
114039.86+414545.9	0.23833	7.87	1.37	<1.00	0.54	44.647	43.770	
114105.71+024117.0	0.09313	7.55	0.47	<1.00	0.06	43.623	43.092	
114327.21+431145.9	0.30505	8.05	3.38	<1.00	0.44	44.447	43.550	
114341.97-014434.5	0.10522	7.52	0.82	<1.00	0.04	43.280	42.444	
114408.90+424357.5	0.27248	7.74	2.20	<1.00	0.58	44.499	43.612	
114559.00+040409.8	0.27359	7.69	1.31	<1.00	0.88	43.916	43.889	
114954.99+044812.9	0.26951	7.53	3.02	1.76	0.62	44.474	43.529	
115105.41+445309.3	0.34394	7.35	1.99	<1.00	0.75	44.455	43.548	
115507.61+520129.6	0.15399	7.30	1.12	2.05	0.58	43.652	42.801	
115549.43+502117.1	0.28443	8.05	3.08	<1.00	0.43	44.386	43.495	
115558.97+593129.3	0.24081	7.98	2.25	<1.00	0.36	44.603	43.579	
115632.24+112653.7	0.22582	7.52	1.54	<1.00	0.38	44.176	43.302	
115758.73-002220.7	0.25984	8.42	3.25	<1.00	0.14	44.034	44.029	
120118.43+060024.1	0.33555	8.79	2.16	<1.00	0.64	44.307	43.402	
120233.08+022559.7	0.27287	8.51	2.85	<1.00	0.37	44.197	43.366	
120347.70+520749.7	0.17760	8.50	4.92	0.68	-0.54	44.681	43.957	
121018.35+015405.9	0.21589	8.40	2.71	<1.00	0.22	43.768	43.174	
122420.28+435157.9	0.32172	8.44	2.39	<1.00	0.69	44.520	43.880	
122549.28+472343.8	0.31883	7.56	2.04	<1.00	0.66	44.294	43.394	
123054.12+110011.1	0.23596	8.29	3.67	<1.00	-0.02	44.503	44.301	
123115.20+590707.2	0.32605	8.05	3.14	<1.00	0.48	44.461	44.272	
123220.11+495721.7	0.26188	7.65	2.94	<1.00	0.50	45.082	43.929	
123234.05+092815.2	0.34665	8.33	2.35	<1.00	0.74	44.614	43.812	
123958.57+490540.1	0.23550	7.04	1.08	<1.00	0.59	43.941	43.308	
124441.41+585626.8	0.19806	7.25	0.75	<1.00	0.54	44.667	43.269	
124635.24+022208.7	0.04818	6.75	0.31	2.23	0.25	44.057	43.159	
124931.72+523039.2	0.16228	8.00	2.66	<1.00	-0.14	44.447	43.272	
125519.69+014412.3	0.34318	8.92	15.52	1.47	0.11	45.350	44.548	
125719.56+442935.4	0.30025	8.25	7.68	<1.00	0.00	44.818	44.084	
125824.57+540429.8	0.34769	8.12	2.17	<1.00	0.76	44.470	43.562	
130250.51+111827.9	0.20284	7.47	1.71	<1.00	0.28	44.424	43.334	
130416.99+020537.1	0.22854	7.67	2.04	<1.00	0.27	45.378	43.966	
130421.84+560817.0	0.29830	7.82	2.17	<1.00	0.62	44.406	43.823	
130604.48+453405.4	0.32709	7.52	2.17	<1.00	0.80	44.683	44.095	
131308.67+542115.5	0.29798	8.45	2.90	<1.00	0.63	44.455	43.872	
131404.97+153054.2	0.16517	7.94	1.27	<1.00	0.22	44.285	43.363	
131651.70+630719.9	0.34724	8.65	2.01	<1.00	0.79	44.110	44.246	
132144.96+033055.7	0.26889	8.70	5.48	<1.00	0.02	44.386	43.765	
132242.47-022522.0	0.12115	7.69	0.68	<1.00	0.18	43.569	43.592	
132447.65+032432.6	0.30578	8.25	4.70	<1.00	0.16	44.358	43.461	
132643.62+015209.4	0.19670	7.68	0.85	<1.00	0.52	44.339	43.680	
132802.54+441805.3	0.23155	8.13	0.77	<1.00	0.84	43.756	42.880	
133051.24+412858.1	0.18174	7.70	1.58	<1.00	0.17	44.429	43.511	
133300.83+451809.0	0.31974	8.36	3.65	<1.00	0.37	44.727	44.267	

Object SDSS	Z (1)	$\log(M_{\text{BH}})$ (M_{\odot}) (2)	$L_{\text{5100}\text{\AA}}$ (10^{44} ergs s $^{-1}$) (3)	$F_{20\text{cm}}$ (mJy) (4)	R_i (5)	$\log(L_{0.1-2.4\text{keV}})$ (ergs s $^{-1}$) (6)	$\log(L_{2-10\text{keV}})$ (ergs s $^{-1}$) (7)	$\log(L_{2-10\text{keV}})$ (ergs s $^{-1}$) (8)
133423.30+434331.7	0.22551	7.65	1.31	<1.00	0.58	44.290	43.751	
133636.65+420934.1	0.22328	8.23	2.46	3.55	0.73	43.559	42.685	
134032.02+052158.5	0.27448	8.73	2.26	<1.00	0.63	44.668	43.892	
134206.57+050523.8	0.26602	7.59	3.48	3.85	0.90	44.468	43.582	
134251.60-005345.2	0.32581	8.37	6.22	<1.00	0.26	44.366	43.364	
134444.17+630337.2	0.29218	7.80	0.98	<1.00	0.85	44.024	43.327	
134459.45-001559.5	0.24490	7.55	1.41	<1.00	0.42	44.472	43.475	
134845.44+451809.5	0.27653	8.90	3.42	<1.00	0.34	44.715	43.930	
135829.58+010908.4	0.24392	8.30	1.76	<1.00	0.60	44.118	43.240	
135946.91+581357.0	0.22391	7.44	0.68	<1.00	0.70	44.096	43.716	
140050.21+532424.5	0.17466	7.25	1.06	<1.00	0.27	43.538	42.678	
140104.18+635234.0	0.34386	8.44	3.47	<1.00	0.48	44.223	42.802	
140604.25+572956.5	0.32581	8.57	2.44	<1.00	0.67	43.943	43.040	
140824.76+543817.7	0.29106	7.63	2.36	<1.00	0.59	44.441	43.548	
140827.51+142233.2	0.31766	7.48	1.62	<1.00	0.84	44.628	43.795	
141046.46+465802.9	0.33409	7.38	2.99	<1.00	0.65	44.873	43.164	
141758.25+360741.4	0.21180	7.62	0.81	<1.00	0.44	44.273	43.558	
141942.46+591259.4	0.31988	8.37	1.59	<1.00	0.89	44.012	43.318	
142424.22+595300.5	0.13495	8.75	2.28	2.51	0.08	44.091	43.537	
142455.53+421407.6	0.31603	8.19	7.32	<1.00	0.17	44.661	43.361	
142734.80+352543.4	0.34038	7.49	1.69	<1.00	0.78	44.084	43.177	
142748.29+050222.0	0.10608	7.24	1.27	<1.00	-0.26	43.657	42.900	
143039.30+493538.9	0.20355	7.82	1.86	<1.00	0.32	44.544	43.403	
143204.60+394439.0	0.34848	9.05	9.25	<1.00	0.14	44.063	43.154	
143704.12+000705.0	0.14036	7.45	0.61	<1.00	0.44	44.165	43.270	
143919.31+551317.8	0.25720	8.52	1.10	<1.00	0.77	44.245	43.807	
143940.27+030528.6	0.26856	7.23	1.75	<1.00	0.64	45.053	43.949	
144012.76+615633.0	0.27547	7.54	4.46	2.88	0.69	44.860	43.701	
144050.76+520445.9	0.31834	7.89	3.73	<1.00	0.46	44.684	43.861	
144202.82+433708.7	0.23146	8.18	3.54	<1.00	0.20	44.584	44.196	
144302.59+404525.1	0.24615	7.79	3.16	<1.00	0.11	44.231	43.673	
144645.93+403505.7	0.26709	8.35	11.37	<1.00	-0.13	44.999	44.073	
144751.79+505328.7	0.30565	7.56	1.41	<1.00	0.86	44.068	43.173	
145006.93+581456.9	0.31520	7.72	3.93	<1.00	0.33	44.714	43.755	
145254.74+395714.6	0.22528	7.31	0.79	<1.00	0.72	43.315	43.261	
145307.63+380319.6	0.26191	7.32	1.51	<1.00	0.61	44.695	43.416	
145439.39+465309.1	0.31645	8.45	1.95	<1.00	0.58	44.502	44.050	
145440.39+455008.5	0.32471	7.92	1.24	<1.00	0.86	44.017	43.115	
145817.42+455514.8	0.28575	7.79	3.12	<1.00	0.41	44.669	43.422	
145824.46+363119.5	0.24637	7.91	3.86	<1.00	0.20	44.384	43.474	
150049.15+343727.1	0.31794	7.52	1.95	<1.00	0.74	44.423	43.423	
150114.45+393927.7	0.23072	7.51	1.68	<1.00	0.49	43.950	43.074	
150155.24+563731.7	0.34338	7.54	2.78	<1.00	0.56	44.370	42.992	
150242.39+395208.5	0.34185	8.19	6.59	<1.00	0.22	44.894	43.976	
150437.67+541149.6	0.30519	8.39	3.17	<1.00	0.45	44.154	43.128	
151304.56+304204.2	0.34801	8.14	5.25	<1.00	0.25	44.886	43.997	
151557.75+302357.1	0.25171	7.60	0.94	<1.00	0.70	44.117	43.235	
151940.87+320157.9	0.30132	8.35	2.40	<1.00	0.58	44.558	43.896	
152153.81+594020.0	0.28599	8.10	2.41	<1.00	0.52	44.260	44.001	
152524.16+523205.7	0.32755	8.28	2.85	<1.00	0.59	44.107	43.125	
152628.19-003809.4	0.12333	7.40	0.57	<1.00	0.46	43.696	42.856	
153941.49+504255.8	0.20315	7.40	2.32	<1.00	0.14	44.766	43.774	

Object SDSS	Z (1)	$\log(M_{\text{BH}})$ (M_{\odot}) (2)	$L_{5100\text{\AA}}$ (10^{44} ergs s $^{-1}$) (4)	$F_{20\text{cm}}$ (mJy) (5)	R_i (6)	$\log(L_{0.1-2.4\text{keV}})$ (ergs s $^{-1}$) (7)	$\log(L_{2-10\text{keV}})$ (ergs s $^{-1}$) (8)
154004.24+355050.1	0.16361	8.16	1.31	<1.00	0.17	43.955	43.126
154344.27-001452.1	0.30189	8.16	3.01	<1.00	0.55	44.512	43.620
154348.62+401324.9	0.31826	8.60	4.93	2.75	0.84	44.885	44.080
155207.18+525347.0	0.33519	8.22	2.50	<1.00	0.65	44.313	43.891
155324.25+490726.9	0.25807	8.19	1.88	<1.00	0.59	44.297	43.414
155328.49+095102.0	0.19174	8.83	3.31	<1.00	-0.04	43.859	42.996
155837.88+282839.0	0.32336	8.54	8.80	<1.00	0.12	45.104	44.163
160518.50+375653.5	0.20087	7.55	1.64	<1.00	0.22	44.229	43.314
161118.82+291932.4	0.29364	8.29	1.68	<1.00	0.74	44.127	43.757
161849.25+442517.2	0.33521	8.46	7.97	3.09	0.63	44.636	44.047
162607.24+335915.2	0.20453	8.10	5.54	3.74	0.44	44.435	43.243
163111.28+404805.2	0.25751	8.03	2.18	<1.00	0.29	44.789	44.152
163631.58+461704.3	0.25170	7.64	1.55	<1.00	0.52	43.988	43.086
164343.24+405654.3	0.34375	7.04	1.34	<1.00	0.95	43.997	43.089
165338.69+634010.6	0.27897	7.54	1.91	<1.00	0.60	44.215	41.701
165352.82+384542.1	0.32302	7.64	1.74	<1.00	0.80	44.414	43.383
165737.30+604939.3	0.31316	7.67	1.48	<1.00	0.92	43.723	43.004
170231.06+324719.6	0.16333	7.76	4.13	1.52	-0.14	44.858	43.879
170302.88+191033.9	0.29045	7.86	4.44	<1.00	0.38	44.980	44.210
170525.54+194722.7	0.19392	8.42	1.33	<1.00	0.48	44.246	43.436
171013.42+334402.6	0.20749	8.67	5.73	4.60	0.40	44.590	43.730
171207.44+584754.5	0.26925	7.69	1.55	<1.00	0.64	44.551	43.446
171601.93+311213.8	0.11015	7.57	1.76	2.42	-0.06	44.301	43.686
171750.59+581514.0	0.31014	8.37	4.19	<1.00	0.42	44.428	43.708
171850.30+304201.6	0.28176	7.39	2.03	0.64	0.43	44.500	43.400
171902.29+593715.8	0.17852	7.49	1.06	<1.00	0.29	44.129	43.018
172711.81+632241.8	0.21736	9.17	2.71	1.23	0.23	44.482	43.647
173229.44+564811.3	0.30320	7.62	1.87	<1.00	0.67	44.309	43.555
211204.85-063535.2	0.20419	7.75	1.71	<1.00	0.34	44.239	43.890
213818.97+011222.4	0.34409	7.99	3.98	<1.00	0.48	44.584	44.853
215010.52-001000.7	0.33481	8.41	1.90	<1.00	0.80	44.719	43.833
215516.14+003250.8	0.27782	7.62	2.11	<1.00	0.59	44.648	43.498
231250.88+001719.0	0.25704	7.56	2.02	<1.00	0.56	44.201	43.319
233512.68-100040.3	0.24288	7.92	0.50	<1.00	0.83	44.095	43.977

Notes.—

Col. 1, object name in J2000.0. Col. 2, redshift given by the SDSS spectroscopic pipeline. Col. 3, logarithm of black hole mass . Col. 4, luminosity at 5100Å in unit of 10^{44} ergs s $^{-1}$. Col. 5, flux density at 20cm, '<1' represents that the object is covered yet not detected by FIRST. Col. 6, the radio-loudness. Col. 7–8, X-ray luminosity of 0.1–2.4 keV and 2–10 keV.