

HIGHEST WEIGHT MODULES OVER QUANTUM QUEER SUPERALGEBRA $U_q(\mathfrak{q}(n))$

DIMITAR GRANTCHAROV¹, JI HYE JUNG^{2,3}, SEOK-JIN KANG² AND MYUNGHO KIM^{2,3}

ABSTRACT. In this paper, we investigate the structure of highest weight modules over the quantum queer superalgebra $U_q(\mathfrak{q}(n))$. The key ingredients are the triangular decomposition of $U_q(\mathfrak{q}(n))$ and the classification of finite dimensional irreducible modules over quantum Clifford superalgebras. The main results we prove are the classical limit theorem and the complete reducibility theorem for $U_q(\mathfrak{q}(n))$ -modules in the category $\mathcal{O}_q^{\geq 0}$.

INTRODUCTION

Since its inception, the representation theory of Lie superalgebras has been known to be much more complicated than the corresponding theory of Lie algebras. One of the Lie superalgebra series attracts special attention due to its resemblance of the Lie algebra \mathfrak{gl}_n on the one hand and because of the unique properties of its structure and representations on the other. This is the so-called *queer* (or *strange*) Lie superalgebra $\mathfrak{q}(n)$ which consists of all endomorphisms of $\mathbb{C}^{n|n}$ that commute with an odd automorphism P of $\mathbb{C}^{n|n}$ such that $P^2 = \text{Id}$. The queer nature of $\mathfrak{q}(n)$ is partly due to the nonabelian structure of its Cartan subsuperalgebra \mathfrak{h} having a nontrivial odd part $\mathfrak{h}_{\bar{1}}$. Another unique property of $\mathfrak{q}(n)$ is that, although it has no invariant bilinear form, it admits an invariant odd bilinear form. Because of the nonabelian structure of \mathfrak{h} , the study of the highest weight modules of $\mathfrak{q}(n)$ requires some tools in addition to the standard technique. For example, the highest weight space \mathbf{v}_λ of an irreducible highest weight $\mathfrak{q}(n)$ -module $V(\lambda)$ has a Clifford module structure. The case when $V(\lambda)$ is a tensor module; i.e., a submodule of some tensor power $V^{\otimes r}$ of the natural $\mathfrak{q}(n)$ -module $V = \mathbb{C}^{n|n}$, was treated first by Sergeev in 1984. In [Se2] Sergeev established several important results, among which are the complete reducibility of $V^{\otimes r}$, a character formula of $V(\lambda)$, and an analog of the fundamental Schur-Weyl duality, often referred as Sergeev duality. The characters of all simple finite-dimensional $\mathfrak{q}(n)$ -modules have been found by Penkov and Serganova in 1996 (see [PS2] and [PS3]) via an algorithm using a supergeometric version of the Borel-Weil-Bott Theorem. In 2004 Brundan, [B], obtained the character formula of Penkov

¹This research was supported by a UT Arlington REP Grant.

²This research was supported by KRF Grant # 2007-341-C00001.

³This research was supported by BK21 Mathematical Sciences Division.

and Serganova using a different approach and formulated a conjecture for the characters of the irreducible modules in the category \mathcal{O} . Important results related to the simplicity of the highest weight $\mathfrak{q}(n)$ -modules were obtained recently by Gorelik in [G].

In this paper we initiate the study of highest weight representations of the quantum superalgebra $U_q(\mathfrak{q}(n))$. The aim of this paper is twofold. We want to study highest weight $U_q(\mathfrak{q}(n))$ -modules on the one hand, and to build the foundations of the crystal bases theory for the tensor modules of $U_q(\mathfrak{q}(n))$ on the other. The latter problem will be treated in a future work.

A quantum deformation of the universal enveloping algebra of $\mathfrak{q}(n)$ was constructed first by Olshanski in [O]. Olshanski's construction is a flat deformation of the universal enveloping algebra $U(\mathfrak{q}(n))$ of $\mathfrak{q}(n)$ and is a quantum enveloping superalgebra in the sense of Drinfeld ([Dr], §7). The idea in [O] is to apply a suitable modification of the procedure used by Faddeev, Reshetikhin, and Takhtajan in [RTF] – using an element S in $\text{End}(\mathbb{C}^{n|n})^{\otimes 2}$ that satisfies the quantum Yang-Baxter equation. However, as pointed out by Olshanski, the r -matrix $r \in \mathfrak{q}(n)^{\otimes 2}$ does not satisfy the classical Yang-Baxter equation. Thus no quantum analogue of $U(\mathfrak{q}(n))$ can be a quasi-triangular Hopf algebra.

In the present paper, based on the description of Olshanski, we give a presentation of $U_q(\mathfrak{q}(n))$ in terms of generators and relations so that the relations are quantum deformations of the relations of $\mathfrak{q}(n)$ obtained in [LS]. Using this presentation, we find a natural triangular decomposition of $U_q(\mathfrak{q}(n))$, and then introduce the notion of highest weight modules and Weyl modules. Similarly to the case of $\mathfrak{q}(n)$, in order to study highest weight modules, one has to describe the modules over the quantum Clifford superalgebra $\text{Cliff}_q(\lambda)$ for a weight λ of $\mathfrak{q}(n)$. These modules, as we show in Section 3, do not have the same structure as the ones over the classical Clifford superalgebra $\text{Cliff}(\lambda)$. For example, the irreducible modules over $\text{Cliff}_q(\lambda)$ are parity invariant for much larger set of weights λ , compared with the irreducibles over $\text{Cliff}(\lambda)$.

In the last two sections of the paper we focus on the category $\mathcal{O}_q^{\geq 0}$ of finite dimensional $U_q(\mathfrak{q}(n))$ -modules all whose weights are of the form $\lambda_1 \epsilon_1 + \dots + \lambda_n \epsilon_n$ ($\lambda_i \in \mathbb{Z}_{\geq 0}$). One of our main results is a classical limit theorem for the irreducible modules in $\mathcal{O}_q^{\geq 0}$. Due to the structure of the quantum Clifford superalgebra, the classical limit theorem is non-standard, as it is not true in general that the classical limit V^1 of an irreducible highest weight $U_q(\mathfrak{q}(n))$ -module $V^q(\lambda)$ is $V(\lambda)$. In fact, as we show in Section 5, if λ has even number of nonzero coordinates $\lambda_1 > \dots > \lambda_{2k}$, then $\text{ch } V^1 = 2 \text{ch } V(\lambda)$. The “queer” version of the classical limit theorems are Theorem 5.14 and Theorem 5.16. With the aid of the classical limit theorems we obtain another important result in the last section: the category $\mathcal{O}_q^{\geq 0}$ is semisimple.

The organization of the paper is as follows. In Section 1 we recall some definitions and basic results about $\mathfrak{q}(n)$. The realization of $U_q(\mathfrak{q}(n))$ and its triangular decomposition is provided in Section 2. Section 3 is devoted to the study of the quantum Clifford superalgebra and its modules. In Section 4 we introduce the notion of highest weight modules and Weyl modules. In particular, we show that every Weyl module $W^q(\lambda)$ has a unique irreducible quotient $V^q(\lambda)$. The classical limit theorem for the category $\mathcal{O}_q^{\geq 0}$ is proved in Section 5 and the complete reducibility of $U_q(\mathfrak{q}(n))$ -modules in $\mathcal{O}_q^{\geq 0}$ is established in the last section.

1. THE LIE SUPERALGEBRA $\mathfrak{q}(n)$ AND ITS REPRESENTATIONS

The ground field in this section will be \mathbb{C} . By $\mathbb{Z}_{\geq 0}$ and $\mathbb{Z}_{> 0}$ we denote the nonnegative integers and strictly positive integers, respectively. We set $\mathbb{Z}_2 = \mathbb{Z}/2\mathbb{Z}$. Every vector space $V = V_{\bar{0}} \oplus V_{\bar{1}}$ over \mathbb{C} is \mathbb{Z}_2 -graded with even part $V_{\bar{0}}$ and odd part $V_{\bar{1}}$. We will write $\dim V = m|n$ if $\dim V_{\bar{0}} = m$ and $\dim V_{\bar{1}} = n$. By Π we denote the parity change functor; i.e., ΠV is a vector space for which $\Pi V_{\bar{0}} = V_{\bar{1}}$ and $\Pi V_{\bar{1}} = V_{\bar{0}}$. The direct sum of r copies of a vector space V will be written as $V^{\oplus r}$.

The Lie subsuperalgebra $\mathfrak{g} = \mathfrak{q}(n)$ of $\mathfrak{gl}(n|n)$ is defined in matrix form by

$$\mathfrak{g} = \mathfrak{q}(n) := \left\{ \left(\begin{array}{cc} A & B \\ B & A \end{array} \right) \mid A, B \in \mathfrak{gl}_n \right\}.$$

By definition, a subsuperalgebra $\mathfrak{h} = \mathfrak{h}_{\bar{0}} \oplus \mathfrak{h}_{\bar{1}}$ of \mathfrak{g} is a *Cartan subsuperalgebra*, if it is a self-normalizing nilpotent subsuperalgebra. Every such \mathfrak{h} has a nontrivial odd part $\mathfrak{h}_{\bar{1}}$. We fix \mathfrak{h} to be the *standard* Cartan subsuperalgebra, namely the one for which $\mathfrak{h}_{\bar{0}}$ has a basis $\{k_1, \dots, k_n\}$ and $\mathfrak{h}_{\bar{1}}$ has a basis $\{k_{\bar{1}}, \dots, k_{\bar{n}}\}$, where $k_i := \begin{pmatrix} E_{i,i} & 0 \\ 0 & E_{i,i} \end{pmatrix}$, $k_{\bar{i}} := \begin{pmatrix} 0 & E_{i,i} \\ E_{i,i} & 0 \end{pmatrix}$ and $E_{i,j}$ is the $n \times n$ matrix having 1 in the (i, j) position and 0 elsewhere. One should note that all Cartan subsuperalgebras of \mathfrak{g} are conjugate to \mathfrak{h} . Let $\{\epsilon_1, \dots, \epsilon_n\}$ be the basis of $\mathfrak{h}_{\bar{0}}^*$ dual to $\{k_1, \dots, k_n\}$. We denote $k_i - k_{i+1}$ by h_i for $i = 1, 2, \dots, n-1$. The root system $\Delta = \Delta_{\bar{0}} \cup \Delta_{\bar{1}}$ of \mathfrak{g} has identical even and odd parts. Namely, $\Delta_{\bar{0}} = \Delta_{\bar{1}} = \{\epsilon_i - \epsilon_j \mid 1 < i \neq j < n\}$. In particular, the root space decomposition $\mathfrak{g} = \bigoplus_{\alpha \in \Delta} \mathfrak{g}_{\alpha}$ has the property that \mathfrak{g}_{α} has dimension 1|1 for every $\alpha \in \Delta$. Set $\alpha_i := \epsilon_i - \epsilon_{i+1}$. Let $Q = \bigoplus_{i=1}^{n-1} \mathbb{Z}\alpha_i$ be the root lattice and $Q_+ = \sum_{i=1}^{n-1} \mathbb{Z}_{\geq 0}\alpha_i$ be the positive root lattice. The notation $Q_- = -Q_+$ will also be used. There is a partial ordering on $\mathfrak{h}_{\bar{0}}^*$ defined by $\lambda \geq \mu$ if and only if $\lambda - \mu \in Q_+$ for $\lambda, \mu \in \mathfrak{h}_{\bar{0}}^*$. The root space \mathfrak{g}_{α_i} is spanned by $e_i := \begin{pmatrix} E_{i,i+1} & 0 \\ 0 & E_{i,i+1} \end{pmatrix}$ and $e_{\bar{i}} := \begin{pmatrix} 0 & E_{i,i+1} \\ E_{i,i+1} & 0 \end{pmatrix}$, while $\mathfrak{g}_{-\alpha_i}$ is spanned

by $f_i := \begin{pmatrix} E_{i+1,i} & 0 \\ 0 & E_{i+1,i} \end{pmatrix}$ and $f_{\bar{i}} := \begin{pmatrix} 0 & E_{i+1,i} \\ E_{i+1,i} & 0 \end{pmatrix}$. Let $P := \bigoplus_{i=1}^n \mathbb{Z}\epsilon_i$ be the weight lattice of \mathfrak{g} and denote by $P^\vee := \bigoplus_{i=1}^n \mathbb{Z}k_i$ the dual weight lattice.

Let $I := \{1, 2, \dots, n-1\}$ and $J := \{1, 2, \dots, n\}$.

Proposition 1.1. [LS] *The Lie superalgebra \mathfrak{g} is generated by the elements $e_i, e_{\bar{i}}, f_i, f_{\bar{i}}$ ($i \in I$), $\mathfrak{h}_{\bar{0}}$ and $k_{\bar{l}}$ ($l \in J$) with the following defining relations:*

$$\begin{aligned}
[h, h'] &= 0 \quad \text{for } h, h' \in \mathfrak{h}_{\bar{0}}, \\
[h, e_i] &= \alpha_i(h)e_i, \quad [h, e_{\bar{i}}] = \alpha_i(h)e_{\bar{i}} \quad \text{for } h \in \mathfrak{h}_{\bar{0}}, \quad i \in I, \\
[h, f_i] &= -\alpha_i(h)f_i, \quad [h, f_{\bar{i}}] = -\alpha_i(h)f_{\bar{i}} \quad \text{for } h \in \mathfrak{h}_{\bar{0}}, \quad i \in I, \\
[h, k_{\bar{l}}] &= 0 \quad \text{for } h \in \mathfrak{h}_{\bar{0}}, \quad l \in J, \\
[e_i, f_j] &= \delta_{ij}(k_i - k_{i+1}), \quad [e_i, f_{\bar{j}}] = \delta_{ij}(k_{\bar{i}} - k_{\overline{i+1}}) \quad \text{for } i, j \in I, \\
[e_{\bar{i}}, f_j] &= \delta_{ij}(k_{\bar{i}} - k_{\overline{i+1}}), \quad [k_{\bar{l}}, e_i] = \alpha_i(k_l)e_{\bar{i}} \quad \text{for } i, j \in I, \quad l \in J, \\
[k_{\bar{l}}, f_i] &= -\alpha_i(k_l)f_{\bar{i}}, \quad [e_{\bar{i}}, f_{\bar{j}}] = \delta_{ij}(k_i + k_{i+1}) \quad \text{for } i, j \in I, \quad l \in J, \\
[k_{\bar{l}}, e_{\bar{i}}] &= \begin{cases} e_i & \text{if } l = i, i+1 \\ 0 & \text{otherwise} \end{cases} \quad \text{for } i \in I, \quad l \in J, \\
[k_{\bar{l}}, f_{\bar{i}}] &= \begin{cases} f_i & \text{if } l = i, i+1 \\ 0 & \text{otherwise} \end{cases} \quad \text{for } i \in I, \quad l \in J, \\
[e_i, e_{\bar{j}}] &= [e_{\bar{i}}, e_{\bar{j}}] = [f_i, f_{\bar{j}}] = [f_{\bar{i}}, f_{\bar{j}}] = 0 \quad \text{for } i, j \in I, \quad |i - j| \neq 1, \\
[e_i, e_j] &= [f_i, f_j] = 0 \quad \text{for } i, j \in I, \quad |i - j| > 1, \\
[e_i, e_{i+1}] &= [e_{\bar{i}}, e_{\overline{i+1}}], \quad [e_i, e_{\overline{i+1}}] = [e_{\bar{i}}, e_{i+1}], \\
[f_{i+1}, f_i] &= [f_{\overline{i+1}}, f_{\bar{i}}], \quad [f_{i+1}, f_{\bar{i}}] = [f_{\overline{i+1}}, f_i], \\
[k_{\bar{i}}, k_{\bar{j}}] &= \delta_{ij}2k_i \quad \text{for } i, j \in J, \\
[e_i, [e_i, e_j]] &= [e_{\bar{i}}, [e_i, e_j]] = 0 \quad \text{for } i, j \in I, \quad |i - j| = 1, \\
[f_i, [f_i, f_j]] &= [f_{\bar{i}}, [f_i, f_j]] = 0 \quad \text{for } i, j \in I, \quad |i - j| = 1.
\end{aligned}$$

Remark. We modified the relations given in [LS]. More precisely, we replaced the relations

$$\begin{aligned}
(1.1) \quad [e_{\bar{i}}, [e_i, e_{\bar{j}}]] &= 0 \quad \text{for } i, j \in I, \quad |i - j| = 1, \\
[f_{\bar{i}}, [f_i, f_{\bar{j}}]] &= 0 \quad \text{for } i, j \in I, \quad |i - j| = 1
\end{aligned}$$

by

$$(1.2) \quad \begin{aligned} [e_i, e_{i+1}] &= [e_{\bar{i}}, e_{\overline{i+1}}], [e_i, e_{\overline{i+1}}] = [e_{\bar{i}}, e_{i+1}], \\ [f_{i+1}, f_i] &= [f_{\overline{i+1}}, f_{\bar{i}}], [f_{i+1}, f_{\bar{i}}] = [f_{\overline{i+1}}, f_{\bar{i}}]. \end{aligned}$$

Since (1.1) can be derived from (1.2) (and other ones), we can easily see that these two presentations are equivalent.

The universal enveloping algebra $U(\mathfrak{g})$ is obtained from the tensor algebra $T(\mathfrak{g})$ by factoring out by the ideal generated by the elements $[u, v] - u \otimes v + (-1)^{\alpha\beta} v \otimes u$, where $\alpha, \beta \in \mathbb{Z}_2$, $u \in \mathfrak{g}_\alpha$, $v \in \mathfrak{g}_\beta$. Let U^+ (respectively, U^0 and U^-) be the subalgebra of $U(\mathfrak{g})$ generated by the elements $e_i, e_{\bar{i}}$ ($i \in I$) (respectively, by $k_i, k_{\bar{i}}$ ($i \in J$) and by $f_i, f_{\bar{i}}$ ($i \in I$)). By the Poincaré-Birkhoff-Witt theorem, the universal enveloping algebra has the triangular decomposition:

$$(1.3) \quad U(\mathfrak{g}) \cong U^- \otimes U^0 \otimes U^+.$$

A \mathfrak{g} -module V is called a *weight module* if it admits a *weight space decomposition*

$$V = \bigoplus_{\mu \in \mathfrak{h}_0^*} V_\mu, \text{ where } V_\mu = \{v \in V \mid hv = \mu(h)v \text{ for all } h \in \mathfrak{h}_0\}.$$

For a weight \mathfrak{g} -module M denote by $\text{wt}(M)$ the set of weights $\lambda \in \mathfrak{h}_0^*$ for which $M_\lambda \neq 0$. Every submodule of a weight module is also a weight module. If $\dim_{\mathbb{C}} V_\mu < \infty$ for all $\mu \in \mathfrak{h}_0^*$, the *character* of V is defined to be

$$\text{ch } V = \sum_{\mu \in \mathfrak{h}_0^*} (\dim_{\mathbb{C}} V_\mu) e^\mu,$$

where e^μ are formal basis elements of the group algebra $\mathbb{C}[\mathfrak{h}_0^*]$ with the multiplication given by $e^\lambda e^\mu = e^{\lambda+\mu}$ for all $\lambda, \mu \in \mathfrak{h}_0^*$.

Denote by \mathfrak{b}_+ the *standard Borel subsuperalgebra* of \mathfrak{g} generated by $k_l, k_{\bar{l}}$ ($l \in J$) and $e_i, e_{\bar{i}}$ ($i \in I$). A weight module V is called a *highest weight module* if it is generated over \mathfrak{g} by a finite dimensional irreducible \mathfrak{b}_+ -submodule (see [PS1, Definition 4]).

Proposition 1.2. [P] *Let \mathfrak{v} be a finite dimensional irreducible \mathbb{Z}_2 -graded \mathfrak{b}_+ -module.*

- (1) *The maximal nilpotent subsuperalgebra \mathfrak{n} of \mathfrak{b}_+ acts on \mathfrak{v} trivially.*
- (2) *For any weight $\mu \in \mathfrak{h}_0^*$, consider the symmetric bilinear form $F_\mu(u, v) := \mu([u, v])$ on $\mathfrak{h}_{\bar{1}}$ and let $\text{Cliff}(\mu)$ be the Clifford superalgebra of the quadratic space $(\mathfrak{h}_{\bar{1}}, F_\mu)$. Then there exists a unique weight $\lambda \in \mathfrak{h}_0^*$ such that \mathfrak{v} is endowed with a canonical \mathbb{Z}_2 -graded $\text{Cliff}(\lambda)$ -module structure and \mathfrak{v} is determined by λ up to Π .*
- (3) *\mathfrak{h}_0 acts on \mathfrak{v} by the weight λ determined in (2).*

From the above proposition, we know that the dimension of the highest weight space of a highest weight \mathfrak{g} -module with highest weight λ is the same as the dimension of an irreducible $\text{Cliff}(\lambda)$ -module. On the other hand all irreducible $\text{Cliff}(\lambda)$ -modules have the same dimension (see, for example, [ABS, Table 2]). Thus the dimension of the highest weight space is constant for all highest weight modules with highest weight λ .

Definition 1.3. Let $\mathbf{v}(\lambda)$ be the irreducible \mathfrak{b}_+ -module determined by λ up to Π . The *Weyl module* $W(\lambda)$ of \mathfrak{g} with highest weight λ is defined to be

$$W(\lambda) := U(\mathfrak{g}) \otimes_{U(\mathfrak{b}_+)} \mathbf{v}(\lambda).$$

Note that the structure of $W(\lambda)$ is determined by λ up to Π .

Remark. One may define the *Verma module* corresponding to λ by $M(\lambda) := U(\mathfrak{g}) \otimes_{U(\mathfrak{b}_+)} \text{Cliff}(\lambda)$. Since the Verma modules are not highest weight modules, they will not be considered in this paper.

We will denote by Λ_0^+ and Λ^+ the set of \mathfrak{gl}_n -dominant integral weights and the set of \mathfrak{g} -dominant integral weights, respectively. These are given by

$$\begin{aligned} \Lambda_0^+ &:= \{\lambda_1 \epsilon_1 + \cdots + \lambda_n \epsilon_n \in \mathfrak{h}_0^* \mid \lambda_i - \lambda_{i+1} \in \mathbb{Z}_{\geq 0} \text{ for all } i \in I\} \\ \Lambda^+ &:= \{\lambda_1 \epsilon_1 + \cdots + \lambda_n \epsilon_n \in \Lambda_0^+ \mid \lambda_i = \lambda_{i+1} \Rightarrow \lambda_i = \lambda_{i+1} = 0 \text{ for all } i \in I\}. \end{aligned}$$

Proposition 1.4. [P]

- (1) For any weight λ , $W(\lambda)$ has a unique maximal submodule $N(\lambda)$.
- (2) For each finite dimensional irreducible \mathfrak{g} -module V , there exists a unique weight $\lambda \in \Lambda_0^+$ such that V is a homomorphic image of $W(\lambda)$.
- (3) $V(\lambda) := W(\lambda)/N(\lambda)$ is finite dimensional if and only if $\lambda \in \Lambda^+$.

Now we restrict our attention to the following subcategory of the category of finite dimensional \mathfrak{g} -modules.

Definition 1.5. Set $P_{\geq 0} := \{\lambda = \lambda_1 \epsilon_1 + \cdots + \lambda_n \epsilon_n \in P \mid \lambda_j \geq 0 \text{ for all } j = 1, \dots, n\}$. The category $\mathcal{O}^{\geq 0}$ consists of finite dimensional $U(\mathfrak{g})$ -modules M with weight space decomposition $M = \bigoplus_{\mu \in P} M_\mu$ satisfying (i) $\text{wt}(M) \subset P_{\geq 0}$, (ii) $k_i|_{M_\mu} = 0$ for $\mu \in P_{\geq 0}$ and $i \in \{1, \dots, n\}$ such that $\langle k_i, \mu \rangle = 0$.

Remark. The reason we impose condition (ii) in Definition 1.5 is that we want the category $\mathcal{O}^{\geq 0}$ and its q -analog $\mathcal{O}_q^{\geq 0}$ (see Definition 6.1) to be completely reducible. If M is a weight module, then one can show that the $\text{Cliff}(\mu)$ -module M_μ is completely reducible if and only if (ii) is satisfied. This follows from the complete reducibility criterion for $\text{Cliff}(\mu)$ -modules. For the q -version of this criterion see Lemma 3.1 and Corollary 3.9. In view of the above,

one can verify that Definition 1.5 is equivalent to the one where (ii) is replaced by: (ii)' M is completely reducible as an \mathfrak{h} -module. The same applies to Definition 6.1 where (ii) can be replaced by the condition that M is a completely reducible U_q^0 -module.

One easily checks that $\mathcal{O}^{\geq 0}$ is closed under finite direct sum, tensor product and taking submodules and quotient modules. As Sergeev showed, [Se2], for any $\lambda \in \Lambda^+ \cap P_{\geq 0}$, $V(\lambda)$ is a submodule of $V^{\otimes |\lambda|}$ where V is the natural representation of \mathfrak{g} and $|(\lambda_1, \dots, \lambda_n)| = \lambda_1 + \dots + \lambda_n$. This, together with the properties of the \mathfrak{gl}_n -module $V(\lambda)$ (see, for example, [HK, Theorem 7.2.3]), implies the following proposition.

Proposition 1.6. *For each $\lambda \in \Lambda^+ \cap P_{\geq 0}$, $V(\lambda)$ is an irreducible $U(\mathfrak{g})$ -module in the category $\mathcal{O}^{\geq 0}$. Conversely, every irreducible $U(\mathfrak{g})$ -module in the category $\mathcal{O}^{\geq 0}$ has the form $V(\lambda)$ for some $\lambda \in \Lambda^+ \cap P_{\geq 0}$.*

In [Se1], Sergeev has presented an explicit set of generators of $Z = \mathcal{Z}(U(\mathfrak{g}))$, the center of $U(\mathfrak{g})$, and showed that each Weyl module $W(\lambda)$ ($\lambda \in \mathfrak{h}_0^*$) admits a central character. Let $\chi_\lambda \in \text{Hom}_{\mathbb{C}}(Z, \mathbb{C})$ be the central character afforded by $W(\lambda)$; i.e., every element $z \in Z$ acts on $W(\lambda)$ as scalar multiplication by $\chi_\lambda(z)$. Following [B, (2.12)], to each weight $\lambda = \lambda_1 \epsilon_1 + \dots + \lambda_n \epsilon_n \in P$, one can assign a formal symbol

$$\delta(\lambda) := \delta_{\lambda_1} + \dots + \delta_{\lambda_n}$$

such that $\delta_0 = 0$ and $\delta_{-i} = -\delta_i$.

Proposition 1.7. [B, Theorem 4.19], [PS2, Proposition 1.1] *For $\lambda, \mu \in P$, $\chi_\lambda = \chi_\mu$ if and only if $\delta(\lambda) = \delta(\mu)$.*

The following proposition will be very useful in Section 5.

Proposition 1.8. *Let V be a finite dimensional highest weight module over \mathfrak{g} with highest weight $\lambda \in \Lambda^+ \cap P_{\geq 0}$. Then V is isomorphic to an irreducible highest weight module $V(\lambda)$.*

Proof. If V is reducible, since it is finite dimensional, it contains a nonzero proper irreducible submodule W . Then W is isomorphic to an irreducible highest weight module $V(\mu)$ for some weight $\mu \in \Lambda^+ \cap P_{\geq 0}$ by Proposition 1.4. We know that $\mu \not\leq \lambda$ and $\chi_\lambda = \chi_\mu$. But, by Proposition 1.7, $\delta(\lambda) = \delta(\mu)$. Since $\lambda, \mu \in \Lambda^+ \cap P_{\geq 0}$, we have $\lambda = \mu$, which is a contradiction. Thus V is irreducible and by Proposition 1.4, it must be isomorphic to the irreducible highest weight module $V(\lambda)$ up to Π . \square

The next proposition gives a sufficient condition for the finite dimensionality of a highest weight \mathfrak{g} -module.

Proposition 1.9. *Let V be a highest weight module over \mathfrak{g} with highest weight $\lambda \in \Lambda^+$. If $f_i^{\lambda(h_i)+1}v = 0$ for all $v \in V_\lambda$ and $i \in I$, then V is finite dimensional.*

Proof. Let $\{x_1, x_2, \dots, x_r\}$ and $\{y_1, y_2, \dots, y_r\}$ be bases of \mathfrak{g}_0 and $\mathfrak{g}_{\bar{1}}$, respectively. Then by the Poincaré-Birkhoff-Witt theorem, $U(\mathfrak{g})$ has a basis consisting of elements of the form $y_1^{\epsilon_1} y_2^{\epsilon_2} \cdots y_r^{\epsilon_r} x_1^{n_1} x_2^{n_2} \cdots x_r^{n_r}$ where $\epsilon_j = 0$ or 1 and $n_j \in \mathbb{N} \cup \{0\}$. Because $\{y_1^{\epsilon_1} y_2^{\epsilon_2} \cdots y_r^{\epsilon_r} \mid \epsilon_j = 0, 1\}$ is a finite set, it is enough to show that $U(\mathfrak{g}_0)V_\lambda$ is finite dimensional. For any $v \in V_\lambda$, we know that $U(\mathfrak{g}_0)v$ is a highest weight module over \mathfrak{g}_0 with highest weight λ satisfying $f_i^{\lambda(h_i)+1}v = 0$ for all $i \in I$. Thus it is finite dimensional. Since $U(\mathfrak{g})V_\lambda \subset \sum_{v \in V_\lambda} U(\mathfrak{g}_0)v$, we have the desired result. \square

We say that a weight $\lambda = \lambda_1 \epsilon_1 + \cdots + \lambda_n \epsilon_n \in \mathfrak{h}_0^*$ is α -typical if $\alpha = \epsilon_i - \epsilon_j$ and $\lambda_i + \lambda_j \neq 0$. In [Se2], Sergeev proved the following character formula for $V(\lambda)$ ($\lambda \in \Lambda^+ \cap P_{\geq 0}$):

$$(1.4) \quad \text{ch } V(\lambda) = \frac{\dim \mathbf{v}_\lambda}{D} \sum_{w \in W} \text{sgn } w \left(e^{\lambda + \rho_0} \prod_{\substack{\alpha \in \Delta_0^+, \\ \lambda \text{ is } \alpha\text{-typical}}} (1 + e^{-\alpha}) \right),$$

where \mathbf{v}_λ is an irreducible Cliff(λ)-module, W is the Weyl group of $\mathfrak{g}_0 = \mathfrak{gl}_n$, $\rho_0 = \frac{1}{2} \sum_{\alpha \in \Delta_0^+} \alpha$ and $D = \sum_{w \in W} \text{sgn } w e^{w(\rho_0)}$ is the Weyl denominator. In [PS2], the formula (1.4) is called the *generic character formula* and an explicit algorithm for computing the character of an arbitrary finite dimensional irreducible \mathfrak{g} -module is presented.

2. THE QUANTUM SUPERALGEBRA $U_q(\mathfrak{q}(n))$

In [O], Olshanski constructed the quantum deformation $U_q(\mathfrak{q}(n))$ of the universal enveloping algebra of $\mathfrak{q}(n)$. The quantum superalgebra $U_q(\mathfrak{q}(n))$ is defined to be the associative algebra over $\mathbb{C}(q)$ generated by L_{ij} , $i \leq j$, with defining relations

$$(2.1) \quad \begin{aligned} L_{ii}L_{-i,-i} &= L_{-i,-i}L_{ii} = 1, \\ (-1)^{p(i,j)p(k,l)} q^{\varphi(j,l)} L_{ij}L_{kl} &+ \{k \leq j < l\} \theta(i,j,k)(q - q^{-1})L_{il}L_{kj} \\ &+ \{i \leq -l < j \leq -k\} \theta(-i,-j,k)(q - q^{-1})L_{i,-l}L_{k,-j} \\ &= q^{\varphi(i,k)} L_{kl}L_{ij} + \{k < i \leq l\} \theta(i,j,k)(q - q^{-1})L_{il}L_{kj} \\ &+ \{-l \leq i < -k \leq j\} \theta(-i,-j,k)(q - q^{-1})L_{-i,l}L_{-k,j}, \end{aligned}$$

where $\varphi(i,j) = \delta_{|i|,|j|} \text{sgn}(j)$, $\theta(i,j,k) = \text{sgn}(\text{sgn}(i) + \text{sgn}(j) + \text{sgn}(k))$, $p(i,j) = \begin{cases} 0 & \text{if } ij > 0 \\ 1 & \text{if } ij < 0, \end{cases}$ for any indices $i \leq j$, $k \leq l$ in $\{\pm 1, \dots, \pm n\}$ and the symbol $\{\dots\}$ (the dots stand for some inequalities) is equal to 1 if all of these inequalities are fulfilled and 0 otherwise.

Following [O, Remark 7.3], we consider the set of generators of $U_q(\mathfrak{g}) = U_q(\mathfrak{q}(n))$ as follows:

$$(2.2) \quad \begin{aligned} q^{k_i} &:= L_{i,i}, \quad q^{-k_i} := L_{-i,-i}, \quad e_i := -\frac{1}{q-q^{-1}}L_{-i-1,-i}, \quad f_i := \frac{1}{q-q^{-1}}L_{i,i+1}, \\ e_{\bar{i}} &:= -\frac{1}{q-q^{-1}}L_{-i-1,i}, \quad f_{\bar{i}} := -\frac{1}{q-q^{-1}}L_{-i,i+1}, \quad k_{\bar{i}} := -\frac{1}{q-q^{-1}}L_{-i,i}. \end{aligned}$$

Our first main result is the following presentation of $U_q(\mathfrak{g})$.

Theorem 2.1. *The quantum superalgebra $U_q(\mathfrak{g})$ is isomorphic to the unital associative algebra over $\mathbb{C}(q)$ generated by the elements $e_i, f_i, e_{\bar{i}}, f_{\bar{i}}$ ($i = 1, \dots, n-1$), $k_{\bar{l}}$ ($l = 1, \dots, n$), and q^h ($h \in P^\vee$), satisfying the following relations*

$$(2.3) \quad \begin{aligned} q^0 &= 1, q^{h_1+h_2} = q^{h_1}q^{h_2} \text{ for } h_1, h_2 \in P^\vee, \\ q^h e_i q^{-h} &= q^{\alpha_i(h)} e_i, q^h f_i q^{-h} = q^{-\alpha_i(h)} f_i \text{ for } h \in P^\vee \\ q^h k_{\bar{i}} q^{-h} &= k_{\bar{i}}, q^h e_{\bar{i}} q^{-h} = q^{\alpha_{\bar{i}}(h)} e_{\bar{i}}, q^h f_{\bar{i}} q^{-h} = q^{-\alpha_{\bar{i}}(h)} f_{\bar{i}} \text{ for } h \in P^\vee \\ e_i f_i - f_i e_i &= \frac{1}{q-q^{-1}} \left(q^{k_i-k_{i+1}} - q^{-k_i+k_{i+1}} \right), \\ qe_{i+1}f_i - f_i e_{i+1} &= e_i f_{i+1} - q f_{i+1} e_i = e_i f_j - f_j e_i = 0 \text{ if } |i-j| > 1, \\ e_i f_{\bar{i}} - f_{\bar{i}} e_i &= q^{-k_{i+1}} k_{\bar{i}} - k_{\bar{i}+1} q^{-k_i}, \\ qe_{i+1}f_{\bar{i}} - f_{\bar{i}} e_{i+1} &= e_i f_{\bar{i}+1} - q f_{\bar{i}+1} e_i = e_i f_{\bar{j}} - f_{\bar{j}} e_i = 0 \text{ if } |i-j| > 1, \\ e_{\bar{i}} f_i - f_i e_{\bar{i}} &= q^{k_{i+1}} k_{\bar{i}} - k_{\bar{i}+1} q^{k_i}, \\ qe_{\bar{i}+1}f_i - f_i e_{\bar{i}+1} &= e_{\bar{i}} f_{i+1} - q f_{i+1} e_{\bar{i}} = e_{\bar{i}} f_j - f_j e_{\bar{i}} = 0 \text{ if } |i-j| > 1, \\ k_{\bar{i}} e_i - q e_i k_{\bar{i}} &= e_{\bar{i}} q^{-k_i}, q k_{\bar{i}} e_{i-1} - e_{i-1} k_{\bar{i}} = -q^{-k_i} e_{\bar{i}-1}, \\ k_{\bar{i}} e_j - e_j k_{\bar{i}} &= 0 \text{ for } j \neq i \text{ and } j \neq i-1, \\ k_{\bar{i}} f_i - q f_i k_{\bar{i}} &= -f_{\bar{i}} q^{k_i}, q k_{\bar{i}} f_{i-1} - f_{i-1} k_{\bar{i}} = q^{k_i} f_{\bar{i}-1}, \\ k_{\bar{i}} f_j - f_j k_{\bar{i}} &= 0 \text{ for } j \neq i \text{ and } j \neq i-1, \\ k_{\bar{i}}^2 &= \frac{q^{2k_i} - q^{-2k_i}}{q^2 - q^{-2}}, k_{\bar{i}} k_{\bar{j}} = -k_{\bar{j}} k_{\bar{i}} \text{ for } i \neq j, \\ e_{\bar{i}} f_{\bar{i}} + f_{\bar{i}} e_{\bar{i}} &= \frac{q^{k_i+k_{i+1}} - q^{-k_i-k_{i+1}}}{q-q^{-1}} + (q-q^{-1}) k_{\bar{i}} k_{\bar{i}+1}, \\ qe_{\bar{i}+1}f_{\bar{i}} + f_{\bar{i}} e_{\bar{i}+1} &= e_{\bar{i}} f_{\bar{i}+1} + q f_{\bar{i}+1} e_{\bar{i}} = e_{\bar{i}} f_{\bar{j}} + f_{\bar{j}} e_{\bar{i}} = 0 \text{ if } |i-j| > 1, \\ k_{\bar{i}} e_{\bar{i}} + q e_{\bar{i}} k_{\bar{i}} &= e_i q^{-k_i}, q k_{\bar{i}} e_{\bar{i}-1} + e_{\bar{i}-1} k_{\bar{i}} = q^{-k_i} e_{i-1}, \\ k_{\bar{i}} e_{\bar{j}} + e_{\bar{j}} k_{\bar{i}} &= 0 \text{ for } j \neq i \text{ and } j \neq i-1, \\ k_{\bar{i}} f_{\bar{i}} + q f_{\bar{i}} k_{\bar{i}} &= f_i q^{k_i}, q k_{\bar{i}} f_{\bar{i}-1} + f_{\bar{i}-1} k_{\bar{i}} = q^{k_i} f_{i-1}, \end{aligned}$$

$$\begin{aligned}
& k_{\bar{i}}f_{\bar{j}} + f_{\bar{j}}k_{\bar{i}} = 0 \text{ for } j \neq i \text{ and } j \neq i-1, \\
& e_i^2 = -\frac{q-q^{-1}}{q+q^{-1}}e_i^2, \quad f_i^2 = \frac{q-q^{-1}}{q+q^{-1}}f_i^2, \\
& e_i e_j - e_j e_i = f_i f_j - f_j f_i = e_{\bar{i}} e_{\bar{j}} + e_{\bar{j}} e_{\bar{i}} = f_{\bar{i}} f_{\bar{j}} + f_{\bar{j}} f_{\bar{i}} = 0 \text{ if } |i-j| > 1, \\
& e_i e_{\bar{j}} - e_{\bar{j}} e_i = f_i f_{\bar{j}} - f_{\bar{j}} f_i = 0 \text{ if } |i-j| \neq 1, \\
& e_i e_{i+1} - e_{i+1} e_i = e_{\bar{i}} e_{\overline{i+1}} + e_{\overline{i+1}} e_{\bar{i}}, \quad f_{i+1} f_i - f_i f_{i+1} = f_i f_{\overline{i+1}} + f_{\overline{i+1}} f_i, \\
& e_i e_{\overline{i+1}} - e_{\overline{i+1}} e_i = e_{\bar{i}} e_{i+1} - e_{i+1} e_{\bar{i}}, \quad f_{\overline{i+1}} f_i - f_i f_{\overline{i+1}} = f_{i+1} f_{\bar{i}} - f_{\bar{i}} f_{i+1}, \\
& q e_i^2 e_{i+1} - (q+q^{-1}) e_i e_{i+1} e_i + q^{-1} e_{i+1} e_i^2 = 0, \\
& q f_i^2 f_{i+1} - (q+q^{-1}) f_i f_{i+1} f_i + q^{-1} f_{i+1} f_i^2 = 0, \\
& q e_i e_{i+1}^2 - (q+q^{-1}) e_{i+1} e_i e_{i+1} + q^{-1} e_{i+1}^2 e_i = 0, \\
& q f_i f_{i+1}^2 - (q+q^{-1}) f_{i+1} f_i f_{i+1} + q^{-1} f_{i+1}^2 f_i = 0, \\
& q e_i^2 e_{\overline{i+1}} - (q+q^{-1}) e_i e_{\overline{i+1}} e_i + q^{-1} e_{\overline{i+1}} e_i^2 = 0, \\
& q f_i^2 f_{\overline{i+1}} - (q+q^{-1}) f_i f_{\overline{i+1}} f_i + q^{-1} f_{\overline{i+1}} f_i^2 = 0, \\
& q e_{\bar{i}} e_{i+1}^2 - (q+q^{-1}) e_{i+1} e_{\bar{i}} e_{i+1} + q^{-1} e_{i+1}^2 e_{\bar{i}} = 0, \\
& q f_{\bar{i}} f_{i+1}^2 - (q+q^{-1}) f_{i+1} f_{\bar{i}} f_{i+1} + q^{-1} f_{i+1}^2 f_{\bar{i}} = 0.
\end{aligned}$$

Proof. Let U be the unital associative algebra over $\mathbb{C}(q)$ generated by the elements $e_i, f_i, e_{\bar{i}}, f_{\bar{i}}$ ($i = 1, \dots, n-1$), $k_{\bar{l}}$ ($l = 1, \dots, n$), and q^h ($h \in P^\vee$) with defining relations given in (2.3). Using (2.1) and (2.2), the relations in (2.3) can be derived easily. Thus there is a well-defined algebra homomorphism $\phi : U \rightarrow U_q(\mathfrak{g})$.

From the relation (2.1), we obtain

$$\begin{aligned}
(2.4) \quad L_{i,i+j} &= (q-q^{-1})q^{-\sum_{h=1}^{j-1} k_{i+h}} \prod_{h=1}^{j-1} \text{ad } f_{i+h}(f_i), \\
L_{-i,i+j} &= -(q-q^{-1})q^{-\sum_{h=1}^{j-1} k_{i+h}} \prod_{h=1}^{j-1} \text{ad } f_{i+h}(f_{\bar{i}}), \\
L_{-i-j,i} &= (-1)^j (q-q^{-1})q^{\sum_{h=1}^{j-1} k_{i+h}} \prod_{h=1}^{j-1} \text{ad } e_{i+h}(e_{\bar{i}}), \\
L_{-i-j,-i} &= (-1)^j (q-q^{-1})q^{\sum_{h=1}^{j-1} k_{i+h}} \prod_{h=1}^{j-1} \text{ad } e_{i+h}(e_i),
\end{aligned}$$

where $\text{ad } b_i(b_j) := b_i b_j - b_j b_i$, $\prod_{h=1}^j \text{ad } b_{i+h}(b_i) := \text{ad } b_{i+j} \cdots \text{ad } b_{i+1}(b_i)$ and $\prod_{h=1}^0 \text{ad } b_{i+h}(b_i) = b_i$ for $b_i = e_i, e_{\bar{i}}, f_i, f_{\bar{i}}$ ($i = 1, \dots, n-1, j > 0$). It follows that the homomorphism ϕ must be surjective.

It remains to prove ϕ is injective. For this purpose, we will show that the relations in (2.1) can be derived from the ones in (2.3). The proof of our assertion is quite lengthy and tedious. But the basic idea is just the case-by-case check-up.

We define the sets

$$\begin{aligned} \Lambda &= \{(i, j) \in \mathbb{Z}/\{0\} \times \mathbb{Z}/\{0\} \mid -n \leq i \leq j \leq n\}, & \Lambda_1 &= \{(i, j) \in \Lambda \mid i > 0, j > 0 \text{ and } i < j\}, \\ \Lambda_2 &= \{(i, j) \in \Lambda \mid i < 0, j > 0 \text{ and } |i| < |j|\}, & \Lambda_3 &= \{(i, j) \in \Lambda \mid i < 0, j > 0 \text{ and } |i| > |j|\}, \\ \Lambda_4 &= \{(i, j) \in \Lambda \mid i < 0, j < 0 \text{ and } |i| > |j|\}, & \Lambda_5 &= \{(i, j) \in \Lambda \mid i < 0, j > 0 \text{ and } |i| = |j|\}. \end{aligned}$$

For $((i, j), (k, l)) \in \Lambda \times \Lambda$, let $a = \min\{|i|, |j|\}$, $b = \max\{|i|, |j|\}$, $c = \min\{|k|, |l|\}$, $d = \max\{|k|, |l|\}$. We list all possible subsets of $\Lambda \times \Lambda$:

$$\begin{aligned} C_1 &= \{((i, j), (k, l)) \in \Lambda \times \Lambda \mid c < d < a < b\}, & C_2 &= \{((i, j), (k, l)) \in \Lambda \times \Lambda \mid c < d = a < b\}, \\ C_3 &= \{((i, j), (k, l)) \in \Lambda \times \Lambda \mid c < a < d < b\}, & C_4 &= \{((i, j), (k, l)) \in \Lambda \times \Lambda \mid c < a < d = b\}, \\ C_5 &= \{((i, j), (k, l)) \in \Lambda \times \Lambda \mid c < a < b < d\}, & C_6 &= \{((i, j), (k, l)) \in \Lambda \times \Lambda \mid c = a < d < b\}, \\ C_7 &= \{((i, j), (k, l)) \in \Lambda \times \Lambda \mid c = a < d = b\}, & C_8 &= \{((i, j), (k, l)) \in \Lambda \times \Lambda \mid c = a < b < d\}, \\ C_9 &= \{((i, j), (k, l)) \in \Lambda \times \Lambda \mid a < c < d < b\}, & C_{10} &= \{((i, j), (k, l)) \in \Lambda \times \Lambda \mid a < c < d = b\}, \\ C_{11} &= \{((i, j), (k, l)) \in \Lambda \times \Lambda \mid a < c < b < d\}, & C_{12} &= \{((i, j), (k, l)) \in \Lambda \times \Lambda \mid a < b = c < d\}, \\ C_{13} &= \{((i, j), (k, l)) \in \Lambda \times \Lambda \mid a < b < c < d\}, & D_1 &= \{((i, j), (k, l)) \in \Lambda_5 \times \Lambda \mid |i| < c < d\}, \\ D_2 &= \{((i, j), (k, l)) \in \Lambda_5 \times \Lambda \mid |i| = c < d\}, & D_3 &= \{((i, j), (k, l)) \in \Lambda_5 \times \Lambda \mid c < |i| < d\}, \\ D_4 &= \{((i, j), (k, l)) \in \Lambda_5 \times \Lambda \mid c < |i| = d\}, & D_5 &= \{((i, j), (k, l)) \in \Lambda_5 \times \Lambda \mid c < d < |i|\}, \\ D_6 &= \{((i, j), (k, l)) \in \Lambda \times \Lambda_5 \mid |k| < a < b\}, & D_7 &= \{((i, j), (k, l)) \in \Lambda \times \Lambda_5 \mid |k| = a < b\}, \\ D_8 &= \{((i, j), (k, l)) \in \Lambda \times \Lambda_5 \mid a < |k| < b\}, & D_9 &= \{((i, j), (k, l)) \in \Lambda \times \Lambda_5 \mid a < b = |k|\}, \\ D_{10} &= \{((i, j), (k, l)) \in \Lambda \times \Lambda_5 \mid a < b < |k|\}. \end{aligned}$$

We consider all cases for $\Lambda_s \times \Lambda_t \cap C_i$ ($1 \leq s, t \leq 4, 1 \leq i \leq 13$) and $\Lambda_s \times \Lambda_t \cap D_i$ ($s = 5, 1 \leq t \leq 4$ or $1 \leq s \leq 4, t = 5$ and $1 \leq i \leq 10$). Since the remaining cases can be checked similarly, we just prove:

$$(2.5) \quad L_{i,i} L_{k,l} L_{i,i}^{-1} = q^{\varphi(l,i) - \varphi(k,i)} L_{k,l} \quad \text{if } (k, l) \in \Lambda_1 \cup \Lambda_2,$$

$$(2.6) \quad L_{i,j} L_{k,l} - L_{k,l} L_{i,j} = 0 \quad \text{if } ((i, j), (k, l)) \in \Lambda_1 \times \Lambda_1 \cap C_1,$$

$$(2.7) \quad L_{i,j} L_{k,l} - L_{k,l} L_{i,j} = (q - q^{-1}) L_{i,l} L_{k,j} \quad \text{if } ((i, j), (k, l)) \in \Lambda_1 \times \Lambda_1 \cap C_2,$$

$$(2.8) \quad (L_{i,j})^2 = \frac{q - q^{-1}}{q + q^{-1}} (L_{-i,j})^2 \quad \text{if } (i, j) \in \Lambda_2.$$

From (2.4), we obtain

$$\begin{aligned} L_{i,j} &= \frac{L_{j-1,j-1}^{-1}}{q - q^{-1}} (L_{j-1,j} L_{i,j-1} - L_{i,j-1} L_{j-1,j}) \quad \text{if } (i, j) \in \Lambda_1 \cup \Lambda_2, \\ L_{i,j} &= \frac{L_{-i-1,-i-1}^{-1}}{q - q^{-1}} (L_{i,i+1} L_{i+1,j} - L_{i+1,j} L_{i,i+1}) \quad \text{if } (i, j) \in \Lambda_3 \cup \Lambda_4. \end{aligned}$$

To prove (2.5), we use induction on $l - k$:

$$\begin{aligned} L_{i,i} L_{k,l} L_{i,i}^{-1} &= \frac{L_{l-1,l-1}^{-1}}{q - q^{-1}} L_{i,i} (L_{l-1,l} L_{k,l-1} - L_{k,l-1} L_{l-1,l}) L_{i,i}^{-1} \\ &= q^{\varphi(l,i) - \varphi(l-1,i) + \varphi(l-1,i) - \varphi(k,i)} \frac{L_{l-1,l-1}^{-1}}{q - q^{-1}} (L_{l-1,l} L_{k,l-1} - L_{k,l-1} L_{l-1,l}) \\ &= q^{\varphi(l,i) - \varphi(k,i)} L_{k,l}. \end{aligned}$$

From (2.3), we know that $f_i f_j - f_j f_i = 0$ if $|i - j| > 1$. By using induction on $j - i$ and (2.5), one can show that $L_{i,j} L_{k,k+1} - L_{k,k+1} L_{i,j} = 0$ when $((i, j), (k, k+1)) \in \Lambda_1 \times \Lambda_1 \cap C_1$. Similarly, one can prove $L_{i,j} L_{k,l} - L_{k,l} L_{i,j} = 0$ by induction on $l - k$. The proof of (2.7) is analogous (we use induction on $l - k$ and (2.5), (2.6)):

$$\begin{aligned} L_{i,j} L_{k,l} &= \frac{L_{l-1,l-1}^{-1}}{q - q^{-1}} L_{i,j} (L_{l-1,l} L_{k,l-1} - L_{k,l-1} L_{l-1,l}) \\ &= \frac{L_{l-1,l-1}^{-1}}{q - q^{-1}} (L_{l-1,l} L_{i,j} L_{k,l-1} + (q - q^{-1}) L_{i,l} L_{l-1,j} L_{k,l-1} - L_{k,l-1} L_{i,j} L_{l-1,l}) \\ &= \frac{L_{l-1,l-1}^{-1}}{q - q^{-1}} (L_{l-1,l} L_{k,l-1} L_{i,j} + (q - q^{-1}) L_{i,l} L_{l-1,j} L_{k,l-1} - L_{k,l-1} L_{l-1,l} L_{i,j} \\ &\quad - (q - q^{-1}) L_{k,l-1} L_{i,l} L_{l-1,j}) \\ &= L_{k,l} L_{i,j} + L_{l-1,l-1}^{-1} L_{i,l} (L_{l-1,j} L_{k,l-1} - L_{k,l-1} L_{l-1,j}) \\ &= L_{k,l} L_{i,j} + (q - q^{-1}) L_{i,l} L_{k,j}. \end{aligned}$$

To verify the relation (2.8), it suffices to show that

$$(L_{j-1,j} L_{i,j-1} - L_{i,j-1} L_{j-1,j})^2 = \frac{q - q^{-1}}{q + q^{-1}} (L_{j-1,j} L_{-i,j-1} - L_{-i,j-1} L_{j-1,j})^2.$$

For this purpose, we need the following formulas for $(i, j) \in \Lambda_2$ which can be derived using induction:

$$L_{j-1,j} L_{i,j-1} L_{j-1,j} = \frac{1}{q + q^{-1}} (q L_{i,j-1} L_{j-1,j}^2 + q^{-1} L_{j-1,j}^2 L_{i,j-1}),$$

$$qL_{-i,j-1}L_{j-1,j}^2 - (q + q^{-1})L_{j-1,j}L_{-i,j-1}L_{j-1,j} + q^{-1}L_{j-1,j}^2L_{-i,j-1} = 0.$$

Using these formulae, we can verify the desired relations

$$\begin{aligned} & (L_{j-1,j}L_{i,j-1} - L_{i,j-1}L_{j-1,j})^2 \\ &= (L_{j-1,j}L_{i,j-1}L_{j-1,j})L_{i,j-1} - \frac{q - q^{-1}}{q + q^{-1}}L_{j-1,j}L_{-i,j-1}^2L_{j-1,j} - L_{i,j-1}L_{j-1,j}^2L_{i,j-1} \\ &\quad + L_{i,j-1}(L_{j-1,j}L_{i,j-1}L_{j-1,j}) \\ &= \frac{q - q^{-1}}{q + q^{-1}} \left(\frac{q^{-1}}{q + q^{-1}}L_{j-1,j}^2L_{-i,j-1}^2 + \frac{q}{q + q^{-1}}L_{-i,j-1}^2L_{j-1,j}^2 - L_{j-1,j}L_{-i,j-1}^2L_{j-1,j} \right) \\ &= \frac{q - q^{-1}}{q + q^{-1}} \left((L_{j-1,j}L_{-i,j-1}L_{j-1,j} - \frac{q}{q + q^{-1}}L_{-i,j-1}L_{j-1,j}^2)L_{-i,j-1} \right. \\ &\quad \left. + L_{-i,j-1}(L_{j-1,j}L_{-i,j-1}L_{j-1,j} - \frac{q^{-1}}{q + q^{-1}}L_{j-1,j}^2L_{-i,j-1}) - L_{j-1,j}L_{-i,j-1}^2L_{j-1,j} \right) \\ &= \frac{q - q^{-1}}{q + q^{-1}}(L_{j-1,j}L_{-i,j-1} - L_{-i,j-1}L_{j-1,j})^2. \end{aligned}$$

□

Set $\deg f_i = \deg \bar{f}_i = -\alpha_i$, $\deg q^h = \deg k_{\bar{i}} = 0$, $\deg e_i = \deg \bar{e}_i = \alpha_i$. Since all the defining relations of the quantum superalgebra $U_q(\mathfrak{g})$ are homogeneous, it has a root space decomposition

$$U_q(\mathfrak{g}) = \bigoplus_{\alpha \in Q} (U_q)_{\alpha},$$

where $(U_q)_{\alpha} = \{u \in U_q(\mathfrak{g}) \mid q^h u q^{-h} = q^{\alpha(h)} u \text{ for all } h \in P^{\vee}\}$.

Remark. If we define

$$F_i = f_i q^{-k_i+1}, \quad E_i = q^{k_i+1} e_i,$$

one can see that the relations involving E_i , F_i and q^h are the same as the standard relations for $U_q(\mathfrak{gl}_n)$ (see, for example, [HK, Definition 7.1.1]). Hence $U_q(\mathfrak{gl}_n)$ is a subalgebra of $U_q(\mathfrak{g})$.

The comultiplication Δ of $U_q(\mathfrak{g})$ is given by the formula

$$(2.9) \quad \Delta(L_{i,j}) = \sum_{k=i}^j L_{i,k} \otimes L_{k,j},$$

(see §4 in [O]). In terms of the new generators we have:

$$\begin{aligned} \Delta(q^h) &= q^h \otimes q^h \text{ for every } h \in P^{\vee}, \\ \Delta(e_i) &= q^{-k_i+1} \otimes e_i + e_i \otimes q^{-k_i}, \end{aligned}$$

$$\begin{aligned}
\Delta(f_i) &= q^{k_i} \otimes f_i + f_i \otimes q^{k_{i+1}}, \\
\Delta(e_{\bar{i}}) &= q^{-k_{i+1}} \otimes e_{\bar{i}} - (q - q^{-1})e_i \otimes k_{\bar{i}} \\
&\quad + (q - q^{-1}) \left(\sum_{j=1}^{i-1} (-1)^{j+1} q^{\sum_{h=1}^j k_{i-j+h}} \prod_{h=1}^j \text{ad } e_{i-j+h}(e_{i-j}) \right. \\
&\quad \left. \otimes q^{-\sum_{h=1}^{j-1} k_{i-j+h}} \prod_{h=1}^{j-1} \text{ad } f_{i-j+h}(f_{i-j}) \right) \\
&\quad + (q - q^{-1}) \left(\sum_{j=1}^{i-1} (-1)^j q^{\sum_{h=1}^j k_{i-j+h}} \prod_{h=1}^j \text{ad } e_{i-j+h}(e_{\bar{i}-j}) \right. \\
&\quad \left. \otimes q^{-\sum_{h=1}^{j-1} k_{i-j+h}} \prod_{h=1}^{j-1} \text{ad } f_{i-j+h}(f_{i-j}) \right) + e_{\bar{i}} \otimes q^{k_i}, \\
\Delta(f_{\bar{i}}) &= q^{-k_i} \otimes f_{\bar{i}} \\
&\quad + (q - q^{-1}) \left(\sum_{j=1}^{i-1} (-1)^j q^{\sum_{h=1}^{j-1} k_{i-j+h}} \prod_{h=1}^{j-1} \text{ad } e_{i-j+h}(e_{i-j}) \right. \\
&\quad \left. \otimes q^{-\sum_{h=1}^j k_{i-j+h}} \prod_{h=1}^j \text{ad } f_{i-j+h}(f_{i-j}) \right) \\
&\quad + (q - q^{-1}) \left(\sum_{j=1}^{i-1} (-1)^{j+1} q^{\sum_{h=1}^{j-1} k_{i-j+h}} \prod_{h=1}^{j-1} \text{ad } e_{i-j+h}(e_{\bar{i}-j}) \right. \\
&\quad \left. \otimes q^{-\sum_{h=1}^j k_{i-j+h}} \prod_{h=1}^j \text{ad } f_{i-j+h}(f_{i-j}) \right) \\
&\quad + (q - q^{-1}) k_{\bar{i}} \otimes f_i + f_{\bar{i}} \otimes q^{k_{i+1}}, \\
\Delta(k_{\bar{i}}) &= q^{-k_i} \otimes k_{\bar{i}} \\
&\quad + (q - q^{-1}) \left(\sum_{j=1}^{i-1} (-1)^j q^{\sum_{h=1}^{j-1} k_{i-j+h}} \prod_{h=1}^{j-1} \text{ad } e_{i-j+h}(e_{i-j}) \right. \\
&\quad \left. \otimes q^{-\sum_{h=1}^{j-1} k_{i-j+h}} \prod_{h=1}^{j-1} \text{ad } f_{i-j+h}(f_{i-j}) \right) \\
&\quad + (q - q^{-1}) \left(\sum_{j=1}^{i-1} (-1)^{j+1} q^{\sum_{h=1}^{j-1} k_{i-j+h}} \prod_{h=1}^{j-1} \text{ad } e_{i-j+h}(e_{\bar{i}-j}) \right.
\end{aligned}$$

$$\otimes q^{-\sum_{h=1}^{j-1} k_{i-j+h}} \prod_{h=1}^{j-1} \text{ad } f_{i-j+h}(f_{i-j}) \Big) + k_{\bar{i}} \otimes q^{k_i}.$$

Let U_q^+ (respectively, U_q^-) be the subalgebra of $U_q(\mathfrak{g})$ generated by the elements $e_i, e_{\bar{i}}$ (respectively, $f_i, f_{\bar{i}}$) for $i = 1, \dots, n-1$, and let U_q^0 be the subalgebra of $U_q(\mathfrak{g})$ generated by q^h ($h \in P^\vee$) and $k_{\bar{l}}$ for $l = 1, \dots, n$. In addition, let $U_q^{\geq 0}$ (respectively, $U_q^{\leq 0}$) be the subalgebra of $U_q(\mathfrak{g})$ generated by U_q^+ and U_q^0 (respectively, by U_q^- and U_q^0). We will show that the quantum superalgebra $U_q(\mathfrak{g})$ has a triangular decomposition. For this purpose, we need the following lemma.

Lemma 2.2.

$$U_q^{\geq 0} \cong U_q^0 \otimes U_q^+, \quad U_q^{\leq 0} \cong U_q^- \otimes U_q^0.$$

Proof. We will prove the second part. Let $\{f_\zeta \mid \zeta \in \Omega\}$ be a basis of U_q^- consisting of monomials in f_i and $f_{\bar{i}}$'s ($i \in I$). Consider a set $\Omega' = \{(a_1, \dots, a_n) \mid a_i = 0 \text{ or } 1 \text{ for all } i \in J\}$. Then $\{q^h k_\eta \mid h \in P^\vee, \eta \in \Omega'\}$ is a basis of U_q^0 , where $k_\eta = k_1^{a_1} \cdots k_n^{a_n}$ for $\eta = (a_1, \dots, a_n)$ by [O, Theorem 6.2]. By the defining relations of $U_q(\mathfrak{g})$, it is easy to see that the elements $f_\zeta q^h k_\eta$ ($\zeta \in \Omega, h \in P^\vee, \eta \in \Omega'$) span $U_q^{\leq 0}$. Thus there is a surjective $\mathbb{C}(q)$ -linear map $U_q^- \otimes U_q^0 \rightarrow U_q^{\leq 0}$ given by $f_\zeta \otimes q^h k_\eta \rightarrow f_\zeta q^h k_\eta$. To show that this map is injective, it suffices to show that the elements $f_\zeta q^h k_\eta$ ($\zeta \in \Omega, h \in P^\vee, \eta \in \Omega'$) are linearly independent over $\mathbb{C}(q)$.

Suppose

$$\sum_{\substack{\zeta \in \Omega, h \in P^\vee, \\ \eta \in \Omega'}} C_{\zeta, h, \eta} f_\zeta q^h k_\eta = 0 \quad \text{for some } C_{\zeta, h, \eta} \in \mathbb{C}(q).$$

We may write

$$\sum_{\beta \in Q_+} \left(\sum_{\substack{\deg f_\zeta = -\beta, \\ h \in P^\vee, \eta \in \Omega'}} C_{\zeta, h, \eta} f_\zeta q^h k_\eta \right) = 0 \quad \text{for some } C_{\zeta, h, \eta} \in \mathbb{C}(q).$$

Since $U_q(\mathfrak{g}) = \bigoplus_{\beta \in Q} (U_q)_\beta$, we have

$$\sum_{\substack{\deg f_\zeta = -\beta, \\ h \in P^\vee, \eta \in \Omega'}} C_{\zeta, h, \eta} f_\zeta q^h k_\eta = 0 \quad \text{for each } \beta \in Q_+.$$

Write $\beta = -\sum_{i=1}^{n-1} m_i \alpha_i$ ($m_i \in \mathbb{Z}_{\geq 0}$), and let $h_\beta = \sum_{i=1}^{n-1} m_i k_{i+1}$. Since f_ζ is a monomial in f_i and $f_{\bar{i}}$'s, the term of degree $(-\beta, 0)$ in $\Delta(f_\zeta)$ is $f_\zeta \otimes q^{h_\beta}$. We consider the terms of degree $(0, 0)$ in $\Delta(k_\eta)$ where $\eta = (a_1, \dots, a_n)$. Then the terms of degree $(0, 0)$ in $\Delta(k_\eta)$ can

be written as

$$\begin{aligned}
& (q^{-k_1} \otimes k_{\bar{1}} + k_{\bar{1}} \otimes q^{k_1})^{a_1} \cdots (q^{-k_n} \otimes k_{\bar{n}} + k_{\bar{n}} \otimes q^{k_n})^{a_n} \\
&= \prod_{i=1}^n \left(\sum_{j=0}^{a_i} q^{-(a_i-j)k_i} k_{\bar{i}}^j \otimes q^{jk_i} k_{\bar{i}}^{a_i-j} \right) \\
&= \sum_{\substack{(j_1, \dots, j_n) \in \Omega' \\ j_i \leq a_i, i \in J}} \prod_{i=1}^n \left(q^{-(a_i-j_i)k_i} k_{\bar{i}}^{j_i} \otimes q^{j_i k_i} k_{\bar{i}}^{a_i-j_i} \right).
\end{aligned}$$

Since the terms of degree $(-\beta, 0)$ of $\sum C_{\zeta, h, \eta} \Delta(f_{\zeta} q^h k_{\eta})$ must sum to zero, we have

$$(2.10) \quad 0 = \sum_{\substack{\eta=(a_1, \dots, a_n) \\ \in \Omega'}} \sum_{\substack{(j_1, \dots, j_n) \in \Omega' \\ j_i \leq a_i, i \in J}} \left(\sum_{\substack{\deg f_{\zeta} = -\beta, \\ h \in P^{\vee}}} C_{\zeta, h, \eta} f_{\zeta} q^h \left(\prod_{i=1}^n q^{-(a_i-j_i)k_i} k_{\bar{i}}^{j_i} \right) \otimes q^{h_{\beta}+h} \left(\prod_{i=1}^n q^{j_i k_i} k_{\bar{i}}^{a_i-j_i} \right) \right).$$

For all $(a_1 - j_1, \dots, a_n - j_n) \in \Omega'$ and $h \in P^{\vee}$, the elements $q^h \left(\prod_{i=1}^n k_{\bar{i}}^{a_i-j_i} \right)$ are linearly independent. Set $\eta_1 := (1, \dots, 1)$. Since there is only one pair of (a_1, \dots, a_n) and (j_1, \dots, j_n) such that $\prod_{i=1}^n k_{\bar{i}}^{a_i-j_i} = k_{\eta_1}$ in the above sum, we obtain

$$\begin{aligned}
0 &= \sum_{h \in P^{\vee}} \sum_{\deg f_{\zeta} = -\beta} C_{\zeta, h, \eta_1} f_{\zeta} q^{h - \sum_{i=1}^n k_i} \otimes q^{h_{\beta}+h} k_{\eta_1} \\
&+ \sum_{\substack{\eta=(a_1, \dots, a_n), \\ (a_1-j_1, \dots, a_n-j_n) \neq \eta_1}} \sum_{\substack{(j_1, \dots, j_n), \\ h \in P^{\vee}, \\ \deg f_{\zeta} = -\beta}} C_{\zeta, h, \eta} f_{\zeta} q^h \left(\prod_{i=1}^n q^{-(a_i-j_i)k_i} k_{\bar{i}}^{j_i} \right) \otimes q^{h_{\beta}+h} \left(\prod_{i=1}^n q^{j_i k_i} k_{\bar{i}}^{a_i-j_i} \right).
\end{aligned}$$

Thus we have

$$\sum_{\deg f_{\zeta} = -\beta} C_{\zeta, h, \eta_1} f_{\zeta} q^{h - \sum_{i=1}^n k_i} = 0 \quad \text{for all } h \in P^{\vee}.$$

Multiplying by $q^{-h + \sum_{i=1}^n k_i}$ from the right we obtain

$$\sum_{\deg f_{\zeta} = -\beta} C_{\zeta, h, \eta_1} f_{\zeta} = 0 \quad \text{for all } h \in P^{\vee}.$$

Using the linear independence of f_{ζ} , we conclude all $C_{\zeta, h, \eta_1} = 0$ for all $\zeta \in \Omega, h \in P^{\vee}$. Now consider general $\eta = (a_1, \dots, a_n) \in \Omega'$. Assume that for all $\eta' = (a'_1, \dots, a'_n)$ such that $a'_i \geq a_i$ for all $i \in J$ and $\eta' \neq \eta$, $C_{\zeta, h, \eta'} = 0$ for all $\zeta \in \Omega, h \in P^{\vee}$. Then there is only one pair of (a_1, \dots, a_n) and (j_1, \dots, j_n) such that $(a_1 - j_1, \dots, a_n - j_n) = \eta$ in (2.10). Repeating the above argument, we conclude $C_{\zeta, h, \eta} = 0$ for all $\zeta \in \Omega, h \in P^{\vee}$.

For example, consider $\eta_2 = (0, 1, \dots, 1)$. Since $C_{\zeta, h, \eta_1} = 0$, there is only one pair of (a_1, \dots, a_n) and (j_1, \dots, j_n) such that $(a_1 - j_1, \dots, a_n - j_n) = (0, 1, \dots, 1)$ in (2.10). Thus we

have

$$\sum_{\deg f_\zeta = -\beta} C_{\zeta, h, \eta_2} f_\zeta q^{h - \sum_{i=2}^n k_i} = 0 \quad \text{for all } h \in P^\vee.$$

Multiplying $q^{-h + \sum_{i=2}^n k_i}$ and using the linear independence of f_ζ , we obtain $C_{\zeta, h, \eta_2} = 0$ for all $\zeta \in \Omega$, $h \in P^\vee$. □

We are now ready to prove the *triangular decomposition* for $U_q(\mathfrak{g})$.

Theorem 2.3. *There is a $\mathbb{C}(q)$ -linear isomorphism*

$$U_q(\mathfrak{g}) \cong U_q^- \otimes U_q^0 \otimes U_q^+.$$

Proof. Let $\{f_\zeta \mid \zeta \in \Omega\}$, $\{q^h k_\eta \mid h \in P^\vee, \eta \in \Omega'\}$, and $\{e_\tau \mid \tau \in \Omega\}$ be monomial bases of U_q^- , U_q^0 and U_q^+ respectively, where Ω and Ω' are the index sets as in the proof for Lemma 2.2. It suffices to show that the elements $f_\zeta q^h k_\eta e_\tau$ ($\zeta, \tau \in \Omega, h \in P^\vee, \eta \in \Omega'$) are linearly independent over $\mathbb{C}(q)$.

Suppose

$$\sum_{\zeta, h, \eta, \tau} C_{\zeta, h, \eta, \tau} f_\zeta q^h k_\eta e_\tau = 0 \quad \text{for some } C_{\zeta, h, \eta, \tau} \in \mathbb{C}(q).$$

The root space decomposition of $U_q(\mathfrak{g})$ yields

$$\sum_{\substack{h, \eta, \\ \deg f_\zeta + \deg e_\tau = \gamma}} C_{\zeta, h, \eta, \tau} f_\zeta q^h k_\eta e_\tau = 0 \quad \text{for all } \gamma \in Q.$$

Using the partial ordering on \mathfrak{h}_0^* , we can choose $\alpha = \deg f_\zeta$ and $\beta = \deg e_\tau$, which are minimal and maximal, respectively, among those for which $\alpha + \beta = \gamma$ and $C_{\zeta, h, \eta, \tau}$ is nonzero. If $\alpha = -\sum m_i \alpha_i$, set $h_\alpha = \sum m_i k_{i+1}$, and if $\beta = \sum n_i \alpha_i$, set $h_\beta = \sum n_i k_{i+1}$. The term of degree $(0, \beta)$ in $\Delta(e_\tau)$ is $q^{-h_\beta} \otimes e_\tau$ and the term of degree $(\alpha, 0)$ of $\Delta(f_\zeta)$ is $f_\zeta \otimes q^{h_\alpha}$.

Since the terms of degree (α, β) of $\sum C_{\zeta, h, \eta, \tau} \Delta(f_\zeta q^h k_\eta e_\tau)$ must sum to zero, we have

$$\sum_{\substack{\deg f_\zeta = \alpha, \\ \deg e_\tau = \beta, \\ h, \eta = (a_1, \dots, a_n)}} \sum_{\substack{(j_1, \dots, j_n) \in \Omega' \\ j_i \leq a_i, i \in J}} C_{\zeta, h, \eta, \tau} f_\zeta q^h \left(\prod_{i=1}^n q^{-(a_i - j_i) k_i - h_\beta} k_i^{j_i} \right) \otimes q^{h_\alpha + h} \left(\prod_{i=1}^n q^{j_i k_i} k_i^{a_i - j_i} \right) e_\tau = 0.$$

The elements $f_\zeta q^h \left(\prod_{i=1}^n k_i^{j_i} \right)$ are linearly independent for $\zeta \in \Omega$, $h \in P^\vee$, $(j_1, \dots, j_n) \in \Omega'$ by Lemma 2.2. By the similar argument in the proof for Lemma 2.2, we obtain

$$\sum_{\deg e_\tau = \beta} C_{\zeta, h, \eta, \tau} e_\tau = 0 \quad \text{for all } h \in P^\vee, \zeta \in \Omega, \text{ and } \eta \in \Omega'.$$

Using the linear independence of e_τ , we conclude that $C_{\zeta, h, \eta, \tau} = 0$ for all $\zeta \in \Omega, h \in P^\vee, \eta \in \Omega',$ and $\tau \in \Omega,$ as desired. \square

3. THE QUANTUM CLIFFORD SUPERALGEBRA $\text{Cliff}_q(\lambda)$

We first introduce some notation that will be used in this section only. Let \mathbb{K} be a field of zero characteristic and A be an associative \mathbb{K} -algebra. Denote by $\text{Mat}_n(A)$ the associative \mathbb{K} -algebra of $n \times n$ matrices with entries in A . If A is a superalgebra, then $\text{Mat}_n(A)$ is a superalgebra as well by setting $\text{Mat}_n(A)_{\bar{i}} = \text{Mat}_n(A_{\bar{i}})$. By $\text{sMat}_{n|n}(\mathbb{K})$ we denote the associative superalgebra of $2n \times 2n$ matrices $\begin{pmatrix} A & B \\ C & D \end{pmatrix}$, where $A, B, C,$ and D are in $\text{Mat}_n(\mathbb{K})$ and

$$\text{sMat}_{n,n}(\mathbb{K})_{\bar{0}} = \left\{ \begin{pmatrix} A & 0 \\ 0 & D \end{pmatrix} \right\}, \quad \text{sMat}_{n,n}(\mathbb{K})_{\bar{1}} = \left\{ \begin{pmatrix} 0 & B \\ C & 0 \end{pmatrix} \right\}.$$

Let $Q_n(\mathbb{K})$ be the subsuperalgebra of $\text{sMat}_{n|n}(\mathbb{K})$ with elements $\begin{pmatrix} A & B \\ B & A \end{pmatrix}$. In particular, $Q_n(\mathbb{K})_{\bar{0}} = Q_n(\mathbb{K})_{\bar{1}} = \text{Mat}_n(\mathbb{K})$. There are \mathbb{K} -superalgebra isomorphisms

$$\text{Mat}_r(\text{sMat}_{1|1}(\mathbb{K})) \cong \text{sMat}_{r|r}(\mathbb{K}), \quad \text{Mat}_r(Q_1(\mathbb{K})) \cong Q_r(\mathbb{K}).$$

Note that if $\mathbb{K} = \mathbb{C}$, then the superalgebra $Q_n(\mathbb{C})$ coincides with \mathfrak{g} as a complex vector space. Another example of a \mathbb{K} -superalgebra is any extension $\mathbb{K}(\alpha)$ of \mathbb{K} of degree 2 considering α as an odd element. If $\alpha^2 = \beta \in \mathbb{K}$ we will denote $\mathbb{K}(\alpha)$ by $\mathbb{K}(\sqrt{\beta})$.

In this section, we set $\mathbb{F} = \mathbb{C}(q)$. For every $\lambda \in P$ we define $I^q(\lambda)$ to be the left ideal of U_q^0 generated by $q^h - q^{\lambda(h)}\mathbf{1}, h \in P^\vee$. Set $\text{Cliff}_q(\lambda) := U_q^0/I^q(\lambda)$. We may consider $\text{Cliff}_q(\lambda)$ as the associative \mathbb{F} -algebra generated by the identity $\mathbf{1} = 1 + I^q(\lambda)$ and $t_{\bar{i}} := k_{\bar{i}} + I^q(\lambda)$ satisfying the relations

$$t_{\bar{i}}t_{\bar{j}} + t_{\bar{j}}t_{\bar{i}} = \delta_{ij} \frac{2(q^{2\lambda_i} - q^{-2\lambda_i})}{q^2 - q^{-2}} \mathbf{1}, \quad i, j = 1, \dots, n.$$

Furthermore, $\text{Cliff}_q(\lambda)$ has an obvious \mathbb{Z}_2 -grading (and thus a superalgebra structure) by assuming that $t_{\bar{i}}$ are odd. More precisely, $\text{Cliff}_q(\lambda)_{\bar{0}}$ is spanned by $\mathbf{1}$ and the monomials $t_{\bar{i}_1} \dots t_{\bar{i}_{2k}}$ of even degree, while $\text{Cliff}_q(\lambda)_{\bar{1}}$ is spanned by those of odd degree. In this section we will describe the structure of $\text{Cliff}_q(\lambda)$ and will classify its irreducible modules. Because of its superalgebra structure, $\text{Cliff}_q(\lambda)$ has both \mathbb{Z}_2 -graded and nongraded modules and both cases will be addressed.

The results in this section may be derived from more general statements about quadratic forms and Clifford superalgebras over arbitrary fields (see, for example, [Lam] and [Sh]). For

the sake of completeness we will give an outline of the proofs. The results and the proofs in this section will also help us to describe explicitly the action of U_q^0 on the highest weight vectors of an irreducible highest weight module over $U_q(\mathfrak{g})$. This is demonstrated in Example 3.10 for the case $n = 3$ and $\lambda = (4, 2, 1)$.

In this section, we fix $V := \bigoplus_{i=1}^n \mathbb{F}t_{\bar{i}}$ and $\Lambda := (\Lambda_1, \dots, \Lambda_n) \in \mathbb{F}^n$ and denote by $B_\Lambda : V \times V \rightarrow \mathbb{F}$ the symmetric bilinear form defined by $B_\Lambda(t_{\bar{i}}, t_{\bar{j}}) = \delta_{ij}\Lambda_i$. Let $\text{Cliff}_q(\Lambda)$ be the unique up to isomorphism Clifford algebra associated to V and B_Λ . If $\Lambda_i = \frac{q^{2\lambda_i} - q^{-2\lambda_i}}{q^2 - q^{-2}}$, then we have $\text{Cliff}_q(\Lambda) \simeq \text{Cliff}_q(\lambda)$.

Define $V(\Lambda) := V / \ker B_\Lambda$, where $\ker B_\Lambda := \{v \in V \mid B_\Lambda(v, u) = 0, \text{ for every } u \in V\}$ and denote by β_Λ the restriction of B_Λ on $V(\Lambda)$. Let $N_\Lambda = \{i \mid \Lambda_i \neq 0\}$, $Z_\Lambda = \{j \mid \Lambda_j = 0\}$, and $|\Lambda| = \#N_\Lambda$. Set $\Lambda_N := (\Lambda_{i_1}, \dots, \Lambda_{i_{|\Lambda|}})$, $0_Z := (\Lambda_{j_1}, \dots, \Lambda_{j_{n-|\Lambda|}}) = (0, \dots, 0)$, where $N_\Lambda = \{i_1, \dots, i_{|\Lambda|}\}$, $Z_\Lambda = \{j_1, \dots, j_{n-|\Lambda|}\}$, and $i_1 < \dots < i_{|\Lambda|}$. It is clear that $\ker B_\Lambda = \bigoplus_{j \in Z_\Lambda} \mathbb{F}t_{\bar{j}}$ and that $\text{Cliff}_q(\Lambda_N) = \bigoplus_{i \in N_\Lambda} \mathbb{F}t_{\bar{i}}$ is the Clifford algebra corresponding to $(V(\Lambda), \beta_\Lambda)$. Furthermore,

$$\text{Cliff}_q(\Lambda) \simeq \text{Cliff}_q(\Lambda_N) \otimes_{\mathbb{F}} \text{Cliff}_q(0_Z) \simeq \text{Cliff}_q(\Lambda_N) \otimes_{\mathbb{F}} \bigwedge \ker B_\Lambda.$$

Here $\bigwedge W$ denotes the exterior algebra of the vector space W . Thanks to the above isomorphisms every $\text{Cliff}_q(\Lambda)$ -module can be considered as a $\text{Cliff}_q(\Lambda_N)$ -module under the embedding $\text{Cliff}_q(\Lambda_N) = \text{Cliff}_q(\Lambda_N) \otimes_{\mathbb{F}} 1 \rightarrow \text{Cliff}_q(\Lambda_N) \otimes_{\mathbb{F}} \text{Cliff}_q(0_Z)$. The class $\overline{\Delta(\Lambda)}$ of $\Delta(\Lambda) = \prod_{i \in N_\Lambda} \Lambda_i$ in \mathbb{F}/\mathbb{F}^2 is called the *discriminant* of (V, B_Λ) .

The following lemma is standard and the proof is left to the reader.

Lemma 3.1. *Let M be an irreducible $\text{Cliff}_q(\Lambda)$ -module. Then M is an irreducible $\text{Cliff}_q(\Lambda_N)$ -module and $t_{\bar{i}}v = 0$ for every $i \in Z_\Lambda$. Conversely, if M_0 is an irreducible $\text{Cliff}_q(\Lambda_N)$ -module then M_0 considered as a $\text{Cliff}_q(\Lambda)$ -module with trivial action of $\text{Cliff}_q(0_Z)$ is irreducible as well.*

Since our goal in this section is to classify the irreducible representations of $\text{Cliff}_q(\Lambda)$, thanks to the above lemma, we may assume that Λ_i are nonzero. So, for simplicity we fix $Z_\Lambda = \emptyset$, and thus $B_\Lambda = \beta_\Lambda$ and $V(\Lambda) = V$, in all statements preceding Corollary 3.9.

Recall that a vector v in V is called β_Λ -isotropic (or simply isotropic) if $\beta_\Lambda(v, v) = 0$. A subspace W of V is β_Λ -isotropic subspace if $\beta_\Lambda(u, w) = 0$ for every u and w in W . A subspace W of V is anisotropic if it contains no nonzero β_Λ -isotropic vector. An isotropic subspace W of V is maximal isotropic if there is no larger β_Λ -isotropic subspace containing W .

Lemma 3.2. *Let W be an isotropic subspace of V . Then there exists an isotropic subspace W^* and a subspace Z of V such that*

$$V = Z \oplus W \oplus W^*, \quad \dim W = \dim W^*,$$

$$\beta_\Lambda(z, w) = \beta_\Lambda(z, w^*) = 0 \quad \text{for every } z \in Z, w \in W, w^* \in W^*.$$

Moreover, there exist bases $\{w_1, \dots, w_m\}$ and $\{w_1^*, \dots, w_m^*\}$ of W and W^* , respectively, such that $\beta_\Lambda(w_i, w_j^*) = \delta_{ij}$.

Proof. The lemma follows by induction on $\dim W$. If $\dim W = 1$, then W^* is spanned by $w_1^* = x - \frac{1}{2}\beta_\Lambda(x, x)w_1$, where $x \in V$ is arbitrarily chosen so that $\beta_\Lambda(w_1, x) = 1$. Then we define Z to be

$$Z = \{z \in V \mid \beta_\Lambda(z, w_1) = \beta_\Lambda(z, w_1^*) = 0\}.$$

For the complete proof, see [Sh, Lemma 1.3]. \square

The decomposition $V = Z \oplus W \oplus W^*$ in Lemma 3.2 is called a *weak Witt decomposition* of V . For any weak Witt decomposition $V = Z \oplus W \oplus W^*$, we denote by $\text{Cliff}(\Lambda_Z)$ the Clifford algebra corresponding to $(Z, \beta_{\Lambda|_Z})$. If $V = Z \oplus W \oplus W^*$ is a weak Witt decomposition for which Z is anisotropic (or, equivalently, W is maximal isotropic) we call it a *Witt decomposition*. We may identify W^* with the dual space of W via the nondegenerate form β_Λ . If $V = Z \oplus W \oplus W^*$ is a Witt decomposition, the dimension of W is an invariant of (V, β_Λ) (see [Sh, Lemma 1.4]) and is known as the *Witt index* of the form β_Λ . We say that the Witt index is *maximal* if $\dim Z \leq 1$. Recall that if the ground field is \mathbb{C} , the Witt index is always maximal. In the case of arbitrary \mathbb{F} though, the Witt index is generally not maximal as we verify in Lemma 3.6. In order to find a Witt decomposition and the Witt index of (V, β_Λ) we need some preparatory statements.

Lemma 3.3. *Let $V = Z \oplus W \oplus W^*$ be a weak Witt decomposition and let $m = 2^{\dim W}$. Then $\text{Cliff}_q(\Lambda) \cong \text{Mat}_m(\text{Cliff}_q(\Lambda_Z))$. Moreover, we have*

$$\text{Cliff}_q(\Lambda)_{\bar{0}} \cong \begin{cases} \text{Mat}_m(\text{Cliff}_q(\Lambda_Z)_{\bar{0}}) & \text{if } Z \neq 0, \\ \text{Mat}_{m/2}(\mathbb{F}) \oplus \text{Mat}_{m/2}(\mathbb{F}) & \text{if } Z = 0. \end{cases}$$

Proof. For the complete proof, see [Sh, Theorem 2.6]. The proof follows by induction on $\dim W$. We sketch the proof for $\dim W = 1$. In this case there is an isomorphism $\Psi : \text{Cliff}_q(\Lambda) \rightarrow \text{Mat}_m(\text{Cliff}_q(\Lambda_Z))$ defined by its restriction $\Psi|_V$ on V :

$$z + rw_1 + sw_1^* \mapsto \begin{pmatrix} z & r \\ s & -z \end{pmatrix}.$$

Notice that if $Z \neq 0$, Ψ is not necessarily parity preserving. In such a case we choose the isomorphism $\Theta : \text{Cliff}_q(\Lambda) \rightarrow \text{Mat}_m(\text{Cliff}_q(\Lambda_Z))$ defined by $\Theta(\alpha) = D^{-1}\Psi(\alpha)D$, where $D = \begin{pmatrix} g & 0 \\ 0 & 1 \end{pmatrix}$ and any $g \in Z$ with $\beta_\Lambda(g, g) \neq 0$. \square

Lemma 3.4. *The nondegenerate Legendre's equation always has a nontrivial solution in \mathbb{F} : for every nonzero A, B, C in \mathbb{F} , there exist $X, Y, Z \in \mathbb{F}$ with $(X, Y, Z) \neq (0, 0, 0)$ such that $AX^2 + BY^2 + CZ^2 = 0$.*

Proof. We modify the proof of the classical Legendre's Theorem (see, for example, [IR, §17.3]). We first assume that A, B, C, X, Y, Z are polynomials in $\mathbb{C}[q]$, where A, B, C are square free. We may fix $C = -1$, since if (X, Y, Z) is a solution of $ACX^2 + BCY^2 = Z^2$ then $(X, Y, \sqrt{-1}\frac{Z}{C})$ is a solution of $AX^2 + BY^2 + CZ^2 = 0$. We prove that $AX^2 + BY^2 = Z^2$ has a nontrivial solution by induction on $N := \max\{\deg A, \deg B\}$.

If $N = 0$; i.e., A and B are constant polynomials, then $AX^2 + BY^2 = Z^2$ has a solution (constant polynomials). Assume that $\deg B \leq \deg A$ and $\deg A \geq 1$. Recall that every polynomial $R \in \mathbb{C}[q]$ is a quadratic residue modulo any square free polynomial S . Indeed, if S is constant, our assertion is obvious. Otherwise, let $S(q) = \prod_{i=1}^r (q - z_i)$ with $z_i \neq z_j$, and let $y_i \in \mathbb{C}$ be such that $y_i^2 = R(z_i)$. Then $y_i^2 \equiv R \pmod{(q - z_i)}$. Using the Chinese Remainder Theorem, we find $y \in \mathbb{C}[q]$ for which $y \equiv y_i \pmod{(q - z_i)}$. But then $y^2 \equiv R \pmod{(q - z_i)}$ and thus $y^2 \equiv R \pmod{S}$.

We fix C_1 with $\deg C_1 < \deg A$ such that $C_1^2 \equiv B \pmod{A}$. Then $C_1^2 - B = AT = AA_1M^2$ for some square free polynomial A_1 . Since $\deg A + \deg A_1 \leq \deg(AA_1M^2) = \deg(C_1^2 - B) < 2\deg A$, we have $0 \leq \deg A_1 < \deg A$. Now we observe that if (X_1, Y_1, Z_1) is a solution of $A_1X^2 + BY^2 = Z^2$, then $(A_1X_1M, C_1Y_1 + Z_1, Z_1C_1 + BY_1)$ is a solution of $AX^2 + BY^2 = Z^2$. Using the induction hypothesis, we complete the proof. \square

Remark. Lemma 3.4 may be proved with a standard algebro-geometric argument using dimensions, see, for example, [Har, Exercise 11.6]. The lemma is also a particular case of the following Theorem of Tsen-Lang: if K is a field of transcendence degree n over an algebraically closed field k , then any quadratic form over K of dimension bigger than 2^n is isotropic. For details, see [Lam, Chapter XI].

In what follows, we assume $\Lambda_i = \frac{q^{2\lambda_i} - q^{-2\lambda_i}}{q^2 - q^{-2}}$. For simplicity, we will write β_λ , $|\lambda|$, and $\Delta(\lambda)$ for β_Λ , $|\Lambda|$, and $\Delta(\Lambda)$, respectively. The following technical lemma can be easily verified.

Lemma 3.5. *Define an equivalence relation \sim in $\{\lambda_i \mid i = 1, \dots, n\}$ by $\lambda_i \sim \lambda_j$ if $\lambda_i^2 = \lambda_j^2$ and denote by $o(\lambda_i)$ the orbit of λ_i relative to \sim . Then $\overline{\Delta(\lambda)} = \bar{1}$ (or, equivalently, $\Delta(\lambda)$ is a square in \mathbb{F}) if and only if the orbit $o(\lambda_i)$ of every $\lambda_i \neq \pm 1$ contains even number of elements.*

Lemma 3.6. *The space V is anisotropic if and only if $\dim V = 1$ or $\dim V = 2$ and $\overline{\Delta(\lambda)} \neq \overline{1}$. If V is isotropic, there is a Witt decomposition $V = Z \oplus W \oplus W^*$ of V such that*

- (1) $\dim W = k$ if $\dim V = 2k + 1$, $k \geq 1$ (maximal Witt index);
- (2) $\dim W = k - 1$ if $\dim V = 2k$ and $\overline{\Delta(\lambda)} \neq \overline{1}$;
- (3) $\dim W = k$ if $\dim V = 2k$ and $\overline{\Delta(\lambda)} = \overline{1}$ (maximal Witt index).

In particular, if $\lambda_1 > \lambda_2 > \dots > \lambda_n > 0$, then $\dim W = \lfloor \frac{n-1}{2} \rfloor$.

Proof. The proof consists of several steps.

Step 1: The case $\dim V = 1$. This case is straightforward.

Step 2: The case $\dim V = 2$. In this case, $v = a_1 t_{\overline{1}} + a_2 t_{\overline{2}}$ is β_λ -isotropic if and only if $a_1^2 \Lambda_1 + a_2^2 \Lambda_2 = 0$. The latter equation has a solution for a_1 and a_2 if and only if $\frac{\Lambda_1}{\Lambda_2}$ is a square (or equivalently, $\Lambda_1 \Lambda_2$ is a square).

Step 3: If $\dim V \geq 3$, then

$$V \cong \mathbb{F}w \oplus \mathbb{F}w^* \oplus \mathbb{F}v_3 \oplus \dots \oplus \mathbb{F}v_n,$$

where

$$\begin{aligned} \beta_\lambda(w, w) &= \beta_\lambda(w^*, w^*) = \beta_\lambda(w, v_i) = \beta_\lambda(w^*, v_i) = 0 \quad \text{for } i \geq 3, \\ \beta_\lambda(w, w^*) &= 1, \quad \beta_\lambda(v_3, v_3) = \Lambda_1 \Lambda_2 \Lambda_3, \quad \beta_\lambda(v_i, v_i) = \Lambda_i \quad \text{if } i \geq 4. \end{aligned}$$

Let us first consider the case $\dim V = 3$. We use Lemma 3.4 to find $w = x_1 t_{\overline{1}} + x_2 t_{\overline{2}} + x_3 t_{\overline{3}}$ such that $\beta_\lambda(w, w) = 0$. Applying Lemma 3.2 to $W = \mathbb{F}w$, we find $w^* = y_1 t_{\overline{1}} + y_2 t_{\overline{2}} + y_3 t_{\overline{3}}$ and $z = z_1 t_{\overline{1}} + z_2 t_{\overline{2}} + z_3 t_{\overline{3}}$ such that

$$\beta_\lambda(w^*, w^*) = \beta_\lambda(w^*, z) = \beta_\lambda(w, z) = 0, \quad \beta_\lambda(w, w^*) = 1.$$

The choice of z is unique up to a multiplication by a nonzero constant in \mathbb{F} . A simple calculation shows that z_i may be chosen as follows

$$\begin{aligned} z_1 &= \sqrt{-1} \Lambda_2 \Lambda_3 (x_2 y_3 - x_3 y_2), \\ z_2 &= \sqrt{-1} \Lambda_1 \Lambda_3 (x_3 y_1 - x_1 y_3), \\ z_3 &= \sqrt{-1} \Lambda_1 \Lambda_2 (x_1 y_2 - x_2 y_1). \end{aligned}$$

Then one can easily verify that $\beta_\lambda(z, z) = \Lambda_1 \Lambda_2 \Lambda_3$.

In the case $\dim V > 3$, write $V = \mathbb{F}t_{\overline{1}} \oplus \mathbb{F}t_{\overline{2}} \oplus \mathbb{F}t_{\overline{3}} \oplus \left(\bigoplus_{i \geq 4} \mathbb{F}t_{\overline{i}} \right)$. Fix $w, w^*, z \in \mathbb{F}t_{\overline{1}} \oplus \mathbb{F}t_{\overline{2}} \oplus \mathbb{F}t_{\overline{3}}$ as above, and set $v_3 = z$ and $v_i = t_{\overline{i}}$ for $i \geq 4$.

Step 4: If $\dim V \geq 3$, then V has a Witt decomposition

$$V \cong Z \oplus W \oplus W^*,$$

where

$$\dim Z = \begin{cases} 0 & \text{if } \dim V \text{ is even and } \Lambda_1 \Lambda_2 \dots \Lambda_n \text{ is a square,} \\ 1 & \text{if } \dim V \text{ is odd,} \\ 2 & \text{if } \dim V \text{ is even and } \Lambda_1 \Lambda_2 \dots \Lambda_n \text{ is not a square.} \end{cases}$$

This follows from an inductive argument using Step 1, Step 2, and Step 3. \square

Lemma 3.7.

(1) Assume that $\dim V = 1$. Then

$$\text{Cliff}_q(\lambda) \cong \begin{cases} Q_1(\mathbb{F}) & \text{if } \overline{\Delta(\lambda)} = \bar{1} \text{ (equivalently, } \Lambda_1 \text{ is a square in } \mathbb{F}), \\ \mathbb{F}(\sqrt{\Lambda_1}) & \text{if } \overline{\Delta(\lambda)} \neq \bar{1} \text{ (equivalently, } \Lambda_1 \text{ is not a square in } \mathbb{F}). \end{cases}$$

(2) Assume that $\dim V = 2$. Then $\text{Cliff}_q(\lambda) \cong \text{Mat}_2(\mathbb{F})$ as (nongraded) algebras and $\text{Cliff}_q(\lambda)_{\bar{0}} \cong \text{Cliff}_q(\Lambda_1 \Lambda_2)$.

Proof. The case (1) corresponds to the ‘‘classical case’’ (Clifford superalgebra over \mathbb{C}) and can be easily verified.

(2) Let $A = \text{Cliff}_q(\lambda)$. Then A is a quaternion algebra over \mathbb{F} . Since it is not a division algebra, by Wedderburn’s Theorem, we have $A \cong \text{Mat}_2(\mathbb{F})$ (see [Lam, Theorem 2.7] for details). The isomorphism $A_{\bar{0}} \cong \text{Cliff}_q(\Lambda_1 \Lambda_2)$ is straightforward. \square

Remark. The superalgebraic structure of $\text{Cliff}_q(\lambda)$ for $\dim V = 2$ is ‘‘explicit’’ only when $\overline{\Delta(\lambda)} = \bar{1}$. In this case, one can show that $\text{Cliff}_q(\lambda) \cong \text{sMat}_{1|1}(\mathbb{F})$.

We are now ready to describe the superalgebra structure of $\text{Cliff}_q(\lambda)$.

Proposition 3.8.

(1) If n is even, then $\text{Cliff}_q(\lambda) \cong \text{Mat}_r(A)$, where $A = \text{Cliff}_q((\Delta(\lambda), 1))$ and $r = 2^{\frac{n}{2}-1}$. Furthermore, $\text{Cliff}_q(\lambda) \cong \text{Mat}_{2r}(\mathbb{F})$ as (nongraded) algebras and

$$\text{Cliff}_q(\lambda)_{\bar{0}} \cong \begin{cases} \text{Mat}_r(\mathbb{F}) \oplus \text{Mat}_r(\mathbb{F}) & \text{if } \overline{\Delta(\lambda)} = \bar{1}, \\ \text{Mat}_r(\mathbb{F}(\sqrt{\Delta(\lambda)})) & \text{if } \overline{\Delta(\lambda)} \neq \bar{1}. \end{cases}$$

(2) If n is odd, then $\text{Cliff}_q(\lambda) \cong \text{Mat}_r(B)$, where $B = \text{Cliff}_q(\Delta(\lambda))$ and $r = 2^{\frac{n-1}{2}}$. Furthermore,

$$\begin{cases} \text{Cliff}_q(\lambda) \cong Q_r(\mathbb{F}), & \text{Cliff}_q(\lambda)_{\bar{0}} \cong \text{Mat}_r(\mathbb{F}) & \text{if } \overline{\Delta(\lambda)} = \bar{1}, \\ \text{Cliff}_q(\lambda) \cong \text{Mat}_r(\mathbb{F}(\sqrt{\Delta(\lambda)})), & \text{Cliff}_q(\lambda)_{\bar{0}} \cong \text{Mat}_r(\mathbb{F}) & \text{if } \overline{\Delta(\lambda)} \neq \bar{1}. \end{cases}$$

In particular, $\text{Cliff}_q(\lambda)$ is a simple superalgebra which is isomorphic to

- a direct sum of two isomorphic simple algebras if n is odd and $\overline{\Delta(\lambda)} = \bar{1}$;

- a simple algebra otherwise.

Proof. We first consider the case when n is even and let $r = 2^{\frac{n}{2}-1}$. If $\Lambda_1 \dots \Lambda_n$ is a square in \mathbb{F} , then (1) is proved by Lemma 3.6 (3) and Lemma 3.3. Now if $\overline{\Delta(\lambda)} \neq \bar{1}$, by Lemma 3.3 and Step 3 in the proof of Lemma 3.6, we have $\text{Cliff}_q(\Lambda) \cong \text{Mat}_r(A)$, where $A = \text{Cliff}_q(\Lambda_1 \dots \Lambda_{n-1}, \Lambda_n)$. We now apply Lemma 3.7 (1),(2) and prove (1).

Next, assume that n is odd and let $r = 2^{\frac{n-1}{2}}$. By Lemma 3.3 and Step 3 in the proof of Lemma 3.6, we have $\text{Cliff}_q(\lambda) \cong \text{Mat}_r(B)$, where B is the 2-dimensional Clifford superalgebra $\text{Cliff}_q(\Lambda_1 \dots \Lambda_n)$. We use Lemma 3.7 (1) to complete the proof. \square

In the statement of the following corollary we allow λ_i to be zero for some i . Recall that $|\lambda|$ is the number of nonzero λ_i . We also set $\lambda_N := (\lambda_{i_1}, \dots, \lambda_{i_{|\lambda|}})$ where $N_\lambda = \{i_1, \dots, i_{|\lambda|}\}$ and $i_1 < \dots < i_{|\lambda|}$.

Corollary 3.9. *Every \mathbb{Z}_2 -graded $\text{Cliff}_q(\lambda_N)$ -module is completely reducible. Furthermore, the superalgebra $\text{Cliff}_q(\lambda)$ has up to isomorphism*

- (1) two simple modules $E^q(\lambda)$ and $\Pi(E^q(\lambda))$ of dimension $2^{k-1}|2^{k-1}$ if $|\lambda| = 2k$ and $\overline{\Delta(\lambda)} = \bar{1}$;
- (2) one simple module $E^q(\lambda) \cong \Pi(E^q(\lambda))$ of dimension $2^k|2^k$ if $|\lambda| = 2k$ and $\overline{\Delta(\lambda)} \neq \bar{1}$ (in particular, if $\lambda_1 > \dots > \lambda_{2k} > 0$);
- (3) one simple module $E^q(\lambda) \cong \Pi(E^q(\lambda))$ of dimension $2^k|2^k$ if $|\lambda| = 2k + 1$.

Proof. Thanks to Lemma 3.1, we may assume that $\lambda_i \neq 0$; i.e., $|\lambda| = n$. The category of all \mathbb{Z}_2 -graded $\text{Cliff}_q(\lambda)$ -modules is equivalent to the category of all nongraded $\text{Cliff}_q(\lambda)_{\bar{0}}$ -modules. Indeed, the reverse correspondence is obtained by

$$V_0 \mapsto \text{Cliff}_q(\lambda) \otimes_{\text{Cliff}_q(\lambda)_{\bar{0}}} V_0.$$

The corollary follows from Proposition 3.8 and the characterization of the simple and indecomposable (nongraded) modules of $\text{Mat}_r(\mathbb{F}) \oplus \text{Mat}_r(\mathbb{F})$, $\text{Mat}_r(\mathbb{F})$, and $\text{Mat}_r(\mathbb{F}(\sqrt{\overline{\Delta(\lambda)}}))$. (This characterization may be found, for example, in [Lang, Chapter XVII].) \square

Example 3.10. Let $n = 3$ and $\lambda = (4, 2, 1)$. We describe the action of $t_{\bar{i}}$ ($i = 1, 2, 3$) on $E^q(\lambda)$. We have

$$\Lambda_1 = (q^2 + q^{-2})(q^4 + q^{-4}), \quad \Lambda_2 = q^2 + q^{-2}, \quad \Lambda_3 = 1.$$

For simplicity, let $t = q^2 + q^{-2}$. We first find a solution of Legendre's equation

$$(3.1) \quad \Lambda_1 X^2 + \Lambda_2 Y^2 + \Lambda_3 Z^2 = 0$$

We follow the proof of Lemma 3.4. Let $Z = tZ'$ and $Y = \sqrt{-1}Y'$. In order to solve the equation $(t^2 - 2)X^2 + tZ'^2 = Y'^2$ we find $C_1 \in \mathbb{C}[t]$ for which $C_1^2 - t$ is a multiple of $t^2 - 2$.

Using the Chinese Remainder Theorem, we choose

$$C_1 = \frac{\sqrt[4]{8}}{4}(1 - \sqrt{-1})t + \frac{\sqrt[4]{2}}{2}(1 + \sqrt{-1}).$$

Then we solve the equation $A_1X_1^2 + BZ_1^2 = Y_1^2$ for $A_1 = -\frac{\sqrt{2}}{4}\sqrt{-1}$ and $B = t$. A solution for this is

$$(X_1, Y_1, Z_1) = \left(1, \frac{\sqrt[4]{8}}{4}(1 - \sqrt{-1}), 0\right).$$

Then (3.1) has a solution

$$(A_1X_1, \sqrt{-1}(Y_1C_1 + BZ_1), t(C_1Z_1 + Y_1)) = \left(-\frac{\sqrt{2}}{4}\sqrt{-1}, \frac{\sqrt{2}}{4}t + \frac{1}{2}\sqrt{-1}, \frac{\sqrt[4]{8}}{4}(1 - \sqrt{-1})t\right).$$

Multiplying by an appropriate constant and changing signs, we fix the following solution of (3.1)

$$w = (X, Y, Z) = (1, \sqrt{-1}t - \sqrt{2}, \sqrt[4]{2}(1 + \sqrt{-1})t).$$

We consider w as an element in V relative to the basis $\{t_{\bar{1}}, t_{\bar{2}}, t_{\bar{3}}\}$. We use Lemma 3.2 to find a Witt decomposition $V = \mathbb{F}w \oplus \mathbb{F}w^* \oplus \mathbb{F}z$. As mentioned in the proof of Lemma 3.2, we find

$$w^* = c(X, Y, -Z),$$

where $c = \frac{\sqrt{-1}}{4\sqrt{2}}t^{-2}$ such that $\beta_\lambda(w, w^*) = 1$ and $\beta_\lambda(w^*, w^*) = 0$. Then, as pointed out in Step 3 of the proof of Lemma 3.6, we can find

$$z = c(tYZ, -t(t^2 - 2)XZ, 0)$$

such that $\beta_\lambda(z, w) = \beta_\lambda(z, w^*) = 0$ and $\beta_\lambda(z, z) = -\frac{1}{4}\Lambda_1\Lambda_2\Lambda_3 = -\frac{1}{4}t^2(t^2 - 2)$. Set $\alpha = \sqrt{\Delta(\lambda)} = \sqrt{t^2(t^2 - 2)}$. Using Lemma 3.3, we define an isomorphism $\Theta : \text{Cliff}_q(\lambda) \rightarrow \text{Mat}_2(\mathbb{F}(\alpha))$ by

$$w \mapsto \begin{pmatrix} 0 & \alpha^{-1} \\ 0 & 0 \end{pmatrix}, \quad w^* \mapsto \begin{pmatrix} 0 & 0 \\ \alpha & 0 \end{pmatrix}, \quad z \mapsto \begin{pmatrix} \alpha & 0 \\ 0 & -\alpha \end{pmatrix}.$$

From Proposition 3.8 and Corollary 3.9 we find that $E^q(\lambda) = \mathbb{F}(\alpha)^{\oplus 2}$. Let v_1 and v_2 be the standard basis vectors of the $\mathbb{F}(\alpha)$ -vector space $E^q(\lambda)$, and let $\bar{v}_i = \alpha v_i$ ($i = 1, 2$). The action of $\text{Cliff}_q(\lambda)$ on $E^q(\lambda)$ is given by

$$\begin{aligned} z(v_1) &= \bar{v}_1, z(v_2) = -\bar{v}_2, z(\bar{v}_1) = t^2(t^2 - 2)v_1, z(\bar{v}_2) = -t^2(t^2 - 2)v_2 \\ w(v_1) &= 0, w(v_2) = (t^2(t^2 - 2))^{-1}\bar{v}_1, w(\bar{v}_1) = 0, w(\bar{v}_2) = v_1 \\ w^*(v_1) &= \bar{v}_2, w^*(v_2) = 0, w^*(\bar{v}_1) = t^2(t^2 - 2)v_2, w^*(\bar{v}_2) = 0 \end{aligned}$$

In order to determine the action of $t_{\bar{i}}$ ($i = 1, 2, 3$) on $E^q(\lambda)$, we need to express $t_{\bar{1}}, t_{\bar{2}}, t_{\bar{3}}$ in terms of z, w, w^* . With simple computations we find:

$$\begin{aligned}
t_{\bar{1}} &= \frac{\sqrt{-1}t^2 - 2}{4\sqrt{2}} \frac{w}{t} + t(t^2 - 2)w^* + \frac{\sqrt[4]{8}(1 - \sqrt{-1})}{2} \frac{\sqrt{-1}t - \sqrt{2}}{t} z, \\
t_{\bar{2}} &= \frac{\sqrt{-1}}{4\sqrt{2}} \frac{\sqrt{-1}t - \sqrt{2}}{t} w + (\sqrt{-1}t^2 - \sqrt{2}t)w^* + \frac{\sqrt[4]{8}(\sqrt{-1} - 1)}{2} \frac{1}{t} z, \\
t_{\bar{3}} &= \frac{1 - \sqrt{-1}}{4\sqrt[4]{2}t} w + \frac{\sqrt{2}(1 + \sqrt{-1})t}{\sqrt[4]{2}} w^*.
\end{aligned}$$

4. HIGHEST WEIGHT REPRESENTATION THEORY OF $U_q(\mathfrak{g})$

A $U_q(\mathfrak{g})$ -module V^q is called a *weight module* if it admits a *weight space decomposition*

$$V^q = \bigoplus_{\mu \in P} V_{\mu}^q, \text{ where } V_{\mu}^q = \{v \in V^q \mid q^h v = q^{\mu(h)} v \text{ for all } h \in P^{\vee}\}.$$

For a weight $U_q(\mathfrak{g})$ -module V^q , we set $\text{wt } V^q = \{\lambda \in P \mid V_{\lambda}^q \neq 0\}$. By the same argument as in [HK, Ch.3], it can be verified that every submodule of a weight $U_q(\mathfrak{g})$ -module is also a weight module. If $\dim_{\mathbb{C}(q)} V_{\mu}^q < \infty$ for all $\mu \in P$, then the *character* of V^q is defined to be

$$\text{ch } V^q = \sum_{\mu \in P} (\dim_{\mathbb{C}(q)} V_{\mu}^q) e^{\mu},$$

where e^{μ} are formal basis elements of the group algebra $\mathbb{C}(q)[P]$ with the multiplication given by $e^{\lambda} e^{\mu} = e^{\lambda + \mu}$ for all $\lambda, \mu \in P$.

A weight module V^q is called a *highest weight module* if it is generated over $U_q(\mathfrak{g})$ by a finite dimensional irreducible $U_q^{\geq 0}$ -module \mathfrak{v}^q . Note that \mathfrak{v}^q also admits a weight space decomposition. We call a vector in \mathfrak{v}^q a *highest weight vector* of V^q . Combining Lemma 2.2 and the triangular decomposition of $U_q(\mathfrak{g})$ (Theorem 2.3), we obtain $V^q = U_q^{-} \mathfrak{v}^q$.

Proposition 4.1. *If \mathfrak{v}^q is a finite dimensional irreducible $U_q^{\geq 0}$ -module with a weight space decomposition $\mathfrak{v}^q = \bigoplus_{\mu \in P} \mathfrak{v}_{\mu}^q$, then \mathfrak{v}^q is irreducible as a U_q^0 -module and $\mathfrak{v}^q = \mathfrak{v}_{\lambda}^q$ for some $\lambda \in P$. Conversely, if \mathfrak{v}^q is an irreducible U_q^0 -module on which the even part of U_q^0 acts by a weight λ , then \mathfrak{v}^q can be endowed with the structure of an irreducible $U_q^{\geq 0}$ -module by letting U_q^{+} act trivially on \mathfrak{v}^q .*

Proof. Because \mathfrak{v}^q is finite dimensional, there exists a weight $\lambda \in P$ such that $\mathfrak{v}_{\lambda}^q \neq 0$ and $\mathfrak{v}_{\lambda + \alpha_i}^q = 0$ for all $i \in I$. Then we have $U_q^{+} \mathfrak{v}_{\lambda}^q = \mathfrak{v}_{\lambda}^q$ and $U_q^0 \mathfrak{v}_{\lambda}^q = \mathfrak{v}_{\lambda}^q$. Thus \mathfrak{v}_{λ}^q is a $U_q^{\geq 0}$ -submodule of \mathfrak{v}^q and hence $\mathfrak{v}_{\lambda}^q = \mathfrak{v}^q$. The other direction is obvious from the defining relations of $U_q(\mathfrak{g})$ in Theorem 2.1. \square

Remark. If \mathfrak{v}^q is a finite dimensional irreducible $U_q^{\geq 0}$ -module which generates a highest weight module V^q of highest weight λ , then, by Proposition 4.1, we know that \mathfrak{v}^q is an

irreducible U_q^0 -module of weight λ . Thus \mathfrak{v}^q is a finite dimensional irreducible module over $\text{Cliff}_q(\lambda) = U_q^0/I^q(\lambda)$. Conversely, if E^q is a finite dimensional irreducible $\text{Cliff}_q(\lambda)$ -module, then it is clear that E^q is an irreducible U_q^0 -module of weight λ .

By Corollary 3.9, we know that, up to isomorphism, $\text{Cliff}_q(\lambda)$ has at most two simple modules: $E^q(\lambda)$ and $\Pi(E^q(\lambda))$. The $U_q(\mathfrak{g})$ -module $W^q(\lambda) = U_q(\mathfrak{g}) \otimes_{U_q^{\geq 0}} E^q(\lambda)$ is called the *Weyl module* of $U_q(\mathfrak{g})$ corresponding to λ (defined up Π).

Proposition 4.2.

- (1) $W^q(\lambda)$ is a free U_q^- -module of rank $\dim E^q(\lambda)$.
- (2) Every highest weight $U_q(\mathfrak{g})$ -module with highest weight λ is a homomorphic image of $W^q(\lambda)$.
- (3) Every Weyl module $W^q(\lambda)$ has a unique maximal submodule $N^q(\lambda)$.

Proof. (1) This is clear from the definition.

(2) Let V^q be a highest weight module with highest weight λ generated by the irreducible $U_q^{\geq 0}$ -module \mathfrak{v}^q . Because \mathfrak{v}^q is irreducible over $\text{Cliff}_q(\lambda)$, it is isomorphic to $E^q(\lambda)$ up to Π . Thus the map $\phi : W^q(\lambda) \rightarrow V^q$ induced by $E^q(\lambda) \rightarrow \mathfrak{v}^q$ is a surjective $U_q(\mathfrak{g})$ -module homomorphism.

(3) Since $E^q(\lambda)$ is an irreducible $\text{Cliff}_q(\lambda)$ -module, any proper submodule N^q of $W^q(\lambda)$ does not contain highest weight vectors (the vectors in $E^q(\lambda)$). That is, N^q must lie in $\bigoplus_{\mu < \lambda} W^q(\lambda)_\mu$. Thus the sum of two proper submodules is again a proper submodule of $W^q(\lambda)$. Then the sum $N^q(\lambda)$ of all proper submodules of $W^q(\lambda)$ is the unique maximal submodule of $W^q(\lambda)$. \square

For $\lambda \in P$, the unique irreducible quotient $V^q(\lambda) := W^q(\lambda)/N^q(\lambda)$ is called the *irreducible highest weight module* over $U_q(\mathfrak{g})$ with highest weight λ (defined up to Π).

We introduce the notation

$$[n]_q := \frac{q^n - q^{-n}}{q - q^{-1}},$$

which is called a *q-integer*. We also define $[0]_q! := 1$ and $[n]_q! := [n]_q \cdot [n-1]_q \cdots [1]_q$. We define the *divided powers* of e_i and f_i as follows:

$$e_i^{(k)} := \frac{e_i^k}{[k]_q!}, \quad f_i^{(k)} := \frac{f_i^k}{[k]_q!}.$$

By a straightforward induction argument, we can prove the following lemma.

Lemma 4.3. *For all $i \in I$ and $k \in \mathbb{Z}_{\geq 0}$, we have*

$$e_i f_i^{(k)} = f_i^{(k)} e_i + f_i^{(k-1)} \frac{q^{h_i} q^{-k+1} - q^{-h_i} q^{k-1}}{q - q^{-1}}.$$

Proposition 4.4. *Let $\lambda \in \Lambda^+$ and $V^q(\lambda)$ be the irreducible highest weight $U_q(\mathfrak{g})$ -module generated by an irreducible finite dimensional $U_q^{\geq 0}$ -module \mathbf{v}^q . Then $f_i^{\lambda(h_i)+1}v = 0$ for all $v \in \mathbf{v}^q$ and $i \in I$.*

Proof. Lemma 4.3 implies

$$e_i f_i^{(k)} v = [\lambda(h_i) - k + 1]_q f_i^{(k-1)} v \quad \text{for all } v \in \mathbf{v}^q.$$

If $k = \lambda(h_i) + 1$, we see that $e_i f_i^{\lambda(h_i)+1} v = 0$. Moreover, for $j \neq i$, we already know $e_j f_i^{\lambda(h_i)+1} v = 0$ and $e_{\bar{j}} f_i^{\lambda(h_i)+1} v = 0$, since $V^q(\lambda) = \bigoplus_{\mu \leq \lambda} V_\mu^q$.

Suppose that $e_{\bar{i}} f_i^{\lambda(h_i)+1} v \neq 0$. We have

$$\begin{aligned} e_i(e_{\bar{i}} f_i^{\lambda(h_i)+1} v) &= e_{\bar{i}}(e_i f_i^{\lambda(h_i)+1} v) = 0, \\ e_{\bar{i}}(e_i f_i^{\lambda(h_i)+1} v) &= -\frac{q - q^{-1}}{q + q^{-1}} e_i^2 f_i^{\lambda(h_i)+1} v = 0. \end{aligned}$$

Also, $e_j(e_{\bar{i}} f_i^{\lambda(h_i)+1} v) = e_{\bar{j}}(e_{\bar{i}} f_i^{\lambda(h_i)+1} v) = 0$ for $j \neq i$, since $V^q(\lambda) = \bigoplus_{\mu \leq \lambda} V_\mu^q$.

If $\lambda(h_i) \geq 1$, then $\text{wt}(e_{\bar{i}} f_i^{\lambda(h_i)+1} v) = \lambda - \lambda(h_i)\alpha_i < \lambda$. Thus $e_{\bar{i}} f_i^{\lambda(h_i)+1} v$ would generate a nontrivial proper submodule of $V^q(\lambda)$, which contradicts the irreducibility of $V^q(\lambda)$.

If $\lambda(h_i) = 0$, then we have $\lambda_i = \lambda_{i+1} = 0$ so that $k_{\bar{i}} v = k_{\bar{i}+1} v = 0$ by Lemma 3.1. From the defining relation of $U_q(\mathfrak{g})$, we know

$$e_{\bar{i}} f_i v = f_i e_{\bar{i}} v + (q^{k_{i+1}} k_{\bar{i}} - q^{k_{\bar{i}}} k_{i+1}) v = 0.$$

Therefore, in any case, $e_{\bar{i}} f_i^{\lambda(h_i)+1} v = 0$ for all $v \in \mathbf{v}^q$.

Similarly, if $f_i^{\lambda(h_i)+1} v \neq 0$, it would generate a nontrivial proper submodule of $V^q(\lambda)$. Hence we conclude $f_i^{\lambda(h_i)+1} v = 0$ for all $v \in \mathbf{v}^q$. \square

5. CLASSICAL LIMITS

Let $\mathbf{A}_1 := \{f/g \in \mathbb{C}(q) \mid f, g \in \mathbb{C}[q], g(1) \neq 0\}$. For an integer $n \in \mathbb{Z}$, we formally define

$$[y; n]_x := \frac{yx^n - y^{-1}x^{-n}}{x - x^{-1}}, \quad (y; n)_x := \frac{yx^n - 1}{x - 1}.$$

For example,

$$[q^h; 0]_q = \frac{q^h - q^{-h}}{q - q^{-1}}, \quad (q^h; 0)_q = \frac{q^h - 1}{q - 1}.$$

Definition 5.1. We define the \mathbf{A}_1 -form $U_{\mathbf{A}_1}$ of the quantum superalgebra $U_q(\mathfrak{g})$ to be the \mathbf{A}_1 -subalgebra of $U_q(\mathfrak{g})$ with 1 generated by the elements $e_i, e_{\bar{i}}, f_i, f_{\bar{i}}, q^h, k_{\bar{l}}$ and $(q^h; 0)_q$ ($i \in I, l \in J, h \in P^\vee$).

We denote by $U_{\mathbf{A}_1}^+$ (respectively, $U_{\mathbf{A}_1}^-$) the \mathbf{A}_1 -subalgebra of $U_q(\mathfrak{g})$ with 1 generated by $e_i, e_{\bar{i}}$ (respectively, $f_i, f_{\bar{i}}$) for $i \in I$, and by $U_{\mathbf{A}_1}^0$ the \mathbf{A}_1 -subalgebra of $U_q(\mathfrak{g})$ with 1 generated by $q^h, k_{\bar{l}}$ and $(q^h; 0)_q$ for $l \in J$, $h \in P^\vee$.

Lemma 5.2.

- (1) $(q^h; n)_q \in U_{\mathbf{A}_1}^0$ for all $n \in \mathbb{Z}$ and $h \in P^\vee$.
- (2) $[q^h; 0]_q \in U_{\mathbf{A}_1}^0$ for all $n \in \mathbb{Z}$ and $h \in P^\vee$.

Proof. Our assertions follow immediately from the following identities:

$$(q^h; n)_q = q^n (q^h; 0)_q + \frac{q^n - 1}{q - 1},$$

$$[q^h; 0]_q = q \frac{q - 1}{q^2 - 1} (1 + q^{-h}) (q^h; 0)_q.$$

□

Note that

$$k_{\bar{i}}^2 = [q^{2k_i}; 0]_{q^2} = q^2 \frac{q^2 - 1}{q^4 - 1} (1 + q^{-2k_i}) \frac{1}{q + 1} (q^{2k_i}; 0)_q.$$

Proposition 5.3. *We have the triangular decomposition of the algebra $U_{\mathbf{A}_1}$. Namely,*

$$U_{\mathbf{A}_1} \cong U_{\mathbf{A}_1}^- \otimes U_{\mathbf{A}_1}^0 \otimes U_{\mathbf{A}_1}^+$$

as \mathbf{A}_1 -modules.

Proof. Recall the canonical isomorphism $U_q(\mathfrak{g}) \xrightarrow{\sim} U_q^- \otimes U_q^0 \otimes U_q^+$ given by Theorem 2.3. The following commutation relations hold:

$$\begin{aligned} e_i(q^h; 0)_q &= (q^h; -\alpha_i(h))_q e_i, & e_{\bar{i}}(q^h; 0)_q &= (q^h; -\alpha_i(h))_q e_{\bar{i}}, \\ (q^h; 0)_q f_i &= f_i(q^h; -\alpha_i(h))_q, & (q^h; 0)_q f_{\bar{i}} &= f_{\bar{i}}(q^h; -\alpha_i(h))_q, \\ e_i f_i &= f_i e_i + [q^{k_i - k_{i+1}}; 0]_q, \\ e_{i+1} f_i &= q^{-1} f_i e_{i+1}, & e_i f_{i+1} &= q f_{i+1} e_i, & e_i f_j - f_j e_i &= 0 \text{ for } |i - j| > 1, \\ e_{\bar{i}} f_{\bar{i}} &= -f_{\bar{i}} e_{\bar{i}} + [q^{k_i + k_{i+1}}; 0]_q + (q - q^{-1}) k_{\bar{i}} k_{\bar{i}+1}, \\ e_{\bar{i}+1} f_{\bar{i}} &= -q^{-1} f_{\bar{i}} e_{\bar{i}+1}, & e_{\bar{i}} f_{\bar{i}+1} &= -q f_{\bar{i}+1} e_{\bar{i}}, & e_{\bar{i}} f_{\bar{j}} - f_{\bar{j}} e_{\bar{i}} &= 0 \text{ for } |i - j| > 1. \end{aligned}$$

Together with Lemma 5.2, one can show that the image of the canonical isomorphism lies inside $U_{\mathbf{A}_1}^- \otimes U_{\mathbf{A}_1}^0 \otimes U_{\mathbf{A}_1}^+$ when restricted to $U_{\mathbf{A}_1}$. Its inverse map is given by multiplication. Hence the two spaces are isomorphic as \mathbf{A}_1 -modules. □

In what follows, V^q is a highest weight module over $U_q(\mathfrak{g})$ with highest weight $\lambda \in P$ generated by a finite dimensional irreducible $U_q^{\geq 0}$ -submodule \mathfrak{v}^q . Then \mathfrak{v}^q is a finite dimensional irreducible $\text{Cliff}_q(\lambda)$ -module. Since it is irreducible, it is generated by a nonzero vector $v \in (\mathfrak{v}^q)_0$; i.e., $\mathfrak{v}^q = \text{Cliff}_q(\lambda)v$. Note that

$$\frac{q^{2n} - q^{-2n}}{q^2 - q^{-2}} = q^{2n-2} + q^{2n-6} + \cdots + q^{-2n+6} + q^{-2n+2} \in \mathbf{A}_1 \quad \text{for } n \in \mathbb{Z}_{>0}.$$

We denote by $\text{Cliff}_{\mathbf{A}_1}(\lambda)$ the \mathbf{A}_1 -subalgebra of $\text{Cliff}_q(\lambda)$ generated by $\{t_{\bar{i}} \mid i \in J\}$.

Definition 5.4. Let V^q be a highest weight $U_q(\mathfrak{g})$ -module generated by a finite dimensional irreducible $U_q^{\geq 0}$ -module \mathfrak{v}^q and let $E^{\mathbf{A}_1}(\lambda)$ be the $\text{Cliff}_{\mathbf{A}_1}(\lambda)$ -submodule of $\mathfrak{v}^q \cong E^q(\lambda)$ generated by a nonzero element $v \in (\mathfrak{v}^q)_0$. The \mathbf{A}_1 -form of V^q is defined to be the $U_{\mathbf{A}_1}$ -submodule $V_{\mathbf{A}_1}$ of V^q generated by $E^{\mathbf{A}_1}(\lambda)$.

In what follows, V^q will denote a highest weight $U_q(\mathfrak{g})$ -module.

Proposition 5.5. $V_{\mathbf{A}_1} = U_{\mathbf{A}_1}^- E^{\mathbf{A}_1}(\lambda)$.

Proof. In view of Proposition 5.3, it suffices to show that $U_{\mathbf{A}_1}^+ E^{\mathbf{A}_1}(\lambda) = E^{\mathbf{A}_1}(\lambda)$ and $U_{\mathbf{A}_1}^0 E^{\mathbf{A}_1}(\lambda) = E^{\mathbf{A}_1}(\lambda)$. The first assertion is clear by the definition of highest weight modules. For the second assertion, we observe that

$$\begin{aligned} q^h w &= q^{\lambda(h)} w, \\ (q^h; 0)_q w &= \frac{q^{\lambda(h)} - 1}{q - 1} w \quad \text{for all } w \in E^{\mathbf{A}_1}(\lambda). \end{aligned}$$

Hence we obtain $V_{\mathbf{A}_1} = U_{\mathbf{A}_1} E^{\mathbf{A}_1}(\lambda) = U_{\mathbf{A}_1}^- E^{\mathbf{A}_1}(\lambda)$. \square

For each $\mu \in P$, let us denote by $(V_{\mathbf{A}_1})_{\mu}$ the space $V_{\mathbf{A}_1} \cap V_{\mu}^q$. The following assertion can be proved using the same arguments as in [HK, Proposition 3.3.6].

Proposition 5.6. $V_{\mathbf{A}_1}$ has the weight space decomposition $V_{\mathbf{A}_1} = \bigoplus_{\mu \leq \lambda} (V_{\mathbf{A}_1})_{\mu}$.

Proposition 5.7. For each $\mu \in P$, the weight space $(V_{\mathbf{A}_1})_{\mu}$ is a free \mathbf{A}_1 -module with $\text{rank}_{\mathbf{A}_1}(V_{\mathbf{A}_1})_{\mu} = \dim_{\mathbb{C}(q)} V_{\mu}^q$. In particular, $\text{rank}_{\mathbf{A}_1} E^{\mathbf{A}_1}(\lambda) = \dim_{\mathbb{C}(q)} E^q(\lambda)$.

Proof. Because \mathbf{A}_1 is a principal ideal domain, every finitely generated torsion free module over \mathbf{A}_1 is free. Furthermore, since $\mathbb{C}(q)$ is the field of quotients of the integral domain \mathbf{A}_1 , a finite subset of a $\mathbb{C}(q)$ -vector space is linearly independent over $\mathbb{C}(q)$ if and only if it is linearly independent over \mathbf{A}_1 . Thus it is enough to show that each V_{μ}^q has a $\mathbb{C}(q)$ -basis which is also contained in $(V_{\mathbf{A}_1})_{\mu}$. The highest weight space $\mathfrak{v}^q = E^q(\lambda)$ has a linearly independent subset of $\{t_1^{\epsilon_1} t_2^{\epsilon_2} \cdots t_n^{\epsilon_n} v \mid \epsilon_j = 0 \text{ or } 1\}$ which generates $E^q(\lambda)$ over $\mathbb{C}(q)$, since $E^q(\lambda) = \text{Cliff}_q(\lambda)v$. By definition, this subset is contained in $E^{\mathbf{A}_1}(\lambda)$. For V_{μ}^q , it is easy to show that there is a

basis of V_μ^q whose elements are of the form $f_\zeta t_1^{\epsilon_1} t_2^{\epsilon_2} \cdots t_n^{\epsilon_n} v$, where f_ζ are monomials in f_i and $f_{\bar{j}}$. This basis is also contained in $(V_{\mathbf{A}_1})_\mu$, which proves the proposition. \square

Corollary 5.8. *The map $\phi : \mathbb{C}(q) \otimes_{\mathbf{A}_1} V_{\mathbf{A}_1} \longrightarrow V^q$ given by $f \otimes v \longmapsto fv$ ($f \in \mathbb{C}(q), v \in V_{\mathbf{A}_1}$) is a $\mathbb{C}(q)$ -linear isomorphism.*

Let \mathbf{J}_1 be the ideal of \mathbf{A}_1 generated by $q - 1$. Then there is a canonical isomorphism of fields

$$\mathbf{A}_1/\mathbf{J}_1 \xrightarrow{\sim} \mathbb{C} \quad \text{given by } f(q) + \mathbf{J}_1 \longmapsto f(1).$$

Define the \mathbb{C} -linear vector spaces

$$\begin{aligned} U_1 &= (\mathbf{A}_1/\mathbf{J}_1) \otimes_{\mathbf{A}_1} U_{\mathbf{A}_1}, \\ V^1 &= (\mathbf{A}_1/\mathbf{J}_1) \otimes_{\mathbf{A}_1} V_{\mathbf{A}_1}. \end{aligned}$$

Then V^1 is naturally a U_1 -module. Note that $U_1 \cong U_{\mathbf{A}_1}/\mathbf{J}_1 U_{\mathbf{A}_1}$ and $V^1 \cong V_{\mathbf{A}_1}/\mathbf{J}_1 V_{\mathbf{A}_1}$. We use the bar notation for the images under these maps. The passage under these maps is referred to as taking the *classical limit*.

Since $V_{\mathbf{A}_1} = U_{\mathbf{A}_1} E^{\mathbf{A}_1}(\lambda)$, we have:

$$V^1 \cong V_{\mathbf{A}_1}/\mathbf{J}_1 V_{\mathbf{A}_1} = U_{\mathbf{A}_1} E^{\mathbf{A}_1}(\lambda)/\mathbf{J}_1 U_{\mathbf{A}_1} E^{\mathbf{A}_1}(\lambda) = (U_{\mathbf{A}_1}/\mathbf{J}_1 U_{\mathbf{A}_1}) \cdot (E^{\mathbf{A}_1}(\lambda)/\mathbf{J}_1 E^{\mathbf{A}_1}(\lambda)).$$

Hence V^1 is generated by $E^{\mathbf{A}_1}(\lambda)/\mathbf{J}_1 E^{\mathbf{A}_1}(\lambda)$ over U^1 . For each $\mu \in P$, denote by V_μ^1 the space $(\mathbf{A}_1/\mathbf{J}_1) \otimes_{\mathbf{A}_1} (V_{\mathbf{A}_1})_\mu \cong (V_{\mathbf{A}_1})_\mu/\mathbf{J}_1 (V_{\mathbf{A}_1})_\mu$.

Proposition 5.9.

- (1) $V^1 = \bigoplus_{\mu \leq \lambda} V_\mu^1$
- (2) For each $\mu \in P$, $\dim_{\mathbb{C}} V_\mu^1 = \text{rank}_{\mathbf{A}_1} (V_{\mathbf{A}_1})_\mu$.

Proof. The first assertion follows from Proposition 5.6. Using the same argument as in [HK, Lemma 3.4.1], we can prove the second assertion. \square

Let $\bar{h} \in U_1$ be the classical limit of $(q^h; 0)_q \in U_{\mathbf{A}_1}$. Using [HK, Lemma 3.4.3], we have:

Lemma 5.10.

- (1) For all $h \in p^\vee$, we have $\overline{q^h} = 1$.
- (2) For any $h, h' \in P^\vee$, $\overline{h + h'} = \bar{h} + \bar{h}'$.

Theorem 5.11.

- (1) The elements $\bar{e}_i, \bar{e}_{\bar{i}}, \bar{f}_i, \bar{f}_{\bar{i}}$, ($i \in I$), \bar{k}_l ($l \in J$) and \bar{h} ($h \in P^\vee$) satisfy the defining relations of $U(\mathfrak{g})$. Hence there exists a surjective \mathbb{C} -algebra homomorphism $\psi : U(\mathfrak{g}) \longrightarrow U_1$ and the U_1 -module V^1 has a $U(\mathfrak{g})$ -module structure.

- (2) For each $\mu \in P$ and $h \in P^\vee$, the element \bar{h} acts on V_μ^1 as scalar multiplication by $\mu(h)$. So V_μ^1 is the μ -weight space of the $U(\mathfrak{g})$ -module V^1 .
- (3) There is an isomorphism $\text{Cliff}(\lambda) \xrightarrow{\sim} \text{Cliff}_1(\lambda) := \text{Cliff}_{\mathbf{A}_1}(\lambda)/\mathbf{J}_1 \text{Cliff}_{\mathbf{A}_1}(\lambda)$.
- (4) As a $U(\mathfrak{g})$ -module, V^1 is a highest weight module or the sum of two highest weight modules with highest weight $\lambda \in P$

Proof. (1) The first relation for $U(\mathfrak{g})$ is trivial. Since

$$\begin{aligned} (q^h; 0)_q e_i - e_i(q^h; 0)_q &= e_i(q^h; \alpha_i(h))_q - e_i(q^h; 0)_q \\ &= \frac{q^{\alpha_i(h)} - 1}{q - 1} e_i q^h, \end{aligned}$$

we obtain $[\bar{h}, \bar{e}_i] = \alpha_i(h) \bar{e}_i$ by letting $q \rightarrow 1$. Similarly,

$$[\bar{h}, \bar{e}_i] = \alpha_i(h) \bar{e}_i, \quad [\bar{h}, \bar{f}_i] = -\alpha_i(h) \bar{f}_i, \quad [\bar{h}, \bar{f}_i] = -\alpha_i(h) \bar{f}_i \quad \text{and} \quad [\bar{h}, \bar{k}_l] = 0.$$

We have

$$e_i f_i - f_i e_i = [q^{h_i}; 0]_q = \frac{q}{q+1} (1 + q^{-h_i}) (q^{h_i}; 0)_q.$$

Taking the classical limit to both sides above leads to $\bar{e}_i \bar{f}_i - \bar{f}_i \bar{e}_i = \frac{1}{2} 2\bar{h}_i = \bar{h}_i$.

Also

$$k_i^2 = [q^{2k_i}; 0]_{q^2} = q^2 \frac{q^2 - 1}{q^4 - 1} (1 + q^{-2k_i}) \frac{1}{q+1} (q^{2k_i}; 0)_q.$$

When we take $q \rightarrow 1$, we obtain $\bar{k}_i^2 = \bar{k}_i$.

Since we can obtain the following relations in $U(\mathfrak{g})$ by the Jacobi identity,

$$[\bar{e}_i, [e_i, e_j]] = [[\bar{e}_i, e_i], e_j] + [e_i, [\bar{e}_i, e_j]] = [e_i, [\bar{e}_i, e_j]], \quad \text{for } |i - j| = 1,$$

in order to prove the corresponding relations in U_1 , it suffices to show that $[\bar{e}_i, [\bar{e}_i, \bar{e}_j]] = 0$. The latter relation can be checked easily by letting $q \rightarrow 1$. The rest of the relations can be derived in a similar manner.

Therefore, there exists a surjective algebra homomorphism $\psi : U(\mathfrak{g}) \rightarrow U_1$ defined by $e_i \mapsto \bar{e}_i, e_{\bar{i}} \mapsto \bar{e}_{\bar{i}}, f_i \mapsto \bar{f}_i, f_{\bar{i}} \mapsto \bar{f}_{\bar{i}}, h \mapsto \bar{h}, k_{\bar{l}} \mapsto \bar{k}_{\bar{l}}$ ($i \in I, l \in J$), which can be used to define a $U(\mathfrak{g})$ -module structure on V^1 .

- (2) For $v \in (V_{\mathbf{A}_1})_\mu$ and $h \in P^\vee$, we have

$$(q^h; 0)_q v = \frac{q^{\mu(h)} - 1}{q - 1} v.$$

Taking the classical limit of both sides yields our assertion.

- (3) Note that $\overline{t_i t_j} + \overline{t_j t_i} = 2\delta_{ij}\lambda_i$ in $\text{Cliff}_1(\lambda)$ and $\text{Cliff}(\lambda)$ is the associative \mathbb{C} -algebra with $\mathbf{1}$ generated by $\{k_i \mid i \in J\}$ with defining relations $k_i k_j + k_j k_i = 2\delta_{ij}\lambda_i$. Thus we have a surjective \mathbb{C} -algebra homomorphism $\text{Cliff}(\lambda) \rightarrow \text{Cliff}_1(\lambda)$. Observe that

$$\begin{aligned} \dim_{\mathbb{C}} \text{Cliff}_1(\lambda) &= \text{rank}_{\mathbf{A}_1} \text{Cliff}_{\mathbf{A}_1}(\lambda) \\ &= \dim_{\mathbb{C}(q)} \text{Cliff}_q(\lambda) \\ &= \dim_{\mathbb{C}} \text{Cliff}(\lambda). \end{aligned}$$

The first two equalities follow by using the same reasoning as in Proposition 5.9 and Proposition 5.7, respectively. It is well known that the dimension of the Clifford algebra associated with a symmetric bilinear form on a vector space of dimension k is 2^k . This result holds for any base field of characteristic different from 2. Thus we proved the last equality.

- (4) V^q is generated by a finite dimensional irreducible $U_q^{\geq 0}$ -submodule $\mathbf{v}^q \cong E^q(\lambda)$ up to Π . By Corollary 3.9

$$\dim E^q(\lambda) = \begin{cases} 2^k & \text{if } |\lambda| = 2k \text{ and } \overline{\Delta(\lambda)} = \bar{1}, \\ 2^{k+1} & \text{if } |\lambda| = 2k \text{ and } \overline{\Delta(\lambda)} \neq \bar{1}, \\ 2^{k+1} & \text{if } |\lambda| = 2k + 1. \end{cases}$$

It is well known that the dimension of the \mathbb{Z}_2 -graded irreducible $\text{Cliff}(\lambda)$ -modules is $2^{\lfloor \frac{|\lambda|-1}{2} \rfloor} |2|^{\lfloor \frac{|\lambda|-1}{2} \rfloor}$ (see, for example, [ABS]). With this in mind we deduce that $E^{\mathbf{A}_1}(\lambda)/\mathbf{J}_1 E^{\mathbf{A}_1}(\lambda)$ is an irreducible $\text{Cliff}(\lambda)$ -module when $|\lambda| = 2k + 1$ or $|\lambda| = 2k$ and $\overline{\Delta(\lambda)} = 1$, and the direct sum of two irreducible $\text{Cliff}(\lambda)$ -modules otherwise. Since $E^q(\lambda)$ is a parity invariant module over $\text{Cliff}_q(\lambda)$ for $|\lambda| = 2k$ and $\overline{\Delta(\lambda)} \neq \bar{1}$, $E^{\mathbf{A}_1}(\lambda)/\mathbf{J}_1 E^{\mathbf{A}_1}(\lambda)$ is a parity invariant $\text{Cliff}(\lambda)$ -module as well. Hence $E^{\mathbf{A}_1}(\lambda)/\mathbf{J}_1 E^{\mathbf{A}_1}(\lambda) = \mathbf{v}(\lambda) \oplus \Pi \mathbf{v}(\lambda)$ for some irreducible $\text{Cliff}(\lambda)$ -module $\mathbf{v}(\lambda)$. By definition, V^1 is a highest weight $U(\mathfrak{g})$ -module generated by $E^{\mathbf{A}_1}(\lambda)/\mathbf{J}_1 E^{\mathbf{A}_1}(\lambda)$ or the sum of two highest weight modules generated by $\mathbf{v}(\lambda)$ and $\Pi \mathbf{v}(\lambda)$ for some irreducible $\text{Cliff}(\lambda)$ -module $\mathbf{v}(\lambda)$. \square

By Propositions 5.7 and 5.9 and Theorem 5.11, we obtain the following identity between the characters of a highest weight $U(\mathfrak{g})$ -module and a highest weight $U_q(\mathfrak{g})$ -module.

Proposition 5.12. $\text{ch } V^1 = \text{ch } V^q$.

Corollary 5.13. $V^q(\lambda)$ is finite dimensional if and only if $\lambda \in \Lambda^+$.

Proof. Let $V^q = V^q(\lambda)$. If $\lambda \in \Lambda^+$, then we have $f_i^{\lambda(h_i)+1} v = 0$ for all $v \in V_\lambda^q$ by Proposition 4.4. Taking the classical limit, we have $\bar{f}_i^{\lambda(h_i)+1} \bar{v} = 0$ for all $\bar{v} \in V_\lambda^1$. Because V^1 is a

highest weight module or the sum of two highest weight modules, it is finite dimensional by Proposition 1.9, and hence V^q is finite dimensional by Proposition 5.12. Conversely, assume that λ is not in Λ^+ . Then V^1 has a submodule which is a highest weight module and whose irreducible quotient is isomorphic to an irreducible highest weight module with highest weight λ . It is not finite dimensional by (2) of Proposition 1.4. Again by Proposition 5.12, V^q cannot be finite dimensional. \square

Theorem 5.14. *If $\lambda \in \Lambda^+ \cap P_{\geq 0}$ and V^q is the irreducible highest weight $U_q(\mathfrak{g})$ -module $V^q(\lambda)$ with highest weight λ , then V^1 is isomorphic to*

- (1) $V(\lambda)$ or $\Pi V(\lambda)$ if $|\lambda| = 2k$ and $\overline{\Delta(\lambda)} = \bar{1}$,
- (2) $V(\lambda) \oplus \Pi V(\lambda)$ if $|\lambda| = 2k$ and $\overline{\Delta(\lambda)} \neq \bar{1}$ (in particular, if $\lambda_1 > \dots > \lambda_{2k} > 0$),
- (3) $V(\lambda) \cong \Pi V(\lambda)$ if $|\lambda| = 2k + 1$.

$$\text{Hence, } \text{ch } V^q(\lambda) = \begin{cases} \text{ch } V(\lambda) & \text{if } |\lambda| = 2k \text{ and } \overline{\Delta(\lambda)} = \bar{1}, \\ 2 \text{ch } V(\lambda) & \text{if } |\lambda| = 2k \text{ and } \overline{\Delta(\lambda)} \neq \bar{1}, \\ \text{ch } V(\lambda) & \text{if } |\lambda| = 2k + 1. \end{cases}$$

Proof. By Theorem 5.11 (4), V^1 is a highest weight module or the sum of two highest weight modules over $U(\mathfrak{g})$ with highest weight λ . By Proposition 1.8, we have

$$V^1 \cong \begin{cases} V(\lambda) \text{ or } \Pi V(\lambda) & \text{if } |\lambda| = 2k \text{ and } \overline{\Delta(\lambda)} = \bar{1}, \\ V(\lambda) \oplus \Pi V(\lambda) & \text{if } |\lambda| = 2k \text{ and } \overline{\Delta(\lambda)} \neq \bar{1}, \\ V(\lambda) \cong \Pi V(\lambda) & \text{if } |\lambda| = 2k + 1. \end{cases}$$

The second assertion follows from Proposition 5.12. \square

Remark. The main reason we restrict our attention in Theorem 5.14 to the dominant set of weights $\Lambda^+ \cap P_{\geq 0}$ is the statement of Proposition 1.8. We still believe that the theorem holds in more general setting and conjecture that it is true for any weight $\lambda \in \Lambda^+$ for which the generic character formula (1.4) holds.

Corollary 5.15. *If V^q is a finite dimensional highest weight module over $U_q(\mathfrak{g})$ with highest weight $\lambda \in \Lambda^+ \cap P_{\geq 0}$, then V^q is isomorphic to $V^q(\lambda)$ up to Π .*

Proof. Note that V^1 is a highest weight module or the sum of two highest weight modules over $U(\mathfrak{g})$ with highest weight λ and it is finite dimensional by Proposition 5.12. From Proposition 1.8, we know that V^1 is an irreducible module or the direct sum of two irreducible modules. Thus we get $\text{ch } V^q = \text{ch } V^1 = \text{ch } V^q(\lambda)$ by Theorem 5.14 and hence $V^q \cong V^q(\lambda)$. \square

Define the subalgebras $U_1^\pm := \mathbf{A}_1/\mathbf{J}_1 \otimes_{\mathbf{A}_1} U_{\mathbf{A}_1}^\pm$ and $U_1^0 := \mathbf{A}_1/\mathbf{J}_1 \otimes_{\mathbf{A}_1} U_{\mathbf{A}_1}^0$ of U_1 .

Theorem 5.16. *The classical limit U_1 of $U_q(\mathfrak{g})$ is isomorphic to the universal enveloping algebra $U(\mathfrak{g})$.*

Proof. By Theorem 5.11 (1), there exists a surjective algebra homomorphism $\psi : U(\mathfrak{g}) \rightarrow U_1$ defined by $e_i \mapsto \bar{e}_i$, $e_{\bar{i}} \mapsto \bar{e}_{\bar{i}}$, $f_i \mapsto \bar{f}_i$, $f_{\bar{i}} \mapsto \bar{f}_{\bar{i}}$, $h \mapsto \bar{h}$, $k_{\bar{l}} \mapsto \bar{k}_{\bar{l}}$ for $i \in I$, $h \in P^\vee$ and $l \in J$. From (1.3), $U(\mathfrak{g}) \cong U^- \otimes U^0 \otimes U^+$.

We first show that U^0 is isomorphic to U_1^0 . Consider the restriction ψ_0 of ψ to U^0 . Note that $\text{Cliff}_{\mathbf{A}_1}(\lambda)$ is a $U_{\mathbf{A}_1}^0$ -module. Indeed, as in the proof of Proposition 5.5, we know that

$$q^h w = q^{\lambda(h)} w,$$

$$(q^h; 0)_q w = \frac{q^{\lambda(h)} - 1}{q - 1} w \quad \text{for all } w \in \text{Cliff}_{\mathbf{A}_1}(\lambda).$$

In particular, the action of $k_{\bar{i}}$ is just the left multiplication by $t_{\bar{i}}$. Let $g \in \ker \psi_0$. By the Poincaré-Birkhoff-Witt theorem, we can write $g = \sum_{i=1}^{2n} g_i k_{\bar{\eta}_i}$, where $k_{\bar{\eta}_i} = k_{\bar{1}}^{a_1} \cdots k_{\bar{n}}^{a_n}$, $0 \leq a_j \leq 1$ for all $j \in J$ and each g_i is a polynomial in k_1, \dots, k_n . For each $\lambda \in P$ we have

$$0 = \psi_0(g) \cdot \bar{\mathbf{1}} = \sum_{i=1}^{2n} \lambda(g_i) \bar{t}_{\bar{\eta}_i} \in \text{Cliff}_1(\lambda),$$

where $\lambda(g_i)$ denotes the polynomial in λ_j corresponding to g_i . Since $\{\bar{t}_{\bar{\eta}_i}\}$ is a linearly independent subset of $\text{Cliff}_1(\lambda) \cong \text{Cliff}(\lambda)$, we have $\lambda(g_i) = 0$ for all $i = 1, \dots, 2n$. Since we may take any integer value for λ_j , g_i must be zero for all $i = 1, \dots, 2n$ and hence g is identically zero. Thus ψ_0 is injective.

Next we show that the restriction of ψ_- of ψ to U^- is an isomorphism of U^- onto U_1^- . Suppose $\ker \psi_- \neq 0$ and $u = \sum b_\zeta f_\zeta \in \ker \psi_-$, where $b_\zeta \in \mathbb{C}$ and f_ζ are monomials in f_i and $f_{\bar{i}}$'s. Let N be the maximal length of the monomials f_ζ in the expression of u , and choose $\lambda \in \Lambda^+ \cap P_{\geq 0}$ satisfying $\lambda(h_i) > N$ and $|\lambda| = 2k$ and $\Delta(\lambda) = \bar{\mathbf{1}}$ or $|\lambda| = 2k + 1$ for all $i \in I$. By Theorem 5.14, the classical limit V^1 of $V^q(\lambda)$ is isomorphic to the irreducible $U(\mathfrak{g})$ -module $V(\lambda)$ when $|\lambda| = 2k$ and $\Delta(\lambda) = \bar{\mathbf{1}}$, or $|\lambda| = 2k + 1$. Set $r = 2^{\lfloor \frac{|\lambda|+1}{2} \rfloor}$. Consider the map $\phi : (U^-)^{\oplus r} \rightarrow V^1$, given by $(x_1, \dots, x_n) \mapsto \sum_{i=1}^r \psi(x_i) \cdot v_i$ for a basis $\{v_i \mid i = 1, \dots, r\}$ of V_λ^1 . Then by Proposition 1.8 and Proposition 1.9, $\ker \phi$ is the left ideal of $(U^-)^{\oplus r}$ generated by $(f_i^{\lambda(h_i)+1}, 0, \dots, 0), \dots, (0, \dots, 0, f_i^{\lambda(h_i)+1})$ for $i \in I$. In particular, $(u, 0, \dots, 0) = (\sum b_\zeta f_\zeta, 0, \dots, 0) \notin \ker \phi$. That is $\psi_-(u)v_1 \neq 0$, which is a contradiction. So $\ker \psi_- = 0$ and U^- is isomorphic to U_1^- .

Similarly, we can show that $U^+ \cong U_1^+$. By the triangular decomposition we have

$$U(\mathfrak{g}) \cong U^- \otimes U^0 \otimes U^+ \cong U_1^- \otimes U_1^0 \otimes U_1^+ \cong U_1.$$

It can be checked easily that this isomorphism is an algebra isomorphism. \square

Theorem 5.17. *Let $\lambda \in P$. If V^q is the Weyl module $W^q(\lambda)$ over $U_q(\mathfrak{g})$ with highest weight λ , then its classical limit V^1 is isomorphic to*

- (1) $W(\lambda)$ or $\Pi W(\lambda)$ if $|\lambda| = 2k$ and $\overline{\Delta(\lambda)} = \bar{1}$,
- (2) $W(\lambda) \oplus \Pi W(\lambda)$ if $|\lambda| = 2k$ and $\overline{\Delta(\lambda)} \neq \bar{1}$ (in particular, if $\lambda_1 > \dots > \lambda_{2k} > 0$),
- (3) $W(\lambda) \cong \Pi W(\lambda)$ if $|\lambda| = 2k + 1$.

Proof. Let $\mathbf{v}(\lambda)$ be a finite dimensional irreducible \mathfrak{b}_+ -module of weight λ which generates $W(\lambda)$. Since $U^- \cong U_1^-$ and $E^{\mathbf{A}_1}(\lambda)/\mathbf{J}_1 E^{\mathbf{A}_1}(\lambda)$ is isomorphic to $\mathbf{v}(\lambda)$ or $\mathbf{v}(\lambda) \oplus \Pi \mathbf{v}(\lambda)$ as a $\text{Cliff}(\lambda)$ -module, it suffices to show that V^1 is a free U_1^- -module whose rank is $\dim_{\mathbb{C}} \mathbf{v}(\lambda)$ or $2 \dim_{\mathbb{C}} \mathbf{v}(\lambda)$.

By Proposition 4.2 we know that $W^q(\lambda)$ is a free U_q^- -module generated by $E^q(\lambda)$. Since $V_{\mathbf{A}_1}$ is a subspace of V^q , taking Proposition 5.7 into account, $V_{\mathbf{A}_1}$ is a free $U_{\mathbf{A}_1}^-$ -module generated by $E^{\mathbf{A}_1}(\lambda)$. Taking the classical limit, we see that $V^1 = U_1^- \cdot \left(E^{\mathbf{A}_1}(\lambda)/\mathbf{J}_1 E^{\mathbf{A}_1}(\lambda) \right)$ and

$$\dim_{\mathbb{C}} E^{\mathbf{A}_1}(\lambda)/\mathbf{J}_1 E^{\mathbf{A}_1}(\lambda) = \dim_{\mathbb{C}(q)} E^q(\lambda) = \dim_{\mathbb{C}} \mathbf{v}(\lambda) \text{ or } 2 \dim_{\mathbb{C}} \mathbf{v}(\lambda).$$

By a similar argument as in [HK, Proposition 3.4.10], we can show that V^1 is a free U_1^- -module. When $|\lambda| = 2k$ and $\overline{\Delta(\lambda)} \neq \bar{1}$, $E^q(\lambda)$ is parity invariant. Hence we have

$$V^1 \cong \begin{cases} W(\lambda) \text{ or } \Pi W(\lambda) & \text{if } |\lambda| = 2k \text{ and } \overline{\Delta(\lambda)} = \bar{1}, \\ W(\lambda) \oplus \Pi W(\lambda) & \text{if } |\lambda| = 2k \text{ and } \overline{\Delta(\lambda)} \neq \bar{1}, \\ W(\lambda) \cong \Pi W(\lambda) & \text{if } |\lambda| = 2k + 1. \end{cases}$$

□

6. COMPLETE REDUCIBILITY OF THE CATEGORY $\mathcal{O}_q^{\geq 0}$

In this section, we prove the complete reducibility theorem for $U_q(\mathfrak{g})$ -modules in the category $\mathcal{O}_q^{\geq 0}$.

Definition 6.1. The category $\mathcal{O}_q^{\geq 0}$ consists of finite dimensional $U_q(\mathfrak{g})$ -modules M with a weight space decomposition $M = \bigoplus_{\mu \in P} M_{\mu}$ satisfying (i) $\text{wt}(M) \subset P_{\geq 0}$, (ii) $k_i|_{M_{\mu}} = 0$ for $\mu \in P_{\geq 0}$ and $i \in \{1, \dots, n\}$ such that $\langle k_i, \mu \rangle = 0$.

Remark. The complete reducibility theorem for $\mathcal{O}_q^{\geq 0}$, which we establish at the end of this section, implies that $\mathcal{O}_q^{\geq 0}$ is isomorphic to the category \mathcal{T}_q of tensor modules; i.e., submodules of a tensor power of the natural representation $\mathbb{C}(q)^{n|n}$. Indeed, using the description of \mathcal{T}_q provided by Olshanski and Sergeev we first check that every simple object of $\mathcal{O}_q^{\geq 0}$ is a tensor module. Then, by the complete reducibility result for \mathcal{T}_q , obtained again by Sergeev and Olshanski, we conclude that the two categories are isomorphic.

Proposition 6.2. (i) The category $\mathcal{O}_q^{\geq 0}$ is closed under taking tensor products.

(ii) For each $\lambda \in \Lambda^+ \cap P_{\geq 0}$, $V^q(\lambda)$ is an irreducible $U_q(\mathfrak{g})$ -module in the category $\mathcal{O}_q^{\geq 0}$. Conversely, every finite dimensional irreducible $U_q(\mathfrak{g})$ -module in the category $\mathcal{O}_q^{\geq 0}$ has the form $V^q(\lambda)$ for some $\lambda \in \Lambda^+ \cap P_{\geq 0}$.

Proof. For (i) we use the comultiplication formula for $k_{\bar{i}}$. For (ii) we use Theorem 5.14 and Proposition 1.6. \square

Let S be the *antipode* on $U_q(\mathfrak{g})$ defined in [O, Section 4]. We have $S(q^h) = q^{-h}$ for all $h \in P^\vee$. Because S is an anti-automorphism on $U_q(\mathfrak{g})$, one can define two $U_q(\mathfrak{g})$ -module structures on the dual vector space of a $U_q(\mathfrak{g})$ -module $V \in \mathcal{O}_q^{\geq 0}$ by

$$\begin{aligned} \langle x \cdot \phi, v \rangle &:= \langle \phi, S(x) \cdot v \rangle \text{ and} \\ \langle x \cdot \phi, v \rangle &:= \langle \phi, S^{-1}(x) \cdot v \rangle \end{aligned}$$

for each $x \in U_q(\mathfrak{g})$ and linear functional ϕ on V . We denote these modules by V^* and V' , respectively. As vector spaces both modules are just $\bigoplus_{\mu \in P} V_\mu^*$, where $V_\mu^* = \text{Hom}_{\mathbb{C}(q)}(V_\mu, \mathbb{C}(q))$. The following lemma is an immediate consequence of the definitions.

Lemma 6.3. Suppose that V is a $U_q(\mathfrak{g})$ -module in the category $\mathcal{O}_q^{\geq 0}$.

- (1) There exist canonical $U_q(\mathfrak{g})$ -module isomorphisms $(V^*)' \cong V \cong (V')^*$.
- (2) The space V_μ^* is a weight space of weight $-\mu$.

Since $q^h S(e_i) q^{-h} = q^{\alpha_i(h)} S(e_i)$, we have $S(e_i)V_\mu \subset V_{\mu+\alpha_i}$, which implies $e_i V_\mu^* \subset V_{\mu-\alpha_i}^*$. By Lemma 6.3, we get $e_i(V^*)_{-\mu} \subset (V^*)_{-\mu+\alpha_i}$. Similarly, we also have $e_{\bar{i}}(V^*)_{-\mu} \subset (V^*)_{-\mu+\alpha_{\bar{i}}}$, $f_i(V^*)_{-\mu} \subset (V^*)_{-\mu-\alpha_i}$, $f_{\bar{i}}(V^*)_{-\mu} \subset (V^*)_{-\mu-\alpha_{\bar{i}}}$ for all $i \in I$ and $k_{\bar{i}}(V^*)_{-\mu} \subset (V^*)_{-\mu}$ for all $i \in J$. A weight module M is called a *lowest weight module* with lowest weight $\lambda \in P$ if it is generated over $U_q(\mathfrak{g})$ by an irreducible finite dimensional $U_q^{\leq 0}$ -module. By a similar argument as in Proposition 4.1, one can show that $(V^q(\lambda)_\lambda)^*$ is an irreducible $U_q^{\leq 0}$ -module so that $V^q(\lambda)^*$ and $V^q(\lambda)'$ are lowest weight modules of lowest weight $-\lambda$.

Suppose that V is a $U_q(\mathfrak{g})$ -module in the category $\mathcal{O}_q^{\geq 0}$. Because V is finite dimensional, we may choose a *maximal* weight $\lambda \in \text{wt}(V)$ with the property that $\lambda + \alpha_i$ is not a weight of V for any $i \in I$. Then the weight space V_λ is a completely reducible U_q^0 -module (see the remark after Definition 1.5). Fix an irreducible summand \mathbf{v} of V_λ and set $L = U_q(\mathfrak{g})\mathbf{v}$. Then L is a highest weight $U_q(\mathfrak{g})$ -module with highest weight λ . By the assumption, $\lambda \in \Lambda^+ \cap P_{\geq 0}$ and from Corollary 5.15 we know $L \cong V^q(\lambda)$ up to Π .

Now consider $\bar{\mathbf{v}} = \text{Hom}_{\mathbb{C}(q)}(\mathbf{v}, \mathbb{C}(q))$. Then it is a U_q^0 -submodule of $(V^*)_\lambda$. Set

$$\bar{L} = U_q(\mathfrak{g})\bar{\mathbf{v}} \subset V^*.$$

It is easy to show that $\bar{\mathbf{v}}$ is an irreducible $U_q^{\leq 0}(\mathfrak{g})$ -module and \bar{L} is a lowest weight module with lowest weight $-\lambda$. Translating Corollary 5.15 to the case of lowest weight modules, we get the following lemma.

Lemma 6.4. *The $U_q(\mathfrak{g})$ -module \bar{L} is isomorphic to the irreducible lowest weight module $V^q(\lambda)^*$ with lowest weight $-\lambda$ and lowest weight space $\bar{\mathbf{v}}$.*

Now we can prove the completely reducibility theorem for $U_q(\mathfrak{g})$ -modules in the category $\mathcal{O}_q^{\geq 0}$.

Theorem 6.5. *Every $U_q(\mathfrak{g})$ -module V in the category $\mathcal{O}_q^{\geq 0}$ is completely reducible.*

Proof. Take a maximal weight λ and consider a submodule of V , say L , generated by an irreducible $U_q^{\geq 0}$ -submodule of V_λ . We want to show $V \cong L \oplus V/L$. Taking dual with respect to S^{-1} of the inclusion $\bar{L} \rightarrow V^*$, we obtain a $U_q(\mathfrak{g})$ -module homomorphism $V \cong (V^*)' \rightarrow (\bar{L})'$. Thus we have a map:

$$\psi : L \hookrightarrow V \rightarrow (\bar{L})'.$$

It is easy to check that ψ is a nontrivial homomorphism. Since both L and $(\bar{L})'$ are irreducible, ψ is an isomorphism by Schur's lemma and we see that the following short exact sequence splits:

$$0 \rightarrow L \rightarrow V \rightarrow V/L \rightarrow 0.$$

Since $V/L \in \mathcal{O}_q^{\geq 0}$, using induction on the dimension of V , we complete the proof. \square

Corollary 6.6. *The tensor product of a finite number of $U_q(\mathfrak{g})$ -modules in the category $\mathcal{O}_q^{\geq 0}$ is completely reducible.*

Remark. The same argument can be applied to prove the completely reducibility of $\mathcal{O}^{\geq 0}$. In that case, the antipode is given by $S(x) = -x$ for all $x \in \mathfrak{g}$ (see [N, Section 4]) and Proposition 1.8 plays the same role as Proposition 5.15.

Acknowledgements. We would like to thank Ivan Penkov and Vera Serganova for the stimulating discussions. D.G. gratefully acknowledges the hospitality and excellent working conditions at the Seoul National University where most of this work was completed.

REFERENCES

- [ABS] M.F. Atiyah, R. Bott, A. Shapiro, Clifford modules, *Topology* **3** (1964), 3–38.
- [B] J. Brundan, Kazhdan-Lusztig polynomials and character formulae for the Lie superalgebra $\mathfrak{q}(n)$, *Adv. Math.* **182** (2004), 28–77.
- [BKM] G. Benkart, S.-J. Kang, D. Melville, Quantized enveloping algebras for Borcherds superalgebras, *Trans. Amer. Math. Soc.* **350** (1998), 3297–3319.

- [Dr] V. Drinfel'd, Quantum groups, *Proceedings of the International Congress of Mathematicians, Vol. 1* (Berkeley, Calif., 1986), 798–820, Amer. Math. Soc., Providence, RI, 1987.
- [G] M. Gorelik, Shapovalov determinants of Q -type Lie superalgebras, *Int. Math. Res. Pap.*, Article ID **96895** (2006), 1–71.
- [Har] J. Harris, Algebraic Geometry, A first course. Corrected reprint of the 1992 original. *Graduate Texts in Mathematics* **133** Springer-Verlag, New York, 1995.
- [HK] J. Hong, S.-J. Kang, Introduction to Quantum Groups and Crystal Bases, *Graduate Studies in Mathematics* **42**, American Mathematical Society, 2002.
- [IR] K. Ireland, M. Rosen, A Classical Introduction to Modern Number Theory, 2nd ed., *Graduate Texts in Mathematics* **84**, Springer-Verlag, New York, 1990.
- [K] V. Kac, Lie superalgebras, *Adv. Math.* **26** (1977), 8–96.
- [Lam] T. Y. Lam, Introduction to Quadratic Forms over Fields, *Graduate Studies in Mathematics* **67**, American Mathematical Society, Providence, RI, 2005.
- [Lang] S. Lang, Algebra, Revised third edition, *Graduate Texts in Mathematics* **211**, Springer-Verlag, New York, 2002.
- [LS] D. Leites, V. Serganova, Defining relations for classical Lie superalgebras I. Superalgebras with Cartan matrix or Dynkin-type diagram, *Proc. Topological and Geometrical Methods in Field Theory (Eds. J. Mickelson, et al)*, World Sci., Singapore, 1992, 194–201.
- [N] M. Nazarov, Capelli identities for Lie superalgebras, *Ann. Sci. Ecole Norm. Sup. (4)* **30**, **6** (1997), 847–872.
- [O] G. Olshanski, Quantized universal enveloping superalgebra of type Q and a super-extension of the Hecke algebra, *Lett. Math. Phys.* **24** (1992), 93–102.
- [P] I. Penkov, Characters of typical irreducible finite-dimensional $\mathfrak{q}(n)$ -modules, *Funct. Anal. Appl.* **20** (1986), 30–37.
- [PS1] I. Penkov, V. Serganova, Generic irreducible representations of finite-dimensional Lie superalgebras, *International Journal of Mathematics* **5** (1994), 389–419.
- [PS2] I. Penkov, V. Serganova, Characters of irreducible G -modules and cohomology of G/P for the Lie supergroup $G = Q(N)$, *J. Math. Sci. (New York)* **84** (1997), 1382–1412.
- [PS3] I. Penkov, V. Serganova, Characters of finite-dimensional irreducible $\mathfrak{q}(n)$ -modules, *Lett. Math. Phys.* **40** (1997), 147–158.
- [RTF] N. Reshetikhin, L. Takhtadzhyan, L. Faddeev, Quantization of Lie groups and Lie algebras. (Russian) *Algebra i Analiz* **1** (1989), 178–206; translation in *Leningrad Math. J.* **1** (1990), 193–225
- [Se1] A. Sergeev, The centre of enveloping algebra for Lie superalgebra $Q(n, C)$, *Lett. Math. Phys.* **7** (1983), 177–179.
- [Se2] A. Sergeev, Tensor algebra of the identity representation as a module over the Lie superalgebras $Gl(n, m)$ and $Q(n)$ (Russian), *Mat. Sb. (N.S.)* **123(165)** (1984), 422–430.
- [Sh] G. Shimura, Arithmetic and Analytic Theories of Quadratic Forms and Clifford Groups, *Mathematical Surveys and Monographs* **109**, American Mathematical Society, Providence, RI, 2004.

DEPARTMENT OF MATHEMATICAL SCIENCES AND RESEARCH INSTITUTE OF MATHEMATICS, SEOUL NATIONAL UNIVERSITY, SAN 56-1 SILLIM-DONG, GWANAK-GU, SEOUL 151-747, KOREA

E-mail address: `jhjung@math.snu.ac.kr`

DEPARTMENT OF MATHEMATICAL SCIENCES AND RESEARCH INSTITUTE OF MATHEMATICS, SEOUL NATIONAL UNIVERSITY, SAN 56-1 SILLIM-DONG, GWANAK-GU, SEOUL 151-747, KOREA

E-mail address: `sjkang@math.snu.ac.kr`

DEPARTMENT OF MATHEMATICAL SCIENCES AND RESEARCH INSTITUTE OF MATHEMATICS, SEOUL NATIONAL UNIVERSITY, SAN 56-1 SILLIM-DONG, GWANAK-GU, SEOUL 151-747, KOREA

E-mail address: `mkim@math.snu.ac.kr`