

# THE EQUIVARIANT EULER CHARACTERISTIC OF MODULI SPACES OF CURVES.

EUGENE GORSKY

ABSTRACT. We derive a formula for the  $S_n$ -equivariant Euler characteristic of the moduli space  $\mathcal{M}_{g,n}$  of genus  $g$  curves with  $n$  marked points.

## 1. INTRODUCTION

Consider the moduli space  $\mathcal{M}_{g,n}$  of algebraic curves of genus  $g$  with  $n$  marked points. The symmetric group  $S_n$  acts naturally on this space. Let  $V_\lambda$  denote the irreducible representation of  $S_n$  corresponding to a Young diagram  $\lambda$ , then one can decompose the cohomology of  $\mathcal{M}_{g,n}$  into isotypic components:

$$H^i(\mathcal{M}_{g,n}) = \bigoplus_{\lambda} a_{i,\lambda} V_\lambda.$$

The  $S_n$ -equivariant Euler characteristic of  $\mathcal{M}_{g,n}$  is defined by the formula

$$\chi^{S_n}(\mathcal{M}_{g,n}) = \sum_{i,\lambda} (-1)^i a_{i,\lambda} s_\lambda,$$

where  $s_\lambda$  denotes the Schur polynomial labeled by the diagram  $\lambda$ . We calculate these equivariant Euler characteristics for all  $g \geq 2$  and  $n$ .

**Theorem 1.1.** *The generating function for the  $S_n$ -equivariant Euler characteristics of  $\mathcal{M}_{g,n}$  has the form*

$$\sum_{n=0}^{\infty} t^n \chi^{S_n}(\mathcal{M}_{g,n}) = \sum_{\underline{k}} c_{k_1, \dots, k_r} \prod_{j=1}^r (1 + p_j t^j)^{k_j},$$

where  $p_j$  are power sums and the coefficients  $c_{k_1, \dots, k_r}$  are defined by the equation (6).

Consider the moduli space  $\mathcal{M}_g(k_1, \dots, k_r)$  of pairs  $(C, \tau)$  where  $C$  is a genus  $g$  curve and  $\tau$  is an automorphism of  $C$  such that for all  $i$  the Euler characteristic of the set of points in  $C$  having the orbit of length  $i$  under the action of  $\tau$  equals  $ik_i$ . The coefficient  $c_{k_1, \dots, k_r}$  can be also defined as the orbifold Euler characteristic of  $\mathcal{M}_g(k_1, \dots, k_r)$ .

This moduli space can be defined for any tuple of integers  $(k_1, \dots, k_r)$  of arbitrary size  $r$ , but we prove that (for a fixed genus  $g$ ) it is non-empty only for a finite number of such tuples. In particular,  $r$  cannot exceed  $4g + 2$ .

**Corollary 1.2.** The generating function  $\sum_{n=0}^{\infty} t^n \chi^{S_n}(\mathcal{M}_{g,n})$  is a rational function in  $t$ . Furthermore, for any  $n$ ,

$$\chi^{S_n}(\mathcal{M}_{g,n}) \in \mathbb{Z}[p_1, \dots, p_{4g+2}].$$

Theorem 1.1 can be compared with the computations of [4], [5], [8] and [10] in genus 2 and with the computations of [1], [2], [9], [17] and [18] in genus 3. A similar generating function for the moduli spaces of hyperelliptic curves was previously obtained in [11]. The non-equivariant Euler characteristics of moduli spaces of curves were computed by Bini and Harer in [3].

The paper is organized as follows. In Section 2 we consider a complex quasi-projective variety  $X$  with an action of a finite group  $G$ . Theorem 2.5 provides a formula for the  $S_n$ -equivariant

Euler characteristic of quotients  $F(X, n)/G$ , where  $F(X, n)$  is a configuration space of  $n$  labeled distinct points on  $X$ . This theorem was previously proved in [10] using the results of Getzler [6, 7] concerning Adams operations over the equivariant motivic rings (see also [12]). The alternative proof presented here uses only the basic properties of Euler characteristic and seems to be more geometric. It also makes the proof of the main result self-contained.

In Section 3 we apply this theorem to the universal family over  $\mathcal{M}_g$ , the moduli space of genus  $g$  curves. This allows us to prove in Theorem 3.3 that the coefficient  $c_{k_1, \dots, k_r}$  is equal to the orbifold Euler characteristic of  $\mathcal{M}_g(k_1, \dots, k_r)$ . These Euler characteristics are then computed in Theorem 3.8 using the results of Harer and Zagier.

#### ACKNOWLEDGEMENTS

The author is grateful to J. Bergström, S. Gusein-Zade, M. Kazaryan and S. Lando for useful discussions. This work was partially supported by the grants RFBR-007-00593, RFBR-08-01-00110-a, NSh-709.2008.1 and the Möbius Contest fellowship for young scientists.

#### 2. EQUIVARIANT EULER CHARACTERISTICS

Let  $X$  be a complex quasi-projective variety with an action of a finite group  $G$ . Let us denote by  $F(X, n)$  the configuration space of ordered  $n$ -tuples of distinct points on  $X$ . For each  $n$ , the action of the group  $G$  on  $X$  can be naturally extended to the action of  $G$  on  $F(X, n)$ , commuting with the natural action of  $S_n$ .

In the computations below we will use the additivity and multiplicativity of the Euler characteristic, as well as the Fubini formula for the integration with respect to the Euler characteristic ([15, 19], see also [16]).

**Lemma 2.1.** *The following equation holds:  $\sum_{n=0}^{\infty} \frac{t^n}{n!} \chi(F(X, n)) = (1+t)^{\chi(X)}$ .*

*Proof.* The map  $\pi_n : F(X, n) \rightarrow F(X, n-1)$ , which forgets the last point in the  $n$ -tuple, has fibers isomorphic to  $X$  without  $n-1$  points. Therefore  $\chi(F(X, n)) = (\chi(X) - n + 1) \cdot \chi(F(X, n-1))$ , and  $\chi(F(X, n)) = \chi(X) \cdot (\chi(X) - 1) \cdot \dots \cdot (\chi(X) - n + 1)$ .  $\square$

Let  $p_k$  denote the  $k$ th power sum and let  $V_\lambda$  denote the irreducible representation of  $S_n$  labelled by the Young diagram  $\lambda$ . We define the  $S_n$ -equivariant Euler characteristic of  $F(X, n)/G$  by the equation

$$\chi^{S_n}(F(X, n)/G) = \sum_{i, \lambda} (-1)^i a_{i, \lambda} s_\lambda,$$

where  $H^i(F(X, n)/G) = \bigoplus_{\lambda} a_{i, \lambda} V_\lambda$  and  $s_\lambda$  is the Schur polynomial.

**Lemma 2.2.** *The following equation holds:*

$$\chi^{S_n}(F(X, n)/G) = \frac{1}{n!} \sum_{\sigma \in S_n} p_1^{k_1(\sigma)} \cdot \dots \cdot p_n^{k_n(\sigma)} \cdot \chi([F(X, n)/G]^\sigma),$$

where  $k_i(\sigma)$  is the number of cycles of length  $i$  in a permutation  $\sigma$ .

*Proof.* It is well known that for every  $i$

$$\sum_{\lambda} a_{i, \lambda} s_\lambda = \frac{1}{n!} \sum_{\sigma \in S_n} p_1^{k_1(\sigma)} \cdot \dots \cdot p_n^{k_n(\sigma)} \cdot \text{Tr}(\sigma)|_{H^i(F(X, n)/G)},$$

hence

$$\chi^{S_n}(F(X, n)/G) = \frac{1}{n!} \sum_i (-1)^i \sum_{\sigma \in S_n} p_1^{k_1(\sigma)} \cdot \dots \cdot p_n^{k_n(\sigma)} \cdot \text{Tr}(\sigma)|_{H^i(F(X, n)/G)}$$

Now the statement follows from the Lefschetz fixed point theorem.  $\square$

**Lemma 2.3.** *Let  $\sigma \in S_n$ . Then*

$$\chi([F(X, n)/G]^\sigma) = \frac{1}{|G|} \sum_{g \in G} \chi\left(F(X, n)^{g^{-1}\sigma}\right).$$

*Proof.* For a point  $\mathbf{y} \in F(X, n)$  whose projection on  $F(X, n)/G$  is  $\sigma$ -invariant there exists an element  $g \in G$  such that  $\sigma\mathbf{y} = g\mathbf{y}$ . Consider the set of pairs

$$S = \{(g, \mathbf{y}) | g \in G, \mathbf{y} \in F(X, n) | \sigma\mathbf{y} = g\mathbf{y}\}$$

and its two-step projection  $S \rightarrow F(X, n) \rightarrow F(X, n)/G$ . The fiber of the first projection over a point  $\mathbf{y}$  is isomorphic to  $G$ -stabiliser of  $\mathbf{y}$  or empty, the fiber of the second projection containing  $\mathbf{y}$  is exactly the orbit of  $\mathbf{y}$ . Therefore the cardinality of every fiber of the composition is equal to  $|G|$ .  $\square$

**Definition 2.4.** For any  $g \in G$  we denote by  $X_k(g)$  the subset of  $X$  consisting of points with  $g$ -orbits of length  $k$ . For example,  $X_1(g)$  is a set of  $g$ -fixed points. Let  $\tilde{X}_k(g) = X_k(g)/(g)$ , where  $(g)$  is a cyclic subgroup in  $G$  generated by  $g$ .

The following theorem was deduced in [10] from the results of Getzler [6, 7], here we would like to present a more geometric and straightforward proof.

**Theorem 2.5.** *The generating function for the  $S_n$ -equivariant Euler characteristics of the quotients  $F(X, n)/G$  is given by the following equation:*

$$(1) \quad \sum_{n=0}^{\infty} t^n \chi^{S_n}(F(X, n)/G) = \frac{1}{|G|} \sum_{g \in G} \prod_{k=1}^{\infty} (1 + p_k t^k)^{\frac{\chi(X_k(g))}{k}}.$$

*Proof.* Since all points in  $X_k(g)$  have  $g$ -orbit of length  $k$ , we have  $\chi(\tilde{X}_k(g)) = \chi(X_k(g))/k$ . From Lemma 2.1 one gets:

$$(1 + p_j t^j)^{\chi(\tilde{X}_j(g))} = \sum_{k_j=0}^{\infty} \frac{p_j^{k_j} t^{j k_j}}{(k_j)!} \chi\left(F\left(\tilde{X}_j(g), k_j\right)\right),$$

Therefore the coefficient at  $t^n$  in the right hand side of (1) equals to:

$$\frac{1}{|G|} \sum_{g \in G} \sum_{\sum_j k_j = n} \prod_j \frac{p_j^{k_j}}{k_j!} \chi\left(F\left(\tilde{X}_j(g), k_j\right)\right).$$

On the other hand, by Lemma 2.2 and Lemma 2.3, the left hand side of (1) can be rewritten as following:

$$\frac{1}{|G|} \sum_{g \in G} \frac{1}{n!} \sum_{\sigma \in S_n} p_1^{k_1(\sigma)} \cdots p_n^{k_n(\sigma)} \cdot \chi([F(X, n)]^{g^{-1}\sigma}).$$

If for a tuple  $\mathbf{y} \in F(X, n)$  we have  $\sigma(\mathbf{y}) = g(\mathbf{y})$ , the action of  $(g)$  at this tuple has  $k_j(\sigma)$  cycles of length  $j$ . Every cycle of length  $j$  corresponds to a point in  $\tilde{X}_j(g)$ , hence for every  $g$  we can define a map

$$\alpha_g : \sqcup_{\sigma \in S_n} [F(X, n)]^{g^{-1}\sigma} \rightarrow \prod_j F(\tilde{X}_j(g), k_j)/S_{k_j}.$$

Given a  $g$ -invariant  $n$ -tuple of distinct points in  $X$ , there are  $n!$  ways to label them and make an ordered tuple  $\mathbf{y}$ . Every such ordering defines a unique permutation  $\sigma$  such that  $\sigma(\mathbf{y}) = g(\mathbf{y})$ , therefore all fibers of  $\alpha_g$  have cardinality  $n!$  and

$$\frac{1}{n!} \sum_{\sigma \in S_n} \chi([F(X, n)]^{g^{-1}\sigma}) = \prod_j \chi\left(F(\tilde{X}_j(g), k_j)/S_{k_j}\right) = \prod_j \frac{\chi\left(F(\tilde{X}_j(g), k_j)\right)}{k_j!}.$$

□

### 3. MODULI SPACES OF CURVES

Let us apply Theorem 2.5 to the study of moduli spaces of curves. Let  $\mathcal{M}_g$  denote the moduli space of genus  $g$  algebraic curves and let  $\mathcal{M}_{g,n}$  denote the moduli space of genus  $g$  algebraic curves with  $n$  parked points (we will always assume  $g \geq 2$ ). Let  $\mathcal{M}_g(k_1, \dots, k_r)$  be the moduli space of pairs  $(C, \tau)$  where  $C$  is a genus  $g$  curve and  $\tau$  is an automorphism of  $C$  such that  $\chi(C_i(\tau)) = ik_i$  for all  $i$ . Since  $g \geq 2$ , every automorphism of  $C$  has finite order, hence one can choose  $r$  such that  $k_r \neq 0$  and  $k_i = 0$  for  $i > r$ .

There is a natural forgetful map  $\pi_{g,\underline{k}} : \mathcal{M}_g(k_1, \dots, k_r) \rightarrow \mathcal{M}_g$  sending  $(C, \tau)$  to  $C$ . For a curve  $C$  we define  $\text{Aut}_{\underline{k}}(C) = \pi_{g,\underline{k}}^{-1}(C) \subset \text{Aut}(C)$ .

**Proposition 3.1.** Suppose that  $\mathcal{M}_g(k_1, \dots, k_r)$  is not empty. Then  $k_r < 0, k_i = 0$  for  $i \nmid r$  and  $k_i \geq 0$  for  $i \mid r, i < r$ . Moreover, we have the following bounds on  $r$  and  $k_i$ :

$$r \leq 4g + 2, |k_r| \leq 2g, \sum_{i=1}^{r-1} k_i \leq 2g + 2.$$

*Proof.* Let  $\tau$  be an automorphism of a genus  $g$  curve  $C$  such that  $\chi(C_i(\tau)) = ik_i$  for all  $i$ . Note that  $C_i(\tau)$  are finite sets for  $i < r$  and

$$(2) \quad \chi(C) = 2 - 2g = \sum_{i=1}^{r-1} ik_i - r|k_r|$$

The quotient  $C_1 = C/\tau$  is a smooth curve of some genus  $h$ , and the Riemann-Hurwitz formula yields its Euler characteristic:

$$(3) \quad \chi(C_1) = 2 - 2h = \sum_{i=1}^{r-1} k_i - |k_r|.$$

The projection of  $C$  to  $C_1$  is a ramified covering of order  $r$  with  $s = \sum_{i=1}^{r-1} k_j$  ramification points. The automorphism  $\tau$  has order  $r$ , so  $i \mid r$ , if  $k_i \neq 0$ . By a theorem of Wiman ([20], see also [14]), the maximal order for an automorphism of a genus  $g$  curve equals  $4g + 2$ , hence  $r \leq 4g + 2$ .

Since proper divisors of  $r$  cannot exceed  $r/2$ , equation (3) implies:

$$\sum_{i=1}^{r-1} ik_i \leq \frac{r}{2} \sum_{i=1}^{r-1} k_i = \frac{r}{2}(2 - 2h + |k_r|),$$

hence by (2):

$$(4) \quad 2g - 2 = r|k_r| - \sum_{i=1}^{r-1} ik_i \geq \frac{r}{2}(2h + |k_r| - 2).$$

Therefore  $|k_r| - 2 \leq 2g - 2$  and  $|k_r| \leq 2g$ . Finally,  $\sum_{i=1}^{r-1} k_i = |k_r| + 2 - 2h \leq 2g + 2$ . □

*Remark 3.2.* The bounds on  $r$  and on  $k_i$  are sharp. Indeed, consider a hyperelliptic curve  $P$  covering  $\mathbb{CP}^1$  with ramifications at the vertices of a regular  $(2g + 1)$ -gon and at its center. The covering can be chosen such that the automorphism of  $P$  induced by the rotation of this polygon acts nontrivially in the fibers and hence has order  $r = 2(2g + 1) = 4g + 2$ .

On the other hand, consider a hyperelliptic curve  $C$  with involution  $\tau$ . We have

$$\chi(C_1(\tau)) = 2g + 2, \chi(C_2(\tau)) = 2 - 2g - (2g + 2) = -4g,$$

hence a pair  $(C, \tau)$  belongs to the moduli space  $\mathcal{M}_g(2g + 2, -2g)$ .

**Theorem 3.3.** *The following equation holds:*

$$(5) \quad \sum_{n=0}^{\infty} t^n \chi^{S_n}(\mathcal{M}_{g,n}) = \sum_{\underline{k}} \chi^{orb}(\mathcal{M}_g(k_1, \dots, k_r)) \cdot \prod_{j=1}^r (1 + p_j t^j)^{k_j}.$$

*Proof.* Consider the forgetful map  $\pi_{g,n} : \mathcal{M}_{g,n} \rightarrow \mathcal{M}_g$ . Its fiber over a point representing a curve  $C$  is isomorphic to  $F(C, n)/\text{Aut}(C)$ , hence one can apply Theorem 2.5 to compute its equivariant Euler characteristic:

$$\begin{aligned} \sum_{n=0}^{\infty} t^n \chi^{S_n}(\pi_{g,n}^{-1}(C)) &= \sum_{n=0}^{\infty} t^n \chi^{S_n}(F(C, n)/\text{Aut}(C)) = \\ &= \frac{1}{|\text{Aut}(C)|} \sum_{\tau \in \text{Aut}(C)} \prod_i (1 + p_i t^i)^{\frac{\chi(C_i(\tau))}{i}} = \sum_{\underline{k}} \frac{1}{|\text{Aut}(C)|} \sum_{\tau \in \text{Aut}_{\underline{k}}(C)} \prod_i (1 + p_i t^i)^{k_i}. \end{aligned}$$

Therefore:

$$\begin{aligned} \sum_{n=0}^{\infty} t^n \chi^{S_n}(\mathcal{M}_{g,n}) &= \int_{\mathcal{M}_g} \sum_{n=0}^{\infty} t^n \chi^{S_n}(\pi_{g,n}^{-1}(C)) d\chi = \\ &= \sum_{\underline{k}} \prod_i (1 + p_i t^i)^{k_i} \int_{\mathcal{M}_g} \frac{|\text{Aut}_{\underline{k}}(C)|}{|\text{Aut}(C)|} d\chi. \end{aligned}$$

On the other hand,

$$\chi^{orb}(\mathcal{M}_g(k_1, \dots, k_r)) = \int_{\mathcal{M}_g} \frac{|\pi_{g,\underline{k}}^{-1}(C)|}{|\text{Aut}(C)|} d\chi = \int_{\mathcal{M}_g} \frac{|\text{Aut}_{\underline{k}}(C)|}{|\text{Aut}(C)|} d\chi \quad \square$$

Using the Proposition 3.1, we conclude that the sum in the right hand side of (5) is finite.

**Corollary 3.4.** The generating function  $\sum_{n=0}^{\infty} t^n \chi^{S_n}(\mathcal{M}_{g,n})$  is a rational function in  $t$ . Furthermore, for any  $n$ ,

$$\chi^{S_n}(\mathcal{M}_{g,n}) \in \mathbb{Z}[p_1, \dots, p_{4g+2}].$$

The orbifold Euler characteristic of  $\mathcal{M}_g(k_1, \dots, k_r)$  can be computed using the combinatorial results of Harer and Zagier [13]. We will denote the greatest common divisor of integers  $a$  and  $b$  by  $(a, b)$ . Let  $\varphi(n)$  and  $\mu(n)$  denote the Euler function and the Möbius function respectively. Define

$$c(k, l, d) := \mu\left(\frac{d}{(d, l)}\right) \frac{\varphi(k/l)}{\varphi(d/(d, l))},$$

**Definition 3.5.** Let  $\lambda = (\lambda_1, \dots, \lambda_s)$  be a partition. We define a number

$$N(r; \lambda) = |\{(x_1, \dots, x_s) \in (\mathbb{Z}/r\mathbb{Z})^s : x_1 + \dots + x_s \equiv 0 \pmod{r}, (x_i, k) = \lambda_i\}|.$$

**Lemma 3.6.** ([13]) *The following equation holds:*

$$N(r; \lambda) = \frac{1}{r} \sum_{d|r} \varphi(d) \prod_{i=1}^s c(k, \lambda_i, d).$$

**Theorem 3.7.** ([13]) *The orbifold Euler characteristic of the moduli space  $\mathcal{M}_{h,s}$  of genus  $h$  curves with  $s$  marked points is given by the formula:*

$$\chi^{orb}(\mathcal{M}_{h,s}) = (-1)^s \frac{(2h-1)B_{2h}}{(2h)!} (2h+s-3)!$$

where  $B_k$  denote Bernoulli numbers.

**Theorem 3.8.** *The generating function for the  $S_n$ -equivariant Euler characteristics of  $\mathcal{M}_{g,n}$  has the form*

$$\sum_{n=0}^{\infty} t^n \chi^{S_n}(\mathcal{M}_{g,n}) = \sum_{\underline{k}} c_{k_1, \dots, k_r} \prod_{j=1}^r (1 + p_j t^j)^{k_j},$$

where  $p_j$  are power sums and the coefficients  $c_{k_1, \dots, k_r}$  are defined by the equation:

$$(6) \quad c_{k_1, \dots, k_r} = \chi^{orb}(\mathcal{M}_{h,s}) r^{2h} \prod_{p|\gamma} (1 - p^{-2h}) \cdot \frac{N(r; \lambda)}{r \prod_{i=1}^{r-1} k_i!}.$$

Here  $h = \frac{1}{2}(1 - \sum_{j=1}^r k_j)$ ,  $s = \sum_{j=1}^{r-1} k_j$ ,  $\gamma = \text{GCD}(i : k_i > 0)$ ,  $\lambda = (1^{k_1} 2^{k_2} \dots (r-1)^{k_{r-1}})$

*Proof.* By Theorem 3.3 one has  $c_{k_1, \dots, k_r} = \chi^{orb}(\mathcal{M}_g(k_1, \dots, k_r))$ . Consider the moduli space  $\mathcal{M}_g(k_1, \dots, k_r)$  of pairs  $(C, \tau)$ . As in Proposition 3.1, to such a pair one can associate a genus  $h$  curve  $C_1 = C/\tau$ . The projection from  $C$  to  $C_1$  is ramified in  $s$  points subdivided into groups of size  $k_1, \dots, k_{r-1}$ . The orbifold Euler characteristic of the moduli space of genus  $h$  curves with such markings equals  $\chi^{orb}(\mathcal{M}_{h,s}) / \prod_{i=1}^{r-1} k_i!$ .

The number of pairs  $(C, \tau)$  associated to a curve  $C_1$  with fixed marked points was computed in [13, pages 478–479] and equals

$$\frac{1}{r} r^{2h} \prod_{p|\gamma} (1 - p^{-2h}) \cdot N(r; \lambda).$$

This completes the proof. □

The non-equivariant Euler characteristic of  $\mathcal{M}_{g,n}$  has been computed in [3, Theorem 4.3]. It can be compared with Theorem 3.8 since

$$\chi(\mathcal{M}_{g,n}) = n! \cdot \chi^{S_n}(\mathcal{M}_{g,n}) [p_1 = 1, p_k = 0 \text{ for } k > 1].$$

**Example 3.9.** The generating function for the  $S_n$ -equivariant Euler characteristics of the moduli spaces of genus 2 curves with marked points has a form [10]:

$$\begin{aligned} \sum_{n=0}^{\infty} t^n \chi^{S_n}(\mathcal{M}_{2,n}) &= -\frac{1}{240} (1 + p_1 t)^{-2} - \frac{1}{240} (1 + p_1 t)^6 (1 + p_2 t^2)^{-4} + \\ &+ \frac{2}{5} (1 + p_1 t)^3 (1 + p_5 t^5)^{-1} + \frac{2}{5} (1 + p_1 t) (1 + p_2 t^2) (1 + p_5 t^5) (1 + p_{10} t^{10})^{-1} + \\ &+ \frac{1}{6} (1 + p_1 t)^2 (1 + p_2 t^2) (1 + p_6 t^6)^{-1} - \frac{1}{12} (1 + p_1 t)^4 (1 + p_3 t^3)^{-2} - \\ &- \frac{1}{12} (1 + p_2 t^2)^2 (1 + p_3 t^3)^2 (1 + p_6 t^6)^{-2} + \frac{1}{12} (1 + p_1 t)^2 (1 + p_2 t^2)^{-2} + \\ &+ \frac{1}{4} (1 + p_1 t)^2 (1 + p_4 t^4) (1 + p_8 t^8)^{-1} - \frac{1}{8} (1 + p_1 t)^2 (1 + p_2 t^2)^2 (1 + p_4 t^4)^{-2}. \end{aligned}$$

These coefficients can be matched with the ones defined in Theorem 3.8.

## REFERENCES

- [1] J. Bergström. Cohomologies of moduli spaces of curves of genus three via point counts. *J. Reine Angew. Math.* **622** (2008), 155–187.
- [2] J. Bergström, O. Tommasi. The rational cohomology of  $\overline{\mathcal{M}}_4$ . *Math. Ann.* **338** (2007), no. 1, 207–239.
- [3] G. Bini, J. Harer. Euler Characteristics of Moduli Spaces of Curves. *J. Eur. Math. Soc. (JEMS)* **13** (2011), no. 2, 487–512.
- [4] G. Bini, G. Gaiffi, M. Polito. A formula for the Euler characteristic of  $\overline{\mathcal{M}}_{2,n}$ . *Math. Z.* **236** (2001) 491–523.
- [5] C. Faber, G. van der Geer. Sur la cohomologie des systèmes locaux sur les espaces de modules des courbes de genre 2 et des surfaces abéliennes. I, II. *C. R. Math. Acad. Sci. Paris* **338** (2004), no. 5, 381–384; no. 6, 467–470.

- [6] E. Getzler. Mixed Hodge structures of configuration spaces. arXiv:math.AG/9510018
- [7] E. Getzler. Resolving mixed Hodge modules on configuration spaces. *Duke Math. J.* **96** (1999), no. 1, 175–203.
- [8] E. Getzler. Euler characteristics of local systems on  $\mathcal{M}_2$ . *Compositio Math.* **132** (2002), 121–135.
- [9] E. Getzler, E. Looijenga. The Hodge polynomial of  $\overline{\mathcal{M}}_{3,1}$ . arXiv:math.AG/9910174
- [10] E. Gorsky. On the  $S_n$ -equivariant Euler characteristic of  $\mathcal{M}_{2,n}$ . arXiv:0707.2662.
- [11] E. Gorsky. On the  $S_n$ -equivariant Euler characteristic of moduli spaces of hyperelliptic curves. *Math. Res. Lett.* **16** (2009), no. 4, 591–603.
- [12] E. Gorsky. Adams operations and power structures. *Mosc. Math. J.* **9** (2009), no. 2, 305–323.
- [13] J. Harer, D. Zagier. The Euler characteristic of the moduli space of curves. *Invent. Math.* **85** (1986), 457–485.
- [14] W. J. Harvey. Cyclic groups of automorphisms of a compact Riemann surface. *Quart. J. Math. Oxford Ser. (2)* **17** (1966), 86–97.
- [15] A. Khovanskii, A. Pukhlikov. Integral transforms based on Euler characteristic and their applications. *Integral Transform. Spec. Funct.* **1** (1993), no. 1, 19–26.
- [16] R. MacPherson. Chern classes for singular algebraic varieties. *Ann. of Math. (2)* **100** (1974), 423–432.
- [17] O. Tommasi. Rational cohomology of the moduli space of genus 4 curves. *Compos. Math.* **141** (2005), no. 2, 359–384.
- [18] O. Tommasi. Rational cohomology of  $\mathcal{M}_{3,2}$ . *Compos. Math.* **143** (2007), no. 4, 986–1002.
- [19] O. Viro. Some integral calculus based on Euler characteristic. *Topology and geometry – Rohlin Seminar*, 127–138, *Lecture Notes in Math.*, **1346**, Springer, Berlin, 1988.
- [20] A. Wiman. Ueber die hyperelliptischen Curven und diejenigen vom Geschlechte  $p = 3$  welche eindeutigen Transformationen in sich zulassen. *Bihang Till. Kongl. Svenska Vetenskaps-Akademiens Handlingar* **21** (1895-6) 1–23.

MATHEMATICS DEPARTMENT, STONY BROOK UNIVERSITY, STONY BROOK NY, 11794-3651, USA

*E-mail address:* egorsky@math.sunysb.edu