# Complex bimatrix variate generalised beta distributions

José A. Díaz-García \*

Department of Statistics and Computation 25350 Buenavista, Saltillo, Coahuila, Mexico E-mail: jadiaz@uaaan.mx

> Ramón Gutiérrez Jáimez Department of Statistics and O.R. University of Granada Granada 18071, Spain

> > E-mail: rgjaimez@ugr.es

#### Abstract

In this paper, the study of bivariate generalised beta type I and II distributions is extended to the complex matrix variate case, for which the corresponding density functions are found. In addition, for complex bimatrix variate beta type I distributions, several basic properties, including the joint eigenvalue density and the maximum eigenvalue distribution, are studied.

## 1 Introduction

Complex matrix variate distributions play an important role in various fields of research. Applications of complex random matrices can be found in multiple time series analysis, nuclear physics, complex multivariate linear model, shape theory, the evaluation of the capacity of multiple-input multiple-output (MIMO) wireless communication systems, see [Mehta \(1991\)](#page-12-0), [Khatri \(1965\)](#page-11-0), [Micheas](#page-12-1) et al. [\(2006\)](#page-12-1) and [Ratnarajah](#page-12-2) et al. [\(2005a\)](#page-12-2) among many others.

Several studies on the distribution of complex random matrices have been made, see [Mathai and Provost \(2005](#page-12-3)). The complex matrix variate Gaussian distribution was introduced by [Wooding \(1956\)](#page-12-4), and further developed by [Turin \(1960\)](#page-12-5) and [Goodman \(1963](#page-11-1)). The complex Wishart distribution was studied by [Goodman \(1963\)](#page-11-1) and [James \(1964\)](#page-11-2), among many others. [James \(1964\)](#page-11-2) and [Khatri \(1965](#page-11-0)) derived both complex central and noncentral matrix variate beta distributions.

When the goal is to generalise the distribution of a random variable to the multivariate case, two options are normally addressed, by which it is extended to either the vectorial or the matrix case, e.g. normal, t or bessel distributions, among many others. However,

<sup>∗</sup>Corresponding author

Key words. Complex random matrices, complex beta distribution, complex bimatrix variate generalised beta.

<sup>2000</sup> Mathematical Subject Classification. 15A52, 60E05, 62E15

some of these generalisations have traditionally been made directly to the matrix case, when such a matrix is symmetric, as in the case of the chi-square and beta distributions, for which the corresponding multivariate distributions are the Wishart and matrix variate beta distributions, respectively. Nevertheless, these latter generalisations are inappropriate in some cases, because, in some applications, the researcher is interested in a vector variate, not in a symmetric matrix, see [Libby and Novick \(1982\)](#page-12-6). In other words, the researcher is interested in a vector, say,  $\mathbf{X} = (x_1, \ldots, x_m)'$ , such that  $x_i$  has a marginal beta type I or II distribution for all  $i = 1, \ldots, m$ . In this respect, [Libby and Novick \(1982\)](#page-12-6) and [Chen and Novick \(1984\)](#page-11-3) proposed a multivariate vector-version of the beta type I and II distributions. Let us consider the following bivariate version, see [Olkin and Liu \(2003\)](#page-12-7).

Let  $X_0, X_1, X_2$  be distributed as independent gamma random variates with parameters  $a = a_0, a_1, a_2$ , respectively (see Definition [2.1](#page-2-0) in Section 2); and define

<span id="page-1-0"></span>
$$
U_1 = \frac{X_1}{X_1 + X_0}, \qquad U_2 = \frac{X_2}{X_2 + X_0}.
$$
 (1)

Clearly,  $U_1$  and  $U_2$  each have a beta type I distribution,  $U_1 \sim \mathcal{B}I_1(a_1, a_0)$  and  $U_2 \sim$  $\mathcal{B}I_1(a_2, a_0)$ , over  $0 \leq u_1, u_2 \leq 1$  (see Subsection 2.1). However, they are correlated such that  $(U_1, U_2)'$  has a bivariate generalised beta type I distribution over  $0 \le u_1, u_2 \le 1$ . The kernel of the joint density function of  $U_1$  and  $U_2$  is

$$
\propto \frac{u_1^{a_1-1} u_2^{a_2-1} (1-u_1)^{a_2+a_0-1} (1-u_2)^{a_1+a_0-1}}{(1-u_1 u_2)^{a_1+a_2+a_0}}, \quad 0 \le u_1, u_2 \le 1.
$$

A similar result is obtained in the case of beta type II. Here it defines

$$
F_1 = \frac{X_1}{X_0}, \qquad F_2 = \frac{X_2}{X_0}.
$$

Once again it is evident that  $F_1$  and  $F_2$  each have a beta type II distribution,  $F_1 \sim$  $\mathcal{B}II_1(a_1, a_0)$  and  $F_2 \sim \mathcal{B}II_1(a_2, a_0)$ , over  $f_1, f_2 \geq 0$ . As in the beta type I case, they are correlated such that  $(F_1, F_2)'$  has a bivariate generalised beta type II distribution over  $f_1, f_2 \geq 0$ . The kernel of the joint density function of  $F_1$  and  $F_2$  is

$$
\propto \frac{f_1^{a_1-1} f_2^{a_2-1}}{(1+f_1+f_2)^{a_1+a_2+a_0}}, \quad f_1, f_2 \ge 0.
$$

Some applications to utility modelling and Bayesian analysis are presented in [Libby and Novick](#page-12-6) [\(1982](#page-12-6)) and [Chen and Novick \(1984](#page-11-3)), respectively.

These ideas can be extended to the matrix variate case. Thus, let us assume a parti-

tioned matrix  $\mathbb{U} = (\mathbf{U}_1 \mathbf{U}_2)' \in \mathfrak{C}^{2m \times m}$ , then under the complex matrix variate versions of the transformations [\(1\)](#page-1-0), we are interested in finding the joint density of  $U_1$  and  $U_2$ , from where it is easy to see that the marginal densities of  $U_1$  and  $U_2$  are complex matrix variate beta type I distributions. In the central and noncentral real cases, the matrix variate joint densities of  $U_1$  and  $U_2$  and of  $F_1$  and  $F_2$ , together with some of their properties, are stud-ied in Díaz-García and Gutiérrez-Jáimez (2009a[,b\)](#page-11-5). These distributions are termed central complex bimatrix variate generalised beta type I and II distributions, respectively.

In the present paper, the bivariate generalised beta type I and II distributions are extended the the complex matrix variate case, see Sections [3](#page-5-0) and [4.](#page-7-0) In Section [5,](#page-8-0) certain basic properties, the joint eigenvalue density and the density of their maximum eigenvalues are studied for the complex bimatrix variate generalised beta type I distribution.

## 2 Preliminary results

The join eigenvalues density can be calculated using complex hypergeometric functions and complex invariant polynomials with matrix arguments. In this section, these and other issues are addressed for the case of the complex multivariate distribution. Let us first establish some notation.

#### 2.1 Notation and matrix variate distributions

Let  $\mathbf{A} = (a_{rs}) = (a_{rs_1} + ia_{rs_2}) = \mathbf{A}_1 + i\mathbf{A}_2$  be an  $m \times n$  of complex numbers,  $\mathbf{A} \in \mathbb{C}^{m \times n}$ , where  $\mathbf{A}_1 = (a_{rs_1})$  and  $\mathbf{A}_2 = (a_{rs_2})$  are real matrices  $m \times n$ ,  $\mathbf{A}_1, \mathbf{A}_2 \in \mathbb{R}^{m \times n}$ , and  $i = \sqrt{-1}$ . Then, A' denotes the transpose of A,  $\overline{A}$  denotes the conjugate of A, and A<sup>H</sup> denotes the conjugate transpose of **A**. For  $n = m$ , let  $tr(A) = a_{11} + \cdots + a_{mm}$ ,  $etr(tr(A)) = exp(tr(A))$ , and then |A| denotes the determinant of **A**.  $\mathbf{A} = \mathbf{A}^H > \mathbf{0}$  is a Hermitian positive definite matrix, and  $A^{1/2}$  denotes the unique Hermitian positive definite square root matrix of  $A = A<sup>H</sup> > 0$ . dA denotes the differential matrix of A and  $(dA) = (dA<sub>1</sub>)(dA<sub>2</sub>)$  denotes the volume element (Lebesgue measure) commonly associated with A. For example if A is a Hermitian matrix, then  $A_1$  is a symmetric matrix  $(A = A')$  and  $A_2$  is a skew-symmetric matrix  $(\mathbf{A} = -\mathbf{A}')$ . In this case

$$
(d\mathbf{A}) = \bigwedge_{r \leq s} da_{rs_1} \bigwedge_{r < s} da_{rs_2}.
$$

The space of all matrices  $\mathbf{G}_1 \in \mathbb{C}^{n \times m}$  ( $m \leq n$ ) with orthonormal columns is termed the **Stiefel manifold**, denoted by  $CV_{m,n}$ . Thus

$$
\mathfrak{CV}_{m,n}=\{\mathbf{G}_1\in \mathfrak{C}^{n\times m}|\mathbf{G}_1^H\mathbf{G}_1=\mathbf{I}_m\}.
$$

From [James \(1964\)](#page-11-2)

$$
\text{Vol}(\mathfrak{C}\mathcal{V}_{m,n}) = \int_{\mathbf{G}_1 \in \mathfrak{C}\mathcal{V}_{m,n}} (\mathbf{G}_1^H d\mathbf{G}_1) = \frac{2^m \pi^{mn}}{\mathfrak{C} \Gamma_m[n]},
$$

where  $\mathfrak{C}\Gamma_m[a]$  denotes the complex multivariate gamma function and is defined as

$$
\mathfrak{C}\Gamma_m[a] = \int_{\mathbf{V}=\mathbf{V}^H>0} \text{etr}(-\mathbf{V})|\mathbf{V}|^{a-m}(d\mathbf{V}) = \pi^{m(m-1)/2}\prod_{j=1}^m \Gamma[a-j+1],
$$

where  $\text{Re}(a) > m - 1$ .

If  $m = n$ , it is have a special case of the Stiefel manifold, termed unitary manifold or unitary group and denoted as  $\mathcal{U}(m) \equiv \mathfrak{CV}_{m,m}$ .

<span id="page-2-0"></span>Definition 2.1 (The complex multivariate beta function). The complex multivariate beta function, denoted as  $\mathfrak{C}\mathfrak{R}_m[a, b]$ , is defined by

$$
\mathfrak{C}\beta_m[b,a] = \int_{0 < \mathbf{S} = \mathbf{S}^H < \mathbf{I}_m} |\mathbf{S}|^{a-m} |\mathbf{I}_m - \mathbf{S}|^{b-m}(d\mathbf{S})
$$
  
\n
$$
= \int_{\mathbf{R} = \mathbf{R}^H > 0} |\mathbf{R}|^{a-m} |\mathbf{I}_m + \mathbf{R}|^{-(a+b)}(d\mathbf{R}), \ \mathbf{R} = (\mathbf{I} - \mathbf{S})^{-1} - \mathbf{I}
$$
  
\n
$$
= \frac{\mathfrak{C} \Gamma_m[a] \mathfrak{C} \Gamma_m[b]}{\mathfrak{C} \Gamma_m[a+b]}.
$$

where  $\text{Re}(a) > m - 1$  and  $\text{Re}(b) > m - 1$ .

We now give definitions of the complex matrix variate gamma, beta type I and II distributions, see [James \(1964\)](#page-11-2), [Khatri \(1965\)](#page-11-0) and [Mathai \(1997\)](#page-12-8).

**Definition 2.2** (Complex matrix variate gamma distribution). It is said that  $A \in \mathbb{C}^{m \times m}$ , a random Hermitian positive definite, has a complex matrix variate gamma distribution with parameters a and a Hermitian positive definite matrix  $\mathbf{\Theta} \in \mathbb{C}^{m \times m}$ , if its density function is

<span id="page-3-2"></span>
$$
\frac{1}{\mathfrak{C}\Gamma_m[a]|\Theta|^a}|\mathbf{A}|^{a-m}\operatorname{etr}(-\Theta^{-1}\mathbf{A})(d\mathbf{A}),\quad \mathbf{A}=\mathbf{A}^H>0,\tag{2}
$$

where  $\text{Re}(a) > m - 1$ . Such a distribution is denoted as  $\mathbf{A} \sim \mathfrak{C} \mathcal{G}_m(a, \mathbf{\Theta})$ .

**Lemma 2.1** (Complex matrix variate beta type I and II distribution). If A and B have a complex matrix variate gamma distribution, i.e.  $\mathbf{A} \sim \mathfrak{C} \mathcal{G}_m(a, \mathbf{I}_m)$  and  $\mathbf{B} \sim \mathfrak{C} \mathcal{G}_m(b, \mathbf{I}_m)$ independently.

1. Then the complex matrix variate beta type I distribution is defined as

$$
\mathbf{U} = \begin{cases} (\mathbf{A} + \mathbf{B})^{-1/2} \mathbf{A} ((\mathbf{A} + \mathbf{B})^{-1/2})', & Definition 1 \text{ or,} \\ \mathbf{A}^{1/2} (\mathbf{A} + \mathbf{B})^{-1} (\mathbf{A}^{1/2})', & Definition 2. \end{cases}
$$
(3)

Thus under definitions 1 and 2 its density function is denoted as

$$
\mathfrak{C}\mathcal{B} I_m(\mathbf{U}; a, b),
$$

and given by

$$
\frac{1}{\mathfrak{C}\beta_m[a,b]}|\mathbf{U}|^{a-m}|\mathbf{I}_m-\mathbf{U}|^{b-m}(d\mathbf{U}), \quad \mathbf{0} < \mathbf{U} = \mathbf{U}^H < \mathbf{I}_m,\tag{4}
$$

this being denoted as  $\mathbf{U} \sim \mathfrak{C} \mathcal{B} I_m(a, b)$  with  $Re(a) > m - 1$  and  $Re(b) > m - 1$ .

2. Then the complex matrix variate beta type II distribution is defined as

$$
\mathbf{F} = \begin{cases} \mathbf{B}^{-1/2} \mathbf{A} (\mathbf{B}^{-1/2})', & Definition 1, \\ \mathbf{A}^{1/2} \mathbf{B}^{-1} (\mathbf{A}^{1/2})', & Definition 2, \end{cases}
$$
(5)

Thus under definitions 1 and 2 its density function is denoted as

$$
\mathfrak{C}\mathcal{B}II_m(\mathbf{U}; a, b),
$$

and given by

$$
\frac{1}{\mathfrak{C}\beta_m[a,b]}|\mathbf{F}|^{a-m}|\mathbf{I}_m + \mathbf{F}|^{-(a+b)}.(d\mathbf{F}), \quad \mathbf{F} = \mathbf{F}^H > \mathbf{0}.\tag{6}
$$

this being denoted as  $\mathbf{F} \sim \mathcal{C} \mathcal{B} \Pi_m(a, b)$  with  $Re(a) > m - 1$  and  $Re(b) > m - 1$ .

#### 2.2 Complex hypergeometric functions and invariant polynomials

The following definitions of hypergeometric functions with a matrix argument are based on [Constantine \(1963\)](#page-11-6) and [Koev and Edelman \(2006\)](#page-12-9).

<span id="page-3-1"></span>Definition 2.3. The hypergeometric functions of a matrix argument are given by

<span id="page-3-0"></span>
$$
{}_{p}F_{q}^{(\alpha)}(a_1,\ldots,a_p;b_1,\ldots,b_q; \mathbf{X}) = \sum_{t=0}^{\infty} \sum_{\tau} \frac{(a_1)_{\tau}^{(\alpha)} \cdots (a_p)_{\tau}^{(\alpha)}}{(b_1)_{\tau}^{(\alpha)} \cdots (b_q)_{\tau}^{(\alpha)}} \frac{C_{\tau}^{(\alpha)}(\mathbf{X})}{t!},\tag{7}
$$

where  $\sum_{\tau}$  denotes the summation over all the partitions  $\tau = (t_1, \ldots, t_m)$ ,  $t_1 \geq \cdots \geq t_m \geq 0$ , of  $t = t_1 + \cdots + t_m$ ,  $C_{\tau}^{(\alpha)}(X)$  is the Jack polynomial of **X** corresponding to  $\tau$  and the generalised hypergeometric coefficient  $(a)_{\tau}^{(\alpha)}$  is given by

$$
(a)_{\tau}^{(\alpha)} = \prod_{j=1}^{m} (a - (i-1)/(\alpha))_{t_i},
$$

where  $(a)_t = a(a+1)(a+2)\cdots(a+t-1)$ ,  $(a)_0 = 1$ . Here  $\mathbf{X} \in \mathfrak{C}^{m \times m}$ , is a complex symmetric matrix and the parameters  $a_i$ ,  $b_j$  are arbitrary complex numbers.

Other characteristics of the parameters  $a_i$  and  $b_j$  and the convergence of [\(7\)](#page-3-0) appear in [Muirhead \(1982](#page-12-10), p. 258), [Gross and Richards \(1987\)](#page-11-7) and [Ratnarajah](#page-12-11) et al. [\(2005b\)](#page-12-11).

**Remark 2.1.** In Definition [2.3,](#page-3-1) when  $\alpha = 1$  and 2 the complex and real cases are obtained, respectively. In this paper it is considered only the complex case. Then, adopting the notation used by [James \(1964\)](#page-11-2), denoting the complex hypergeometric function, generalised hypergeometric coefficient and zonal polynomials as  ${}_{p}\widetilde{F}_{q}^{(\alpha)} \equiv {}_{p}F_{q}^{(1)}$ ,  $[a]_{\tau} \equiv (a)_{\tau}^{(1)}$ , and  $\widetilde{C}_{\tau}(\cdot) \equiv C_{\tau}^{(1)}(\cdot)$ , respectively

A special case of [\(7\)](#page-3-0) is

$$
{}_{1}\widetilde{F}_{0}(a; \mathbf{X}) = \sum_{t=0}^{\infty} \sum_{\tau} [a]_{\tau} \frac{\widetilde{C}_{\tau}(\mathbf{X})}{t!}, \quad (||\mathbf{X}|| < 1)
$$

$$
= |\mathbf{I}_{m} - \mathbf{X}|^{-a},
$$

where  $\|\mathbf{X}\|$  denotes the maximum of the absolute values of the eigenvalues of X.

From [Ratnarajah](#page-12-11) et al. [\(2005b](#page-12-11)), is known that,

$$
\begin{aligned} \n\widetilde{F}_1(a;c; \mathbf{X}) &= \frac{1}{\mathfrak{C}\beta_m[a,c-a]} \\ \n&\times \int_{\mathbf{0} < \mathbf{Y} = \mathbf{Y}^H < \mathbf{I}_m} \n\widetilde{F}_0(\mathbf{X}\mathbf{Y}) |\mathbf{Y}|^{a-m} |\mathbf{I} - \mathbf{Y}|^{c-a-m}(d\mathbf{Y}), \n\end{aligned}
$$

and

$$
{}_{2}\widetilde{F}_{1}(a,a_{1};c;\mathbf{X}) = \frac{1}{\mathfrak{C}\beta_{m}[a,c-a]} \times \int_{0<\mathbf{Y}=\mathbf{Y}^{H}<\mathbf{I}_{m}} {}_{1}\widetilde{F}_{0}(a_{1};\mathbf{X}\mathbf{Y})|\mathbf{Y}|^{a-m}|\mathbf{I}-\mathbf{Y}|^{c-a-m}(d\mathbf{Y}). \tag{8}
$$

Thus we have

<span id="page-4-0"></span>**Lemma 2.2.** Let  $X = X^H \in \mathbb{C}^{m \times m}$ , with  $\text{Re}(X) < I$ ,  $\text{Re}(a) > m - 1$ ,  $\text{Re}(c) > m - 1$  and  $Re(c - a) > m - 1$ . Then

$$
{}_{p+1}\widetilde{F}_{q+1}(a,a_1,\ldots,a_p;c,b_1,\ldots,b_q; \mathbf{X}) = \frac{1}{\mathfrak{C}\beta_m[a,c-a]}\\ \times \int_{\mathbf{0} < \mathbf{Y} = \mathbf{Y}^H < \mathbf{I}_m} {}_p\widetilde{F}_q(a_1\cdots a_p; b_1\cdots b_q; \mathbf{XY}) |\mathbf{Y}|^{a-m} |\mathbf{I} - \mathbf{Y}|^{c-a-m}(d\mathbf{Y}).
$$

Proof. First, an expansion is applied in terms of complex zonal polynomials

$$
{}_{p}\widetilde{F}_{q}(a_1,\ldots,a_p;b_1,\ldots,b_q; \mathbf{XY})=\sum_{t=0}^{\infty}\sum_{\tau}\frac{[a_1]_{\tau}\cdots[a_p]_{\tau}}{[b_1]_{\tau}\cdots[b_q]_{\tau}}\frac{\widetilde{C}_{\tau}(\mathbf{XY})}{t!}.
$$

Then, after integrating term by term, see [Ratnarajah](#page-12-11) et al. [\(2005b\)](#page-12-11), we have that

$$
\int_{0<\mathbf{Y}=\mathbf{Y}^H<\mathbf{I}_m} p\widetilde{F}_q(a_1\cdots a_p; b_1\cdots b_q; \mathbf{XY})|\mathbf{Y}|^{a-m}|\mathbf{I}-\mathbf{Y}|^{c-a-m}(d\mathbf{Y})
$$
\n
$$
=\sum_{t=0}^{\infty}\sum_{\tau}\frac{[a_1]_{\tau}\cdots[a_p]_{\tau}}{[b_1]_{\tau}\cdots[b_q]_{\tau}}\int_{0<\mathbf{Y}=\mathbf{Y}^H<\mathbf{I}_m} |\mathbf{Y}|^{a-m}|\mathbf{I}-\mathbf{Y}|^{c-a-m}\widetilde{C}_{\tau}(\mathbf{XY})(d\mathbf{Y})
$$
\n
$$
=\mathfrak{C}\beta_m[a,c-a]\sum_{t=0}^{\infty}\sum_{\tau}\frac{[a]_{\tau}}{[c]_{\tau}}\frac{[a_1]_{\tau}\cdots[a_p]_{\tau}}{[b_1]_{\tau}\cdots[b_q]_{\tau}}\widetilde{C}_{\tau}(\mathbf{X})
$$
\n
$$
=\mathfrak{C}\beta_m[a,c-a]_{p+1}\widetilde{F}_{q+1}(a,a_1\cdots a_p;c,b_1\cdots b_q;\mathbf{X}),
$$

and the required result follows.

The use of complex zonal polynomials and the hypergeometric function with a matrix argument has only recently been extended; to a large extent this is derived from the work of [Koev and Edelman \(2006\)](#page-12-9), who in [Koev \(2004](#page-11-8)) provided a program in MatLab with a very efficient algorithm for calculating Jack polynomials (in particular complex zonal polynomials) and the complex hypergeometric function with a matrix argument.

This section concludes by establishing the following two properties of a class of homogeneous polynomials  $\widetilde{C}^{\kappa,\tau}_\phi(\mathbf{R},\mathbf{\mathbf{S}})$  of degrees  $k$  and  $t$  in the eigenvalues of the Hermitian matrices  $\mathbf{R}, \mathbf{S} \in \mathfrak{C}^{m \times m}$ , respectively, see [Davis \(1980](#page-11-9)) and [Ratnarajah](#page-12-2) *et al.* [\(2005a\)](#page-12-2). These properties generalise the incomplete beta function equation (61) of [Constantine \(1963\)](#page-11-6). The first is proposed by [Davis \(1979,](#page-11-10) eq. (3.3)) and the second is obtained using the complex versions of [Chikuse \(1980,](#page-11-11) eq. (3.33)) and the review version of [Chikuse \(1980](#page-11-11), eq. (3.11)) given in Chikuse and Davis  $(1986, eq. (2.7)).$ 

**Lemma 2.3.** Let  $\mathbf{R}, \mathbf{S}, \mathbf{\Omega}, \Xi \in \mathfrak{C}^{m \times m}$  Hermitian matrices. Then

<span id="page-5-2"></span>
$$
\int_{0}^{\Delta=\Delta^{H}} |\mathbf{R}|^{a-m} |\mathbf{I} - \mathbf{R}|^{b-m} \widetilde{C}_{\tau}(\Omega \mathbf{R}) (d\mathbf{R})
$$
  

$$
= \mathfrak{C} \beta_{m}[a, m] |\Delta|^{a} \sum_{k=0}^{\infty} \sum_{\kappa; \phi \in \kappa. \tau} \frac{[a]_{\phi} [-b + m]_{\kappa} \theta_{\phi}^{\kappa, \tau} \widetilde{C}_{\phi}^{\kappa, \tau}(\Delta, \Omega \Delta)}{k! [a + m]_{\phi}}
$$
(9)

and

$$
\int_{0}^{\Delta=\Delta^{H}} |\mathbf{R}|^{a-m} |\mathbf{I}-\mathbf{R}|^{b-m} \widetilde{C}_{\phi}^{\kappa,\tau}(\mathbf{\Xi},\mathbf{\Omega}\mathbf{R}) (d\mathbf{R})
$$
  

$$
= \mathfrak{C} \beta_{m}[a,m] |\Delta|^{a} \sum_{s=0}^{\infty} \sum_{\sigma;\phi_{1}\in\sigma,\phi} \frac{[a]_{\phi_{1}}[-b+m]_{\sigma} \pi^{\sigma,\kappa,\tau;\phi_{1}}_{\sigma,\phi} \widetilde{C}_{\phi_{1}}^{\sigma,\kappa,\tau}(\Delta,\mathbf{\Xi},\mathbf{\Omega}\Delta)}{s! [a+m]_{\phi_{1}}} \qquad (10)
$$

<span id="page-5-3"></span>where, from [Chikuse and Davis \(1986,](#page-11-12) Lemma 2.2(i) and (ii), respectively) it is known that

$$
\theta^{\kappa,\tau}_\phi = \frac{\widetilde{C}^{\kappa,\tau}_\phi(\mathbf{I},\mathbf{I})}{\widetilde{C}_\phi(\mathbf{I})} \quad \text{and} \quad \pi^{\sigma,\kappa,\tau;\phi_1}_{\sigma,\phi} = \sum_{\phi_1' \equiv \phi_1} \gamma^{\sigma,\kappa,\tau;\phi_1}_{\sigma;\phi_1'} \overline{\alpha^{\sigma^*,\kappa,\tau;\phi_1'}_{\phi}}.
$$

### <span id="page-5-0"></span>3 Bimatrix variate generalised beta type I distribution

Let **A**, **B** and **C** be independent complex random matrices, such that  $\mathbf{A} \sim \mathfrak{C} \mathcal{G}_m(a, \mathbf{I}_m)$ ,  $\mathbf{B} \sim \mathfrak{C} \mathcal{G}_m(b, \mathbf{I}_m)$  and  $\mathbf{C} \sim \mathfrak{C} \mathcal{G}_m(c, \mathbf{I}_m)$  with  $\text{Re}(a) > m-1$ ,  $\text{Re}(b) > m-1$  and  $\text{Re}(c) > m-1$ and let us define

<span id="page-5-1"></span>
$$
U_1 = (A + C)^{-1/2} A(A + C)^{-1/2} \text{ and } U_2 = (B + C)^{-1/2} B(B + C)^{-1/2}. \tag{11}
$$

 $\Box$ 

Of course,  $U_1 \sim \mathfrak{C} \mathcal{B} I_m(a, c)$  and  $U_2 \sim \mathfrak{C} \mathcal{B} I_m(b, c)$ . However, they are correlated such that the distribution of  $\mathbb{U} = (\mathbf{U}_1 \mathbf{U}_2)' \in \mathfrak{C}^{2m \times m}$  can be termed a complex bimatrix variate generalised beta type I distribution, denoted as

$$
\mathbb{U} \sim \mathfrak{CBGBI}_{2m \times m}(a, b, c).
$$

<span id="page-6-1"></span>**Theorem 3.1.** Assume that  $\mathbb{U} \sim \mathcal{BGBI}_{2m \times m}(a, b, c)$ . Then its density function is

<span id="page-6-0"></span>
$$
\frac{|\mathbf{U}_1|^{a-m}|\mathbf{U}_2|^{b-m}|\mathbf{I}_m - \mathbf{U}_1||^{b+c-m}|\mathbf{I}_m - \mathbf{U}_2|^{a+c-m}}{\mathfrak{C}\beta_m^*[a, b, c]|\mathbf{I}_m - \mathbf{U}_1\mathbf{U}_2|^{a+b+c}}(d\mathbb{U}),\tag{12}
$$

 $0 < U_1 = U_1^H < I_m$ ,  $0 < U_2 = U_2^H < I_m$ , where the measure

$$
(d\mathbb{U})=(d\mathbf{U}_1)\wedge (d\mathbf{U}_2),
$$

and

$$
\mathfrak{C} \beta^*_m [a, b, c] = \frac{\mathfrak{C} \Gamma_m [a] \mathfrak{C} \Gamma_m [b] \mathfrak{C} \Gamma_m [c]}{\mathfrak{C} \Gamma_m [a + b + c]}.
$$

and  $\text{Re}(a) > m - 1$ ,  $\text{Re}(b) > m - 1$  and  $\text{Re}(c) > m - 1$ .

Proof. The joint density of A, B and C is

$$
\frac{|\mathbf{A}|^{a-m}|\mathbf{B}|^{b-m}|\mathbf{C}|^{c-m}}{\mathfrak{C}\Gamma_m[a]\mathfrak{C}\Gamma_m[b]\mathfrak{C}\Gamma_m[c]} \operatorname{etr}(-(\mathbf{A}+\mathbf{B}+\mathbf{C}))(d\mathbf{A})(d\mathbf{B})(d\mathbf{C}).
$$

By effecting the change of variable [\(11\)](#page-5-1), and taking into account [Mathai \(1997,](#page-12-8) Theorems 3.5 and 3.8, pp. 183 and 190, respectively) we have

$$
(d\mathbf{A})(d\mathbf{B})(d\mathbf{C}) = |\mathbf{C}|^{2m} |\mathbf{I}_m - \mathbf{U}_1|^{-2m} |\mathbf{I}_m - \mathbf{U}_2|^{-2m} (d\mathbf{U}_1)(d\mathbf{U}_2)(d\mathbf{C}).
$$

The joint density of  $U_1$ ,  $U_2$  and  $C$  is

$$
\frac{|\mathbf{U}_1|^{a-m}|\mathbf{U}_2|^{b-m}}{\mathfrak{C}\Gamma_m[a]\mathfrak{C}\Gamma_m[b]\mathfrak{C}\Gamma_m[c]|\mathbf{I}_m-\mathbf{U}_1|^{a+m}|\mathbf{I}_m-\mathbf{U}_2|^{b+m}}|\mathbf{C}|^{a+b+c-m}
$$
  
×  $\text{etr}\left[-(\mathbf{I}_m-\mathbf{U}_2)^{-1}(\mathbf{I}_m-\mathbf{U}_1\mathbf{U}_2)(\mathbf{I}_m-\mathbf{U}_1)^{-1}\mathbf{C}\right] (d\mathbf{C})(d\mathbf{U}_1)(d\mathbf{U}_2).$ 

Integrating with respect to C using

$$
\int_{\mathbf{C}=\mathbf{C}^{H}>0} |\mathbf{C}|^{a+b+c-m} \operatorname{etr} \left[ -(\mathbf{I}_{m} - \mathbf{U}_{2})^{-1} (\mathbf{I}_{m} - \mathbf{U}_{1} \mathbf{U}_{2}) (\mathbf{I}_{m} - \mathbf{U}_{1})^{-1} \mathbf{C} \right] (d\mathbf{C})
$$
\n
$$
= \mathfrak{C} \Gamma [a+b+c] \frac{|\mathbf{I}_{m} - \mathbf{U}_{1}|^{a+b+c} |\mathbf{I}_{m} - \mathbf{U}_{2}|^{a+b+c}}{|\mathbf{I}_{m} - \mathbf{U}_{1} \mathbf{U}_{2}|^{a+b+c}},
$$

(from [\(2\)](#page-3-2)) gives the stated marginal density function for  $(\mathbf{U}_1:\mathbf{U}_2)'$ .

$$
\mathbf{L}^{\prime}
$$

As in the real case (Díaz-García and Gutiérrez-Jáimez, [2009a\)](#page-11-4), the joint density  $(12)$  can be represented as a mixture. Let us first note that

$$
|\mathbf{I}_{m} - \mathbf{U}_{1} \mathbf{U}_{2}|^{-(a+b+c)} = 1 \widetilde{F}_{0}(a+b+c; \mathbf{U}_{1} \mathbf{U}_{2})
$$
  
= 
$$
\sum_{t=0}^{\infty} \sum_{\tau} [a+b+c]_{\tau} \frac{\widetilde{C}_{\tau}(\mathbf{U}_{1} \mathbf{U}_{2})}{t!}.
$$

By substituting in [\(12\)](#page-6-0) it is obtained that the joint density function of  $(\mathbf{U}_1 \cdot \mathbf{U}_2)'$  is

<span id="page-7-1"></span>
$$
\sum_{t=0}^{\infty} \sum_{\tau} \frac{[a+b+c]_{\tau}}{\mathfrak{C} \beta_m^*[a,b,c]} |\mathbf{U}_1|^{a-m} |\mathbf{U}_2|^{b-m} |\mathbf{I}_m - \mathbf{U}_1| |_{\mathcal{M}}^{b+c-m} \times |\mathbf{I}_m - \mathbf{U}_2|^{a+c-m} \frac{\widetilde{C}_{\tau}(\mathbf{U}_1 \mathbf{U}_2)}{t!}.
$$
 (13)

Moreover

$$
\sum_{t=0}^{\infty}\sum_{\tau}\frac{[a+b+c]_{\tau}\mathfrak{C}\Gamma_{m}[b+c]\mathfrak{C}\Gamma_{m}[a+c]}{\mathfrak{C}\Gamma_{m}[a+b+c]\mathfrak{C}\Gamma_{m}[c]}\mathfrak{C}\mathcal{B}I_{m}(\mathbf{U}_{1};a,b+c)\frac{\times}{\mathfrak{C}\mathcal{B}I_{m}(\mathbf{U}_{2};b,a+c)}\frac{\widetilde{C}_{\tau}(\mathbf{U}_{1}\mathbf{U}_{2})}{t!}.
$$

## <span id="page-7-0"></span>4 Bimatrix variate generalised beta type II distribution

Let **A**, **B** and **C** ∈  $\mathbb{C}^{m \times m}$  be independent, where  $\mathbf{A} \sim \mathfrak{C} \mathcal{G}_m(a, \mathbf{I}_m)$ ,  $\mathbf{B} \sim \mathfrak{C} \mathcal{G}_m(b, \mathbf{I}_m)$  and  $\mathbf{C} \sim \mathfrak{C} \mathcal{G}_m(c, \mathbf{I}_m)$  with  $\text{Re}(a) > m - 1$ ,  $\text{Re}(b) > m - 1$  and  $\text{Re}(c) > m - 1$  and let us define

$$
\mathbf{F}_1 = \mathbf{C}^{-1/2} \mathbf{A} \mathbf{C}^{-1/2} \quad \text{and} \quad \mathbf{F}_2 = \mathbf{C}^{-1/2} \mathbf{B} \mathbf{C}^{-1/2}.
$$
 (14)

Clearly,  $\mathbf{F}_1 \sim \mathcal{C} \mathcal{B} I I_m(a,c)$  and  $\mathbf{F}_2 \sim \mathcal{C} \mathcal{B} I I_m(b,c)$ . But they are correlated and so the distribution of  $\mathbb{F} = (\mathbf{F}_1 \cdot \mathbf{F}_2)' \in \mathbb{R}^{2m \times m}$  can be termed a complex bimatrix variate generalised beta type II distribution, which is denoted as  $\mathbb{F} \sim \mathfrak{C}\mathcal{B}\mathcal{B}II_{2m \times m}(a, b, c)$ .

**Theorem 4.1.** Assume that  $\mathbb{F} \sim \mathcal{BGBI}_{2m \times m}(a, b, c)$ . Then its density function is

$$
\frac{|\mathbf{F}_1|^{a-m}|\mathbf{F}_2|^{b-m}}{\mathfrak{C}\beta_m^*[a,b,c]|\mathbf{I}_m + \mathbf{F}_1 + \mathbf{F}_2|^{a+b+c}}(d\mathbb{F}),\tag{15}
$$

 $\mathbf{F}_1 = \mathbf{F}_1^H > \mathbf{0}, \ \mathbf{F}_2 = \mathbf{F}_2^H > \mathbf{0}, \ where \ the \ measure$ 

$$
(d\mathbb{F}) = (d\mathbf{F}_1) \wedge (d\mathbf{F}_2).
$$

and  $\text{Re}(a) > m - 1$ ,  $\text{Re}(b) > m - 1$  and  $\text{Re}(c) > m - 1$ .

*Proof.* As an alternative to proceeding as in Theorem [3.1,](#page-6-1) let us recall that if  $\mathbf{U} \sim \mathfrak{C} \mathcal{B} I_m(a, b)$ , then  $(\mathbf{I}_m-\mathbf{U})^{-1}-\mathbf{I}_m \sim \mathfrak{C}\mathcal{B}II_m(a, b)$ , see [Srivastava and Khatri \(1979\)](#page-12-12) and Díaz-García and Gutiérrez-Jáimez [\(2007](#page-11-13)). Then

$$
\mathbb{F} = \left( \begin{array}{c} \mathbf{F}_1 \\ \mathbf{F}_2 \end{array} \right) = \left( \begin{array}{c} (\mathbf{I}_m - \mathbf{U}_1)^{-1} - \mathbf{I}_m \\ (\mathbf{I}_m - \mathbf{U}_2)^{-1} - \mathbf{I}_m \end{array} \right),
$$

with the Jacobian given by (see [Mathai \(1997](#page-12-8), Theorem 3.8, p. 190))

$$
(d\mathbf{U}_1)(d\mathbf{U}_2) = |\mathbf{I}_m + \mathbf{F}_1|^{-2m} |\mathbf{I}_m + \mathbf{F}_2|^{-2m} (d\mathbf{F}_1)(d\mathbf{F}_2).
$$

Also, note that

$$
\mathbf{I}_m - (\mathbf{I}_m + \mathbf{F}_1)^{-1} = (\mathbf{I}_m + \mathbf{F}_1)^{-1} ((\mathbf{I}_m + \mathbf{F}_1) - \mathbf{I}_m) = (\mathbf{I}_m + \mathbf{F}_1)^{-1} \mathbf{F}_1 \n\mathbf{I}_m - (\mathbf{I}_m + \mathbf{F}_2)^{-1} = (\mathbf{I}_m + \mathbf{F}_2)^{-1} \mathbf{F}_2,
$$

Then the joint density of  $(\mathbf{F}_1 \cdot \mathbf{F}_2)'$  is

$$
\frac{|\mathbf{F}_1|^{a-m}|\mathbf{F}_2|^{b-m}|\mathbf{I}_m+\mathbf{F}_1|^{-(a+b+c)}|\mathbf{I}_m+\mathbf{F}_2|^{-(a+b+c)}}{\mathfrak{C}\beta_m^*[a,b,c]|\mathbf{I}_m-(\mathbf{I}_m+\mathbf{F}_1)^{-1}\mathbf{F}_1\mathbf{F}_2(\mathbf{I}_m+\mathbf{F}_2)^{-1}|^{a+b+c}}(d\mathbb{F}).
$$

The desired results then follows, noting that

$$
\frac{|\mathbf{I}_m + \mathbf{F}_1|^{-1}|\mathbf{I}_m + \mathbf{F}_2|^{-1}}{|\mathbf{I}_m - (\mathbf{I}_m + \mathbf{F}_1)^{-1}\mathbf{F}_1\mathbf{F}_2(\mathbf{I}_m + \mathbf{F}_2)^{-1}|} = |\mathbf{I}_m + \mathbf{F}_1 + \mathbf{F}_2|^{-1}.
$$

Other properties of the distribution  $\mathcal{L}\mathcal{B}\mathcal{B}II_{2m\times m}(a, b, c)$  can be found in a similar way.

## <span id="page-8-0"></span>5 Properties

In this section, several basic properties and eigenvalue distributions are studied.

#### 5.1 Basic properties

As direct consequences of Lemma [2.2](#page-4-0) the following basic properties are derived: the moments  $E(|\mathbf{U}_1|^r|\mathbf{U}_2|^s)$ , the distributions of the product  $\mathbf{Z} = \mathbf{U}_2^{1/2}\mathbf{U}_1\mathbf{U}_2^{1/2}$  and the inverse

 $(\mathbf{U}_1^{-1}$  $\cdot :$ U<sub>2</sub><sup>-1</sup>). Their proofs are similar to those given for the real case, see Díaz-García and Gutiérrez-Jáimez [\(2009a](#page-11-4)).

**Theorem 5.1.** Assume that  $(U_1:U_2) \sim \mathcal{C} \mathcal{B} \mathcal{G} \mathcal{B} I_{2m \times m}(a, b, c)$  then

$$
E(|\mathbf{U}_1|^r |\mathbf{U}_2|^s) = \frac{\mathfrak{C}\beta_m[a+r,b+c]\mathfrak{C}\beta_m[b+s,a+c]}{\mathfrak{C}\beta_m^*[a,b,c]} \times {}_3\widetilde{F}_2(a+r,b+s,a+b+c;a+b+c+r,a+b+c+s;\mathbf{I}_m),
$$

with  $\text{Re}(b + r) > m - 1$ , and  $\text{Re}(a + c) > m - 1$ .

**Theorem 5.2.** Consider that  $(U_1:U_2) \sim \mathcal{C} \mathcal{B} \mathcal{B} I_{2m \times m}(a,b,c)$ . Then the density function of  $\mathbf{Z} = \mathbf{Z}^H = \mathbf{U}_2^{1/2} \mathbf{U}_1 \mathbf{U}_2^{1/2} \in \mathfrak{C}^{m \times m}$  is

$$
\frac{\mathfrak{C}\beta_m[a+c,b+c]|{\bf Z}|^{a-m}|{\bf I}_m-{\bf Z}|^{c-m}}{\mathfrak{C}\beta^*_m[a,b,c]}\widehat{F}_1(a+c,a+c;a+b+2c;{\bf I}_m-{\bf Z})(d{\bf Z}),
$$

and

$$
E(|\mathbf{Z}|^r)=\frac{\mathfrak{C}\pmb{\beta}_m[a+c,b+c] \pmb{\beta}_m[a+r,c]}{\mathfrak{C}\pmb{\beta}_m^*[a,b,c]} {}_3\widetilde{F}_2(c,a+c,a+c;a+c+r,a+b+2c;\mathbf{I}_m),
$$

with  $0 < \text{Re}(\mathbf{Z}) < \mathbf{I}_m$ ,  $\text{Re}(a+b) > m-1$  and  $\text{Re}(b+c) > m-1$ .

**Theorem 5.3.** Let  $(\mathbf{U}_1 \cdot \mathbf{U}_2)' \sim \mathcal{C} \mathcal{B} \mathcal{G} \mathcal{B} I_{2m \times m}(a, b, c)$ . Then the density function of  $\mathbb{V} =$  $({\bf V}_1.{\bf V}_2)'=({\bf U}_1^{-1}$  $\therefore$ U<sub>2</sub><sup>-1</sup>)'  $\in \mathfrak{C}^{2m \times m}$  is

$$
\frac{|\mathbf{V}_1|^{-a-m}|\mathbf{V}_2|^{-b-m}|\mathbf{I}_m-\mathbf{V}_1^{-1}||^{b+c-m}|\mathbf{I}_m-\mathbf{V}_2^{-1}|^{a+c-m}}{\mathfrak{C}\pmb{\beta}_m^*[a,b,c]|\mathbf{I}_m-(\mathbf{V}_1\mathbf{V}_2)^{-1}|^{a+b+c}}(d\mathbb{V}),
$$

 $0 < V_1 = V_1^H < I_m$ ,  $0 < V_2 = V_2^H < I_m$ , where the measure

$$
(d\mathbb{V}) = (d\mathbf{V}_1) \wedge (d\mathbf{V}_2).
$$

and  $\text{Re}(a) > m - 1$ ,  $\text{Re}(b) > m - 1$  and  $\text{Re}(c) > m - 1$ .

#### 5.2 Joint eigenvalue distribution

Many statistics of the multivariate test hypothesis are functions of the eigenvalues or of the maximum eigenvalue. In these final two subsections, we find the eigenvalues distribution. The following result is needed, see [James \(1964\)](#page-11-2).

<span id="page-9-1"></span>**Lemma 5.1.** If  $f_{\mathbf{A}}(\mathbf{A})$  (d $\mathbf{A}$ ) is the density function of a Hermitian matrix variate  $\mathbf{A} \in$  $\mathfrak{C}^{m \times m}$ , then the distribution of the diagonal matrix  $\Lambda = \text{diag}(\lambda_1, \ldots, \lambda_m)$ ,  $\lambda_1 > \cdots > \lambda_m >$ 0, of the eigenvalues of **A**, where  $\mathbf{A} = \mathbf{G}\Lambda\mathbf{G}^H$ ,  $\mathbf{G} \in \mathcal{U}(m)$ , is the eigendecomposition of **A**, is

$$
f_{\mathbf{\Lambda}}(\mathbf{\Lambda}) = \frac{\pi^{m(m-1)}}{\mathfrak{C}\Gamma_m[m]} \prod_{r < s} (\lambda_r - \lambda_s)^2 \int_{\mathcal{U}(m)} f_{\mathbf{\Lambda}}(\mathbf{G}\mathbf{\Lambda}\mathbf{G}^H)(d\mathbf{G}),
$$

where  $(d\mathbf{G})$  is the invariant measure on the unitary group  $\mathcal{U}(m)$  normalised, given as

<span id="page-9-2"></span>
$$
(d\mathbf{G}) = \frac{\mathfrak{C}\Gamma_m[m]}{2^m \pi^{m^2}} (\mathbf{G} d\mathbf{G}^H), \quad \text{such that} \quad \int_{\mathcal{U}(m)} (d\mathbf{G}) = 1. \tag{16}
$$

**Theorem 5.4.** Assume that  $(U_1:U_2) \sim \mathfrak{C} \mathcal{B} \mathcal{G} \mathcal{B} I_{2m \times m}(a, b, c)$  and let

<span id="page-9-0"></span>
$$
\mathbb{U} = \left(\begin{array}{c} \mathbf{U}_1 \\ \mathbf{U}_2 \end{array}\right) = \left(\begin{array}{c} \mathbf{ED}_{\lambda} \mathbf{E}^H \\ \mathbf{GD}_{\delta} \mathbf{G}^H \end{array}\right). \tag{17}
$$

The spectral decomposition of  $U_1$  and  $U_2$ , with  $E, G \in \mathcal{U}(m)$  and  $D_\lambda = \text{diag}(\lambda_1, \ldots, \lambda_m)$ ,  $1 > \lambda_1 > \ldots, \lambda_m > 0$  and  $\mathbf{D}_{\delta} = \text{diag}(\delta_1, \ldots, \delta_m)$ ,  $1 > \delta_1 > \ldots, \delta_m > 0$ . Then the joint density function of  $\lambda_1, \ldots, \lambda_m, \delta_1, \ldots, \delta_m$  is

$$
\frac{\pi^{2m(m-1)}}{(\mathfrak{C}\Gamma_m[m])^2 \mathfrak{C}\beta^*[a,b,c]} \prod_{r=1}^m \left(\lambda_r^{a-m}(1-\lambda_r)^{b+c-m}\right) \prod_{e=1}^m \left(\delta_e^{b-m}(1-\delta_e)^{a+c-m}\right)
$$

$$
\times \prod_{r
$$

with  $\text{Re}(a) > m - 1$ ,  $\text{Re}(b) > m - 1$  and  $\text{Re}(c) > m - 1$ .

Proof. . The desired result follows by making the transformation [\(17\)](#page-9-0), from Lemma [5.1,](#page-9-1) equation [\(16\)](#page-9-2) and observing that, see [Chikuse \(2004](#page-11-14))

$$
\int_{\mathbf{G}\in\mathcal{U}(m)}\int_{\mathbf{E}\in\mathcal{U}(m)}\widetilde{C}_{\kappa}(\mathbf{E}\mathbf{D}_{\lambda}\mathbf{E}^{H}\mathbf{G}\mathbf{D}_{\delta}\mathbf{G}^{H})(d\mathbf{E})(d\mathbf{G})=\frac{\widetilde{C}_{\kappa}(\mathbf{D}_{\lambda})\widetilde{C}_{\kappa}(\mathbf{D}_{\delta})}{\widetilde{C}_{\kappa}(\mathbf{I})}.\quad \Box
$$

#### 5.3 Joint distribution of  $\lambda_{max}$  and  $\delta_{max}$

In this subsection, we derive the distribution of the largest eigenvalues,  $\lambda_{max}$  and  $\delta_{max}$  of a complex bimatrix variate beta type I matrix. With this aim, let us consider the following theorem.

<span id="page-9-3"></span>Theorem 5.5. Assume that  $(U_1:U_2) \sim \mathcal{C} \mathcal{B} \mathcal{G} \mathcal{B} I_{2m \times m}(a,b,c)$  and let  $\Delta_1, \Delta_2 \in \mathfrak{C}^{m \times m}$ be Hermitian positive definite matrices,  $0 < \Delta_1, \Delta_2 < I$ . Then the probability  $P(U_1 <$  $\Delta_1, U_2 < \Delta_2$ ) is given by

$$
\begin{split} \frac{\mathfrak{C}\beta_{m}[a,m]\mathfrak{C}\beta_{m}[b,m]}{\mathfrak{C}\beta_{m}^{*}[a,b,c]}|\Delta_{1}|^{a}|\Delta_{2}|^{b}\sum_{k,t,s=0}^{\infty}\sum_{\kappa,\tau,\sigma;\phi_{1}\in\kappa.\tau;\phi_{2}\in\sigma^{*}.\phi_{1}^{*}}\frac{[a]_{\phi_{2}}[b]_{\phi_{1}}[a+b+c]_{\kappa}}{k! \;t! \;s!}\\ \times\frac{[-(a+c)+m]_{\tau}[-(b+c)+m]_{\sigma}}{[a+m]_{\phi_{2}}[b+m]_{\phi_{1}}} \theta_{\phi_{1}}^{\kappa,\tau,\pi,\sigma,\tau,\kappa;\phi_{2}}\widetilde{C}_{\phi_{2}}^{\sigma,\tau,\kappa}(\Delta_{1},\Delta_{2},\Delta_{1}\Delta_{2})\\ \sum_{\kappa,\tau,\sigma;\phi_{1}\in\kappa.\tau;\phi_{2}\in\sigma^{*}.\phi_{1}^{*}}=\sum_{\kappa,\tau,\sigma}\sum_{\phi_{1}\in\kappa.\tau}\sum_{\phi_{2}\in\sigma^{*}.\phi_{1}^{*}} \end{split}
$$

where

and  $\text{Re}(a) > m - 1$ ,  $\text{Re}(b) > m - 1$  and  $\text{Re}(c) > m - 1$ .

*Proof.* From [\(12\)](#page-6-0) the probability  $P(\mathbf{U}_1 < \mathbf{\Delta}_1, \mathbf{U}_2 < \mathbf{\Delta}_2)$  is

<span id="page-10-0"></span>
$$
\int_{\mathbf{O}}^{\mathbf{\Delta}_1} \int_{\mathbf{O}}^{\mathbf{\Delta}_2} \frac{|\mathbf{U}_1|^{a-m} |\mathbf{U}_2|^{b-m} |\mathbf{I}_m - \mathbf{U}_1|^{b+c-m} |\mathbf{I}_m - \mathbf{U}_2|^{a+c-m}}{\mathfrak{C} \beta_m^*[a, b, c] |\mathbf{I}_m - \mathbf{U}_1 \mathbf{U}_2|^{a+b+c}} (d\mathbf{U}_1)(d\mathbf{U}). \tag{18}
$$

□

By rewritten [\(18\)](#page-10-0) as in [\(13\)](#page-7-1), and integrating with respect to  $U_2$  using [\(9\)](#page-5-2), it is obtained that

$$
\frac{\mathfrak{C}\beta_m[b,m]}{\mathfrak{C}\beta_m^*[a,b,c]} |\mathbf{\Delta}_1|^a \sum_{k,t=0}^{\infty} \sum_{\kappa,\tau;\phi_1 \in \kappa.\tau} \frac{[b]_{\phi_1}[a+b+c]_{\kappa}[-(a+c)+m]_{\tau}\theta_{\phi_1}^{\kappa,\tau}}{k! \ t! [b+m]_{\phi_1}} \\ \times \int_{\mathbf{O}}^{\mathbf{\Delta}_1} |\mathbf{U}_1|^{a-m} |\mathbf{I}_m - \mathbf{U}_1| |^{b+c-m} \widetilde{C}_{\phi_1}^{\tau,\kappa}(\mathbf{\Delta}_2, \mathbf{\Delta}_2 \mathbf{U}_1)(d\mathbf{U}_1).
$$

The desired result follows by integrating with respect to  $U_1$  using [\(10\)](#page-5-3).

The following result is obtained from Theorem [5.5.](#page-9-3)

Corollary 5.1. Let  $(U_1:U_2) \sim \mathfrak{C} \mathcal{B} \mathcal{B} I_{2m \times m}(a,b,c)$ , Re $(a) > m-1$ , Re $(b) > m-1$  and  $\text{Re}(c) > m - 1$ . If  $\lambda_{max}$  and  $\delta_{max}$  are the largest eigenvalues of (U<sub>1</sub> and U<sub>2</sub>, respectively, then their joint distribution function,  $P(\lambda_{max} < x, \delta_{max} < y)$  is given by

$$
\frac{\mathfrak{C}\beta_{m}[a,m]\mathfrak{C}\beta_{m}[b,m]}{\mathfrak{C}\beta_{m}^{*}[a,b,c]} \sum_{k,t,s=0}^{\infty} \sum_{\kappa,\tau,\sigma;\phi_{1}\in\kappa.\tau;\phi_{2}\in\sigma^{*}.\phi_{1}^{*}} \frac{x^{am+s+k}y^{bm+t+k}[a]_{\phi_{2}}[b]_{\phi_{1}}}{k! \; t! \; s!} \times \frac{[-(a+c)+m]_{\tau}[-(b+c)+m]_{\sigma}}{[a+b+c]_{\kappa}[a+m]_{\phi_{2}}[b+m]_{\phi_{1}}} \theta_{\phi_{1}}^{\kappa,\tau}\pi_{\sigma,\phi_{1}}^{\sigma,\tau,\kappa;\phi_{2}}\theta_{\phi_{2}}^{\kappa,\tau,\sigma}\widetilde{C}_{\phi_{2}}^{\sigma,\tau,\kappa}(\mathbf{I}).
$$

*Proof.* . The inequalities  $\lambda_{max} < x$  and  $\delta_{max} < y$  are equivalent to  $U_1 < xI$  and  $U_2 < yI$ . Therefore, the result follows by letting  $\Delta_1 = xI$  and  $\Delta_2 = yI$  in Theorem [5.5,](#page-9-3) and taking into account that from Davis  $(1979, \text{ eqs. } (2.1) \text{ and } (2.7))$ 

$$
\widetilde{C}^{\sigma,\tau,\kappa}_{\phi_2}(x\mathbf{I},y\mathbf{I},xy\mathbf{I})=x^{s+k}y^{t+k}\widetilde{C}^{\sigma,\tau,\kappa}_{\phi_2}(\mathbf{I}).\qquad \qquad \Box
$$

## 6 Conclusions

Random matrices play a prominent role because of their deep mathematical structure. They have arisen in a number of fields (statistics, graph theory, stochastic linear algebra, physics, signal processing, etc.), often independently. In particular, the statisticians are interested in studying the Normal (Gaussian), Wishart, manova, and circular random matrices, which, from the point of view of random matrix theory are termed Hermite, Laguerre, Jacobi, and Fourier ensembles, [Mehta \(1991\)](#page-12-0) and [Edelman and Rao \(2005\)](#page-11-15).

The results obtained in the present work can be considered as a generalisation of the Jacobi ensemble for the case in which there are two correlated Jacobi ensembles, and with all their potential applications, see [Edelman and Sutton \(2008\)](#page-11-16).

Another potential use appears in the context of complex shape theory, see [Micheas](#page-12-1) *et al.* [\(2006](#page-12-1)). Specifically, in the approach known as affine shape or configuration densities, see [Caro-Lopera](#page-11-17) et al. [\(2009\)](#page-11-17).

## Acknowledgments

This research work was partially supported by CONACYT-México, Research Grant No. 81512 and IDI-Spain, Grants No. FQM2006-2271 and MTM2008-05785. This paper was written during J. A. Díaz- García's stay as a visiting professor at the Department of Statistics and O. R. of the University of Granada, Spain.

#### References

- <span id="page-11-17"></span>F. J. Caro-Lopera, J. A. Díaz-García G. González-Farías, Inference in statistical shape theory: Elliptical configuration densities, J. Stat. Res. In press.
- <span id="page-11-3"></span>J. J. Chen, M. R. Novick, Bayesian analysis for binomial models with generalized beta prior distributions, J. Educational Statist. 9 (1984) 163–175.
- <span id="page-11-11"></span>Y. Chikuse, Invariant polynomials with matrix arguments and their applications, In: Gupta R P (ed.) Multivariate Statistical Analysis, North-Holland Publishing Company 1980, 53–68.
- <span id="page-11-12"></span>Y. Chikuse, W. Davis, Some properties of invariant polynomials with matrix arguments and their applications in econometrics, Ann. Inst. Statist. Math. Part A 38 (1986)109–122.
- <span id="page-11-14"></span>Y. Chikuse, Hermite and Lauguerre polynomials with complex matrix arguments Linear Algebra Appl. 388 (2004) 91–105.
- <span id="page-11-6"></span>A. G. Constantine, Some noncentral distribution problems in multivariate analysis, Ann. Math. Statist. 34 (1963) 1270–1285.
- <span id="page-11-10"></span>A. W. Davis, Invariant polynomials with two matrix arguments extending the zonal polynomials: Applications to multivariate distribution theory, Ann. Math. Statist. 31 (1979) 465–485.
- <span id="page-11-9"></span>A. W. Davis, Invariant polynomials with two matrix arguments, extending the zonal polynomials, In Multivariate Analysis V. (P. R. Krishnaiah, Ed.), North-Holland Publishing Company 1980, pp. 287–299..
- <span id="page-11-13"></span>J. A. Díaz-García, R. Gutiérrez-Jáimez, Noncentral, nonsingular matrix variate beta distribution. Brazilian J. Prob. Statist. 21 (2007) 175–186.
- <span id="page-11-4"></span>J. A. Díaz-García, R. Gutiérrez-Jáimez, Bimatrix variate generalised beta distributions, (2009a) http://arxiv.org/abs/0904.1830. Also submitted.
- <span id="page-11-5"></span>J. A. Díaz-García, R. Gutiérrez-Jáimez, Noncentral bimatrix variate generalised beta distributions, (2009b) Submitted.
- <span id="page-11-15"></span>A. Edelman, N. R. Rao, Random matrix Theory, Acta Numerica 14 (2005) 233-297.
- <span id="page-11-16"></span>A. Edelman, B. D. Sutton, The beta-Jacobi matrix model, the CS decomposition, and generalized singular value problem Found. Comput. Math. 2 (2008) 259-285.
- <span id="page-11-1"></span>N. R. Goodman, Statistical analysis based on a certain multivariate complex Gaussian distribution (an introdiction), Ann. Math. Statist. 34 (1963) 152–177.
- <span id="page-11-7"></span>K. I. Gross, D. St. P. Richards, Special functions of matrix argument. I: Algebraic induction, zonal polynomials, and hypergeometric functions, Trans. Amer. Math. Soc. 301 (1987) 781–811.
- <span id="page-11-2"></span>A. T. James, Distributions of matrix variate and latent roots derived from normal samples, Ann. Math. Statist. 35 (1964) 475–501.
- <span id="page-11-0"></span>C. G. Khatri, Classical statistical analysis based on a certain multivariate complex Gaussian distribution, Ann. Math. Statist. 37 (1965) 98–114.
- <span id="page-11-8"></span>Koev, P., 2004. [http://www.math.mit.edu/˜plamen.](http://www.math.mit.edu/~plamen)
- <span id="page-12-9"></span>P. Koev, A. Edelman, The efficient evaluation of the hypergeometric function of a matrix argument, Math. Comp. 75 (2006) 833–846.
- <span id="page-12-6"></span>D. L. Libby, M. R. Novick, Multivariate Generalized beta distributions with applications to utility assessment, J. Educational Statist. 7 (1982) 271–294.
- <span id="page-12-8"></span>A. M. Mathai, Jacobians of matrix transformations and functions of matrix argument, World Scientific, London, 1997.
- <span id="page-12-3"></span>A. m. Mathai, S. B. Provost, Some complex matrix-variate statistical distributions on rectangular matrices, Linear Algebra Appl. 410 (2005) 198–216.
- <span id="page-12-0"></span>M. Metha, Random matrices. Academic Press, San Diego, 1991.
- <span id="page-12-1"></span>A. C. Micheas, D. K. Dey, K. V. Mardia, Complex elliptical distribution with application to shape theory, J. Statist. Plann. Infer. 136 (2006) 2961-2982.
- <span id="page-12-10"></span>R. J. Muirhead, Aspects of Multivariate Statistical Theory, John Wiley & Sons, New York, 1982.
- <span id="page-12-7"></span>I. Olkin, R. Liu, A bivariate beta distribution, Statist. Prob. Letters, 62 (2003) 407–412.
- <span id="page-12-2"></span>T. Ratnarajah, R. Villancourt, M. Alvo, Complex random matrices and Rician channel capacity, Problems of Information transmission 41 (2005a) 1–22.
- <span id="page-12-11"></span>T. Ratnarajah, R. Villancourt, M. Alvo, Eigenvalues and condition numbers of complex random matrices, SIAM J. Matrix Anal. Appl. 26 (2005b) 441–456.
- <span id="page-12-12"></span>M. S. Srivastava, C. G. Khatri, An Introduction to Multivariate Analysis. North-Holland Publ., Amsterdam, 1979.
- <span id="page-12-5"></span>G. L. Turin, The characteristic function of Hermitian quadratic forms in complex normal variables, Biometrika 47 (1960) 199–201.
- <span id="page-12-4"></span>R. A. Wooding, The multivariate distribution of complex normal variables. Biometrika 43 (1956) 212–215.