

# THE BOUNIAKOWSKY CONJECTURE AND THE DENSITY OF POLYNOMIAL ROOTS TO PRIME MODULI

Timothy Foo

*Division of Mathematical Sciences, Nanyang Technological University, Singapore*  
S080074@ntu.edu.sg

## Abstract

We establish a result linking the Bouniakowsky conjecture and the density of polynomial roots to prime moduli.

**Mathematics Subject Classification (2000):** 11B05, 11C08, 11K06, 11N32

**Keywords:** Bouniakowsky conjecture, Schinzel Hypothesis, density, polynomial roots, prime moduli

## Introduction

In this paper, we study roots of irreducible polynomials to prime moduli. We think of  $\mathbf{Z}/p\mathbf{Z}$  as the set  $0, 1, 2, \dots, p-1$  and hence we think of the root of our polynomial as a number in that set. When the root  $z$  is divided by  $p$ , we naturally have a number in  $(0, 1)$ . If we fix a polynomial  $f(x)$  of degree  $n \geq 2$  which is irreducible in  $\mathbf{Z}[x]$ , we can consider the set

$$A_f = \cup_p \left\{ \frac{z}{p} : f(z) \equiv 0 \pmod{p}, 1 \leq z \leq p-1 \right\}$$

The aim of this paper is to prove that if a certain conjecture called the Bouniakowsky conjecture is true, then the set  $A_f$  is dense in  $(0, 1)$ . We stress that our result is conditional. Results that are not dependent on open conjectures have been proven about roots of polynomials to various moduli. Hooley [H] proved that the roots of an irreducible polynomial, considered over the ring  $\mathbf{Z}/n\mathbf{Z}$ ,  $n$  not necessarily prime, when suitably normalized by dividing by  $n$  and considered over all  $n$ , are in fact equidistributed in  $(0, 1)$ . Duke, Friedlander and Iwaniec [DFI] proved equidistribution for quadratic polynomials of negative discriminant, to prime moduli. Toth [T] proved equidistribution for quadratic polynomials of positive discriminant, to prime moduli. We now state the main theorem of our paper.

**Theorem** If the Bouniakowsky conjecture is true, the set  $A_f = \cup_p \left\{ \frac{z}{p} : f(z) \equiv 0 \pmod{p}, 1 \leq z \leq p-1 \right\}$  is dense in  $(0, 1)$ .

## The Bouniakowsky conjecture

We now discuss the Bouniakowsky conjecture to give some background.

**Conjecture** Bouniakowsky Conjecture: Let  $f(x)$  be a polynomial that is irreducible in  $\mathbf{Z}[x]$ . Let  $r_f = \gcd(\{f(x) : x \in \mathbf{Z}\})$ . Then  $\frac{f(x)}{r_f}$  is prime infinitely often.

It is easy to construct polynomials which are always divisible by a given prime  $q$ . We know by Fermat's little theorem that the prime  $q$  always divides  $x^q - x$ . Therefore, all we have to do is choose a value  $k$  so that  $x^q - x + qk$  is irreducible in  $\mathbf{Z}[x]$ . It then follows that  $q$  divides all the values of this polynomial.

## The Result

We first begin by considering a subset of  $(0, 1)$  which we will prove to be dense. We are then going to use this set to help prove the density of  $A_f$ . Here, we let  $n$  be the degree of  $f$  and  $c$  be the leading coefficient of  $f$ .

Let  $B_f = \{\frac{a}{b} : 1 \leq a < b, b \text{ odd prime}, (cr_f, b) = 1, acx^{n-1} = -r_f \pmod{b} \text{ has a solution}\}$

**Lemma 1**  $B_f$  is dense in  $(0, 1)$ .

*Proof*

Case 1:  $n$  is even. Consider the map  $x \rightarrow x^{n-1}$  on  $(\mathbf{Z}/b\mathbf{Z})^*$ . This map is injective and surjective if  $(n-1, b-1) = 1$ . For such  $b$ , we can in fact solve  $acx^{n-1} \equiv -r_f \pmod{b}$  for all  $a \in (\mathbf{Z}/b\mathbf{Z})^*$ . Since  $b$  is prime, we can pick  $b$  larger than  $cr_f$  to ensure  $(b, cr_f) = 1$ . We can also pick infinitely many such  $b$  with  $(n-1, b-1) = 1$ . It thus follows that  $B_f$  is dense in this case.

Case 2:  $n$  is odd. Since  $n-1$  is even, let  $n-1 = 2^e h$ ,  $h$  odd. The map  $x \rightarrow x^{n-1}$  on  $(\mathbf{Z}/b\mathbf{Z})^*$  is therefore a composition of the maps  $x \rightarrow x^2$  applied  $e$  times and  $x \rightarrow x^h$ .  $x \rightarrow x^h$  is a permutation of  $(\mathbf{Z}/b\mathbf{Z})^*$  if  $(b-1, h) = 1$ . Also, if  $b \equiv 3 \pmod{4}$ ,  $x \rightarrow x^2$  is a permutation of the squares in  $(\mathbf{Z}/b\mathbf{Z})^*$ , so by choosing  $b \equiv 3 \pmod{4}$  and  $(b-1, h) = 1$ , we can ensure that the image of  $x \rightarrow x^{n-1}$  is the squares. We also want  $(b, cr_f) = 1$ . We have infinitely many primes  $b$  satisfying these conditions, and for such  $b$ , the numerator of the fractions  $\frac{a}{b}$  ranges over either only the squares or only the nonsquares in  $(\mathbf{Z}/b\mathbf{Z})^*$ . By a result of Brauer, the maximum number of consecutive squares or nonsquares in  $(\mathbf{Z}/b\mathbf{Z})^*$  is less than  $b^{0.5}$  when  $b \equiv 3 \pmod{4}$ . [B] This ensures that  $B_f$  is dense in this case.

We will now show how  $\frac{z}{p}$  is related to the values in  $B_f$ . To do this, first consider the original polynomial  $f$ . From  $f = \sum_i c_i x^i$ , we can construct a polynomial  $g(x, y) = \sum_i c_i x^i y^{n-i}$ . Now for any prime  $b$  with  $(b, cr_f) = 1$  we

have a polynomial in one variable  $g(bw + t, b)$  where  $w$  is the variable and  $t \in (\mathbf{Z}/b\mathbf{Z})^*$ . Since we can vary  $b$  and  $t$ , we have many such polynomials associated to  $f$ . We will show that the gcd of the values of all these polynomials is also  $r_f$  and that they are also irreducible in  $\mathbf{Z}[w]$ . It is these polynomials that we apply the Bouniakowsky conjecture to. If the Bouniakowsky conjecture is true, then there are infinitely many primes  $p$  with  $r_f p = g(bw + t, b)$  as  $w \rightarrow \infty$ . Furthermore, for these primes  $p$ , we can construct a root  $z$  of  $f \bmod p$  such that  $\frac{z}{p}$  is “close” to  $\frac{a}{b}$  where  $a$  is chosen so that  $\frac{ap+bw+t}{b}$  is an integer and  $\frac{a}{b} \in (0, 1)$ . This is the same as choosing  $1 \leq a < b$  and  $a$  such that  $act^{n-1} \equiv -r_f \pmod{b}$ . We thus see the relation to the set  $B_f$ . We then let  $z = \frac{ap+bw+t}{b}$  and show that  $z$  is a root of  $f \bmod p$ .

**Lemma 2** The polynomial  $g(bw + t, b)$ , where  $w$  is the variable,  $b$  is prime,  $(b, cr_f) = 1$ ,  $1 \leq t < b$  is irreducible in  $\mathbf{Z}[w]$ .

*Proof*  $g(bw + t, b)$  is related in a simple way to the original polynomial  $f$ .  $g(bw+t, b) = \sum_i c_i (bw+t)^i b^{n-i} = b^n \sum_i c_i (w+\frac{t}{b})^i = b^n g(w+\frac{t}{b}, 1) = b^n f(w+\frac{t}{b})$ . Since a polynomial is irreducible in  $\mathbf{Z}[x]$  if and only if it is irreducible in  $\mathbf{Q}[x]$ , the lemma follows.

**Lemma 3** Let  $b$  be prime,  $(b, cr_f) = 1$ , and  $1 \leq t < b$ . Then  $\gcd(\{g(bw+t, b) : w \in \mathbf{Z}\}) = r_f$

*Proof* Let  $r = r_f$ . Since  $f$  has integer coefficients, we can think of  $f$  as a polynomial in  $(\mathbf{Z}/r\mathbf{Z})[x]$ . But since  $r$  divides all the values of  $f$ , it follows that  $f(x) = 0$  in  $(\mathbf{Z}/r\mathbf{Z})[x]$ . We showed in the proof of Lemma 2 that  $g(bw + t, b) = b^n f(w + \frac{t}{b})$  in  $\mathbf{Q}[x]$ . Since  $(b, r_f) = 1$ ,  $b$  has an inverse mod  $r$  and hence the rational number  $\frac{t}{b}$  can be thought of as an element in  $\mathbf{Z}/r\mathbf{Z}$ . Hence  $g(bw + t, b) = b^n f(w + \frac{t}{b}) = 0$  in  $(\mathbf{Z}/r\mathbf{Z})[x]$ . Therefore, for each such  $b$  and  $t$ , we have that  $r$  divides  $\gcd(\{g(bw + t, b) : w \in \mathbf{Z}\})$ . Conversely, let  $r_{b,t} = \gcd(\{g(bw + t, b) : w \in \mathbf{Z}\})$ . We have  $g(bw + t, b) = 0$  in  $(\mathbf{Z}/r_{b,t}\mathbf{Z})[w]$ . But  $f(w) = (b^n)^{-1} g(b(w - \frac{t}{b}) + t, b)$ , so  $f(w) = 0$  in  $(\mathbf{Z}/r_{b,t}\mathbf{Z})[w]$ . Therefore  $r_{b,t}$  divides  $r$  for each such  $b$  and  $t$ . It follows that the polynomials  $g(bw + t, b)$  have the same gcd as  $f$ .

**Lemma 4** If  $a$  is chosen such that  $z = \frac{ap+bw+t}{b}$  is an integer, then  $z$  is a root of the polynomial  $f \bmod p$ .

*Proof*  $b^n f(z) = b^n f(\frac{ap+bw+t}{b}) = b^n \sum_i c_i (\frac{ap+bw+t}{b})^i = \sum_i c_i (ap + bw + t)^i b^{n-i} \equiv \sum_i c_i (bw + t)^i b^{n-i} = g(bw + t, b) = r_f p \equiv 0 \pmod{p}$ . Since  $(b, p) = 1$ , the lemma is proven.

Having proven these lemmas, we know that  $\frac{z}{p}$  is close to  $\frac{a}{b}$ . Assuming the Bouniakowsky conjecture, we can let  $w \rightarrow \infty$  and obtain infinitely many primes  $p$  and a root  $z$  for each prime. As  $w \rightarrow \infty$ ,  $\frac{z}{p}$  is arbitrarily close to

$\frac{a}{b}$ , since  $n \geq 2$ . Since we showed in Lemma 1 that  $B_f$  is dense in  $(0, 1)$ , the theorem is now proved.

### Acknowledgements

Thanks to Professor Zhengyu Mao for suggesting the problem to me and for many helpful discussions.

### References

- [B] A. Brauer, Über die Verteilung der Potenzreste, Math Z. 35 (1932) 39-50.
- [DFI] W. Duke, J.B. Friedlander, H. Iwaniec, Equidistribution of roots of a quadratic congruence to prime moduli, Annals of Mathematics (2) 141 (1995), no. 2, 423-441.
- [H] C. Hooley, On the distribution of the roots of a polynomial congruences, Mathematika 11 (1964) 39-49
- [T] A. Toth, Roots of quadratic congruences, IMRN 2000, no. 14, 719-739