# Counterflow and paired superfluidity in one-dimensional Bose mixtures in optical lattices

Anzi Hu<sup>1</sup>, L. Mathey<sup>1</sup>, Ippei Danshita<sup>2,3</sup>, Eite Tiesinga<sup>1</sup>, Carl J. Williams<sup>1</sup> and Charles W. Clark<sup>1</sup>

"Joint Quantum Institute,

University of Maryland and National Institute of Standards and Technology,

Gaithersburg, MD 20899

"Department of Physics, Faculty of Science,

Tokyo University of Science,

Shinjuku-ku, Tokyo 162-8601, Japan

"Department of Physics,

Boston University, Boston, MA 02215

We study the quantum phases of mixtures of ultra-cold bosonic atoms held in an optical lattice that confines motion or hopping to one spatial dimension. The phases are found by using Tomonaga-Luttinger liquid theory as well as the numerical method of time evolving block decimation (TEBD). We consider a binary mixture with repulsive intra-species interactions, and either repulsive or attractive inter-species interaction. For a homogeneous system, we find paired- and counterflow-superfluid phases at different filling and hopping energies. We also predict parameter regions in which these types of superfluid order coexist with charge density wave order. We show that the Tomonaga-Luttinger liquid theory and TEBD qualitatively agree on the location of the phase boundary to superfluidity. We then describe how these phases are modified and can be detected when an additional harmonic trap is present. In particular, we show how experimentally measurable quantities, such as time-of-flight images and the structure factor, can be used to distinguish the quantum phases. Finally, we suggest applying a Feshbach ramp to detect the paired superfluid state, and a  $\pi/2$  pulse followed by Bragg spectroscopy to detect the counterflow superfluid phase.

#### I. INTRODUCTION

Bose-Einstein condensation [1] is a fascinating manybody phenomenon. It demonstrates the significance of quantum statistics at low temperature. Identical bosons can occupy the same single particle state and are in fact more likely to do so than classical particles. At a critical temperature, a gas of bosons undergoes a phase transition towards a state in which a macroscopic fraction of the particles occupy the lowest energy state, creating a condensate. Such a state was realized in ultra-cold atom systems in [2], demonstrating that the technology of cooling and manipulating atoms had reached a level of control with which novel states of matter could be generated and studied.

In the case of a Fermi gas, the Pauli exclusion principle prevents such a phenomenon to occur, because no single particle state can be more than singly occupied. However, the phenomenon of condensation can still occur in Fermi systems via a different mechanism: fermions can form pairs to create composite bosons. The bosonic particles then form a condensate of pairs. Conventional superconductors, for example, were understood as a condensate of electron pairs [3]. In ultra-cold atoms, fermionic condensates of this type were created in [4].

Interestingly, this mechanism of condensation of pairs is not limited to fermionic systems but can occur in bosonic systems as well. In fermionic systems, formation of Bosonic pairs necessarily occurs before condensation. In bosonic systems this mechanism can be favored ener-

getically, and will typically be in competition with single particle condensation.

In [5, 6], two types of composite bosons were predicted for a binary Bose mixture in a optical lattice: pairs and anti-pairs. For attractive mutual interactions, a bosonic mixture can form pairs of atoms which then form a paired superfluid (PSF) state, as is visualized in Fig. 1. For repulsive interactions, at special fillings, the atoms can form anti-pairs, which can be interpreted as pairs of one atom of one species and one hole of the other species. These anti-pairs can then generate a counterflow superfluid (CFSF) state, visualized in Fig. 2. Most of their simulations were performed for two dimensional systems.

Quantum phases of atoms in optical lattices have been experimentally studied. Following the prediction by Jaksch et al. in [7], the Mott insulator (MI) to superfluid (SF) transition was realized in Ref. [8] in a three dimensional lattice. In [9] this transition was achieved in 1D. More recently, Ref. [10] observed the two dimensional (2D) transition.

In one-dimensional gases quantum phases have quasilong range order (QLRO), rather than true long range order. QLRO of an operator O(x) is defined as follows: The correlation function  $R(x) = \langle O^{\dagger}(x)O(0) \rangle$  falls off algebraically as  $R(x) \sim |x|^{\alpha-2}$  as  $|x| \to \infty$  with  $\alpha > 0$ . Various order parameters O(x) will be defined in the text. In contrast in higher dimensional bosonic systems correlation functions can have true long range order, where correlation functions approach a finite value. Power-law scaling in a 1D optical lattice has been observed in [11].




Figure 1: Sketch of a condensate of pairs. Atoms of each species (red/green) pair together and form a paired superfluid (PSF) state.




Figure 2: Sketch of a condensate of anti-pairs. Here, atoms of one species are strongly anti-correlated with atoms of the other species, creating a counterflow superfluid (CFSF) state. These composite bosons can also be thought of as a pair of one atom of one species and one hole of the other species.

They observed the Tonks-Girardeau regime of strongly interacting bosons.

In this paper we consider a two-component Bose mixture held in an optical lattice that only allows atoms to hop in one spatial dimension. We ask the question of how the superfluid as well as other phases or orders can be realized. We assume that the two species of the mixture have the same filling  $\nu$ , restricted to the range  $0 < \nu < 1$ . The phase diagram of these mixtures is determined using Tomonaga-Luttinger liquid theory [13], which gives the universal phase diagram in terms of a few effective parameters. Based on the univerisal phase diagram, we generate the numerical phase diagram using the time-evolving block decimation (TEBD) method [14, 16, 17, 18]. With these two approaches we find that CFSF can exist for  $\nu = 1/2$  (half-filling) and repulsive interaction, whereas PSF can exist for  $\nu < 1$ and attractive interaction (see also [19]).

We also find that charge density wave (CDW) quasiorder can coexist with both PSF and CFSF, as well as single particle superfluidity (SF). The regimes in which CDW and SF quasi-order coexist constitute a quasisupersolid phase [20, 21]. Similarly, the regimes where CDW and PSF quasi-order coexist is a quasi-supersolid of pairs and in the case of CFSF, a quasi-supersolid of anti-pairs. Previous work has predicted coexistence of CDW and PSF for 1D Bose mixtures [20, 22] and bilayer 2D lattice bosons with long-range interactions [23], and that of CDW and CFSF for 1D Bose-Fermi mixtures [21, 24].

We then address the question whether PSF and CFSF can be realized and detected in experiment. To simulate the effect of a global trap, we numerically study a mixture confined by a harmonic trap and find that PSF and CFSF can indeed exist in such trapped systems. Their existence can be detected through various measurements. The PSF phase can be detected by using a Feshbach ramp, similar to what has been used in BEC-BCS experiments [4], which generates a quasi-condensate signal in the resulting molecules. The CFSF phase can be detected by applying a  $\pi/2$  pulse followed by Bragg spectroscopy. This generates a quasi-condensate signal in the structure factor. Time-of-flight expansion can also be used to show the absence of single particle superfluidity in PSF and CFSF. Measuring the structure factor via Bragg spectroscopy can be one way of detecting CDW order.

This paper is organized as follows: in Section II, we introduce the model that is used to describe the system; in Section III, we use Tomonaga-Luttinger liquid theory to derive the phase diagram. The numerical approach and results are discussed in Section IV. Specifically, phase diagrams of the homogeneous system are presented in Section IV A, and the realization and detection of PSF and CFSF are discussed in Section IV B. We conclude in Section V.

#### II. HAMILTONIAN

Ultra-cold bosonic atoms in optical lattices can be well described by Bose Hubbard models [7]. Here, we consider a mixture of two types of atoms confined to a one-dimensional lattice system. The Hamiltonian of such a system is given by

$$H = -t \sum_{a=1,2} \sum_{i=1}^{N-1} (b_{a,i}^{\dagger} b_{a,i+1} + h.c.) + U_{12} \sum_{i=1}^{N} n_{1,i} n_{2,i} + \frac{U}{2} \sum_{a=1,2} \sum_{i=1}^{N} n_{a,i} (n_{a,i} - 1).$$

$$(1)$$

We denote the different types of atoms with index a=1,2, and the lattice site with index i. We assume that the two species have equal particle density  $\nu \leq 1$ , the same intra-species interaction U>0 and hopping parameter t>0. The inter-species interaction is given by  $U_{12}$ . The operators  $b_{a,i}^{\dagger}$  and  $b_{a,i}$  are the creation and annihilation operators for atoms of type a and site i and  $n_{a,i}=b_{a,i}^{\dagger}b_{a,i}$  are the number operators.

# III. TOMONAGA-LUTTINGER LIQUID APPROACH

The universal behavior of this system can be found within a Tomonaga-Luttinger liquid description [13]. In this paper, we are interested in the phase diagram of the

system at various densities and interactions. First, we switch to a continuum description,  $b_{a,i} \to b_a(x)$ , and express the operators  $b_a(x)$  through a bosonization identity, according to Haldane [25, 26]:

$$b_a(x) = [n + \Pi_a(x)]^{1/2} \sum_{m} e^{2mi\Theta_a(x)} e^{i\phi_a(x)}, \qquad (2)$$

where the real-space density of each species is  $n = \nu/a_L$  and  $a_L$  is the lattice constant. The lattice sites are at positions  $x = ia_L$ . This expression is a phase-density representation of the Bose operators, in which the square root of the density operator has been written in an intricate way. The fields  $\Pi_{1,2}(x)$  describe the small amplitude and the long wave length density fluctuations. The fields  $\Theta_{1,2}(x)$  are given by  $\Theta_{1,2}(x) = \pi nx + \theta_{1,2}(x)$ , where  $\theta_{1,2}(x) = \pi \int^x dy \Pi_{1,2}(y)$ . The fields  $\phi_{1,2}(x)$  describe the phase, and are conjugate to the density fluctuations  $\Pi_{1,2}(x)$ .

The contact interactions between the densities in 1 written in Haldane's representation generate an infinite series of terms that contain  $\exp(2m_1i(\pi nx + \theta_1) + 2m_2i(\pi nx + \theta_2))$ , where  $m_1$  and  $m_2$  are some integers. A term of this form can only drive a phase transition, if the oscillatory part  $2\pi m_1 nx + 2\pi m_2 nx$  vanishes for all lattice sites. This leads to the requirement  $m_1\nu + m_2\nu = m_3$ , with  $m_3$  another integer [22]. As a further requirement, small integers  $m_1$  and  $m_2$  are necessary, because the scaling dimension of the term scales quadratically in  $m_1$  and  $m_2$ .

For the range  $0 < \nu \le 1$ , we find that there are three different cases: unit-filling  $(\nu=1)$ , half-filling  $(\nu=1/2)$ , and non-commensurate filling  $(\nu\ne 1)$  and  $\nu\ne 1/2$ . It can be checked, using renormalization group arguments as below, that higher forms of commensurability do not generate new phases, but that either phase separation or collapse is reached first. Our numerical findings are consistent with this.

Non-commensurate filling. The action of the system, assuming a short-range spatial cut-off  $r_0$ , at non-commensurate filling is given by [13, 20, 26]:

$$S = \int d^{2}r \left[ \sum_{j=1,2} \frac{1}{2\pi K} \left( \left( \partial_{v\tau} \theta_{j} \right)^{2} + \left( \partial_{x} \theta_{j} \right)^{2} \right) + \frac{U_{12} a_{L}}{\pi^{2} v \hbar} \partial_{x} \theta_{1} \partial_{x} \theta_{2} + \frac{2g_{\sigma}}{(2\pi r_{0})^{2}} \cos(2\theta_{1} - 2\theta_{2}) \right] (3)$$

The first line of the action is characterized by a Luttinger parameter K and a velocity v, contained in  $\mathbf{r} = (v\tau, x)$ . This part of the action, without the coupling between the two fields  $\theta_a(x)$ , generates a linear dispersion  $\omega = v|k|$ , where . v should therefore be interpreted as the phonon velocity. The Luttinger parameter K is a measure of the intra-species interaction U. We will be interested in the regime  $U \gtrsim t$ , in which we have approximately [27]

$$K \approx 1 + \frac{8t}{U} \frac{\sin \pi \nu}{\pi}.$$
 (4)

The velocity v can also be related to the parameters of the underlying Hubbard model by

$$v \approx v_F (1 - 8t\nu \cos \pi \nu / U) \tag{5}$$

where  $v_F$  is the 'Fermi velocity' of an identical system of fermions,  $v_F = 2(a_L t/\hbar) \sin \pi \nu$ , and  $k_F$  is the 'Fermi wave vector',  $k_F = \pi n$ . Here,  $\hbar$  is the Planck constant.

The two fields  $\theta_a(x)$  are coupled by the inter-species interaction. The interaction term  $U_{12}n_1n_2$  in the underlying Hubbard model generates both the term containing  $\partial_x\theta_1\partial_x\theta_2$ , as well as the backscattering term [13, 22] containing  $\cos(2\theta_1-2\theta_2)$ . The action S is only well-defined with a short-range cut-off  $r_0$ . It is proportional to 1/n. At this scale,  $g_\sigma$  is approximately given by

$$g_{\sigma} = U_{12}a_L/(v\hbar). \tag{6}$$

We diagonalize the quadratic part of the action by switching to the symmetric and antisymmetric combinations  $\theta_{S/A} = \frac{1}{\sqrt{2}}(\theta_1 \pm \theta_2)$ . For the two sectors we find

$$K_{S/A} = (1/K^2 \pm U_{12}a_L/(v\hbar\pi K))^{-1/2}$$
 (7)

as effective Luttinger parameters. To lowest order in  $U_{12}$  this gives  $K_{S/A} \approx K \mp U_{12} a_L K^2/(2\pi v \hbar)$ . The effective velocities are  $v_{S/A} = v \sqrt{1 \pm U_{12} a_L K/(\pi v \hbar)}$ . Collapse (phase separation) of the superfluid phase is when  $v_{S/A}$  is imaginary. We note that  $K_S$  diverges when collapse (CL) is approached, and that  $K_A$  diverges as the system approaches phase separation (PS).

The anti-symmetric sector contains the nonlinear backscattering term  $\cos(2\sqrt{2}\theta_A)$ . To study its effect, we use an RG approach. We renormalize the short-range cut-off  $r_0$  to a slightly larger value, and correct for it at one-loop order. The resulting flow equations are given by [13]:

$$\frac{dg_{\sigma}}{dl} = (2 - 2K_A)g_{\sigma} \tag{8}$$

$$\frac{dK_A}{dl} = -\frac{g_\sigma^2}{2\pi^2} K_A^3 \tag{9}$$

The flow parameter l is given by

$$l = \log_e \left(\frac{r_0'}{r_0}\right),\tag{10}$$

where  $r_0'$  is the new cut-off that has been created in the RG process.

The flow equations 8 and 9 have two qualitatively different fixed points: Either  $g_{\sigma}$  diverges, which in turn renormalizes  $K_A$  to zero, or  $g_{\sigma}$  is renormalized to zero for finite  $K_A = K_A^*$ . In the latter case, the action S is quadratic in  $\theta_S$  and  $\theta_A$ . For the parameter  $K_S$ , we use the bare value given in Eq. 7.

As mentioned in the introduction, we can determine the phase diagram by studying the long-range scaling behavior of correlation functions,  $\langle O^{\dagger}(x)O(y)\rangle$ , of various order parameters O(x). In particular, the singleparticle superfluid order parameter is  $O_{SF} = b_a(x)$  with




Figure 3: Phase diagram of a bosonic mixture at non-unit and non-half-filling. For attractive interactions  $U_{12}$  and K < 2 the system can form a paired superfluid state, in the regime labeled PSF and PSF(CDW). This phase can coexist with CDW order for weaker interactions. For large repulsive (attractive) interactions  $U_{12}$  the system phase separates (PS) (collapses (CL)). For the remaining regime the system shows single particle superfluidity (SF). This can coexist with CDW order, resulting in a quasi-supersolid (SS) regime.

a=1,2. The CDW order is related to the  $2k_F$  wavevector component of the density operator,  $O_{CDW} = n_a$ . PSF is described by  $O_{PSF} = b_1(x)b_2(x)$ , and CFSF by  $O_{CFSF} = b_1^{\dagger}(x)b_2(x)$ . In the homogeneous system, it suffices to study

$$G(x) = \langle b_a^{\dagger}(x)b_a(0)\rangle, a = 1, 2 \tag{11}$$

$$R_{n,a}(x) = \langle n_a(x)n_a(0)\rangle, a = 1, 2$$
 (12)

$$R_S(x) = \langle b_1^{\dagger}(x)b_2^{\dagger}(x)b_1(0)b_2(0)\rangle$$
 (13)

$$R_A(x) = \langle b_1^{\dagger}(x)b_2(x)b_1(0)b_2^{\dagger}(0)\rangle.$$
 (14)

We find that away from collapse (CL) and phase separation (PS), the correlation functions scale either algebraically or exponentially. For algebraic scaling, we have

$$G(x) \sim |x|^{\alpha_{SF}-2},$$
  
 $\alpha_{SF} = 2 - 1/(4K_S) - 1/(4K_A) \quad (15)$   
 $R_{n,a}(x) \sim \cos(2k_F x)|x|^{\alpha_{CDW}-2},$ 

$$\alpha_{CDW} = 2 - K_S - K_A \tag{16}$$

$$R_S(x) \sim |x|^{\alpha_{PSF}-2}, \alpha_{PSF} = 2 - 1/K_S$$
 (17)

$$R_A(x) \sim |x|^{\alpha_{CFSF}-2}, \alpha_{CFSF} = 2 - 1/K_A.$$
 (18)

where the scaling exponents  $\alpha_O$  are determined by  $K_S$ and  $K_A$  after the RG flow. For the case that  $g_{\sigma}$  diverges in Eqs. 8 and 9 and  $K_A$  is undefined, these expressions can still be used. We set  $K_A$  to zero, and find that  $\alpha_{CDW}$ and  $\alpha_{PSF}$  are well defined. Hence  $R_{n,a}$  and  $R_S$  still show algebraic scaling. On the other hand,  $\alpha_{CFSF}$  and  $\alpha_{SF}$ become  $-\infty$  and G and  $R_A$  scale exponentially.

We can identify regimes where different scaling exponents are positive based on the relationship between the scaling exponents and  $K_{S/A}$  after the flow. This determines the different quasi-long range orders that are present. The resulting phase diagram is shown in Fig. 3, as a function K and  $U_{12}a_L/(v\hbar)$ , as appearing in the action in Eq. 3. These two parameters determine the initial values of the flow equations through equations 7 and 6.

We can estimate the phase boundary between PSF and SF. For small  $U_{12}a_L/(v\hbar)$  this boundary is near the point  $K_A = 1$  and  $g_{\sigma} = 0$ . For that limit, Eq. (9) can be linearized to

$$\frac{dK_A}{dl} = -\frac{g_\sigma^2}{2\pi^2} \tag{19}$$

and the expression  $A = \pi^2 (1 - K_A)^2 - g_\sigma^2/4$  becomes an invariant of the flow. From the properties of the RG flow of a Berezinskii-Kosterlitz-Thouless transition (see e.g. [13, 29]), the phase boundary is given by A=0 and  $g_{\sigma} < 0$ . Using the expressions of  $K_A$  and v in terms of the Hubbard parameters, we estimate the critical interaction  $U_{12}$  for PSF to occur at

$$\frac{U_{12}}{U}\bigg|_{2} = -32\frac{t^2}{U^2}\sin^2(\pi\nu). \tag{20}$$

The phase boundary between supersolid (SS) and SF has been derived in Ref. [20].

Half-filling. In the case of half-filling, another nonlinear term has to be introduced in the action

$$S_{uk} = \frac{2g_{uk}}{(2\pi r_0)^2} \int d^2r \cos(2\theta_1 + 2\theta_2).$$
 (21)

This term describes Umklapp scattering. At the initial cut-off  $r_0 \sim 1/n$ ,  $g_{uk}$  is approximately given by  $U_{12}a_L/v$ . In addition to the RG flow in the antisymmetric sector we now also have

$$\frac{dg_{uk}}{dl} = (2 - 2K_S)g_{uk} \tag{22}$$

$$\frac{dK_S}{dl} = -\frac{g_{uk}^2}{2\pi^2} K_S^3 \tag{23}$$

in the symmetric sector. Proceeding along the same lines as for the non-commensurate case, we find the phase diagram shown in Fig. 4.

We estimate the SF-CFSF phase boundary in the same way as the PSF-SF boundary. We find

$$\frac{U_{12}}{U}\bigg|_{c} = 32\frac{t^2}{U^2}\sin^2(\pi\nu). \tag{24}$$

*Unit-filling*. At unit-filling we have to introduce a term of the form

$$S_1 = \frac{2g_1}{(2\pi r_0)^2} \int d^2r \left(\cos(2\theta_1) + \cos(2\theta_2)\right).$$
 (25)

The resulting RG flow for this system is given by

$$\frac{dg_{uk}}{dl} = (2 - 2K_S)g_{uk} + \alpha_3 \frac{g_1^2(K_A - K_S)}{2\pi}$$
 (26)




Figure 4: Phase diagram of a bosonic mixture at half-filling. In addition to the phases that appear in Fig. 3, the system now develops a counterflow superfluid (CFSF) phase, which can coexist with CDW order.

$$\frac{dg_{\sigma}}{dl} = (2 - 2K_A)g_{\sigma} + \alpha_3 \frac{g_1^2(K_S - K_A)}{2\pi}$$

$$\frac{dg_1}{dl} = (2 - \frac{K_S + K_A}{2} + \alpha_3 \frac{g_{uk}K_S + g_{\sigma}K_A}{\pi})g_1(28)$$

$$\frac{dK_A}{dl} = -\frac{g_\sigma^2}{2\pi^2} K_A^3 - \frac{g_1^2}{16\pi^2} (K_S + K_A) K_A^2$$
 (29)

$$\frac{dK_S}{dl} = -\frac{g_{uk}^2}{2\pi^2} K_S^3 - \frac{g_1^2}{16\pi^2} (K_S + K_A) K_S^2 \tag{30}$$

where  $\alpha_3$  is some non-universal parameter [28]. The behavior of this set of equations depends strongly on the initial value of  $g_1$ . For small values of  $g_1$ , four phases can be stable: Single-particle superfluidity, CFSF, PSF and a Mott phase. For large values only single-particle SF and MI are stable. We determine with our numerical approach, that the Hubbard model falls into the second category, i.e. there is only a single-particle SF and a Mott state at unit-filling.

Having established the universal behavior of the system from Tomonaga-Luttinger liquid theory, we now want to connect the phase diagram with the parameters in the Hubbard model. The expressions 4 and 5, which relate the Luttinger parameter K and the velocity v to microscopic parameters of the Hubbard model, are only approximate, no full analytic expression is known. In addition, only some phase boundaries are predicted reliably, because we use perturbative RG in the  $g_{\sigma}$ . We expect that the analytic calculation only predicts the general structure of the phase diagram, as well as the decay behavior of the correlation functions. To obtain the phase diagram in terms of the parameters in the Hubbard model, we need to use numerical methods. The next section describes the numerical determination of the phase diagram.

	$R_S(x)$	$R_A(x)$	G(x)
MI	Exp.	Exp.	Exp.
SF	Alg.	Alg.	Alg.
CFSF	Exp.	Alg.	Exp.
PSF	Alg.	Exp.	Exp.
$\mathrm{CL/PS}$	$R_S(x)$ , $R_A(x)$ undefined		

Table I: Definitions of Mott insulator (MI), superfluid (SF), counterflow superfluid (CFSF) and paired superfluid (PSF) orders in terms of the long-range behavior of the correlation functions  $R_S(x)$ ,  $R_A(x)$ , and G(x). Each of these can either show algebraic (Alg.) or exponential (Exp.) decay when the system is away from collapse (CL) or phase separation (PS). From the Tomonaga-Luttinger liquid theory,  $R_S(x)$  and  $R_A(x)$  approach a constant (or  $K_{S/A}$  diverges) when the system approaches CL/PS regime. For the numerical calculation in the CL/PS regimes, the behavior of the correlation functions is inconclusive and we assign the phase from additional observables as discussed in the text.

## IV. NUMERICAL APPROACH

We use the time-evolving-block-decimation (TEBD) method [14] to study our discrete one-dimensional two-species Hubbard Hamiltonian. With this method, explained in Appendix A, we obtain an approximate ground state solution. We consider N lattice sites with hard-wall boundary conditions and express the Hubbard parameters in units of the intra-species interaction U. The number of sites N is equal to 80, unless otherwise noted. In our numerical analysis, we limit the particle number on each site and each species to two for filling  $\nu \leqslant 0.8$  and four otherwise. Once we obtain the ground state, we calculate the energy, density distributions, correlation functions, and the structure factor to identify the quasi-long range order and other properties of the ground state.

For example, to determine whether a SF, PSF, or CFSF is present, we study the decay behavior of the correlation functions, G(x),  $R_A(x)$ , and  $R_S(x)$ , defined in Eqs. 11, 14, and 13, respectively. If both  $R_A$  and  $R_S$  decay algebraically, the system is in a single-particle superfluid (SF) state. If both are exponential, the system is in a Mott insulator(MI) state. If  $R_S$  or  $R_A$  decays algebraically, the system is in the PSF or CFSF state, respectively. These relationships are summarized in Table I.

In Fig. 5(a) and (b), we show the decay behavior of the correlation functions in the PSF and CFSF phase, respectively. As the Hamiltonian is discrete, the correlation functions are calculated as discrete functions:  $G(i,j) = \langle b_{a,i}^{\dagger} b_{a,j} \rangle$ ,  $R_S(i,j) = \langle b_{1,i}^{\dagger} b_{2,i}^{\dagger} b_{1,j} b_{2,j} \rangle$ , and  $R_A(i,j) = \langle b_{1,i}^{\dagger} b_{2,i} b_{1,j} b_{2,j}^{\dagger} \rangle$ . For the PSF phase,  $R_A(i,j)$  decays exponentially, while  $R_S$  decays algebraically. It is also worthwhile to notice that the single-particle Green's function decays exponentially, implying the absence of single-particle superfluidity. For the CFSF phase,  $R_A$  decays algebraically while  $R_S$  decays exponentially. Single-




Figure 5: The correlation functions  $R_A$ ,  $R_S$ , and G on a logarithmic scale as a function of distance |i - j|. The index i is 40, the center of the 80 lattice sites. The squares are the numerical data. The blue lines are exponential fits to the data and red dotted lines are algebraic fits. Note that the scale of the vertical axis of the graphs differs by orders of magnitude. In (a), we show an example for the paired superfluid phase at  $\nu = 0.3$ , t = 0.02U, and  $U_{12} = -0.16U$ .  $R_A$  decays exponentially and  $R_S$  decays algebraically. The single-particle correlation function decays exponentially, implying the absence of single-particle superfluidity. In (b), we show an example for the counterflow superfluid phase at  $\nu = 0.5$ , t = 0.02U, and  $U_{12} = 0.2U$ . The anti-pair correlation function  $R_A$  decays algebraically, while the pair correlation function decays exponentially. Single-particle superfluidity is again absent. The algebraic fits deviate from the data around  $|i-j| \approx 40$ , due to the boundary conditions of our numerical calculations.

particle superfluidity is again absent.

Behavior of  $K_S$  and  $K_A$ : We study the decay behavior of  $R_S$  and  $R_A$  in more detail. Using the fit function,  $c \cdot |i-j|^{\alpha-2}$ , where c and  $\alpha$  are the fitting parameters, we obtain the power-law exponent  $\alpha$  and, hence, the Luttinger parameters  $K_S$  and  $K_A$  based on Eqs. 17 and 18. In Fig. 6(a), we show these  $K_S$  and  $K_A$  as a function of  $U_{12}$ , for non-commensurate filling. A Luttinger parameter is formally set to zero when its correlation function decays exponentially.

For  $U_{12} < -0.06U$ ,  $R_A$  decays exponentially, while for  $U_{12} > -0.06U$ ,  $R_A$  decays algebraically, and  $K_A$  increases as  $U_{12}$  increases. The system undergoes a PSF to SF transition at  $U_{12} = -0.06U$ . On the other hand,  $K_S$  decreases monotonically for  $U_{12} > -0.6U$ . For  $U_{12} < -0.6U$  the numerics failed to converge to a homogeneous

state. This indicates that the system collapses, and we therefore cannot extract a Luttinger liquid parameter. We can observe charge density wave (CDW) order for a range of  $U_{12}/U$  in Fig. 6. According to Eq. 16, this order exists when  $K_S+K_A<2$ . In fact, it co-exists with the SF, PSF or CFSF order. At half-filling,  $K_S$  will go to zero at a critical, positive value of  $U_{12}$ . This indicates the transition from the SF to CFSF phase.

Finite-size effect: The behavior of  $K_{A/S}$  stated above is affected by the size of the system. Finite size effects can 'smooth out' a sudden change in  $K_{A/S}$  at the phase transition. This effect can be estimated from the RG flow calculation by integrating Eqs. 8 and 9 out to a finite value l rather than to infinity. In Fig. 6(b), we show an example of a finite-l RG calculation in the vicinity of the PSF-to-SF transition. We see that as l increases,  $K_A$ dramatically changes for the attractive  $U_{12}$ . In the limit of  $l \to \infty$ ,  $K_A$  becomes discontinuous and 'jumps' from 0 to 1 at  $U_{12} \approx -0.01U$ . This is where the PSF-to-SF transition occurs. This transition is a Berezinskii-Kosterlitz-Thouless transition [13, 29]. In order to compare the RG result with our TEBD result, we associate the system size N with the flow parameter l, based on the relation in Eq. 10. The cut-off  $r_0$  is the lattice constant  $a_L$  and  $r'_0 = Na_L$ . For N = 80 we have l = 4.4 and we find that the RG and TEBD are in good agreement. The regime between  $U_{12}/U \approx -0.06$  and -0.01 is a cross-over regime due to the finite size of the system.

Collapse and phase separation: For large  $|U_{12}|$ , the system approaches collapse or phase separation. According to Tomonaga-Luttinger liquid theory,  $K_S \to \infty$  as the system approaches collapse and  $K_A \to \infty$  as the system approaches phase separation. As seen in Fig. 6, we indeed find such a tendency in our TEBD calculations. For  $U_{12} > 0.8U$  (not shown),  $K_A$  increases rapidly to values around 10, indicating a possible phase separation. For  $U_{12} < -0.6U$ , due to the slow decay of the correlation function  $R_S$  and the finite-size of our system, we are unable to extract an accurate  $K_S$  from the numerical result. On the other hand, we observe a peaked density distribution for  $U_{12} < -0.6U$ , indicating a collapse. In the phase separation regime, G(x) has algebraic decay except for  $\nu = 0.5$  or 1, where it has exponential decay. An algebraic decay implies two spatially-separated single-species superfluids while the exponential decay implies two spatially-separated Mott insulators. [30].

# A. Phase diagram

We study the phase diagram as a function of filling  $\nu$  and parameters of the Hubbard Hamiltonian. Assuming a positive U, the system can be fully characterized in terms of  $\nu$ , t/U, and  $U_{12}/U$ . Our results are shown in Fig. 7 for a fixed hopping parameter and in Fig. 8 for half filling.




Figure 6: (a)  $K_S$  and  $K_A$  as a function of  $U_{12}$  as extracted from the fit of the correlation functions,  $R_S$  and  $R_A$ . The filling  $\nu$  is 0.7 and t/U is 0.02. Around  $U_{12}/U \approx -0.06$ , the anti-pair correlation function changes from algebraic to exponential decay. This corresponds to the transition from the PSF to SF phase. When  $R_A$  decays exponentially,  $K_A$  is formally set to zero. For  $K_a + K_s \lesssim 2$ , the system has CDW order. Error bars are one standard deviation uncertainties obtained from the power-law fit to the numerical data. (b) A comparison of  $K_A$  obtained from our RG and TEBD calculations. The red square connected by lines are the TEBD results while all other lines are determined from the RG flow with flow parameter l = 3, 4, 7, and 10, where l is defined in Eq. 10. The error bars are as in panel (a). The PSF-to-SF transition obtained from TEBD is around  $U_{12}/U \approx -0.06$ , while the RG calculation shows that for l = 10, the transition occurs near  $U_{12}/U \approx -0.01$ . We interpret the regime between  $U_{12}/U \approx -0.06$  and -0.01 the cross-over region.

0.3

0.4

# 1. Phase diagram at a fixed hopping parameter

In Fig. 7 we show the phase diagram for filling fractions between 0 and 1 and the interaction  $U_{12}/U$  between -1.1 and 1.1. The symbols correspond to numerical data points at which the phases have been characterized. Different markers represent the different orders. The orders are determined from the decay behavior of the three correlation functions  $R_A$ ,  $R_S$ , and G.

For weak attractive inter-species interaction,  $-0.06 < U_{12}/U < 0$ , the system is in a SF state. As  $U_{12}$  grows more attractive, paired superfluidity (PSF) occurs. The critical  $U_{12}$  is largest,  $\sim -0.08U$ , at half-filling and grad-

ually decreases away from half-filling. This phase boundary differs from that predicted by our RG calculation (Eq. 20), plotted as the dotted line in Fig. 7. This discrepancy is the result of the finite-size effect discussed in Fig. 6(b). In the SF to PSF cross-over regime, charge density wave (CDW) order can coexist. According to the phase diagram Fig. 3, for attractive interaction, CDW order can co-exist only with PSF order. In our numerical work, we observed the CDW order slightly outside the numerical phase boundary of PSF but within the RG phase boundary of PSF. The sub-regime where CDW and PSF coexist ends when  $U_{12}/U \lesssim -0.4$ . When the inter-species attraction is comparable to the intra-species repulsion,  $U_{12} \lesssim -U$ , the system collapses (CL) and no long-range order is present.

For repulsive inter-species interaction and  $U_{12} < U$ , the system is in a SF state for all non-commensurate fillings. Within the SF regime, there is a smaller parameter region where CDW order coexist with the SF order. This subregime is a quasi-supersolid regime. The boundary between a normal superfluid and a quasi-supersolid is estimated by RG calculation in Ref. [20]. At half-filling, counterflow superfluidity (CFSF) occurs when  $0.08 \lesssim U_{12}/U \lesssim 1$ . Within the CFSF regime, the CDW order can coexist, forming a quasi-supersolid of anti-pairs. It also worthwhile to point out that at half-filling, CDW order only exists within the PSF and CFSF regimes.

At unit filling, our numerical results do not show evidence of PSF or CFSF for any  $U_{12}$ . We find a Mott insulator (MI) state for  $|U_{12}| < U$ .

### 2. Phase diagram at half-filling

In Fig. 8, we show the phase diagram at half filling as a function of  $U_{12}/U$  and t/U. From this diagram, we find that the border between PSF and SF and the border between PSF and CL approach each other as t increases. Similarly, the border between the CFSF and SF and the border between CFSF and PS approach each other. In fact, the PSF and CFSF phases end around  $t \sim 0.16U$ . Within the PSF and CFSF regimes, CDW order can co-exist. In the phase separated regime, the separated single-species ensembles form two individual Mott insulating states for  $t \lesssim 0.14U$  and two individual SF states for t > 0.14U.

We can compare this phase diagram with the half-filling phase diagram in Fig. 4 obtained from Tomonaga-Luttinger liquid theory. Especially, we can compare the location of the phase boundary between SF and PSF(CFSF). To do so, we plot the RG phase boundaries, described by Eqs. 20 and 24, onto our phase diagram. The area near the two boundaries is interpreted as the cross-over regime where finite-size effects modify the phase boundary.




Figure 7: Phase diagram for a homogeneous system with 80 sites and the hopping parameter  $t=0.02\mathrm{U}$  as a function of filling  $\nu$  and inter-species interaction  $U_{12}/U$ . The horizontal axis shows three disconnected regions in  $U_{12}/U$ . The solid lines are the estimated phase boundaries based on the TEBD results and the dotted line is the PSF-to-SF phase boundary predicted by our RG calculation (see Eq. 20). For attractive interaction  $U_{12} \lesssim -0.06U$ , the system forms a paired-superfluid (PSF). The state collapses(CL) for  $U_{12} \lesssim -0.7U$ . For  $U_{12} \gtrsim -0.06$  and  $U_{12} \lesssim U$  the system shows single-particle superfluidity (SF). The system phase-separates (PS) for  $U_{12} \gtrsim 1$  and forms two single-particle superfluids (SF). Open circles are the points where  $K_S + K_A < 2$  and charge density wave (CDW) order coexists with a superfluid phase (SF,PSF, or CFSF). At half and unit filling there exist special phases. For repulsive interaction  $U_{12} \gtrsim 0.08U$  and half-filling, the system forms a counterflow superfluid (CFSF). For unit filling, we find a Mott-Insulator (MI) phase for interactions  $|U_{12}| \lesssim U$ . Finally, in the PS region at half- and unit-filling, the system forms two individual MI states.

#### B. Realization and detection

Having established the phase diagram for the homogeneous system, we now discuss how to realize and detect the PSF and CFSF phases. First, we need to modify the Hubbard Hamiltonian in Eq. 1 because in any ultra-cold atom experiment an additional trapping potential is present. We add a harmonic potential,  $\Omega(j-j_c)^2(n_{1,j}+n_{2,j})$ , where j is the site index and  $j_c$  is the index at the center of the system. The TEBD method is used to find the ground state. We consider a system of 80 lattice sites and adjust the total number of particles and the trap frequency so that the number of particles is negligible at the edge of the lattice.

We again determine the orders of the system by studying the correlation functions in Table I. We find that, in spite of the presence of the trap, the correlation functions still show exponential or algebraic scaling away from the edge of the lattice. In fact, a correlation function can have different decay behavior in different parts of the trap. We also find that SF, PSF, and CFSF still exist. The remainder of this article focusses on experimental signatures that distinguish between these orders by cal-

culating the density distibution, the time-of-flight image after an expansion, or the structure factor for Bragg spectroscopy.

Density distribution: We find that in a trapped system PSF and CFSF can only exist when the density distribution satisfies certain conditions. For PSF, the density of each species at the center of the trap,  $n_{\rm center}$ , must be less than one atom per site or equivalently per lattice constant  $a_L$ . (The density is largest at the center.) For CFSF,  $n_{\rm center}$  must satisfy  $n_{\rm center}a_L=1/2$ . Once such conditions are satisfied, the critical value of  $U_{12}$  for PSF and CFSF is close to the one for a homogeneous system (See Figs. 7 and 8).

In Fig. 9(a) we show density distributions for three attractive interactions  $U_{12}$  and a hopping parameter equal to the one used for Fig. 7. For all attractive interactions, the density distributions of each species are the same. For more attractive inter-species interaction, the density distribution concentrates near the center of the trap. There is no discontinuous change in the density distribution when the system goes from SF to PSF.

In Fig. 9(b) we show the density distribution for  $U_{12} = 0.2U$ . In this case in the center of the trap, where the




Figure 8: Phase diagram at half-filling as a function of  $U_{12}/U$  and t/U. The solid lines are estimated phase boundaries from the TEBD calculation and the dotted lines are the phase boundaries predicted by the RG calculation (see Eqs. 20 and 24). For large repulsive interaction, the system phase separates (PS) and for large attractive interaction, the system collapses (CL). For moderate interactions and for  $t/U \lesssim 0.2$ , the system shows paired superfluidity (PSF) on the attractive side and counterflow superfluidity on the repulsive side. Both PSF and CFSF can coexist with charge density wave (CDW) order when  $t \lesssim 0.1U$ .

density distribution is constant or has a "plateau", the system is in a CFSF state. The "plateau" is at half-filling consistent with predictions from a local density approximation and noting that in Fig. 7 CFSF only occurs at  $\nu=1/2$ . Towards the edge, where the density is decreasing sharply, it is in a SF state. The plateau implies that the system is incompressible in the center.

Time of flight measurement: A widely used measurement technique in the field of ultra-cold atoms is measuring the density of atoms after a time-of-flight (TOF) expansion. The 1D optical lattice potential and the harmonic trap are abruptly turned off at time T=0 and the atoms expand freely afterwards. We calculate the density at time T, according to

$$n_a(x,T) = \langle c_a^{\dagger}(x,T)c_a(x,T)\rangle$$
 (31)

with a = 1, 2. The operators  $c_a(x, T)$  are related to the lattice operator  $b_{a,j}$  according to

$$b_a(x,T) = \sum_{j=1}^{N} w(x - r_j, T) b_{a,j},$$
 (32)

where  $w(x,T)=\sqrt{d/\sqrt{2\pi}\Delta(T)^2\exp(-x^2/(4\Delta(T)^2))}$  describes the free expansion from the initial Gaussian wavefunction of an atom in a lattice site and  $\Delta(T)^2=d^2+iT\hbar/(2m)$ . The parameter d is the width of the initial Gaussian state and m is the atomic mass. The

density distribution  $n_a(x,T)$  is then given by

$$n_a(x,T) = \sum_{j_1,j_2=1}^{N} w^*(x - r_{j_1}, T)w(x - r_{j_2}, T)G(j_1, j_2),$$

where  $G(j_1, j_2)$  is the single-particle Green's function. In Fig. 10 we show examples of TOF expansions of PSF, CFSF, and SF order. For the SF phase, we find a strongly peaked interference pattern, reflecting the single-particle quasi-long range order. For both PSF and CFSF phases, the TOF density shows a broad Lorentzian distribution, which is due to the exponential decay of the single-particle Green's function.

Feshbach ramp: In order to detect the superfluidity of pairs, we consider applying a Feshbach ramp to pairwise project the atoms onto molecules formed by one atom from each species, which is similar to detection of fermionic pairs in the BCS regime [4]. In those experiments, a fast ramp across a Feshbach resonance was used, followed by a time-of-flight expansion. The density distribution of the molecules showed the superfluidity of fermionic pairs. We propose a similar detection for bosonic pairs in PSF.

To give a simple estimate of a TOF image after a Feshbach ramp, we imagine that bosons of different species on the same lattice site are converted into molecules. This leads to the replacement  $b_{1,j}b_{2,j} \to M_j$ , where  $M_j$  is the molecule annihilation operator. A TOF density of the




Figure 9: Density distribution of a trapped system for t=0.02U. (a) Attractive interaction  $U_{12}$ . The trap frequency is  $\Omega=1\times 10^{-5}U$  and the number of atoms is 20 for each species. For attractive interactions, the density distributions of the two species are identical. For  $U_{12}=-0.01U$  (curve I) the system is superfluid. For  $U_{12}=-0.11U$  (curve II) and  $U_{12}=-0.21U$  (curve III), the system is in the paired superfluid (PSF) state. As  $U_{12}$  becomes more negative the distribution gradually shrinks in size. (b) Repulsive interaction  $U_{12}=0.2U$  with  $\Omega=8\times 10^{-5}U$  and 30 atoms of each species. The red and green curves correspond to the species, respectively. The density distribution has a 'plateau' with half-filling in the center of the trap. The system is in a counter-flow superfluid (CFSF) state. The two species have weak interlocked density modulations around half filling.




Figure 10: Density distribution after a time-of-flight expansion. We assume <sup>87</sup>Rb atoms and use an expansion time of 0.03s. The hopping energy is t=0.02U. Panel (a): For attractive interaction  $U_{12}$ , we show the TOF expansion of a SF state at  $U_{12}=-0.01U$  (red line) and of a PSF state at  $U_{12}=-0.21U$  (green line). The two curves correspond to the expansion of the densities shown as curve I and III in Fig. 9(a) The trap frequency is  $\Omega=1\times 10^{-5}U$ . Panel (b): For repulsive interaction, we show a TOF expansion of a SF state at  $U_{12}=0.01U$  and of a CFSF state at  $U_{12}=0.21U$ . The trap frequency is  $\Omega=8\times 10^{-5}U$ .

molecules at position x and time T is given by

$$n_M(x,T) = \sum_{j_1,j_2=1}^{N} w^*(x - r_{j_1}, T)w(x - r_{j_2}, T)R_s(j_1, j_2).$$
(33)

In the expanding wave function w(x,T), the mass m is replaced by the mass of the molecule. We assume the

same initial width d. In a more realistic estimate, the conversion efficiency to molecules would not be 100%, but approximately given by the square of the overlap of the molecular wave function and the single-atom wave functions. This leads to a reduced signal. The spatial dependence, however, remains the same. In Fig. 11, we see an example of the density of molecules after TOF and, for comparison, the atomic density after TOF for the PSF state. The strongly peaked molecular distribution indicates the quasi-condensate of the bosonic pairs. The single-atom density is a broad Lorentzian distribution, indicating the absence of single-particle SF.

Bragg spectroscopy: To detect the presence of CDW order, one can use Brag spectroscopy [31, 32]. The quantity that is measured in those experiments is either the dynamic or static structure factor. Here we calculate the static structure factor  $S_a(k)$  for species a=1,2. It is defined as

$$S_a(k) = \frac{1}{N} \sum_{j_1, j_2} e^{-ika_L(j_1 - j_2)} (\langle n_a(j_1) n_a(j_2) \rangle - \langle n_a(j_1) \rangle \langle n_a(j_2) \rangle). \tag{34}$$

For wavevectors k near twice the "Fermi wavevector"  $k_F$ , the structure factor  $S(k) \sim ||k| - 2k_F|^{1-\alpha_{CDW}}$  with  $\alpha_{CDW} = 2 - K_S - K_A$  [13]. In our system,  $K_S + K_A$  is always larger than 1 and, thus,  $1 - \alpha_{CDW}$  is positive. Consequently, the structure factor does not diverge. In the CDW regime with  $K_S + K_A < 2$  the power  $1 - \alpha_{CDW}$ , however, is less than one. This gives S(k) cusps at  $\pm 2k_F$  when CDW quasi-long range order is present. In Fig. 12 we show examples of S(k) for a case with and without CDW.

Bragg Spectroscopy preceded by a  $\pi/2$  pulse: To de-




Figure 11: Density distribution of molecules after time-of-flight expansion of state III in Fig. 9(a). The expansion time is 0.03s. We assume two hyperfine states of  $^{87}Rb$ . These are converted into Feshbach molecules at T=0 via a fast ramp across a resonance. We assume a complete conversion. The strongly peaked interference pattern of molecules indicates the presence of a quasi-condensate of pairs. For comparison, we also show the TOF expansion of atoms in the PSF phase for the same parameters. The broad Lorentzian distribution demonstrates the absence of single-particle SF.




Figure 12: Structure factor at filling  $\nu=0.3$ . For  $U_{12}=-0.01U$  the system is in the SF regime (dashed line) and for  $U_{12}=-0.07U$  the system is in the PSF regime (continuous line). Cusps at  $|k|=2\pi\nu$  only occur for  $U_{12}=-0.07U$  indicating the coexistence of CDW with PSF order.

tect CFSF order, we propose the following detection method. It applies to the case that the mixture is composed of atoms in different internal states rather than different atomic species. First, we apply a  $\pi/2$  pulse, which transfers the atoms into the superpositions  $b_{1/2,i} \to b_{\pm,i} = (b_{1,i} \pm b_{2,i})/\sqrt{2}$ . We then measure the structure factor, which now corresponds to the Fourier transform of the density correlations  $R_{n\pm}(i,j) = \langle n_{\pm,i} n_{\pm,j} \rangle - \langle n_{\pm,i} \rangle \langle n_{\pm,j} \rangle$ . In terms of the original  $b_{1/2,i}$  operators these density correlations are given by

$$R_{n\pm}(i,j) = \frac{1}{4} \langle (n_{1,i} + n_{2,i})(n_{1,j} + n_{2,j}) \rangle - \frac{1}{4} (\langle n_{1,i} \rangle + \langle n_{2,i} \rangle)(\langle n_{1,j} \rangle + \langle n_{2,j} \rangle) + \frac{1}{2} \langle b_{1,i}^{\dagger} b_{2,i} b_{2,j}^{\dagger} b_{1,j} \rangle$$
(35)

The last term in the above equation is the correlation function  $R_a(i,j)$  of the order parameter of CFSF,  $b_{1,j}b_{2,j}^{\dagger}$ . In Fig. 13, we show the structure factor  $S_+(k)$ , the Fourier transform of Eq. 35, as well as the Fourier transform of  $R_a(i,j)$ . Both  $S_+(k)$  and the Fourier transform of  $R_a(i,j)$  have a cusp around k=0. The cusp is due to the long-range correlations of the anti-pairs in the CFSF. The two functions are nearly identical near k=0, indicating that the momentum distibution of anti-pairs can be measured by determining the structure factor  $S_+(k)$ .

#### V. SUMMARY

We have studied ground state properties of onedimensional Bose mixtures in an optical lattice using both Tomonaga-Luttinger liquid theory and the timeevolving block decimation method. We first discussed the zero-temperature phase diagram in a homogeneous system at different filling fractions and different parameter regimes. We have shown that 1D Bose mixtures in an optical lattice can have quasi-long range orders that include superfluid, paired superfluid (PSF), counterflow superfluid (CFSF), and Mott insulator. We also found that each type of superfluid order can coexist with charge density wave (CDW) order and that in both PSF and CFSF phases single particle superfluidity (SF) is absent.

In addition, we discussed ways of realizing and detecting these phases experimentally. We propose using a Feshbach ramp to probe the momentum distribution of pairs in the PSF, which shows signatures of the quasicondensate of pairs. To detect the CFSF for a mixture composed of two atomic hyperfine states, we propose to measure the static structure factor by using Bragg spectroscopy preceded by a  $\pi/2$  pulse. A sharp peak in the structure factor was shown to be dominated by the contribution from the momentum distribution of anti-pairs in the CFSF phase. Finally, we suggest to detect CDW order with Bragg spectroscopy.

This work was supported by NSF under Physics Frontier Grant PHY-0822671. L.M. acknowledges support from an NRC/NIST fellowship. I.D. acknowledges support from a Grant-in-Aid from JSPS.




Figure 13: Structure factor  $S_+(k)$  (blue line) after applying a  $\pi/2$  pulse in the CFSF phase. The quasi-condensate of antipairs generates an algebraic peak at k=0. The cusp also appear in the Fourier transform of the anti-pair correlation function  $R_a(i,j) = \langle b_{1,i}^{\dagger} b_{2,i} b_{2,j}^{\dagger} b_{1,j} \rangle$  (red dashed line).

- S. Bose, Z. Phys. 26, 178 (1924); A. Einstein, Sitzungsber. K. Preuss. Akad. Wiss., Phys. Math. Kl., 261 (1924); Sitzungsber. K. Preuss. Akad. Wiss., Phys. Math. Kl., 3 (1925).
- [2] K. B. Davis, M. -O. Mewes, M. R. Andrews, N. J. van Druten, D. S. Durfee, D. M. Kurn and W. Ketterle, *Phys. Rev. Lett.* **75** 3969 (1995); M.H. Anderson, J. R. Ensher, M. R. Matthews, C. E. Wieman and E. A. Cornell, *Science* **269** 198 (1995); C. C. Bradley, C. A. Sackett, J. J. Tollett, R. G. Hulet, *Phys. Rev. Lett.* **75** 1687 (1995).
- [3] L.N. Cooper, Phys. Rev. 104 1189 (1956); J. Bardeen,
   L. N. Cooper and J. R. Shrieffer, Phys. Rev. 106 162 (1957); Phys. Rev. 108 1175 (1957).
- [4] C. A. Regal, M. Greiner and D. S. Jin, Phys. Rev. Lett. 92 040403 (2004); M.W. Zwierlein, C. A. Stan, C. H. Schunck, S. M. F. Raupach, A. J. Kerman and W. Ketterle, Phys. Rev. Lett. 92 120403 (2004); M. Bartenstein, A. Altmeyer, S. Riedl, S. Jochim, C. Chin, J. Hecker Denschlag and R. Grimm, Phys. Rev. Lett. 92 120401 (2004); T. Bourdel, L. Khaykovich, J. Cubizolles, J. Zhang, F. Chevy, M. Teichmann, L. Tarruell, S. J. J. M. F. Kokkelmans and C. Salomon Phys. Rev. Lett. 93 050401 (2004).
- [5] A. B. Kuklov and B. V. Svistunov, Phys. Rev. Lett. 90, 100401 (2003).
- [6] A. Kuklov, N. Prokof'ev and B. Svistunov, Phys. Rev. Lett. 92, 050402 (2004); Phys. Rev. Lett. 92, 030403 (2003).
- [7] D. Jaksch, Phys. Rev. Lett. 81, 3108 (1998)
- [8] M. Greiner, O. Mandel, T. Esslinger, T. W. Hänsch and I. Bloch, Nature 415, 39 (2002).
- [9] T. Stöferle, H. Moritz, M. Köhl and T. Esslinger, Phys. Rev. Lett. 92,130403(2004)
- [10] I. B. Spielman, W. D. Phillips, and J. V. Porto , Phys. Rev. Lett. 98, 080404 (2007)
- [11] B. Paredes, A. Widera, V. Murg, O. Mandel, S. Foelling, I. Cirac, G. Shlyapnikov, T. W. Hansch and I. Bloch, Nature 429, 277 (2004).
- [12] A. Widera, S. Trotzky, P. Cheinet, S. Fölling, F. Gerbier, and I. Bloch, Phys. Rev. Lett. 100, 140401 (2008)
- [13] T. Giamarchi, Quantum Physics in one dimension, (Oxford Univ. Press, Oxford, UK, 2004)
- [14] G. Vidal, Phys. Rev. Lett 98, 070201 (2007); G. Vidal,
   ibid. 91, 147902 (2003); ibid. 93, 040502 (2004); S.R.
   White and A. E. Feiguin, ibid. 93, 076401 (2004);
- [15] Y.-Y. Shi, L.-M. Duan, and G. Vidal, Phys. Rev. A 74, 022320 (2006).
- [16] A. J. Daley, S. R. Clark, D. Jaksch, and P. Zoller, Phys. Rev. A 72, 043618 (2005); A. J. Daley, Ph.D. thesis, Leopold-Franzens-Universität Innsbruck, 2005 (unpublished).
- [17] I. Danshita, J. E. Williams, C. A. R. Sá de Melo, and C. W. Clark, *Phys. Rev. A* 76, 043606 (2007); I. Danshita, C. A. R. Sá de Melo, and C. W. Clark, *Phys. Rev. A* 77, 063609 (2008).
- [18] I. Danshita and P. Naidon, *Phys. Rev. A* **79**, 043601 (2009).
- [19] A. Argüelles and L. Santos, Phys. Rev. A 75, 053613 (2007); ibid 77, 059904 (2008).
- [20] L. Mathey, I. Danshita and C. W. Clark, *Phys. Rev. A* **79**, 011602(R) (2009)
- [21] G. G. Batrouni, F. Hébert and R. T. Scalettar, Phys. Rev.

- Lett. 97, 087209 (2006); V. W. Scarola and S. Das Sarma, ibid. 95, 033003 (2005); P. Sengupta1, L. P. Pryadko1, F. Alet, M. Troyer and Guido Schmid, ibid. 94, 207202 (2005); S. Wessel and M. Troyer, ibid. 95, 127205 (2005); D. Heidarian and K. Damle, ibid 95, 127206 (2005); R. G. Melko, A. Paramekanti, A. A. Burkov, A. Vishwanath, D. N. Sheng, and L. Balents, ibid. 95, 127207 (2005); H.P. Büchler and G. Blatter, ibid. 91, 130404 (2004); M. Boninsegni and N. Prokof'ev, ibid. 95, 237204 (2005); M. Boninsegni, J. Low. Temp. Phys. 132, 39 (2005); D. L. Kovrizhin, G. Venketeswara Pai and S. Sinha, Euro. Phys. Lett. 72, 162 (2005); F. Karim Pour, M. Rigol, S. Wessel, and A. Muramatsu, Phys. Rev. B 75, 161104 (2007).
- [22] L. Mathey, Phys. Rev. B 75, 144510 (2007).
- [23] C. Trefzger, C. Menotti, M. Lewenstein, arXiv:0904.1552.
- [24] A. Masaki, S. Tsukada, and H. Mori, J. Phys. Conf. Ser. 150, 032050 (2009).
- [25] F. D. M. Haldane, Phys. Rev. Lett. 47, 1840 (1981).
- [26] M.A. Cazalilla, J. Phys. B: At. Mol. Opt. Phys. 37, S1 (2004).
- [27] M.A. Cazalilla, Phys. Rev. A 70, 041604(R) (2004).
- [28] J. B. Kogut, Rev. Mod. Phys. 51, 659 (1979).
- [29] J. M. Kosterlitz and D. J. Thouless, J. Phys. C 6, 1181 (1973); V. L. Berezinskii, Zh. Eksp. Teor. Fiz. 6, 907 (1970) [Sov. Phys. JETP 32, 493 (1971)].
- [30] T. Mishra, R. V. Pai and B. P. Das, Phys. Rev. A 76, 013604 (2007).
- [31] J. Stenger, S. Inouye, A. P. Chikkatur, D. M. Stamper-Kurn, D. E. Pritchard, and W. Ketterle, *Phys. Rev. Lett.* 82, 4569 (1999); J. Steinhauer, R. Ozeri, N. Katz, and N. Davidson, *ibid.* 88, 120407 (2002).
- [32] D. Clément, N. Fabbri, L. Fallani, C. Fort, and M. Inguscio, Phys. Rev. Lett. 102, 155301 (2009)
- [33] D.S. Hall, M. R. Matthews, C. E. Wieman, and E. A. Cornell, Phys. Rev. Lett. 81, 1543 (1998)
- [34] G. Thalhammer, G. Barontini, L. De Sarlo, J. Catani, F. Minardi, and M. Inguscio, Phys. Rev. Lett. 100, 210402 (2008)

# Appendix A: TEBD METHOD FOR TWO-SPECIES MANY-BODY SYSTEMS

In this appendix, we briefly review the time-evolving block decimation (TEBD) method [14] used in Sec. IV and explain an efficient way to apply the TEBD to a two-species Bose-Hubbard model. We use the number-conserving version of the TEBD method [16].

The TEBD determines the ground state via an imaginary time evolution for one-dimensional (1D) quantum lattice systems. In this method the Hilbert space  ${\bf H}$  is decomposed as

$$\mathbf{H} = \otimes_{l=1}^{M} \mathbf{H}_{l}. \tag{A1}$$

Here, l refers to the lth lattice site, M is the number of sites, and  $\mathbf{H}_l$  is the local Hilbert space at site l with local dimension d, independent of l. Any state  $|\Psi\rangle$  in  $\mathbf{H}$  is

represented as

$$|\Psi\rangle = \sum_{j_1, j_2, \dots, j_M = 1}^{d} c_{j_1, j_2, \dots, j_M} |j_1\rangle |j_2\rangle \cdots |j_M\rangle. \quad (A2)$$

In the TEBD algorithm, coefficients  $c_{j_1,j_2,...,j_M}$  are decomposed as

$$c_{j_{1},j_{2},...,j_{M}} = \sum_{\alpha_{1}=1}^{\chi_{1}} \sum_{\alpha_{2}=1}^{\chi_{2}} \cdots \sum_{\alpha_{M-1}=1}^{\chi_{M-1}} \Gamma_{\alpha_{1}}^{[1]j_{1}} \lambda_{\alpha_{1}}^{[1]} \Gamma_{\alpha_{1}\alpha_{2}}^{[2]j_{2}} \lambda_{\alpha_{2}}^{[2]} \cdots \times \lambda_{\alpha_{M-2}}^{[M-2]} \Gamma_{\alpha_{M-2}\alpha_{M-1}}^{[M-1]j_{M-1}} \lambda_{\alpha_{M-1}}^{[M-1]} \Gamma_{\alpha_{M-1}}^{[M]j_{M}}.$$
(A3)

The variables  $\lambda_{\alpha_l}^{[l]}$  and  $\chi_l$  are the Schmidt coefficients and rank of the Schmidt decomposition of  $|\Psi\rangle$  with respect to the bipartite splitting of the system into  $[1,\ldots,l-1,l]$ :  $[l+1,l+2,\ldots,M]$ ,

$$|\Psi\rangle = \sum_{\alpha_l=1}^{\chi_l} \lambda_{\alpha_l}^{[l]} |\Phi_{\alpha_l}^{[1,\dots,l-1,l]}\rangle |\Phi_{\alpha_l}^{[l+1,l+2,\dots,M]}\rangle. \tag{A4}$$

We take  $\lambda_{\alpha}^{[l]} > \lambda_{\beta}^{[l]}$  for all  $\alpha < \beta$ . In one dimension, the rank  $\chi_l$  at the center of the system must be of the order  $d^{M/2}$  in order to express arbitrary states. However, since it is empirically known that the Schmidt coefficients  $\lambda_{\alpha}^{[l]}$  decrease rapidly with index  $\alpha$  for the ground and lowlying excited states, we set  $\chi_l$  to a relatively small number  $\chi$  for all l.

To efficiently simulate the two-species Bose-Hubbard model (Eq. 1 in the main text), we map it onto the onespecies Hamiltonian

$$H = -t \sum_{l=1}^{2N-2} (b_l^{\dagger} b_{l+2} + \text{h.c.}) + U_{12} \sum_{\text{odd } l} n_l n_{l+1} + \frac{U}{2} \sum_{l=1}^{2N} n_l (n_l - 1),$$
(A5)

where N is the number of sites in the original two-species Hamiltonian. In this one-species Hamiltonian, there are 2N sites, each of which is indexed by l. The odd sites l correspond to species 1 and the even sites to species 2. Hopping between neighboring sites  $-t \, b_{a,i}^{\dagger} b_{a,i+1}$  in Eq. 1 is mapped onto a next-nearest-neighbor hopping  $-t \, b_l^{\dagger} b_{l+2}$  in Eq. A5. Similarly, the inter-species

onsite-interaction  $U_{12}n_{1,i}n_{2,i}$  is mapped onto the nearest-neighbor interaction  $U_{12}n_ln_{l+1}$ . This type of mapping has been successfully applied to treat the two-legged Bose-Hubbard model [17].

We map the two-species Bose-Hubbard Hamiltonian Eq. 1 onto the one-species Hamiltonian because it reduces computational cost dramatically. This cost in TEBD [14] scales as  $Md^3\chi^3$ . For the two-species system with N sites we must define a dimension of the local Hilbert space for each species, say D. Hence, at each site there are  $D^2$  basis functions and the cost scales as  $ND^6$ . On the other hand, for the mapped Hamiltonian with 2N sites and a local dimension D the cost only scales as  $2ND^3$ . In our calculation, we set d=3 for the filling factor  $\nu \leq 0.8$  and d=5 for  $\nu=0.9,1$ . In this case, the mapping makes the computation five to ten times faster.

Imaginary time evolution of any state to the ground state is given by repeated application of  $e^{-iH\delta}$  on  $|\Psi\rangle$ , where  $\delta$  is a small imaginary time step. To apply this operator we first split the Hamiltonian into three parts as  $H=H_{\rm int}+H_{\rm hop}^{\rm odd}+H_{\rm hop}^{\rm even}$ , where

$$H_{\text{int}} = \sum_{m=1}^{N} [U_{12}n_{2m-1}n_{2m} + Un_{2m-1}(n_{2m-1} - 1) + Un_{2m}(n_{2m} - 1)], \qquad (A6)$$

$$H_{\text{hop}}^{\text{odd}} = -t \sum_{\text{odd } m} (b_{2m-1}^{\dagger}b_{2m+1} + b_{2m}^{\dagger}b_{2m+2} + \text{h.c.}),$$

$$H_{\text{hop}}^{\text{even}} = -t \sum_{\text{even } m} (b_{2m-1}^{\dagger}b_{2m+1} + b_{2m}^{\dagger}b_{2m+2} + \text{h.c.}).$$

Subsequently, we use the second-order Suzuki-Trotter expansion to decompose  $e^{-i\hat{H}\delta}$  as

$$e^{-iH\delta} = e^{-iH_{\rm int}\delta/2} e^{-iH_{\rm hop}^{\rm odd}\delta/2} e^{-iH_{\rm hop}^{\rm even}\delta} e^{-iH_{\rm hop}^{\rm odd}\delta/2} \times e^{-iH_{\rm int}\delta/2} + O(\delta^3), \tag{A7}$$

Each of the operators  $e^{-iH_{\rm int}\delta/2}$ ,  $e^{-iH_{\rm hop}^{\rm odd}\delta/2}$ , and  $e^{-i\hat{H}_{\rm hop}^{\rm even}\delta}$  can be decomposed into a product of two-site operators, which can be efficiently applied to the matrix product state  $|\Psi\rangle$  [14, 15, 18]. We use swapping techniques to apply the next-nearest-neighbor operators  $e^{-iH_{\rm hop}^{\rm odd}\delta/2}$  and  $e^{-i\hat{H}_{\rm hop}^{\rm even}\delta}$  [15, 18].