Strong Lensing as a Probe of the Mass Distribution Beyond the Einstein Radius

Mass & Light in SL2S J08544-0121, a Galaxy Group at z = 0.35

Marceau Limousin^{1,2,3}, Eric Jullo^{4,1}, Johan Richard^{5,6}, Rémi Cabanac²,

Sherry H. Suyu⁷, Aleksi Halkola⁸, Jean-Paul Kneib¹, Raphael Gavazzi^{9,10} & Geneviève Soucail¹¹ *

- ¹ Laboratoire d'Astrophysique de Marseille, UMR 6610, CNRS-Université de Provence, 38 rue Frédéric Joliot-Curie, 13 388 Marseille Cedex 13, France
- ² Laboratoire d'Astrophysique de Toulouse-Tarbes, Université de Toulouse, CNRS, 57 avenue d'Azereix, 65 000 Tarbes, France
- ³ Dark Cosmology Centre, Niels Bohr Institute, University of Copenhagen, Juliane Maries Vej 30, 2100 Copenhagen, Denmark
- ⁴ Jet Propulsion Laboratory, Caltech, MS 169-327, 4800 Oak Grove Dr, Pasadena, CA 91109, USA
- ⁵ Durham University, Physics and Astronomy Department, South Road, Durham DH3 1LE, UK
- ⁶ Department of Astronomy, California Institute of Technology, 105-24, Pasadena, CA91125, USA
- ⁷ Argelander-Institut für Astronomie, Universität Bonn, Auf dem Hügel 71, 53121 Bonn, Germany
- ⁸ Excellence Cluster Universe, Technische Universität München, Boltzmannstr. 2, 85748 Garching, Germany
- ⁹ CNRS, UMR 7095, Institut d'Astrophysique de Paris, F-75014, Paris, France
- ¹⁰ UPMC Université Paris 06, UMR 7095, Institut d'Astrophysique de Paris, F-75014, Paris, France
- ¹¹ Laboratoire d'Astrophysique de Toulouse-Tarbes, Université de Toulouse, CNRS, 14 avenue Edouard Belin, 31 400 Toulouse, France

Preprint online version: August 10, 2022

ABSTRACT

Strong lensing is to provide accurate mass measurements within Einstein radii. However, precise modelling of strong lensing systems can be affected by external mass distributions, e.g. the group or cluster within which the lens is embedded. In this article, we propose to turn this limitation to our advantage and to use precise strong lensing modelling to probe external mass distributions surrounding the lens. We consider SL2S J08544-0121, a galaxy group at z = 0.35 that contains a strong lensing system. A simple elliptical isothermal potential cannot reproduce satisfactorily the strong lensing constraints. We include an external mass perturbation corresponding to the group within which the lens is embedded. The lensing properties of this perturbation are parametrised by its total mass M and a smoothing scale s that quantifies the characteristic scale over which M is distributed. For a range of these parameters, we are able to reproduce accurately the observations. This suggests that light is a good tracer of mass. Interestingly, this also shows that a localised strong lensing analysis (on scales of ~ $10^{\prime\prime}$) allows us to constrain global properties of the group as a whole (on scales of ~ $100^{\prime\prime}$). Indeed, we constrain the group mass-to-light ratio to be M/L = 98 ± 27 (*i* band, solar units, not corrected for evolution) and $s = 20'' \pm 9$ $(2 \sigma \text{ confidence level})$. We demonstrate that these strong lensing *only* constraints are due to the perturbed strong lensing configuration, where the main arc is located at $\sim 5''$ from the galaxy, whereas its counter-image is found at $\sim 8''$. To test independently our resulting strong lensing model, we pursue an independent weak lensing analysis of the group and find a mass-to-light ratio in the range 66-146 $(1-\sigma \text{ confidence level})$. We conclude that strong lensing can be used to probe mass distributions beyond their Einstein radius, and we characterise the kind of perturbed strong lensing systems allowing such an analysis: a non dominant strong lensing system used as a particle test probing the main potential. This kind of analysis needs to be validated further on other systems and may constitute a quick method to probe mass distributions. This is particularly relevant in the context of forthcoming wide field surveys which will yield thousands of strong lenses, some of them being perturbed enough to pursue the analysis proposed in this paper.

Key words. Gravitational lensing: strong lensing - Galaxies: groups -

1. Introduction

Gravitational lensing allows one to measure line-of-sight projected mass distributions. When the surface mass density of a lens is larger than a critical threshold, the light from a background source galaxy is lensed into multiple images, a regime called strong lensing (SL). These multiple images provide strong observational constraints on the projected mass distribution of the lens within the Einstein radius. Since the discovery of the first gravitational arc in the core of massive galaxy clusters twenty years ago (Lynds & Petrosian 1986; Soucail et al. 1987), strong lensing is widely used to probe mass distribution of structures on different scales: galaxies (see, *e.g.* the SLACS sur-

Send offprint requests to: marceau.limousin@oamp.fr

^{*} Based on observations obtained with MegaPrime/MegaCam, a joint project of CFHT and CEA/DAPNIA, at the Canada-France-Hawaii Telescope (CFHT) which is operated by the National Research Council (NRC) of Canada, the Institut National des Sciences de l'Univers of the Centre National de la Recherche Scientifique (CNRS) of France, and the University of Hawaii. This work is based in part on data products produced at TERAPIX and the Canadian Astronomy Data Centre as part of the Canada-France-Hawaii Telescope Legacy Survey, a collaborative project of NRC and CNRS. Also based on HST data, program 10876 and Keck telescope data.

vey, Koopmans et al. 2006), galaxy clusters (see, *e.g.* Halkola et al. 2006) and recently galaxy groups (Cabanac et al. 2007; Limousin et al. 2009; Belokurov et al. 2009).

1.1. Perturbing a Strong Lensing System

Because most of the galaxies in the Universe are part of larger structures, either groups or clusters, many SL systems are also part of larger structures. (see, e.g. Kundic et al. 1997; Fassnacht & Lubin 2002; Faure et al. 2004; Morgan et al. 2005; Williams et al. 2006; Momcheva et al. 2006; Auger et al. 2007; Tu et al. 2008; Auger et al. 2008; Grillo et al. 2008; Treu et al. 2009; Inada et al. 2009). Any mass at a small angular distance of a strong lens may induce measurable perturbations in the lensing signal. Not considering this external perturbation can seriously bias the results inferred from the SL modelling as shown by Keeton & Zabludoff (2004): they found that SL modelling of dual-image lenses, which neglect environment, grossly overestimate both the ellipticity of the lens galaxy ($\Delta e/e \sim 0.5$) and the Hubble constant ($\Delta h/h \sim 0.22$). Besides, standard models of four-image lenses, which approximate the environment as a tidal shear, recover the ellipticity reasonably well but still overestimate the Hubble constant ($\Delta h/h \sim 0.15$). They argue that most of the biases are caused by neglecting convergence from nearby massive neighbours. More generally, the topic of modelling a lens with an external mass perturbation has been addressed by different authors (see, e.g Keeton et al. 1997; Kochanek et al. 2001; Keeton & Zabludoff 2004; Oguri et al. 2005, 2008; Dye et al. 2007, and references therein).

To summarise, a precise SL modelling can be *affected* by some external mass distribution, and people have tried to take this bias into account in order to improve the SL modelling. As observations become more and more accurate, we can expect to be more and more sensitive to external mass distributions near strong lenses. In this article, we propose to *turn this bias to our advantage* and to use the perturbations measured in SL modelling as *probes of the external mass distribution*. First, we remind the reader of such an attempt to probe a galaxy cluster potential using a perturbed SL system.

1.2. The "Ring" Test in Abell 1689

In the core of galaxy cluster Abell 1689, Limousin et al. (2007) reported SL systems ("rings") formed around three elliptical galaxies located 100" away from the cluster centre, i.e. between the strong and weak lensing regime regions. These SL systems should be sensitive to the external shear and convergence produced by their parent cluster (Kochanek & Blandford 1991). Based on simulations, Tu et al. (2008) showed that such strong lenses could be used to probe the cluster potential locally. Tu et al. (2008) applied this method to the three rings discovered in Abell 1689. They found that modelling these three rings only (i.e. without including any other multiply-imaged systems also produced by the cluster) provides strong evidence for bimodality of the cluster core; it is not possible to model simultaneously the three rings assuming a unimodal mass distribution for the cluster. This bimodality confirms previous parametric SL studies of Abell 1689 (Miralda-Escude & Babul 1995; Halkola et al. 2006; Limousin et al. 2007; Leonard et al. 2007; Saha et al. 2007; Okura et al. 2008). More importantly, this result shows that SL features of 1-2"-wide Einstein rings actually contain information on the mass distribution of the parent cluster, i.e. on a much larger scale than their Einstein radii. In other words, this study suggests that strong lenses can be used to probe mass distributions *beyond* their Einstein radius. In this article, we develop this idea on another perturbed SL system located in a galaxy group, SL2S J08544-0121.

All results are scaled to the flat, Λ CDM cosmology with $\Omega_{\rm M}$ =0.3, Ω_{Λ} =0.7 and a Hubble constant H₀=70 km s⁻¹ Mpc⁻¹. In such cosmology, at z = 0.35, 1" corresponds to physical transverse distance 4.94 kpc . All images are aligned with the WCS coordinates, i.e. north is up, east is left. Magnitudes are given in the AB system. Ellipticities are expressed as $(a^2 - b^2)/(a^2 + b^2)$, and position angles are given counter-clock wise with respect to the west. Luminosities are given for the *i* band, in solar units, not corrected for passive evolution. Shear and convergence are computed for a source redshift z_s =1.268.

2. SL2S J08544-0121: Presentation & Data

SL2S J08544-0121 is part of the Strong Lensing Legacy Survey (SL2S, Cabanac et al. 2007), which collects SL systems in the Canada France Hawaii Telescope Legacy Survey (CFHTLS). SL2S J08544-0121 is a galaxy group at z = 0.35 presented by Limousin et al. (2009) which contains a SL system (Fig. 1).

2.1. Ground-Based Imaging

SL2S J08544-0121 has been observed in five bands as part of the CFHTLS. The *i*-band data are used to build luminosity maps from isophotal magnitudes of elliptical group members and to perform a weak-lensing analysis.

Fig. 1 bottom panel shows a 10' × 10' CFHTLS *i*-band image, The white cross gives the location of the strong lens. We draw luminosity isodensity contours of 10^5 , 3×10^5 , 10^6 , 3×10^6 and $10^7 L_{\odot} \text{ kpc}^{-2}$. Fig. 1 top-right panel also shows a CFHTLS 1-arcmin² gri colour image centred on the lens.

2.2. Space-Based Imaging

The strong lensing features detected from ground-based images have been observed with the Hubble Space Telescope (HST). Observations were done in snapshot mode (C15, P.I. Kneib, ID 10876) in three bands with the ACS camera (F814, F606 and F475). Fig. 1 shows a colour image of the strong lens based on these observations. We report two multiply-imaged systems: the first system is bright and forms a typical cusp configuration perturbed by a satellite galaxy (labelled Dwarf on Fig. 1). The second system is a very faint arc located West of the lens at a larger radius. It is not possible to reliably separate or identify individual images on the faint arc that may be singly imaged only. Moreover, given its faintness, spectroscopy is hopeless with current facilities, as the surface brightness is ca. 31 mag $\operatorname{arcsec}^{-2}$. Therefore, this faint arc is not used in the following analysis. As can be appreciated on Fig. 1, the HST data brings significant additional information on the lensed features.

2.3. Spectroscopy

We have used the Low Resolution Imager and Spectrograph (LRIS, Oke et al. 1995) on the Keck telescope to measure the spectroscopic redshift of both the lens and the brightest arc of the SL2S J08544-0121 system. On January 14 2007, we obtained 300 seconds on the lensing galaxy and 4 exposures of 900 seconds on the arc, using a 1.0" width slit. A 600 lines mm⁻¹ grism blazed at 4000 Å and a 400 lines mm⁻¹ grating blazed at

8500 Å were used in the blue and red channels of the instrument, both light paths being separated by a dichroic at 5600 Å. The corresponding dispersions are 0.6/1.85 Å and resolutions are 4.0/6.5 Å in the blue/red channel. The resulting extracted spectra are shown Fig. 2. The lens presents a typical elliptical spectrum at $z = 0.3530 \pm 0.0005$ with prominent H and K CaII lines, 4000 Å break, and G band. The spectrum of the arc shows a strong emission line at 8454 Å, resolved in a doublet separated by ~ 5 Å . This gives us an unambiguous redshift determination of $z = 1.2680 \pm 0.0003$.

2.4. Global Properties

The luminosity contours of SL2S J08544-0121 are elongated in the east-west direction. The SL deflector is populated by a single bright galaxy. Note (Fig. 1) that the innermost luminosity isodensity contour at $10^7 L_{\odot} \text{ kpc}^{-2}$ encompasses the SL system but also two bright galaxies located ~ 54" east from the SL system, making this light distribution bimodal. This is the only group of the sample presented by Limousin et al. (2009) for which the luminosity isodensity distribution is not clearly dominated by the lens, making this configuration rather exceptional: the large Einstein radius (~ 5") points towards a significant mass concentration associated with this lens, but the luminosity isodensity distribution is actually bimodal.




Fig. 2. 1-D spectra of the lens (*up*) and of the bright arc (*low*). The green shaded region masks the residuals of a strong OH atmospheric emission

3. Modelling the Lens

In this Section, we attempt to reproduce the SL multiple images using a single elliptical isothermal potential centred on the bright galaxy. All optimisations are performed in the *image plane* using the LENSTOOL software (Jullo et al. 2007). We quantify the goodness of the fit by using the image plane RMS and the corresponding χ^2 . When necessary, we compare the fits using the Bayesian Evidence.

3.1. Observational Constraints

As explained in Section 2.2, we do not use the faint arc in the analysis and focus on the bright multiply-imaged system. This system is composed of 4 main images: 1.1, 1.2, 1.3 and 1.4. Additional images are produced by the satellite (dwarf) galaxy. These are not considered in the analysis because we do not want to complicate the modelling by adding a sub-halo for the satellite galaxy. We also make the assumption that, given its small size, the satellite galaxy does not influence images 1.1 to 1.4. That assumption will be discussed in Section 8 and addressed further in a forthcoming publication. Since the merging arc composed by images 1.2 and 1.3 is well resolved, we can safely associate two other images on this arc, namely 2.2 and 2.3 (Fig. 1). Their counter-images expected near images 1.1 and 1.4 are not safely identified, therefore we do not use them in the analysis. Indeed, parametric strong lensing analyses are highly sensitive to images misidentifications and we prefer to use only images we are confident in. This gives us a total of 8 observational constraints.

3.2. Shape of the Bright Galaxy

In this subsection, we describe the properties of the light distribution of the bright galaxy populating the strong lensing deflector. We use the IRAF task *ellipse* to measure the shape of its isophotes. We find an ellipticity of e = 0.206 and a position angle of 39 ± 5 degrees at 2" from the centre. A closer inspection of the galaxy centre clearly reveals a double core even though the outer isophotes are elliptical. The above measurements therefore correspond to the superposition of the light from each component. The spectrum of the galaxy presented in Fig. 2 do not show features of another galaxy at a different lens redshift along the line of sight. Given the similar colours of these two components, the bright galaxy may be the result of a recent merger.

3.3. Positional Uncertainty

The sizes of the multiple images have been estimated using the IRAF task *imexamine*. They range from 0.11'' to 0.15'', with a mean of 0.13''. Therefore, the positional uncertainty is set to 0.13''. We note that this positional uncertainty sounds large for HST data but we stress that we are not in the case of point-like objects (quasar lensing) where the astrometric precision can reach 0.01''. In our case, the images are extended and the depth of the snapshot observations does not allow to resolve better the conjugated points. We note that large positional uncertainties are often used in the case of extended images (see, *e.g.* Oguri 2010).




Fig. 1. Group SL2S J08544-0121 at $z_{spec} = 0.353$. Upper Left: composite CFHTLS gri colour image (1 arcmin² = 297 × 297 kpc²). Upper Right: composite HST/ACS F814W-F606W-F475W colour image (24" × 24" = 118 × 118 kpc²). We show the proposed multiple-image identification. The dwarf galaxy and the main extra image it produces are labelled. Lower: CFHTLS *i* band (10' × 10' = 2969 × 2969 kpc²). Luminosity isodensity contours of 10⁵, 3 10⁵, 10⁶, 3 10⁶ and 10⁷ L_☉ kpc⁻² are drawn (continuous black line), and the white cross shows the location of the SL system.

3.4. Mass Model

The lens potential is parametrised by a dual Pseudo Isothermal Elliptical Mass Distribution (dPIE, see Elíasdóttir et al. 2007). The 3D density distribution of the dPIE is:

$$\rho(r) = \frac{\rho_0}{(1 + r^2/r_{\rm core}^2)(1 + r^2/r_{\rm cut}^2)}; \quad r_{\rm cut} > r_{\rm core}.$$
 (1)

This distribution represents a spherical system with scale radius r_{cut} , core radius r_{core} and central density ρ_0 .

This profile is formally the same as the Pseudo Isothermal Mass Distribution (PIEMD) described in Limousin et al. (2005). Its scale radius is set to 250 kpc, i.e. larger than the range where the observational constraints are found. Allowing r_{core} to vary produces models with core radii much smaller than the radius over which we have observational constraints. Therefore, we can set r_{core} =0, and the dPIE profile becomes close to isothermal in the constrained region. The remaining free parameters of the dPIE profile are:

- the halo centre position (X,Y) allowed to vary within 3" of the light distribution centre

- the halo ellipticity *e*, forced to be smaller than 0.6, as suggested by numerical simulations (Jing & Suto 2002)

- its position angle θ allowed to vary between 0 and 180 degrees

- The fiducial velocity dispersion¹ allowed to vary between 200 and 900 km/s.

We emphasise that this fiducial velocity dispersion is not the Spherical Isothermal Sphere velocity dispersion. It is usually smaller, see Elíasdóttir et al. (2007) for a self-contained description of the dPIE profile.

3.5. Results: Bad Optimisation

Results of the optimisation are given in Table 1. The fit is bad, the RMS error of image positions is 0.38" in the image plane (i.e. significantly larger than the assumed positional uncertainty of 0.13") and a reduced χ^2 of 29. The halo position is found coincident to the light distribution centre within error bars. The halo ellipticity is at the higher edge of the input prior, and the position angle is equal to ~18 deg. Only when we allow the halo ellipticity to reach values as high as 0.9, are we able to reproduce the observational constraints (RMS equals to 0.06" for e = 0.9, $\theta \sim 19.5$ degrees, and the halo centre is offset from the light distribution centre by one arcsecond.

We conclude that a single potential does not satisfactorily reproduce the observational constraints. In the rest of the paper, we add the contribution of the external mass distribution in the lens modelling. Finally, we have used the non-parametric method of Suyu et al. (2006) in parallel to our method and found that the observational constraints used in this work require an external shear component in order to be properly reproduced.

4. An External Mass Perturbation Based on the Light Distribution: Does Light traces Mass?

The large scale properties of SL2S J08544-0121 shown on Fig. 1 together with the failed modelling attempted in the last Section suggest the need to take into account an external mass perturbation. In order to test the hypothesis that light traces mass, this

¹ linked to the central density by:
$$\sigma_0^2 = \frac{4}{3}G\pi\rho_0 \frac{r_{core}^2 r_{cut}^2}{(r_{cut} - r_{core})(r_{cut} + r_{core})^2}$$

external perturbation will be mapped from the known light distribution properties.

4.1. Luminosity Maps

The first step is to build luminosity maps of SL2S J08544-0121 from which we will derive the external mass perturbation properties. Group members have been colour selected. All galaxies having a r - i colour difference smaller than 0.15 with the bright galaxy deflector are selected as group members (Limousin et al. 2009). Because we want to describe the perturbation of the galaxy group on the SL system, this luminosity map should not take into account the light coming from the galaxies populating the SL deflector. Therefore, we select all group members excepting the bright galaxy populating the deflector and the associated satellite galaxy. This partial group luminosity is referred to as Lext in the following. From this catalogue, we generate luminosity maps as described below. An important ingredient of this procedure is the smoothing scale. Indeed, the shape of the luminosity maps strongly depends on the smoothing scale. Mechanically that has an influence on the properties of the derived mass maps. Therefore, this smoothing scale is a free parameter describing the external mass perturbation.

We use the following smoothing scheme: the $10' \times 10'$ CFHTLS *i*-band image is divided into cells of size *c* pixels, which translates into $c \times 0.186''$. We compute the rest-frame *i*band luminosity *L* of each galaxy located in a given cell with

$$L = 10^{(M_{\odot} - M + \text{DM} + \text{k})/2.5}$$
(2)

where *M* is the *i*-band isophotal magnitude of the galaxy, M_{\odot} is the solar absolute magnitude in the *i* band; DM is the distance modulus, and *k* the k-correction factor, estimated from elliptical templates by Bruzual & Charlot (2003) using single-burst stellar formation models. Then we sum up the luminosities of all galaxies per cell to get the total luminosity within each cell. The resulting luminosity isodensity map is then convolved with a Gaussian kernel of sigma *w*. This gives an angular smoothing scale *s* equals to $c \times 0.186 \times w''$ (since the pixel size equals 0.186''). Fig. 3 shows three luminosity maps, where we distribute the same total luminosity contours of 10^7 and $10^8 L_{\odot} \operatorname{arcsec}^{-2}$. One can appreciate how the smoothing scale *s* influences the shape of the luminosity isodensity contours.

4.2. From Light Map to Mass Map

Once a luminosity map with a given smoothing scale *s* is obtained, we assume a mass-to-light ratio $M_{ext}/L_{ext} = 1$ to convert it into a mass map. Because we omit the bright galaxy populating the deflector when building the luminosity map, this mass map can be considered as the external mass perturbation. Therefore we refer to the mass contained in this map as M_{ext} . Then, we use the algorithm developed by Jullo & Kneib (2009) to interpolate this mass map into a grid of analytic circular dPIE potentials, supported by LENSTOOL.

We model the mass distribution of SL2S J08544-0121 with a 5' hexagonal grid of dPIE potentials. In order to build an adaptive grid where resolution follows the 2-D mass density, we recursively split the input mass map into equilateral triangles until the mean surface density per triangle is lower than 10^7 M_{\odot} arcsec⁻². Then we place a dPIE potential at each node of the grid with the following parameters: core radii $r_{\rm core}$ equal to the local grid resolution and cut-off radii $r_{\rm cut} = 3 \times r_{\rm core}$. In Jullo

Model	$\delta(x)$	$\delta(y)$	е	θ	$\sigma_0 (\mathrm{kms^{-1}})$	RMS	χ^2	log(Evidence)	Prior
1 Lens	-0.24±0.12	-0.02 ± 0.10	$0.597^{+0.002}_{-0.038}$	17.9±1.6	476±7	0.38"	86/3	-62	<i>e</i> < 0.6
Ext. Perturb.	0.01 ± 0.05	-0.26±0.08	0.499 ± 0.04	21.5±0.9	447±3	0.05"	0.96	-20	<i>e</i> < 0.6

Table 1. Parameters of the lens inferred from two optimisations: *First line:* A single halo models the lens potential. *Second line:* We add an external mass perturbation on top of the halo lens potential. Coordinates are given in arcseconds with respect to the centre of the galaxy deflector. e is the central halo ellipticity. Error bars are given for 1σ confidence level.




Fig. 3. Three luminosity maps of the same luminosity L_{ext} at different smoothing scales as indicated on each panel. We draw in white luminosity isodensity contours of 10⁷ and 10⁸ L_{\odot} arcsec⁻². We can appreciate how the smoothing scale *s* influences the shape of the luminosity isodensity contours, and by construction the shape of the resulting mass distribution, hence its lensing properties.

& Kneib (2009), we found that such values of r_{cut} was ensuring a smooth and extended density profile. We estimate the dPIE central velocity dispersions σ_i by inverting the equation

$$\sigma_i^2 = M_{i,j} \Sigma_j \tag{3}$$

where Σ_j is the surface density at the grid nodes location. $M_{i,j}$ is a mapping matrix whose coefficients depend of the dPIE core and cut-off radii (see Jullo & Kneib 2009). In order to prevent negative σ_i^2 , we invert equation 3 by a mean-square minimisation technique. Finally, each luminosity map is scaled to the corresponding mass map by multiplying all dPIE potential velocity dispersions by a constant mass-to-light ratio M_{ext}/L_{ext} . This mass-to-light ratio M_{ext}/L_{ext} is the second free parameter describing the perturbation produced by the galaxy group. The density threshold controls the grid resolution, and as such might be considered a critical parameter but we have tried to use smaller thresholds, down to $10^5 \text{ M}_{\odot} \operatorname{arcsec}^{-2}$, and the results were unchanged. We therefore keep the $10^7 \text{ M}_{\odot} \operatorname{arcsec}^{-2}$ threshold because the corresponding mass maps require less mass clumps. We also force the algorithm to stop after 4 levels of splitting. On average, a grid contains about 200 dPIE potentials.

4.3. Modelling the Lens Accounting for the External Perturbation

We now come back to modelling the SL system, taking into account the external mass perturbation parametrised by a smoothing scale *s* and a mass-to-light ratio M_{ext}/L_{ext} . We generated mass maps with smoothing scales *s* ranging from 1 to 40" by steps of 2.5" and mass-to-light ratios from 10 to 190 by steps of 20. Each mass map is then included in the modelling of the SL system. This modelling is performed in the image plane. Parameters of the lens potential are taken from Section 3.4. For

each set of parameters (s, M_{ext}/L_{ext}), we quantify the goodness of the SL modelling using the image plane RMS, the corresponding χ^2 and the Bayesian Evidence.

In the following, because our goal is to constrain the galaxy group as a whole, we use M/L corresponding to the *total* mass-to-light ratio of the group; i.e. M (resp. L) is the sum of the external mass (luminosity) perturbation *and* the mass (luminosity) of the lens. We checked that degeneracies of each mass component near the lens are small. For the range of parameters (s, M_{ext}/L_{ext}) investigated in this work, we compute M_{ext}/M_{lens} in a circle of radius 10" centred on the lens. This ratio falls between 10⁻⁷ and 10⁻⁵.

Total masses and luminosities are computed within a region of $10' \times 10'$ centred on the lens. At the redshift of the group, it corresponds to ~ 3×3 Mpc² (Fig. 1).

5. Results: Properties of SL2S J08544-0121

For a certain range of parameters characterising the external mass perturbation we obtain excellent fits for the observed constraints. We present first the best-fit model, and then the derived constraints on the galaxy group properties. We emphasise that what we achieve here is to constrain the properties of the galaxy group as a *whole* (on scales of 100'') based on a local SL analysis *only* (on scales of 10'').

5.1. Best-Fit Model for the Lens

Results are given in Table 1. The best fit is found for a total mass-to-light ratio ≈ 75 (*i* band, solar units, not corrected for evolution) and a smoothing scale $\approx 20''$ (Fig. 4). The RMS error between observed and modelled image positions in the image plane is 0.05", yielding a reduced χ^2 of 0.96. This is a sig-




Fig. 4. Results: constraints on the galaxy group mass-to-light ratio M/L and smoothing scale *s* that characterises the size of dark matter clumps. Vertical dashed lines are lower-higher constraints (1 σ error bars) on group mass-to-light ratio from an independent weak-lensing analysis (Section 7).

nificant improvement compared to the modelling without external mass perturbation, which had RMS = 0.38": the difference in log(Evidence) values, which accounts fully for the additional complexity of the new model (extra parameters and their prior PDFs) is 42. The log(Evidence) difference between two models compares the relative probability of getting a set of data explained by the associated model. Here the data set associated with the perturbed model is e^{42} times more probable than the data set associated with the simple model.

We find the position of the halo to coincide with the centre of the light distribution. The modelled position angle of the halo is 21.5 deg. Comparing this value to the position angle of the light distribution is complicated by the bimodal light distribution of the bright galaxy (Section 3.2). In particular, a merger will affect the light and mass distributions so that agreement may be not necessarily expected.

5.2. Constraints on the Group as a Whole

 χ^2 differences between each model are translated into confidence levels and drawn on Fig. 4. Considering the 2σ contour, we find M/L = 98±27 (*i* band, solar units, not corrected for evolution) and $s = 20'' \pm 9$.

5.3. Is Mass Traced by Light ?

We are able to reproduce accurately the observational constraints when considering an external mass perturbation drawn from the light distribution. Because our SL analysis is sensitive to the mass, this finding is consistent with the hypothesis that light is a good tracer of mass. We note, however, that we have not demonstrated the uniqueness of the smoothed light model.

6. Local Effect of Large Scale Perturbation

In this Section, we propose to explain why a *local* SL analysis is able to constrain *global* properties of the galaxy group hosting the lens. First, we investigate the lensing impact of the external

perturbation on the local SL modelling (i.e. on the local image positions).

6.1. Lensing Properties of the External Perturbation

The lensing properties of a mass distribution are commonly parametrised by a shear γ and a convergence κ (see, *e.g.* Schneider et al. 1992, for the definition of these quantities). Here we estimate the mean shear and convergence experienced locally by the lens for each set of parameters (s, M_{ext}/L_{ext}) by averaging γ and κ generated by the perturbation within a 7" square encompassing all the multiple images.

Fig. 5 shows κ and γ maps generated by an external perturbation of fixed mass (5.7×10¹⁴ M_☉) for different values of *s* (reported on each panel). These maps have been generated for a source redshift equal to 1.268. Red crosses indicate the lens position. One can appreciate how the experienced shear and convergence are correlated with the smoothing scale.

6.2. Shape of the Constraints

Fig. 6 shows lines of constant κ and γ on the results of Fig 4. We see that the constraints inferred from the SL analysis do not follow κ lines but do follow γ lines. In particular, the best model generates a shear of ~ 0.075. We interpret the shape of the constraints as follows: one needs to generate a shear value of ~ 0.075 with a mass distribution parametrised by a smoothing scale and a total mass. For a given total mass, the smoother this mass distribution (the higher *s*), the smaller the generated shear. Therefore, the smoother the mass distribution, the higher the total mass in order to generate a given shear level.

6.3. Why Closed Contours ? Shear, Convergence, and Beyond

Fig. 6 suggests that the observational constraints require locally an external shear component of ~0.075. However, our analysis rejects some external mass perturbations characterised by such a shear (Fig. 4). This suggests, as found by Keeton & Zabludoff (2004), that the shear approximation fails to capture all of the environmental effects. In other words, the shape of the constraints follow constant shear lines. However, the contours do close, which means that the constraints are sensitive to more than the shear; most probably higher order terms beyond shear that are naturally provided by the modelling proposed in this work.

Besides, we estimate the shear experienced by images 1.1, 1.2, 1.3 and 1.4. To do so, we consider all models falling in the 1σ contour. For each model, we compute the shear experienced by the images, and from these numbers, we estimate the mean shear and the associated standard deviation. They are equal to 0.075, 0.074, 0.073 and 0.073 respectively, with a typical uncertainty equal to 0.01. Therefore, each image does experience the same shear value within the error bars. We will investigate further the differences between our approach and a constant external shear approach further in Appendix.

7. Mass-to-Light Ratio from Weak Lensing

We have presented constraints on the mass-to-light ratio of galaxy group SL2S J08544-0121 based on a local SL analysis. In this Section, we aim to constrain the mass-to-light ratio of SL2S J08544-0121 from an independent weak lensing (WL) analysis, which is intrinsically more sensitive to the projected




Fig. 5. Shear (upper panels) and convergence (lower panels) maps generated by the external perturbation and experienced by the SL system whose centre is given by the red crosses. The total mass is fixed to the same value $(5.7 \ 10^{14} \ M_{\odot})$ in all panels. Panel sizes are 600×600 square arcseconds, and the smoothing scales *s* vary as indicated on each panel. White contours correspond to shear levels of 0.1 and convergence levels of 0.1 and 0.2. One can appreciate how the shear and convergence generated by the group are correlated with the smoothing scale.

mass distribution on large scales. The goal is to check whether M/L inferred from the two techniques are consistent.

For a detailed description of our WL methodology, see Limousin et al. (2009). Here we just give a brief reminder. We select as background sources all galaxies with *i*-band magnitudes in the range 21.5-24. Their density is 13.5 arcmin⁻². The completeness magnitude in this band is 23.91. The seeing is 0.51". A Bayesian method, implemented in the IM2sHAPE software (Bridle et al. 2002), is used to fit the shape parameters of the faint background galaxies and to correct for PSF smearing. From the catalogue of background galaxies, Limousin et al. (2009) performed a one-dimensional WL analysis. They fit a Singular Isothermal Sphere model (SIS) to the reduced shear signal between 150 kpc and 1.2 Mpc from the group centre, finding an Einstein radius of $5.4\pm2.1''$. In order to relate the strength of the WL signal to a physical velocity dispersion characterising the group potential, Limousin et al. (2009) estimate the mean geometrical factor using the photometric redshift catalogue from the T0004 release of the CFHTLS-Deep survey ² (Ienna & Pelló 2006). They find $\sigma_{\text{SIS}} = 658^{+119}_{-146} \text{ km s}^{-1}$. This translates into a total mass within the considered square of $5.3\pm2.0\ 10^{14}\ M_{\odot}$. Because the total luminosity is $5 \, 10^{12} L_{\odot}$, we find a mass-to-light ratio equals to 106 ± 40 (*i* band, solar units, not corrected for passive evolution).

This is comparable to the M/L constrained by SL *only*. The good agreement (Fig. 4) between the two methods gives support to the SL only analysis.

8. Discussion

8.1. Mass is Traced by Light

An external mass perturbation derived from the light of the group members allows us to fit accurately the observed constraints. Because observed constraints are sensitive to mass and not to light, this suggests that light is a good tracer of mass. We note that this assumption constitutes an efficient way of taking into account an external mass perturbation in SL modelling. Indeed, this perturbation is fully described with only two parameters, the mass-to-light ratio and the smoothing scale. In contrast, describing this perturbation parametrically using a mass clump would require at least three parameters (position and velocity dispersion), unless independent data motivate additional priors (see, *e.g.* Tu et al. 2009, where X-ray observations allow one to constrain the group centre).

8.2. What is the source of the Constraints ?

Why our SL analysis did allow us to infer some constraints on the whole galaxy group? We claim that this is due to the perturbed state of the SL system of SL2S J08544-0121. Most of the

² http://www.ast.obs-mip.fr/users/roser/CFHTLS_T0004/




Fig. 6. As in Fig. 4, constraints obtained on the galaxy group as a whole derived from the *local* SL analysis are shown as dashed contours. Solid lines corresponds to lines of constant κ (left) and γ (right) generated by the external mass perturbation and experienced locally by the lens, their values are labelled on each line.

perturbed signal of the multiply imaged system comes from image 1.4, because it is located much further from the lens centre (~ 8") than images 1.1, 1.2 and 1.3 (~ 5"). If we remove image 1.4 from the set of observational constraints, we are able to fit very well the remaining images without considering any external mass perturbation (the lens being modelled as in Section 3.4.) In that case, we get RMS = 0.03" and a reduced χ^2 equals to 0.03. Therefore, ignoring image 1.4 prevents us to put any constraints on the external mass perturbation, i.e. the host galaxy group. This shows that image 1.4 yields the constraints presented in this work. This finding will help us diagnose the type of the SL systems favouring this kind of analysis (see Section 8.5).

We note that the SL analysis presented here is very simple since we just conjugate a couple of images with each other. In particular, we do not use the constraints coming from the whole Einstein ring. More sophisticated methods fully take into account arc brightness constraints (see, *e.g.* Warren & Dye 2003; Suyu et al. 2006; Barnabè & Koopmans 2007). We are aware that we ignore some informations that could allow us to put stronger constraints on the galaxy group. One the other hand, the basic level of the SL analysis done here emphasises even more the prospects of this method.

8.3. The Satellite Galaxy

We have assumed that the satellite galaxy does not produce significant shear on the images used as constraints. However, one could argue that neglecting this satellite galaxy effectively produces the claimed constraints from the SL analysis. This is not likely, because of the location of the satellite galaxy with respect to the multiple images (Fig. 1). The dwarf galaxy may produce a marginal shear on on images 1.1 1.2 1.3, but is unlikely to have any significant influence on image 1.4, the one image yielding most of the constraints. Indeed, the distance between the satellite galaxy and image 1.4 is ~ 13". We note that we do not quantify the bias that could result under our working assumption.

Besides, a paper focusing on the properties of the satellite galaxy (Suyu & Halkola, submitted) shows that even with the satellite galaxy included in the lens model, an external shear of approximately the same magnitude is needed to fit the observed constraints.

8.4. Choice of the Lens' Scale Radius

The dPIE scale radius is where the logarithmic slope of the 3D density profile smoothly decreases from -2 to -4. The scale radius of the lens is set to 250 kpc in the present analysis. We have also done a complete analysis for a scale radius of 400 kpc as a sanity check and found that the results inferred for the group do not change significantly. To understand why, we superimposed critical lines of the best-fit parameters of Table 1, for a source redshift of 1.268 (without external perturbation), and the critical lines of the best-fit parameters of Section 5.1 (with external mass perturbation). We find that the external mass perturbation generates a critical line shift of 1.3''. In parallel, we investigated the critical lines shifts between various scale radii; increasing from 250 kpc to 400 kpc and decreasing from 250 kpc to 100 kpc. The shifts are 0.12'', an order of magnitude smaller than the shift due to the external mass perturbation.

8.5. Looking for Perturbed SL Systems

We propose to characterise the kind of perturbed SL systems one should look for to be able to perform analyses similar to the one presented in this work. From the ring test done in Abell 1689 (Tu et al. 2008, see Section 1.3) and the analysis presented in this paper, we hint the need for *a non-dominant SL system used as a particle test probing the main potential.*

This is linked to the global geometry of the structure hosting the SL system: in order to be perturbed, a SL system should not be at the centre of the structure. Indeed, if the lens studied in this paper would have dominated the whole group potential, image 1.4 would have been located at a similar distance from the lens centre as images 1.1, 1.2 and 1.3.

The Cosmic Horseshoe (Belokurov et al. 2007; Dye et al. 2008) illustrates this point: it is an almost complete Einstein ring of radius 5" containing a luminous red galaxy in its centre. This galaxy is the brightest object in the group of ~ 26 members as revealed by the SDSS photometry and it dominates the group

light distribution. No external shear is required in the model of the Cosmic Horseshoe SL system, which is already suggested by the nearly perfect circle outlined by the ring.

To summarise, we should look for multiply imaged systems where one of the images is found at a larger radial distance than the other images of the SL system.

8.6. SL2S J08544-0121: Further Evidences for a Bimodal Mass Distribution from Spectroscopy of Group Members

We have shown that the strong lensing modelling *only* provides strong hints for a bimodal mass distribution: a first one is clearly associated with the strong lensing deflector, and the second one perturbing the strong lensing configuration seems, to first order, associated with the second light peak of the bimodal light map. This suggests a dynamically young structure in the process of formation. Additional data support this hint. We measured redshift for 36 galaxies along the direction of SL2S J08544-0121 by using spectroscopic data acquired with FORS2 at the ESO Very Large Telescope (VLT), and confirmed the presence of a high concentration of galaxies at $z \sim 0.35$ (Muñoz et al., in preparation). A careful analysis of the redshift distribution of galaxies around this peak reveals two close structures with a radial velocity difference of $V_r = 1\,180\,\mathrm{km\,s^{-1}}$, maybe bounded-incoming structures. This result is in agreement with our strong lensing only hints.

9. Conclusion

We propose a method to constrain the dark matter distribution of galaxy groups and clusters. Exploiting information contained in perturbed SL systems, we use the SL geometry to probe the main potential of the host structure responsible of that perturbed state.

We show that the SL *only* constraints on the mass-to-light ratio of SL2S J08544-0121 are in good agreement with WL constraints obtained independently, comforting the reliability of the proposed method. Moreover, the SL *only* analysis provides strong hints for a bimodal mass distribution, which is confirmed by spectroscopy of galaxy group members.

We advocate the need for a dedicated search of perturbed SL systems in the HST archive in order to test and validate further this method, particularly promising in the light of future large surveys that will yield thousands of SL systems, some of them being perturbed enough to perform the test presented in this paper.

Acknowledgement

ML acknowledges Bernard Fort, Masamune Oguri & Phil Marshall for related discussions. ML acknowledges the anonymous referee for a detailed report, and Christopher Kochanek for insightful comments on the submitted version of this paper. ML acknowledges the Centre National d'Etudes Spatiales (CNES) and the Centre National de la Recherche Scientifique (CNRS) for their support. ML est bénéficiaire d'une bourse d'accueil de la Ville de Marseille. The Dark Cosmology Centre is funded by the Danish National Research Foundation. We thank the Danish Centre for Scientific Computing at the University of Copenhagen for providing a generous amount of time on its supercomputing facility. JR acknowledges support from an EU Marie-Curie fellowship. SHS is supported in part through the Deutsche Forschungsgemeinschaft (DFG) under project SCHN 342/7-1, and AH by the DFG cluster of excellence 'Origin and Structure of the Universe'. JPK acknowledges CNRS for its support.

References

- Auger, M. W., Fassnacht, C. D., Abrahamse, A. L., Lubin, L. M., & Squires, G. K. 2007, AJ, 134, 668
- Auger, M. W., Fassnacht, C. D., Wong, K. C., et al. 2008, ApJ, 673, 778
- Barnabè, M. & Koopmans, L. V. E. 2007, ApJ, 666, 726
- Belokurov, V., Evans, N. W., Hewett, P. C., et al. 2009, MNRAS, 392, 104
- Belokurov, V., Evans, N. W., Moiseev, A., et al. 2007, ApJ, 671, L9
- Bridle, S., Kneib, J.-P., Bardeau, S., & Gull, S. 2002, in The shapes of galaxies and their dark halos, Proceedings of the Yale Cosmology Workshop , New Haven, Connecticut, USA, 28-30 May 2001. Edited by Priyamvada Natarajan., ed. P. Natarajan, 38–+
- Bruzual, G. & Charlot, S. 2003, MNRAS, 344, 1000
- Cabanac, R. A., Alard, C., Dantel-Fort, M., et al. 2007, A&A, 461, 813
- Dye, S., Evans, N. W., Belokurov, V., Warren, S. J., & Hewett, P. 2008, MNRAS, 388, 384
- Dye, S., Smail, I., Swinbank, A. M., Ebeling, H., & Edge, A. C. 2007, MNRAS, 379, 308
- Elíasdóttir, Á., Limousin, M., Richard, J., et al. 2007, ArXiv e-prints, 710
- Fassnacht, C. D. & Lubin, L. M. 2002, AJ, 123, 627
- Faure, C., Alloin, D., Kneib, J. P., & Courbin, F. 2004, A&A, 428, 741
- Grillo, C., Lombardi, M., Rosati, P., et al. 2008, A&A, 486, 45
- Halkola, A., Seitz, S., & Pannella, M. 2006, MNRAS, 372, 1425
- Ienna, F. & Pelló, R. 2006, in SF2A-2006: Semaine de l'Astrophysique Francaise, ed. D. Barret, F. Casoli, G. Lagache, A. Lecavelier, & L. Pagani, 347-+
- Inada, N., Oguri, M., Shin, M., et al. 2009, AJ, 137, 4118
- Jing, Y. P. & Suto, Y. 2002, ApJ, 574, 538
- Jullo, E. & Kneib, J. 2009, MNRAS, 395, 1319
- Jullo, E., Kneib, J.-P., Limousin, M., et al. 2007, New Journal of Physics, 9, 447
- Keeton, C. R., Kochanek, C. S., & Seljak, U. 1997, ApJ, 482, 604
- Keeton, C. R. & Zabludoff, A. I. 2004, ApJ, 612, 660
- Kochanek, C. S. & Blandford, R. D. 1991, ApJ, 375, 492
- Kochanek, C. S., Keeton, C. R., & McLeod, B. A. 2001, ApJ, 547, 50
- Koopmans, L. V. E., Treu, T., Bolton, A. S., Burles, S., & Moustakas, L. A. 2006, ApJ, 649, 599
- Kundic, T., Hogg, D. W., Blandford, R. D., et al. 1997, AJ, 114, 2276
- Leonard, A., Goldberg, D. M., Haaga, J. L., & Massey, R. 2007, ApJ, 666, 51
- Limousin, M., Cabanac, R., Gavazzi, R., et al. 2009, A&A, 502, 445
- Limousin, M., Kneib, J.-P., & Natarajan, P. 2005, MNRAS, 356, 309
- Limousin, M., Richard, J., Jullo, E., et al. 2007, ApJ, 668, 643
- Lynds, R. & Petrosian, V. 1986, in Bulletin of the American Astronomical Society, 1014-+
- Miralda-Escude, J. & Babul, A. 1995, ApJ, 449, 18
- Momcheva, I., Williams, K., Keeton, C., & Zabludoff, A. 2006, ApJ, 641, 169
- Morgan, N. D., Kochanek, C. S., Pevunova, O., & Schechter, P. L. 2005, AJ, 129, 2531

Oguri, M. 2010, ArXiv e-prints

- Oguri, M., Inada, N., Blackburne, J. A., et al. 2008, MNRAS, 391, 1973
- Oguri, M., Keeton, C. R., & Dalal, N. 2005, MNRAS, 364, 1451
- Oke, J. B., Cohen, J. G., Carr, M., et al. 1995, PASP, 107, 375
- Okura, Y., Umetsu, K., & Futamase, T. 2008, ApJ, 680, 1
- Saha, P., Williams, L. L. R., & Ferreras, I. 2007, ApJ, 663, 29
- Schneider, P., Ehlers, J., & Falco, E. E. 1992, Gravitational Lenses (Berlin: Springer-Verlag)
- Soucail, G., Fort, B., Mellier, Y., & Picat, J. P. 1987, A&A, 172, L14
- Suyu, S. H., Marshall, P. J., Hobson, M. P., & Blandford, R. D. 2006, MNRAS, 371, 983
- Treu, T., Gavazzi, R., Gorecki, A., et al. 2009, ApJ, 690, 670
- Tu, H., Gavazzi, R., Limousin, M., et al. 2009, A&A, 501, 475
- Tu, H., Limousin, M., Fort, B., et al. 2008, MNRAS, 1169
- Warren, S. J. & Dye, S. 2003, ApJ, 590, 673
- Williams, K. A., Momcheva, I., Keeton, C. R., Zabludoff, A. I., & Lehár, J. 2006, ApJ, 646, 85

Appendix A: Taking an External Mass Perturbation Into Account: Comparison with Other Approaches

We have proposed in this article a way of taking into account an external mass perturbation in a strong lensing (SL) modelling. Here we try other possible and more conventional approaches: i) a constant external shear profile and ii) a Singular Isothermal Sphere centred on the second high luminosity peak, which, by construction, is the main mass concentration perturbing the SL in the method proposed in this work.

A.1. A Constant External Shear

Although unphysical (any mass distribution will not generate shear only but also convergence), the external shear model is widely used and is often a good approximation. Here we address the modelling of the SL system with a constant external shear component parametrised by a position angle and a strength (γ_{Kst}). This modelling is performed in the image plane. Parameters of the potential describing the lens are set as in Section 3.4. The external shear strength is allowed to vary between 0 and 0.3. The upper limit corresponds to a very strong shear value: for comparison, the massive galaxy cluster Abell 1689 (Limousin et al. 2007), produces an average shear value of 0.23 at 50" away from its centre.

We are able to get a very good fit, with $\chi^2 < 1$. The best model corresponds to a circular halo for the lens (e = 0.036) centred on the bright galaxy (X = -0.36", Y = 0.07"), making its position angle (95 degrees) irrelevant. The lens fiducial velocity dispersion equals to 450 ± 8 km s⁻¹ (1 σ). The external shear is described by $\gamma_{Kst} = 0.19$, more than twice the one derived in Section 6.2 (~ 0.075), and a position angle equal to 19.8 degrees.

The LENSTOOL software does explore the parameter space using a MCMC sampler (Jullo et al. 2007). Therefore, we can use these MCMC realisations in order to investigate the degeneracies between the different parameters. The following figures have been generated this way.

Fig. A.1 shows that there is a strong degeneracy between *e* and γ_{Kst} . We see that the solution derived in Section 5.1 (i.e. an external shear of ~0.075 and an ellipticity of ~0.5) is included in the 1- σ contour. On the other hand, the position angle of the external shear is very well constrained to be ~20 degrees. This position angle points towards the second high luminosity light clump. This suggests that, to first order, the external mass perturbation is dominated by this component. We note that the best-fit model needs an external shear of order 0.18, which is a pretty unlikely value in our case because it is comparable to what would be experienced at ~ 100" from the centre of Abell 1689.

A.2. An SIS profile

The first order mass perturbation is associated to a second peak of high luminosity. We put an SIS mass distribution at the location of this second luminosity peak (X,Y = -53,10" wrt the lens). We allow its velocity dispersion to vary up to 800 km s⁻¹, an upper limit motivated by the WL analysis of the full group (Section 7), and do the SL modelling with parameters set as in Section 3.4. We are able to get a very good fit, with $\chi^2 < 1$. The lens halo is centred on the bright galaxy. Its ellipticity equals $0.43^{+0.01}_{-0.12}$ and its position angle 27 ± 2 deg. The lens fiducial velocity dispersion equals to 441 ± 7 km s⁻¹ (1 σ). The external shear and convergence generated by the SIS profile at the lo-




Fig. A.1. Results of a constant external shear model. *Top:* degeneracy between the halo ellipticity *e* and the strength of the external shear $\gamma_{\text{Kst.}}$ *Bottom:* posterior probability distribution for the position angle of the external shear.

cation of the multiple images are equal by definition, between 0.07 and 0.10.

We show on Fig. A.2 the degeneracies between the lens halo ellipticity and the SIS profile velocity dispersion, related to the strength of the external shear experienced by the multiple images. We see that the solution derived in Section 5.1 (i.e. an external shear of ~0.075, corresponding to $\sigma_{\text{SIS}} \sim 700 \,\text{km s}^{-1}$ and an ellipticity of ~0.5) is included in the 1- σ contour.

A.3. Discussion

In each cases investigated in this Appendix, we find that the solution we have derived in Section 5.1 using our original method is consistent with solutions derived with more conventional methods.

We note that conventional methods exhibit strong degeneracies between the lens halo ellipticity and the strength of the external shear. These degeneracies are smaller in the case of the SIS profile (lens ellipticity is constrained between 0.3 and 0.6) compared to the case of a constant shear profile (ellipticity un-




Fig. A.2. Degeneracies between the lens halo ellipticity and the SIS profile velocity dispersion, related to the strength of the external shear experienced by the multiple images.

constrained between the allowed priors: 0 and 0.6). The main difference between the SIS profile and the constant shear profile is that the SIS profile generates both shear and convergence.

Concerning the lens itself, we note that all fitted fiducial velocity dispersions are consistent, whatever the method used to take into account the external mass perturbation. They fall between 433 km s⁻¹ and 458 km s⁻¹. This translates into a projected mass computed in a radius of 10" between 0.93 and 1.04×10^{13} M_{\odot}. This is expected because the mass of the lens within this radius is set by the location of the SL constraints and therefore does not depend much on the external mass perturbation (see also Section 4.3).