

Spectral and temporal changes associated with flux enhancement in 4U 1626–67

Chetana Jain^{1,2*}, Biswajit Paul² and Anjan Dutta¹

¹*Department of Physics and Astrophysics, University of Delhi, Delhi 110007, India*

²*Raman Research Institute, Sadashivnagar, C. V. Raman Avenue, Bangalore 560080, India*

30 October 2018

ABSTRACT

4U 1626–67 is an accretion powered X-ray pulsar that shows remarkably stable X-ray luminosity above hours timescale and gradual intensity variation on a few years timescale. Unlike the other high magnetic field binary X-ray pulsars, the spin-up or spin-down rate of the neutron star is also very stable upto several years. Here we report a significant increase in the X-ray intensity in the long term *RXTE*-ASM light curve of 4U 1626–67. Similar enhancement in the X-ray flux has also been detected in the *Swift*-BAT light curve. The increase in the X-ray flux took place over a long period of about 100 days and there appears to be two episodes of flux enhancement. We have investigated the spectral and timing features of 4U 1626–67 during its current state of enhanced flux emission with data obtained from the Proportional Counter Array and the High-Energy X-ray Timing Explorer on board the *Rossi X-ray Timing Explorer*. We report the detection of a torque reversal to spin-up in 4U 1626–67. The source has entered a new spin-up phase with a spin-up rate of $4.02(5) \times 10^{-13} \text{ Hz s}^{-1}$. The present spin-up rate is almost half of the earlier spin-up and spin-down trends. A significant excess in soft X-ray photon emission is observed during the enhanced flux state, which is similar to the energy spectrum obtained during the spin-up era of the pulsar before 1990. 4U 1626–67 is a unique accretion powered X-ray pulsar in which quasi periodic oscillations have been consistently observed over the past ~ 20 years. But during the recent observations, we did not detect a QPO at frequencies as observed in earlier observations. Instead, we report detection of a significant broadening in the wings of the 130 mHz peak and a change in the shape of the continuum of the power spectrum. These results indicate that the flux enhancement is not a simple case of increased mass accretion rate, but there is also a change in the accretion geometry in the vicinity of the neutron star.

Key words: X-ray: Neutron Stars - pulsars: individual (4U 1626–67) - X-ray: binaries

1 INTRODUCTION

4U 1626–67 is an accretion powered X-ray pulsar which was discovered with the *Uhuru* satellite in 1972 (Giacconi et al. 1972). The 7.7s X-ray pulsations were discovered by Rappaport et al. (1977) during SAS3 observations and have been extensively monitored since then. Initially the pulsar was found to be spinning up with a characteristic timescale of ~ 5000 years. It underwent an abrupt torque reversal in 1990 and has been steadily spinning down since then (Chakrabarty et al. 1997, Krauss et al. 2007). Optical counterpart of the pulsar was identified as KZ TrA, a faint blue star with little reddening (McClintock et al. 1977; Bradt &

McClintock 1983). Pulsations at 130 mHz have also been detected from the reprocessed optical emission (Middleditch et al. 1981). Multiple sidebands separated by about 0.4 mHz, are present in the power spectrum of the optical light curve, which was interpreted as signature of an orbital period of ~ 42 min (Middleditch et al. 1981; Chakrabarty et al. 2001). But despite extensive searches, an orbital motion of the neutron star (usually observed in binary pulsars in the form of Doppler shift in the pulse period, or delay in the pulse arrival time) has never been detected in this source (Rappaport et al. 1977; Levine et al. 1988; Jain et al. 2007). It is believed that the neutron star most probably, has a low mass binary companion and is in a nearly face-on orbit.

The X-ray spectrum of 4U 1626–67 was first studied in detail with the *ASCA* spectrometers (Angelini et al. 1995)

* E-mail: chetanajain11@gmail.com

and later with instruments on board *Beppo-SAX* (Owens et al. 1997; Orlandini et al. 1998), *Chandra* (Schulz et al. 2001) and *XMM-Newton* (Krauss et al. 2007). The X-ray continuum spectrum is evidently correlated with the torque state. During the 1977–1990 spin-up phase, the spectrum was well described by an absorbed blackbody, a power law and a high energy cutoff (Pravdo et al. 1979; Kii et al. 1986). But after the torque reversal in 1990, the time-averaged X-ray spectrum was found to be relatively harder (Vaughan & Kitamoto 1997, Yi & Vishniac 1999). Orlandini et al. (1998) reported an absorption feature at ~ 37 keV which they attributed to be the cyclotron absorption line. The spectrum is rich in bright hydrogen-like and helium-like oxygen and neon emission lines. The blue and red Doppler shifted emission lines in the X-ray spectrum indicate that the lines originate in the accretion disk. However, presence of the red and blue Doppler shifted emission lines from the accretion disk is difficult to explain if the binary system is face-on. Presence of a metal rich accretion disk and absence of orbital motion led Jain et al. (2007) to suggest that 4U 1626–67 is a candidate for a neutron star with a supernova fallback accretion disk.

An important signature of the presence of an accretion disk is the detection of Quasi Periodic Oscillations (QPOs) in the power spectrum of the X-ray pulsar. QPOs are believed to arise due to inhomogeneities in the inner accretion disk. This aperiodic X-ray variability has been detected in several X-ray sources, such as, Cen X-3 (Takeshima et al. 1991; Raicher & Paul 2008), XTE J1858+034 (Paul & Rao 1998; Mukherjee et al. 2006), EXO 2030+375 (Angelini, Stella & Parmar 1989), XTE J0111.2–7317 (Kaur et al. 2007) and 3A 0535+262 (Finger, Wilson & Harmon 1996). 4U 1626–67 is a unique X-ray pulsar in which QPOs have been consistently detected in the X-ray and optical observations. The QPO frequency has been evolving over the past 22 years (Kaur et al. 2008). The QPO central frequency was found to be increasing during the spin-up phase of the pulsar, whereas, a negative trend was observed after the torque reversal. During the spin-up phase of the source, QPOs were detected at a frequency of 36 mHz with the EXOSAT data (Kaur et al. 2008) and at 40 mHz from the *Ginga* observations (Shinoda et al. 1990). During the spin-down era of the source, QPOs have been reported to occur at a higher frequency of about 48 mHz from *ASCA*, *Beppo-SAX*, *RXTE*, and *XMM-Newton* (Angelini et al. 1995; Owens et al. 1997; Kommers et al. 1998; Krauss et al. 2007). Large amplitude 0.3–1.2 mHz QPOs were also detected in reprocessed optical emission from both ground based and Hubble Space Telescope observations (Chakrabarty et al. 2001).

Recent observations with the monitoring X-ray instruments on board the *RXTE* and *Swift* satellites, show an increase in the X-ray intensity of 4U 1626–67. In the present work, we have investigated the spectral and temporal properties of 4U 1626–67 during the increased flux state. In particular, we have searched for pulsations and the QPO feature in the power spectrum of 4U 1626–67 which was persistently present in the earlier observations. We have also studied the energy spectrum of the source during the current phase and have compared it with the earlier known spectral features.




Figure 1. *RXTE*-ASM and *Swift*-BAT light curves of 4U 1626–67, binned with 30 d. The 15–50 keV *Swift*-BAT light curve is shown with a thick line over the 1.5–12 keV *RXTE*-ASM light curve (thin line). The inset figure shows the expanded view near the onset of the flux enhancement which occurred around MJD 54500 (thick line is the *Swift*-BAT light curve in the same units as in the main panel).

2 OBSERVATIONS AND ANALYSIS

4U 1626–67 is a medium intensity persistent X-ray source among the few hundred bright X-ray sources regularly monitored by the All Sky Monitor (ASM) on board *Rossi X-ray Timing Explorer (RXTE)*. Figure 1 (thin line) shows the long term 1.5–12 keV ASM light curve of 4U 1626–67, binned with 30 d. The data used here covered the time range between MJD 50088 to MJD 54909. The X-ray intensity was gradually decreasing until MJD 54500, when an increase in the X-ray flux was noticed. The inset in Figure 1 shows the expanded view near the onset of the flux enhancement. There seems to be two episodes of X-ray flux enhancement, both on timescales of ~ 100 days.

4U 1626–67 was also regularly monitored by the Burst Alert Telescope (BAT; Barthelmy et al. 2005) on board the *Swift* observatory (Gehrels et al. 2004). The long term 15–50 keV *Swift*-BAT light curve, binned with 30 d is also shown in Figure 1 (thick line) alongwith the ASM light curve. The observations covered the time range from MJD 53414 to MJD 54920. A sudden increase in the X-ray flux was first reported by Krimm et al. (2008). The inset in Figure 1 (thick line) shows the expanded view (in the same units) near the onset of the flux enhancement. The flux change on timescales of ~ 100 days is also clear in the *Swift*-BAT light curve.

We have also analysed data obtained from the Proportional Counter Array (PCA) and the High Energy X-ray Timing Explorer (HEXTE) on board the *RXTE*. The PCA consists of five xenon/methane proportional counter units (PCUs) and is sensitive in the energy range of 2–60 keV with an effective area of 1300–6500 cm², depending on the number of PCUs ON (Jahoda et al. 1996). The HEXTE operates in the energy range 15–250 keV and consists of two

Table 1. Log of *RXTE* observations analyzed in the present work

Year	Observation ID	Number of pointings	Total Exposure (ks)
1996	P10101	3	153
	P10144	1	10
	P20146	2	2
1997	P20146	12	7
1998	P30058	2	40
	P30060	7	53
2008	P93431	2	7
2009	P94423	3	9

clusters of phoswich scintillation detectors, each having a collecting area of 800 cm^2 (Gruber et al. 1996).

Table 1 gives a log of the *RXTE* observations used for the present analysis. We have studied the temporal and spectral variations in 4U 1626–67, using data collected in the Standard-1 and the Standard-2 mode of the *RXTE*-PCA, respectively. The archive mode data of the HEXTE was used for the spectral analysis.

2.1 Timing Analysis

We searched for pulsations using data obtained in the Standard 1 mode of the *RXTE*-PCA. The background counts were simulated using the *FTOOL* - *RUNPCABACKEST* and subtracted from the source light curve. The photon arrival times were converted to the solar system barycenter using the *FTOOL* - *FAXBARY*. We searched for the spin period of the neutron star using the *EFSEARCH* tool of the *HEASoft* analysis package, *FTOOLS* ver 6.5.1. This tool folds the light curve with a large number of trial periods around an approximate period and determines the best period by the χ^2 maximization technique. Using this method, we determined a pulse period of 7.67941(1) s at MJD 54530.4, 7.67945(7) s at MJD 54538.1 and 7.67848(2) at MJD 54984.5 (Jain & Paul 2009). This gives a spin-up rate of $4.02(5) \times 10^{-13} \text{ Hz s}^{-1}$. Similar behaviour based on the Fermi/GBM data has also been reported by Camero-Arranz et al. (2009).

Figure 2 shows the normalized pulse profile of 4U 1626–67 binned into 64 phasebins. The top panel shows the pulse profile generated from data obtained from *RXTE*-PCA observations made in 1996 (ObsID 10101-01-01-00) and folded with a period of 7.66735 s (Jain et al. 2007). The bottom panel shows the normalized pulse profile of 2008 data (ObsID 93431-01-01-00) folded with a period of 7.67941 s. Although the two profiles resemble the characteristic bi-horned profile of the source (Levine et al. 1988), the pulse fraction is significantly different in the two cases. Moreover, the peaks in the 2008 pulse profile are significantly sharper with a higher amplitude variation than those in the 1996 pulse profile.

We searched for QPOs in the 2–60 keV energy band, using the data collected in the Standard 1 mode of PCA, which has a time resolution of 0.125 s. We created the power density spectrum (PDS) using the *FTOOL* - *POWSPEC* for all the sufficiently long *RXTE* observations (See Table 1). The PDS were normalized such that their integral gives the squared rms fractional variability and the white noise level




Figure 2. The 2–60 keV normalized pulse profile of 4U 1626–67 generated from observations made with *RXTE*-PCA during 1996 (ObsID 10101-01-01-00) and 2008 (ObsID 93431-01-01-00). The background subtracted light curves were folded into 64 phasebins with a period of 7.66735 s and 7.67941 s, respectively.

was subtracted. Figure 3 shows some representative power spectra generated from some of the observations mentioned in Table 1. The dates of observation are mentioned in each panel above their respective observation IDs.

The narrow main peak at around 0.130 Hz corresponds to the spin frequency of the neutron star. Multiple harmonics are also seen in the PDS of the source. In addition to the main peak, a QPO feature is also seen at $\sim 48 \text{ mHz}$. However, the QPO feature is not present in the PDS generated from observations made in 2008 and 2009 (Figure 3 (9), (10), (11), (12)). In all the observations prior to 2008, the power spectrum consists of a broken power-law. As opposed to this, there is a remarkable difference in the power spectra generated from 2008 and 2009 observations.

Figure 4 shows the power spectrum generated from the 2008 observations (ObsID 93431-01-01-00; Figure 3-panel 9) overlaid with a typical power spectrum from a 1996 observation (ObsID 10101-01-01-00; Figure 3-panel 1). In the 2008 power spectrum, absence of a QPO at $\sim 48 \text{ mHz}$ is clearly seen. One more feature that stands out is the substantial broadening of the wings of the pulse peak in the power spectrum during the 2008 observation. The r.m.s variability in the low frequency noise component of the power spectrum was 14.6%, whereas, the r.m.s variability in the broad wings of the fundamental pulse peak and the fundamental pulse peak itself was 6.9% and 5.2%. This broad feature at $\sim 130 \text{ mHz}$ was never detected before and hence provides the first instance of a strong coupling between the aperiodic and periodic X-ray variability in the source.




Figure 3. Some representative X-ray power spectra of 4U 1626–67 generated from the *RXTE*-PCA observations. The date of observation is mentioned in each panel above the respective observation ID.

2.2 Spectral Analysis

The pulse-phase averaged spectra of 4U 1626–67 were generated from the data collected in the Standard 2 mode of the *RXTE*-PCA and the archive mode of the *RXTE*-HEXTE. The background counts for the PCA data were simulated using the *FTOOL* - *RUNPCABACKEST* and subtracted from the source spectrum. In case of the HEXTE data, the source and the background spectra were generated using the *FTOOL* - *HXTBACK*. These were then corrected for the detector dead time using the *FTOOL* - *HXTDEAD*. In order to bring out the significant evolution in the X-ray spectrum, we have generated the spectrum from one of the 1996 observation (ObsID 10101–01–00) and the 2008 observation (ObsID 93431–01–01–00).

Figure 5 shows the 3–20 keV and 15–50 keV energy spectrum of the source from the 1996 *RXTE*-PCA and *RXTE*-HEXTE observations. A simultaneous fit of the two spectrum consists of an intrinsic absorption, a powerlaw, a gaussian, a high energy cutoff and cyclotron resonance feature (CRF). The bottom panel of Figure 5 shows the residual of the fit. The best fit spectral parameters (with 1% systematic errors) are given in Table 2. The spectrum is similar to the broadband continuum spectrum obtained from the *Beppo*-SAX observations made after the torque reversal in 1990 (Owens et al. 1997; Orlandini et al. 1998). The *Beppo*-SAX energy spectrum was well fit by a model consisting of a low energy absorption, a blackbody, a powerlaw and a high energy cutoff. However, the best fit blackbody temperature was ~ 0.3 keV which is much lower than the sensitivity of




Figure 4. X-ray power spectra of 4U 1626–67 from the *RXTE* observations of 1996 and 2008. The upper, dotted curve is the X-ray spectrum of 1996 observation (ObsID 10101-01-01-00; Figure 3-panel 1) and the lower, solid curve is from 2008 observation (ObsID 93431-01-01-00; Figure 3-panel 9).




Figure 5. The X-ray spectrum of the 1996 *RXTE*-PCA and *RXTE*-HEXTE observations of 4U 1626–67. The model components are intrinsic absorption, powerlaw, a gaussian and a high energy cutoff. The bottom panel shows the residual of the fit.

the *RXTE*-PCA detector. We have obtained a high energy cutoff of 6.45 keV which is also quite small as compared to ~ 20 keV obtained from the *Beppo-SAX* observations.

The 3–20 keV *RXTE*-PCA and the 15–50 keV *RXTE*-HEXTE energy spectrum obtained from the 2008 observations are shown in Figure 6. The spectra were simultaneously fitted with a model consisting of a powerlaw, a blackbody, a gaussian, high energy cutoff and cyclotron resonance feature (CRF). In Table 2, we have given the best fit spectral parameters (with 1 % systematic errors). The observations made with *HEAO-1* (Pravdo et al. 1979) and X-ray astronomy satellite *Tenma* (Kii et al. 1986) during the earlier spin-up phase of 4U 1626–67 also showed similar energy spectrum. Figure 7 shows the evolution of the 3–50 keV X-ray




Figure 6. Energy spectrum of 4U 1626–67 obtained from the 2008 *RXTE*-PCA observations. The model consists of a blackbody, a powerlaw and a gaussian component.

Table 2. Best fit spectral parameters of 4U 1626–67, obtained from the *RXTE*-PCA and *RXTE*-HEXTE observations.

Parameter	Value	
	1996	2008
N_H (10^{22} cm $^{-2}$)	0.65	0.82
Photon index (Γ)	0.59(4)	0.74(7)
norm (photons keV $^{-1}$ cm $^{-2}$ s $^{-1}$)	0.0079(6)	0.012(2)
Gaussian line (keV)	6.56(40)	6.33(39)
Gaussian width (keV)	1.15(21)	1.43(38)
Gaussian norm (photons cm $^{-2}$ s $^{-1}$)	0.00054(32)	0.0019(18)
Blackbody, kT (keV)	-	0.66(12)
Blackbody, norm (km 2 kpc $^{-2}$)	-	44(7)
cutoff E (keV)	6.45775(95)	17.7(8)
foldE (keV)	32(5)	26(12)
Energy $_{(CRF)}$ (keV)	39.5(1.8)	40.3(1.2)
Width $_{(CRF)}$ (keV)	13.8(3.0)	4.8(4.5)
Depth $_{(CRF)}$	2.02(28)	2.9(1.7)
Reduced χ^2 (d.o.f.)	0.59 (57)	0.61 (43)

spectrum from 1996 observation to the 2008 observation. In the figure, the best fit spectrum of the 1996 observation (dotted curve) is shown with the spectrum generated from the 2008 observations. The bottom panel in the figure gives the ratio between the two spectra. It is clear that there is an excess soft-photon emission in the spectrum generated from the 2008 observations. The powerlaw photon index has also increased during the recent observation.

3 DISCUSSION

4U 1626–67 is a unique system in which the X-ray flux has been gradually decreasing over the past ~ 30 years. Since the discovery of X-ray pulsations in 1977, the source has shown a smooth spin evolution history. Initially, it was found to be spinning up with a characteristic timescale of 5000




Figure 7. The best fit spectrum of 4U 1626–67, generated from the 1996 (dotted line) and 2008 (solid line) *RXTE* observations. The bottom panel shows the ratio of the two spectra.

yr, but after an abrupt torque reversal, the neutron star started spinning down at about the same rate (Chakrabarty et al. 1997; Krauss et al. 2007). The recent flux enhancement has opened up many questions regarding the stability of the accretion disk. We have found a sudden increase in the X-ray flux of 4U 1626–67, which could have also caused significant changes in the geometry of the accretion column.

The power spectra of X-ray pulsars consist of narrow peaks originating from the periodic signal and its harmonic lines which are often accompanied with aperiodic features like broad bumps, wiggles and steep low frequency excess (Wijnands & van der Klis 1999). The periodic variabilities arise due to the rotation induced motion of the accretion column, through which the infalling matter is funneled towards the magnetic poles of the neutron star. Whereas, any instability in the emissivity of the accretion column, can give rise to an aperiodic variability (Burderi et al. 1997). If the inhomogeneities occur far away from the surface of the neutron star, then the periodic and aperiodic variations occur independent of each other (Angelini et al. 1989). But if the aperiodic variability occurs near the accretion column, then they are modulated at the pulsar spin frequency (Makishima et al. 1988). Lazzati & Stella (1997) found a significant coupling between the aperiodic and periodic variabilities in Vela X-1 and 4U 1145–62. A significant coupling was also reported in the accretion powered millisecond binary pulsar SAX J1808.4–3658 (Menna et al. 2003), Her X-1 (Moon & Eikenberry 2001) and Cen X-3 (Raichur & Paul 2008). The coupling between the aperiodic and periodic variations is expected to alter the shape of the power spectrum because of the convolution of the two signals (Burderi, Robba & Cusumano 1993). The coupling results in a broadening of the base of the power spectrum peaks, as if, the peaks are riding on a QPO.

X-ray spectrum and rapid X-ray variability of accreting compact objects have a common origin and are therefore expected to be correlated. In the case of 4U 1626–67, Yi & Vishniac (1999) had found that the torque reversal in 1990

was accompanied by spectral transition and a change in the luminosity. During the spin-up phase of 4U 1626–67, the phase averaged spectrum was well fit by a blackbody temperature kT of ~ 0.6 and a powerlaw photon index of ~ 1 (Pravdo et al. 1979; Kii et al. 1986). Vaughan & Kitamoto (1997) found that the spin transition had a significant effect on the 2–10 keV energy spectrum. The powerlaw photon index changed from 1.6 (Pravdo et al. 1979) to ~ 0.6 (Owens et al. 1997). The blackbody temperature also decreased from ~ 0.6 keV to ~ 0.3 keV. We have found similar spectral transition in the present high flux state of the source. The spectral parameters derived during the present high flux state are similar to those determined during the earlier spin-up phase.

ACKNOWLEDGMENTS

We thank the anonymous referee for some useful comments on improving this paper. This research has made use of the data obtained through the High Energy Astrophysics Science Archive Research Center Online Service, provided by the NASA/Goddard Space Flight Center. In particular, we thank the *RXTE*-ASM teams at MIT for provision of the ASM data.

REFERENCES

- Angelini L., Stella L., Parmar A. N., 1989, *ApJ*, 346, 906
 Angelini L., White N. E., Nagase F., Kallman T. R., Yoshida A., Takeshima T., Becker C., Paerels F., 1995 *ApJL*, 449, 41
 Barthelmy S. D., Barbier L. M., Cummings J. R. et al., 2005, *Space Science Reviews*, 120, 143
 Bradt H. V. D., McClintock J. E., 1983, *ARA&A*, 21, 13
 Burderi L., Robba N. R., Cusumano G., 1993, *Advances in Space Research*, 13, 291
 Burderi L., Robba N. R., La Barbera N., Guainazzi M., 1997, *ApJ*, 481, 943
 Camero Arranz A., Finger M. H., Ikhsanov N. R., Wilson-Hodge C. A., Beklen E., 2009, arXiv0906.4224
 Chakrabarty D., Bildsten L., Grunsfeld J. M. et al., 1997, *ApJ*, 474, 414
 Chakrabarty D., Homer L., Charles P. A., O’Donoghue D., 2001, *ApJ*, 562, 985
 Finger M. H., Wilson R. B., Harmon B. A., 1996, *ApJ*, 459, 288
 Gehrels N., Chincarini G., Giommi P. et al., 2004, *ApJ*, 611, 1005
 Giacconi R., Murray S., Gursky H., Kellogg E., Schreier E., Tananbaum H., 1972, *ApJ*, 178, 281
 Gruber D. E., Blanco P. R., Heindl W. A., Pelling M. R., Rothschild R. E., Hink P. L., 1996, *A&AS*, 120, 641
 Jahoda K., Swank J. H., Giles A. B., Stark M. J., Strohmayer T., Zhang W., Morgan E. H., 1996, *SPIE*, 2808, 59
 Jain C., Paul B., Joshi K., Dutta A., Raichur H., 2007, *JA&A*, 28, 175
 Jain C., Paul B., 2009, *ATel* 2095
 Kaur R., Paul B., Raichur H., Sagar R., 2007, *ApJ*, 660, 1409

- Kaur R., Paul B., Kumar B., Sagar, R., 2008, *ApJ*, 676, 1184
- Kii T., Hayakawa S., Nagase F., Ikegami T., Kawai N., 1986, *PASJ*, 38, 751
- Kommers J. M., Chakrabarty D., Lewin W. H. G., 1998, *ApJL*, 497, 33
- Krauss M. I., Schulz N. S., Chakrabarty D., Juett A. M., Cottam J. 2007, *ApJ*, 660, 605
- Krimm H. A., Barthelmy S. D., Baumgartner W. et al., 2008, *ATel*, 1426, 1
- Lazzati D., Stella L. 1997, *ApJ*, 476, 267
- Levine A., Ma C. P., McClintock J., Rappaport S., van der Klis M., Verbunt F., 1988, *ApJ*, 327, 732
- Makishima K., 1988, in *Physics of Neutron Stars and Black Holes*, ed. Y. Tanaka (Tokyo: Universal Acad.), 175
- McClintock J. E., van Paradijs J., Hidajat B., Hendricks H., 1977, *IAUC*, 3084, 3
- Menna M. T., Burderi L., Stella L., Robba N., van der Klis M., 2003, *ApJ*, 589, 503
- Middleditch J., Mason K. O., Nelson J. E., White N. E., 1981, *ApJ*, 244, 1001
- Moon Dae-Sik, Eikenberry S. S., 2001, *ApJL*, 552, 135
- Mukherjee U., Bapna S., Raichur H., Paul B., Jaaffrey S. N. A., 2006, *JA&A*, 27, 25
- Orlandini M., Fiume D. D., Frontera F. et al., 1998, *ApJL*, 500, 163
- Owens A., Oosterbroek T., Parmar A. N., 1997, *A&A*, 324, L9
- Paul B., Rao A. R., 1998, *A&A*, 337, 815
- Pravdo S. H., White N. E., Boldt E. A. et al., 1979, *ApJ*, 231, 912
- Raichur H., Paul B., 2008, *ApJ*, 685, 1109
- Rappaport S., Markert T., Li F. K., Clark G. W., Jernigan J. G., McClintock J. E., 1977, *ApJL*, 217, 29
- Schulz N. S., Chakrabarty D., Marshall H. L., Canizares C. R., Lee J. C., Houck J., 2001, *ApJ*, 563, 941
- Shinoda K., Kii T., Mitsuda K., Nagase F., Tanaka Y., Makishima K., Shibasaki N., 1990, *PASJ*, 42, L27
- Takeshima T., Dotani T., Mitsuda K., Nagase F., 1991, *PASJ*, 43, L43
- Vaughan B. A., Kitamoto S., 1997, *astro-ph/9707105*
- Wijnands R., van der Klis M., 1999, *ApJ*, 514, 939
- Yi I., Vishniac E. T., 1999, *ApJL*, 516, 87