

# A SHORT PROOF OF GAMAS'S THEOREM

ANDREW BERGET

ABSTRACT. If  $\chi^\lambda$  is the irreducible character of  $\mathfrak{S}_n$  corresponding to the partition  $\lambda$  of  $n$  then we may symmetrize a tensor  $v_1 \otimes \cdots \otimes v_n$  by  $\chi^\lambda$ . Gamas's theorem states that the result is not zero if and only if we can partition the set  $\{v_i\}$  into linearly independent sets whose sizes are the parts of the transpose of  $\lambda$ . We give a short and self-contained proof of this fact.

## 1. INTRODUCTION

Let  $\lambda$  be a partition of a positive integer  $n$  and let  $\chi^\lambda$  be the irreducible character of the symmetric group  $\mathfrak{S}_n$  corresponding to  $\lambda$ . There is a right action of  $\mathfrak{S}_n$  on  $V^{\otimes n}$ , where  $V$  is a finite-dimensional complex vector space, by permuting tensor positions. Let  $T_\lambda$  be the endomorphism of  $V^{\otimes n}$  given by

$$(v_1 \otimes \cdots \otimes v_n)T_\lambda = \frac{\chi^\lambda(1)}{n!} \sum_{\sigma \in \mathfrak{S}_n} \chi_\lambda(\sigma) v_{\sigma(1)} \otimes \cdots \otimes v_{\sigma(n)}.$$

Our goal is to prove the following result of Carlos Gamas [3].

**Theorem 1** (Gamas's Theorem). Let  $v_1, \dots, v_n$  be vectors in  $V$ . Then

$$(v_1 \otimes \cdots \otimes v_n)T_\lambda \neq 0$$

if and only if it is possible to partition the set  $\{v_i\}$  into linearly independent sets whose sizes are the parts of the transpose of  $\lambda$ .

If  $\{v_1, \dots, v_n\}$  is a collection of vectors satisfying the condition of the theorem we will say that it satisfies "Gamas's Condition for  $\lambda$ ". The theorem is a generalization of the well known fact that the exterior product of a set of vectors is nonzero if and only if the set of vectors is linearly independent.

In addition to Gamas's proof of this result there was a second one given by Pate in 1990 [4] using results he obtained in [5]. The benefit of our proof is that it is self-contained and short. It relies on standard facts from the representation theory of  $\mathrm{GL}(V)$ , namely, Schur-Weyl duality and the Pieri Rule. We refer to Fulton and Harris's book [2] for the needed background and notation.

## 2. PRELIMINARIES AND PROOF

Let  $V$  be a finite dimensional complex vector space. The general linear group  $\mathrm{GL}(V)$  acts diagonally on  $V^{\otimes n}$ . Let  $w \in V^{\otimes n}$  be any tensor. Define  $G(w)$  to be the  $\mathrm{GL}(V)$ -module spanned by

$$\mathrm{GL}(V)w = \{g \cdot w : g \in \mathrm{GL}(V)\}.$$

---

The author is grateful to Victor Reiner for listening to the many incarnations of this proof. His support from NSF grant DMS 0601010 was also greatly appreciated.  
<http://dx.doi.org/10.1016/j.laa.2008.09.027>.

We are interested in which irreducible  $\mathrm{GL}(V)$ -modules appear in  $G(w)$ . Since  $G(w) \subset V^{\otimes n}$  is a polynomial representation, the isomorphism type of the irreducible  $\mathrm{GL}(V)$ -modules which can appear in  $G(w)$  are indexed by partitions  $\lambda$  of  $n$  with at most  $\dim V$  parts. If  $\lambda$  is such a partition, we will say that  $\lambda$  *appears in*  $G(w)$  if this module has a highest weight vector of weight  $\lambda$  (see [2, Chapter 15]). We will write  $\ell(\lambda)$  for the number of parts of  $\lambda$ .

**Proposition 2.** If  $\lambda$  is a partition of  $n$ , then  $\lambda$  appears in  $G(w)$  if and only if  $wT_\lambda \neq 0$ .

*Proof.* Note that  $\lambda$  appears in  $G(w)$  if and only if the projection of  $G(w)$  onto its  $\lambda$ -th isotypic component is not zero. By Schur-Weyl duality (see [2, Lemma 6.22])  $T_\lambda$  is this projector, since it is the projector of  $\mathbb{C}\mathfrak{S}_n$  onto its  $\lambda$ -th isotypic component. Since  $T_\lambda$  commutes with the  $\mathrm{GL}(V)$  action, this isotypic component is zero if and only if  $G(wT_\lambda) = 0$ , which happens if and only if  $wT_\lambda = 0$ .  $\square$

The following corollary is immediate from Proposition 2.

**Corollary 3.** Suppose that  $W$  is a subspace of  $V$  and  $w \in W^{\otimes n} \subset V^{\otimes n}$ . The shape  $\lambda$  appears in  $\mathrm{span} \mathrm{GL}(V)w$  if and only if it appears in  $\mathrm{span} \mathrm{GL}(W)w$ .

*Proof of Gamas's Theorem.* Assume that the vectors  $\{v_1, \dots, v_n\}$  span  $V$ , as we may by Corollary 3. Suppose that  $\{v_i\}$  satisfy Gamas's condition for  $\lambda$ . We prove the result by induction on  $n + \ell(\lambda)$ . Write  $v^\otimes$  for the tensor  $v_1 \otimes \dots \otimes v_n$ .

If  $\lambda$  has one part  $\chi^\lambda$  is the trivial character and  $v^\otimes T_\lambda$  is a scalar multiple of the fully symmetrized tensor  $v_1 \cdots v_n$  in  $\mathrm{Sym}^n(V)$ . This is not zero since none of the  $v_i$  are zero.

If  $\ell(\lambda) < \dim V$  let  $A \in \mathrm{End}(V)$  be a generic projection to a subspace  $W \subset V$  with dimension equal to the length of  $\lambda$ . Since  $A$  is generic, the collection  $\{Av_1, \dots, Av_n\}$  still satisfies Gamas's condition for  $\lambda$ . It follows by induction that  $\lambda$  appears in the span of  $\mathrm{GL}(W)(A \cdot v^\otimes)$  and hence it also appears in  $G(A \cdot v^\otimes)$ . Since  $A$  is a limit of elements of  $\mathrm{GL}(V)$  we have  $G(A \cdot v^\otimes) \subset G(v^\otimes)$  and hence  $\lambda$  appears in  $G(v^\otimes)$ .

If  $\ell(\lambda) = \dim V$  then we consider a Young tableau of shape  $\lambda$  whose columns index independent subsets of the set  $v = \{v_1, \dots, v_n\}$ . Let  $B$  be the set of numbers in the first column of the tableau. The map

$$b_B = \sum_{\sigma \in \mathfrak{S}_B} (-1)^\sigma \sigma \in \mathbb{C}\mathfrak{S}_n = \mathrm{End}_{\mathrm{GL}(V)}(V^{\otimes n})$$

is a map of  $\mathrm{GL}(V)$ -modules and hence we have a surjection of  $\mathrm{GL}(V)$ -modules  $G(v^\otimes) \rightarrow G(v^\otimes b_B)$ . Without loss of generality, we write  $B = \{1, \dots, k\}$  where  $k = \dim V$ , so that

$$G(v^\otimes b_B) = \det_V \otimes G(v_{k+1} \otimes \dots \otimes v_n).$$

Here  $\det_V$  is the one dimensional representation  $g \mapsto \det(g)$  of  $\mathrm{GL}(V)$ . For example, if  $\dim V = 2$  and  $B = \{1, 2\}$  then

$$(v_1 \otimes v_2 \otimes v_3 \otimes v_4 \otimes v_5) b_B = (v_1 \otimes v_2 - v_2 \otimes v_1) \otimes v_3 \otimes v_4 \otimes v_5$$

Since  $v_1$  and  $v_2$  are a basis of  $V$ , we see that  $g \in \mathrm{GL}(V)$  acts by its determinant on the exterior power  $\bigwedge^2 V$  and hence on  $v_1 \wedge v_2 = v_1 \otimes v_2 - v_2 \otimes v_1$ .

Denote by  $\lambda^-$  the shape obtained from  $\lambda$  by removing the first column. Then  $\{v_{k+1}, \dots, v_n\}$  satisfies Gamas's condition for  $\lambda^-$ . By induction we know that  $\lambda^-$  appears in  $G(v_{k+1} \otimes \dots \otimes v_n)$ . By Pieri's Rule (see [2, Equation 6.9]) it follows that  $\lambda$  appears in

$$\det_V \otimes G(v_{k+1} \otimes \dots \otimes v_n)$$

and, hence, in  $G(v^\otimes)$  since it appears in its homomorphic image  $G(v^\otimes b_B)$ . This completes the more difficult implication of Gamas's Theorem.

Although our proof of the converse was essentially known to Pate [4], we include it to keep this paper self-contained. We will need the standard construction of the irreducible  $\mathrm{GL}(V)$  and  $\mathbb{C}\mathfrak{S}_n$  modules via Young symmetrizers. To this end let  $T$  be a tableau of shape  $\lambda$ ,  $a_T$  its row symmetrizer, and  $b_T$  its column antisymmetrizer. These are given by

$$\sum_{\sigma \in \mathrm{Row}(T)} \sigma, \quad \sum_{\sigma \in \mathrm{Col}(T)} \mathrm{sign}(\sigma)\sigma,$$

respectively. For example, using cycle notation for permutations in  $\mathfrak{S}_n$ , if

$$T = \begin{array}{|c|c|c|} \hline 2 & 3 & 4 \\ \hline 1 & 5 & \\ \hline \end{array}$$

then  $b_T = (1 - (12))(1 - (35))$  while

$$a_T = (1 + (23) + (24) + (34) + (234) + (243))(1 + (15)).$$

A product  $b_T a_T$  is called a *Young symmetrizer* and the right ideal in  $\mathbb{C}\mathfrak{S}_n$  generated by a Young symmetrizer is an irreducible  $\mathbb{C}\mathfrak{S}_n$ -module with character  $\chi^\lambda$  while the image of  $b_T a_T$  on  $V^{\otimes n}$  is zero, or irreducible with highest weight  $\lambda$  (see [2, Chapters 4 and 15]). It is clear that  $v^{\otimes} b_T$  is not zero if and only if the sets of vectors indexed by the columns of the tableau  $T$  are linearly independent.

It follows from Schur-Weyl duality that if  $\lambda$  appears in  $G(v^{\otimes})$  then there is an element  $c \in \mathrm{End}_{\mathrm{GL}(V)}(V^{\otimes n}) = \mathbb{C}\mathfrak{S}_n$  such that  $G(v^{\otimes} c)$  equals the irreducible  $\mathrm{GL}(V)$ -module  $V^{\otimes n} b_T a_T$ . It then follows that the right  $\mathbb{C}\mathfrak{S}_n$ -module generated by  $v^{\otimes} c$  is isomorphic to both  $c\mathbb{C}\mathfrak{S}_n$  and  $b_T a_T \mathbb{C}\mathfrak{S}_n$ , in particular the latter two modules are isomorphic. We conclude that  $c$  can be written as a sum  $c = \sum_{\sigma \in \mathfrak{S}_n} x_\sigma \sigma b_T a_T$ ,  $x_\sigma \in \mathbb{C}$ , and hence one of these terms  $x_\sigma \sigma b_T a_T$  applied to  $v^{\otimes}$  is not zero. Finally, since  $v^{\otimes} \sigma b_T$  is not zero Gamas's Condition holds for  $\lambda$ , the shape of  $T$ .  $\square$

Define a sequence of integers  $\rho_i$  by the condition that

$$\rho_1 + \cdots + \rho_k$$

is the size of the largest union of  $k$  linearly independent subsets of  $\{v_i\}$ . The sequence  $\rho$  is called the *rank partition* of  $\{v_i\}$  and was introduced by Dias da Silva in [1]. In our language, Dias da Silva proved the following strengthening of Gamas's Theorem.

**Theorem 4** (Dias da Silva). The partition  $\lambda$  appears in  $G(v^{\otimes})$  if and only if  $\lambda$  is larger (in dominance order) than the transposed rank partition of  $\{v_i\}$ .

The extent to which one can further predict the irreducible  $\mathrm{GL}(V)$ -decomposition of  $G(v^{\otimes})$  is the subject of the author's Ph.D. thesis.

#### REFERENCES

- [1] J.A. Dias da Silva. On the  $\mu$ -colorings of a matroid. *Linear and Multilinear Algebra*, 27(1):25–32, 1990.
- [2] William Fulton and Joe Harris. *Representation theory*, volume 129 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 1991. A first course, Readings in Mathematics.
- [3] Carlos Gamas. Conditions for a symmetrized decomposable tensor to be zero. *Linear Algebra Appl.*, 108:83–119, 1988.
- [4] Thomas H. Pate. Immanants and decomposable tensors that symmetrize to zero. *Linear and Multilinear Algebra*, 28(3):175–184, 1990.
- [5] Thomas H. Pate. Partitions, irreducible characters, and inequalities for generalized matrix functions. *Trans. Amer. Math. Soc.*, 325(2):875–894, 1991.

SCHOOL OF MATHEMATICS, UNIVERSITY OF MINNESOTA, MINNEAPOLIS, MN 55455  
*E-mail address:* berget@math.umn.edu