An Upper Bound on the Minimum Weight of Type II \mathbb{Z}_{2k} -Codes

Masaaki Harada*and Tsuyoshi Miezaki[†]

August 30, 2018

Abstract

In this paper, we give a new upper bound on the minimum Euclidean weight of Type II \mathbb{Z}_{2k} -codes and the concept of extremality for the Euclidean weights when k = 3, 4, 5, 6. Together with the known result, we demonstrate that there is an extremal Type II \mathbb{Z}_{2k} -code of length $8m \ (m \leq 8)$ when k = 3, 4, 5, 6.

Key Words: Type II code, Euclidean weight, extremal code, theta series 2000 *Mathematics Subject Classification*. Primary 94B05; Secondary 11F03.

1 Introduction

Let \mathbb{Z}_{2k} be the ring of integers modulo 2k, where k is a positive integer. In this paper, we take the set of elements of \mathbb{Z}_{2k} to be either $\{0, 1, \ldots, 2k-1\}$ or $\{0, \pm 1, \ldots, \pm (k-1), k\}$. A \mathbb{Z}_{2k} -code C of length n (or a code C of length n over \mathbb{Z}_{2k}) is a \mathbb{Z}_{2k} -submodule of \mathbb{Z}_{2k}^n . The Euclidean weight of a codeword x = (x_1, x_2, \ldots, x_n) is $\sum_{i=1}^n \min\{x_i^2, (2k - x_i)^2\}$. The minimum Euclidean weight $d_E(C)$ of C is the smallest Euclidean weight among all nonzero codewords of C.

^{*}Department of Mathematical Sciences, Yamagata University, Yamagata 990–8560, Japan and PRESTO, Japan Science and Technology Agency, Kawaguchi, Saitama 332–0012, Japan. email: mharada@sci.kj.yamagata-u.ac.jp. This work was partially supported by JSPS KAKENHI (20540103).

[†]Research Fellow of the Japan Society for the Promotion of Science and Department of Mathematics, Hokkaido University, Hokkaido 060–0810, Japan. email: miezaki@math.sci.hokudai.ac.jp

A \mathbb{Z}_{2k} -code C is self-dual if $C = C^{\perp}$ where the dual code C^{\perp} of C is defined as $C^{\perp} = \{x \in \mathbb{Z}_{2k}^n \mid x \cdot y = 0 \text{ for all } y \in C\}$ under the standard inner product $x \cdot y$. As described in [14], self-dual codes are an important class of linear codes for both theoretical and practical reasons. It is a fundamental problem to classify self-dual codes of modest lengths and determine the largest minimum weight among self-dual codes of that length.

A binary doubly even self-dual code is often called Type II. For \mathbb{Z}_4 -codes, Type II codes were first defined in [2] as self-dual codes containing a (± 1) vector and with the property that all Euclidean weights are divisible by eight. Then it was shown in [11] that, more generally, the condition of containing a (± 1) -vector is redundant. For general k, Type II \mathbb{Z}_{2k} -codes were defined in [1] as a self-dual code with the property that all Euclidean weights are divisible by 4k. It is known that a Type II \mathbb{Z}_{2k} -code of length n exists if and only if n is divisible by eight [1].

The aim of this paper is to show the following theorem.

Theorem 1. Let C be a Type II \mathbb{Z}_{2k} -code of length n. If $k \leq 6$ then the minimum Euclidean weight $d_E(C)$ of C is bounded by

$$d_E(C) \le 4k \left\lfloor \frac{n}{24} \right\rfloor + 4k. \tag{1}$$

Remark 2. The upper bound (1) is known for the cases k = 1 [13] and k = 2 [2]. For $k \ge 3$, the bound (1) is known under the assumption that $\lfloor n/24 \rfloor \le k-2$ [1].

We say that a Type II \mathbb{Z}_{2k} -code meeting the bound (1) with equality is *extremal* for $k \leq 6$. For the following cases

(k,m) = (4,7), (4,8), (5,6), (5,7), (5,8), (6,4), (6,5), (6,6), (6,7)and (6,8), (6,7), (

an extremal Type II \mathbb{Z}_{2k} -code of length 8m is constructed for the first time in Section 4. Together with the known results on the existences of extremal Type II codes, we have the following theorem.

Theorem 3. If $k \leq 6$ then there is an extremal Type II \mathbb{Z}_{2k} -code of length 8m for $m \leq 8$.

The existences of a binary extremal Type II code of length 72 and a 72dimensional extremal even unimodular (Type II) lattice are long-standing open questions. In this paper, we propose the following new question.

Question. Is there an extremal Type II \mathbb{Z}_{2k} -code of length 72 for $k \leq 6$?

We remark that if there is an Type II \mathbb{Z}_{2k} -code of length 72 (k = 4, 5, 6) then a 72-dimensional extremal even unimodular lattice can be obtained by Construction A.

All computer calculations in this paper were done by MAGMA [3].

2 Preliminaries

An *n*-dimensional (Euclidean) lattice Λ is a subset of \mathbb{R}^n with the property that there exists a basis $\{e_1, e_2, \ldots, e_n\}$ of \mathbb{R}^n such that $\Lambda = \mathbb{Z}e_1 \oplus \mathbb{Z}e_2 \oplus \cdots \oplus \mathbb{Z}e_n$, i.e., Λ consists of all integral linear combinations of the vectors e_1, e_2, \ldots, e_n . The dual lattice Λ^* of Λ is the lattice $\{x \in \mathbb{R}^n \mid \langle x, y \rangle \in \mathbb{Z} \text{ for all } y \in \Lambda\}$ where $\langle x, y \rangle$ is the standard inner product. A lattice with $\Lambda = \Lambda^*$ is called unimodular. The norm of a vector x is $\langle x, x \rangle$. A unimodular lattice with even norms is said to be *even*. A unimodular lattice containing a vector of odd norm is said to be *odd*. An *n*-dimensional even unimodular lattice exists if and only if $n \equiv 0 \pmod{8}$ while an odd unimodular lattice exists for every dimension. The minimum norm $\min(\Lambda)$ of Λ is the smallest norm among all nonzero vectors of Λ . For Λ and a positive integer m, the shell Λ_m of norm m is defined as $\{x \in \Lambda \mid \langle x, x \rangle = m\}$. Two lattices L and L' are *isomorphic*, denoted $L \simeq L'$, if there exists an orthogonal matrix Awith $L' = L \cdot A = \{xA \mid x \in L\}$. Two lattices L and L' are *neighbors* if both lattices contain a sublattice of index 2 in common.

The theta series $\Theta_{\Lambda}(q)$ of Λ is the following formal power series

$$\Theta_{\Lambda}(q) = \sum_{x \in \Lambda} q^{\langle x, x \rangle} = \sum_{m=0}^{\infty} |\Lambda_m| q^m.$$

For example, when Λ is the E_8 -lattice

$$\Theta_{\Lambda}(q) = E_4(q) = 1 + 240 \sum_{m=1}^{\infty} \sigma_3(m) q^{2m}$$
$$= 1 + 240q^2 + 2160q^4 + 6720q^6 + \cdots$$

where $\sigma_3(m)$ is a divisor function $\sigma_3(m) = \sum_{0 < d \mid m} d^3$. Moreover the following theorem is known (see [6, Chap. 7]).

Theorem 4. If Λ is an even unimodular lattice then

$$\Theta_{\Lambda}(q) \in \mathbb{C}[E_4(q), \Delta_{24}(q)],$$

where $\Delta_{24}(q) = q^2 \prod_{m=1}^{\infty} (1 - q^{2m})^{24}$.

We now give a method to construct even unimodular lattices from Type II codes, which is called Construction A [1]. Let ρ be a map from \mathbb{Z}_{2k} to \mathbb{Z} sending $0, 1, \ldots, k$ to $0, 1, \ldots, k$ and $k + 1, \ldots, 2k - 1$ to $1 - k, \ldots, -1$, respectively. If C is a self-dual \mathbb{Z}_{2k} -code of length n, then the lattice

$$A_{2k}(C) = \frac{1}{\sqrt{2k}} \{\rho(C) + 2k\mathbb{Z}^n\}$$

is an n-dimensional unimodular lattice, where

$$\rho(C) = \{ (\rho(c_1), \dots, \rho(c_n)) \mid (c_1, \dots, c_n) \in C \}.$$

The minimum norm of $A_{2k}(C)$ is $\min\{2k, d_E(C)/2k\}$. Moreover, if C is Type II then the lattice $A_{2k}(C)$ is an even unimodular lattice [1].

The symmetrized weight enumerator of a \mathbb{Z}_{2k} -code C is

$$\operatorname{swe}_{C}(x_{0}, x_{1}, \dots, x_{k}) = \sum_{c \in C} x_{0}^{n_{0}(c)} x_{1}^{n_{1}(c)} \cdots x_{k-1}^{n_{k-1}(c)} x_{k}^{n_{k}(c)},$$

where $n_0(c)$, $n_1(c)$, ..., $n_{k-1}(c)$, $n_k(c)$ are the numbers of $0, \pm 1, \ldots, \pm (k-1)$, k components of c, respectively [1]. Then the theta series of $A_{2k}(C)$ can be found by replacing x_1, x_2, \ldots, x_k by

$$f_0 = \sum_{x \in 2k\mathbb{Z}} q^{x^2/2k}, f_1 = \sum_{x \in 2k\mathbb{Z}+1} q^{x^2/2k}, \dots, f_k = \sum_{x \in 2k\mathbb{Z}+k} q^{x^2/2k},$$

respectively.

3 Proof of Theorem 1

In this section, we give a proof of Theorem 1. Our proof is an analogue of that of [2, Corollary 13] (see also [12]). We remark that in the proof of [2, Corollary 13] $(\Delta/E_8^3 4)$ (p. 973, right, l. -7) should be (tE_4^3/Δ) and $(4\mathbb{Z})^8/2$ (p. 973, right, l. -5) should be $2\mathbb{Z}^8$.

Proof. Let C be a Type II \mathbb{Z}_{2k} -code of length n. Then the even unimodular lattice $A_{2k}(C)$ contains the sublattice $\Lambda_0 = \sqrt{2k}\mathbb{Z}^n$ which has minimum norm 2k. We set $\Theta_{\Lambda_0}(q) = \theta_0$, n = 8j and $j = 3\mu + \nu$ ($\nu = 0, 1, 2$), that is, $\mu = \lfloor n/24 \rfloor$. In this proof, we denote $E_4(q)$ and $\Delta_{24}(q)$ by E_4 and Δ , respectively. By Theorem 4, the theta series of $A_{2k}(C)$ can be written as

$$\Theta_{A_{2k}(C)}(q) = \sum_{s=0}^{\mu} a_s E_4^{j-3s} \Delta^s = \sum_{r \ge 0} |A_{2k}(C)_r| q^r = \theta_0 + \sum_{r \ge 1} \beta_r q^r.$$

Suppose that $d_E(C) \ge 4k(\mu+1)$. We remark that a codeword of Euclidean weight 4km gives a vector of norm 2m in $A_{2k}(C)$. Then we choose the a_0, a_1, \ldots, a_μ so that

$$\Theta_{A_{2k}(C)}(q) = \theta_0 + \sum_{r \ge 2(\mu+1)} \beta_r^* q^r.$$

Here, we set b_{2s} as $E_4^{-j}\theta_0 = \sum_{s=0}^{\infty} b_{2s} (\Delta/E_4^3)^s$. That is, $\theta_0 = \sum_{s=0}^{\infty} b_{2s} E_4^{j-3s} \Delta^s$. Then

$$\sum_{s=0}^{\mu} a_s E_4^{j-3s} \Delta^s = \Theta_{A_{2k}(C)}(q) = \sum_{s=0}^{\infty} b_{2s} (\Delta/E_4^3)^s + \sum_{r \ge 2(\mu+1)} \beta_r^* q^r$$

Comparing the coefficients of q^i $(0 \le i \le 2\mu)$, we get $a_s = b_{2s}$ $(0 \le s \le \mu)$. Hence we have

$$-\sum_{r \ge (\mu+1)} b_{2r} E_4^{j-3r} \Delta^r = \sum_{r \ge 2(\mu+1)} \beta_r^* q^r.$$
(2)

In (2), comparing the coefficient of $q^{2(\mu+1)}$, we have

$$\beta_{2(\mu+1)}^* = -b_{2(\mu+1)}$$

All the series are in $q^2 = t$, and Bürman's formula shows that

$$b_{2s} = \frac{1}{s!} \frac{d^{s-1}}{dt^{s-1}} \left(\left(\frac{d}{dt} (E_4^{-j} \theta_0) \right) (t E_4^3 / \Delta)^s \right)_{\{t=0\}}$$

Using the fact that $\theta_0 = \theta_1^j$ where θ_1 is the theta series of the lattice $\sqrt{2k\mathbb{Z}^8}$,

$$b_{2s} = \frac{-j}{s!} \frac{d^{s-1}}{dt^{s-1}} \left(E_4^{3s-j-1} \theta_1^{j-1} (\theta_1 E_4' - \theta_1' E_4) (t/\Delta)^s \right)_{\{t=0\}}$$

where f' is the derivation of f with respect to $t = q^2$.

The condition that there is a codeword of Euclidean weight $4k(\mu + 1)$ is equivalent to the condition $\beta_{2(\mu+1)}^* > 0$. It is sufficient to show that the coefficients of $\theta_1^{j-1}(\theta_1 E'_4 - \theta'_1 E_4)$ are positive up to the exponent μ since E_4 and $1/\Delta$ have positive coefficients.

By Proposition 3.4 in [1], there exists a Type II \mathbb{Z}_{2k} -code of length 8 for every k. Hence let C_8 be a Type II \mathbb{Z}_{2k} -code of length 8. Then $A_{2k}(C_8)$ is the E_8 -lattice. In addition, we can write

$$E_4 = \text{swe}_{C_8}(f_0, f_1, \dots, f_k) \text{ and } \theta_1 = f_0^8$$

Deriving

$$E_4/\theta_1 = \operatorname{swe}_{C_8}(1, f_1/f_0, \dots, f_k/f_0),$$

we find

$$\theta_1^{j-1}(\theta_1 E'_4 - \theta'_1 E_4) = \frac{\partial \operatorname{swe}_{C_8}(f_0, f_1, \dots, f_k)}{\partial x_1} f_0^{8j-1}(f_0 f'_1 - f'_0 f_1) + \dots + \frac{\partial \operatorname{swe}_{C_8}(f_0, f_1, \dots, f_k)}{\partial x_k} f_0^{8j-1}(f_0 f'_k - f'_0 f_k)$$

Hence it is sufficient to show that $f_0^{8j-1}(f_0f'_1 - f'_0f_1), \ldots, f_0^{8j-1}(f_0f'_k - f'_0f_k)$ have positive coefficients up to μ . We only consider the case $f_0^{8j-1}(f_0f'_1 - f'_0f_1)$ and the other cases are similar. We have that

$$t(f_0f_1' - f_0'f_1) = \sum_{x,y \in \mathbb{Z}} \frac{(1+2ky)^2 - (2kx)^2}{4k} t^{((1+2ky)^2 + (2kx)^2)/4k},$$

then

$$tf_0^s(f_0f_1' - f_0'f_1) = \sum_{x,y,x_1,\dots,x_s \in \mathbb{Z}} \frac{(1+2ky)^2 - (2kx)^2}{4k} \cdot t^{((1+2ky)^2 + (2kx)^2 + (2kx_1)^2 + \dots + (2kx_s)^2)/4k}.$$
 (3)

Fix one of the choices $y, x, x_1, \ldots x_s \in \mathbb{Z}$ and define l as follows:

$$l = (1 + 2ky)^{2} + (2kx)^{2} + (2kx_{1})^{2} + \dots + (2kx_{s})^{2}.$$
 (4)

Consider all permutations on the set $\{x, x_1, \ldots, x_s\}$. As the sum of coefficients of $t^{l/4k}$ in the right hand side of (3) under these cases, we have that some positive constant multiple by

$$\frac{(s+1)(1+2ky)^2 - (2kx)^2 - (2kx_1)^2 - \dots - (2kx_s)^2}{4k} = \frac{(s+2)(1+2ky)^2 - l}{4k}.$$
 (5)

If l < s + 2 then (5) is positive. Since we consider the case s = 8j - 1, l < n + 1. Hence if the exponent l/4k of t is less than (n + 1)/4k then (5) is positive. This means that if $\mu < (n + 1)/4k$ then (5) is positive. This condition $\mu < (n + 1)/4k$ is satisfied since $k \leq 6$. Thus for any choice y, $x, x_1, \ldots x_s$, (5) is positive. The coefficient of $t^{l/4k}$ in the right hand side of (3) is the sum of those coefficients (5), that is, positive. This completes the proof of Theorem 1.

4 Extremal Type II \mathbb{Z}_{2k} -codes

An extremal Type II \mathbb{Z}_{2k} -code of length 8m is currently known for the cases (k, m) listed in the second column in Table 1. In this section, an extremal Type II \mathbb{Z}_{2k} -code of length 8m is constructed for the first time for the cases (k, m) listed in the last column in Table 1.

Table 1: Existence of extremal Type II \mathbb{Z}_{2k} -codes of length 8m

k	m (known cases)			m (new cases)
1	$1, 2, \ldots, 8, 10, 11, 13, 14, 17$	[6, p. 194], [8]		
2	$1, 2, \dots, 8$	[2], [4], [9]		
3	$1, 2, \dots, 8$	[5], [10]		
4	$1, 2, \ldots, 6$	[5], [7]	7, 8	$(C_{8,56}, C_{8,64})$
5	$1, 2, \ldots, 5$	[5], [7]	6, 7, 8	(Proposition 6, $C_{10,56}$, $C_{10,64}$)
6	1, 2, 3	[5]	4,5,6,7,8	(Proposition 7)

Let A and B be $n\times n$ negacirculant matrices, that is, A and B have the following form

$$\begin{pmatrix} r_0 & r_1 & r_2 & \cdots & r_{n-1} \\ -r_{n-1} & r_0 & r_1 & \cdots & r_{n-2} \\ -r_{n-2} & -r_{n-1} & r_0 & \cdots & r_{n-3} \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ -r_1 & -r_2 & -r_3 & \cdots & r_0 \end{pmatrix}$$

If $AA^T + BB^T = -I_n$, then it is trivial that

$$\begin{pmatrix} & A & B \\ & I_{2n} & -B^T & A^T \end{pmatrix}$$
(6)

generates a self-dual code where I_n denotes the identity matrix of order n and A^T is the transpose of A.

Codes	r_A	r_B
$C_{8,56}$	(0, 0, 4, 3, 4, 1, 6, 3, 1, 1, 1, 1, 1, 2)	(1, 1, 0, 0, 0, 0, 0, 3, 2, 0, 0, 0, 0, 4)
$C_{8,64}$	(0, 0, 0, 2, 0, 7, 3, 2, 0, 0, 5, 3, 1, 4, 0, 2)	(0, 0, 1, 0, 0, 0, 0, 1, 7, 1, 3, 0, 1, 2, 2, 0)
$C_{10,56}$	(0, 0, 0, 2, 5, 1, 4, 1, 2, 0, 5, 0, 4, 1)	(0, 0, 0, 0, 0, 1, 0, 5, 7, 0, 3, 9, 1, 0)
$C_{10,64}$	(0, 0, 4, 3, 2, 0, 0, 1, 9, 0, 0, 0, 9, 1, 2, 0)	(1, 3, 0, 1, 0, 6, 9, 4, 6, 2, 0, 5, 0, 0, 2, 3)
$C_{12,32}$	(0, 0, 7, 6, 0, 1, 7, 10)	(0, 1, 0, 4, 1, 0, 3, 11)
$C_{12,40}$	(0, 0, 0, 2, 1, 10, 5, 9, 2, 10)	(0, 1, 0, 1, 0, 0, 11, 1, 0, 4)
$C_{12,56}$	(2, 11, 2, 2, 4, 11, 0, 5, 0, 0, 6, 1, 5, 7)	(1, 0, 5, 3, 0, 8, 0, 2, 0, 7, 7, 0, 0, 4)

Table 2: New extremal Type II \mathbb{Z}_{2k} -codes

Using the above construction method, we have found extremal Type II \mathbb{Z}_8 -codes $C_{8,56}$ and $C_{8,64}$ of lengths 56 and 64, respectively, and extremal Type II \mathbb{Z}_{10} -codes $C_{10,56}$ and $C_{10,64}$ of lengths 56 and 64, respectively. The

first rows r_A and r_B of the matrices A and B in their generator matrices (6) are listed in Table 2. Hence we have the following:

Proposition 5. For lengths 56 and 64, there is an extremal Type II \mathbb{Z}_{2k} -code when k = 4 and 5.

An *n*-dimensional even unimodular lattice is called *extremal* if it has minimum norm $2\lfloor n/24 \rfloor + 2$. The existence of an extremal Type II \mathbb{Z}_{10} -code of length 48 is established by considering the existence of a 10-frame in some extremal even unimodular lattice. Recall that a set $\{f_1, \ldots, f_n\}$ of *n* vectors f_1, \ldots, f_n in an *n*-dimensional unimodular lattice *L* with $\langle f_i, f_j \rangle = \ell \delta_{i,j}$ is called an ℓ -frame of *L*, where $\delta_{i,j}$ is the Kronecker delta. It is known that an even unimodular lattice *L* contains a 2*k*-frame if and only if there is a Type II \mathbb{Z}_{2k} -code *C* such that $A_{2k}(C) \simeq L$.

Proposition 6. There is an extremal Type II \mathbb{Z}_{10} -code of length 48.

Proof. Let $C_{5,48}$ be the \mathbb{F}_5 -code with generator matrix (6) where the first rows r_A and r_B of the matrices A and B are

 $r_A = (2, 3, 0, 2, 2, 3, 2, 2, 3, 2, 2, 0)$ and $r_B = (3, 0, 4, 4, 0, 1, 0, 0, 4, 0, 0, 1)$,

respectively. Then this code $C_{5,48}$ is a self-dual code and the lattice $A_5(C_{5,48}) = \frac{1}{\sqrt{5}} \{x \in \mathbb{Z}^{48} \mid x \pmod{5} \in C_{5,48}\}$ is an odd unimodular lattice. The lattice has theta series $1 + 393216q^5 + 26201600q^6 + \cdots$. We have verified that $A_5(C_{5,48})$ has an even unimodular neighbor L_{48} which is extremal. Clearly the lattice $A_5(C_{5,48})$ contains the 5-frame $\{\sqrt{5}e_1, \sqrt{5}e_2, \ldots, \sqrt{5}e_{48}\}$ where e_i $(i = 1, 2, \ldots, 48)$ denotes the *i*-th unit vector $(\delta_{i,1}, \delta_{i,2}, \ldots, \delta_{i,48})$ of length 48. Then the set $F = \{\sqrt{5}(e_{2i-1} \pm e_{2i}) \mid i = 1, 2, \ldots, 24\}$ is a 10-frame of the even sublattice of $A_5(C_{5,48})$. Hence F is also a 10-frame of the extremal even unimodular neighbor L_{48} . Therefore there is a Type II \mathbb{Z}_{10} -code $C_{10,48}$ of length 48 such that $A_{10}(C_{10,48}) \simeq L_{48}$. Moreover, the code $C_{10,48}$ must be extremal since the lattice L_{48} is extremal.

Similar to the above proposition, the existence of 12-frames in extremal even unimodular lattices yields that of some extremal Type II \mathbb{Z}_{12} -codes.

Proposition 7. There is an extremal Type II \mathbb{Z}_{12} -code of length 8m for m = 4, 5, 6, 7 and 8.

Proof. It is known that there is an extremal Type II \mathbb{Z}_6 -code of length 8m for m = 4, 5, 6, 7 and 8 (see Table 1). We denote these codes by $C_{6,8m}$ (m = 4, 5, 6, 7 and 8), respectively. Since $C_{6,8m}$ is an extremal Type II code, the lattice $A_6(C_{6,8m})$ is an extremal even unimodular lattice for

m = 4, 5, 6, 7 and 8. Moreover, clearly the lattice $A_6(C_{6,8m})$ contains the 6-frame $\{\sqrt{6}e_1, \sqrt{6}e_2, \ldots, \sqrt{6}e_{8m}\}$ where e_i denotes the *i*-th unit vector of length 8m. Then the set $\{\sqrt{6}(e_{2i-1} \pm e_{2i}) \mid i = 1, 2, \ldots, 4m\}$ is a 12-frame of $A_6(C_{6,8m})$. Hence there is a Type II \mathbb{Z}_{12} -code N_{8m} of length 8m such that $A_{12}(N_{8m}) \simeq A_6(C_{6,8m})$. Moreover, the code N_{8m} must be extremal since the lattice $A_6(C_{6,8m})$ is extremal.

Remark 8. Similar to Proposition 5, by considering generator matrices (6), we have found extremal Type II \mathbb{Z}_{12} -codes $C_{12,32}$, $C_{12,40}$ and $C_{12,56}$ of lengths 32, 40 and 56, respectively where the first rows r_A and r_B of the matrices A and B in (6) are listed in Table 2.

Together with the known results on the existences of extremal Type II codes (see Table 1), Propositions 5, 6 and 7 give Theorem 3.

Acknowledgment. The authors would like to thank Kenichiro Tanabe for useful discussions, and Koichi Betsumiya for helpful conversations on the construction of extremal Type II codes.

References

- E. Bannai, S.T. Dougherty, M. Harada and M. Oura, Type II codes, even unimodular lattices, and invariant rings, *IEEE Trans. Inform. Theory* 45 (1999), 1194–1205.
- [2] A. Bonnecaze, P. Solé, C. Bachoc and B. Mourrain, Type II codes over Z₄, IEEE Trans. Inform. Theory 43 (1997), 969–976.
- [3] W. Bosma and J. Cannon, Handbook of Magma Functions, Department of Mathematics, University of Sydney, Available online at http://magma.maths.usyd.edu.au/magma/.
- [4] A.R. Calderbank and N.J.A. Sloane, Double circulant codes over \mathbb{Z}_4 and even unimodular lattices, J. Algebraic Combin. 6 (1997), 119–131.
- [5] R. Chapman, Double circulant constructions of the Leech lattice, J. Aust. Math. Soc. Ser. A 69 (2000), 287–297.
- [6] J.H. Conway and N.J.A. Sloane, Sphere Packing, Lattices and Groups (3rd ed.), Springer-Verlag, New York, 1999.
- [7] T.A. Gulliver and M. Harada, Extremal self-dual codes over Z₆, Z₈ and Z₁₀, AKCE Int. J. Graphs Comb. 2 (2005), 11−24.

- [8] M. Harada, An extremal doubly even self-dual code of length 112, Electron. J. Combin. 15 (2008), Note 33, 5 pp.
- [9] M. Harada, Extremal Type II Z₄-codes of lengths 56 and 64, (submitted).
- [10] M. Harada, M. Kitazume and M. Ozeki, Ternary code construction of unimodular lattices and self-dual codes over Z₆, J. Algebraic Combin. 16 (2002), 209–223.
- [11] M. Harada, P. Solé and P. Gaborit, Self-dual codes over \mathbb{Z}_4 and unimodular lattices: a survey, Algebras and combinatorics (Hong Kong, 1997), 255–275, Springer, Singapore, 1999.
- [12] C.L. Mallows, A.M. Odlyzko and N.J.A. Sloane, Upper bounds for modular forms, lattices, and codes, J. Algebra 36 (1975), 68–76.
- [13] C.L. Mallows and N.J.A. Sloane, An upper bound for self-dual codes, Inform. Control 22 (1973), 188–200.
- [14] E. Rains and N.J.A. Sloane, "Self-dual codes," Handbook of Coding Theory, V.S. Pless and W.C. Huffman (Editors), Elsevier, Amsterdam 1998, pp. 177–294.