

An algebraic geometric model of an action of the face monoid associated to a Kac-Moody group on its building

Claus Mokler

Abstract

We described in [M1] a monoid \widehat{G} , the face monoid, acting on the integrable irreducible highest weight modules of a symmetrizable Kac-Moody algebra. It has similar structural properties as a reductive algebraic monoid whose unit group is a Kac-Moody group G . We found in [M5] two natural extensions of the action of the Kac-Moody group G on its building Ω to actions of the face monoid \widehat{G} on the building Ω . Now we give an algebraic geometric model of one of these actions of the face monoid \widehat{G} on Ω , where the building Ω is obtained as a part of the \mathbb{F} -valued points of the spectrum of all homogeneous prime ideals of the Cartan algebra CA of the Kac-Moody group G . We describe the spectrum of all homogeneous prime ideals of the Cartan algebra CA and determine its \mathbb{F} -valued points.

Introduction

The face monoid \widehat{G} is an infinite-dimensional algebraic monoid. It has been obtained in [M1] by a Tannaka reconstruction from categories determined by the integrable irreducible highest weight representations of a symmetrizable Kac-Moody algebra. By its construction and by the involved categories it is a very natural object. Its Zariski open dense unit group G coincides, up to a slightly extended maximal torus, with the special Kac-Moody group defined representation theoretically in [KP1].

The face monoid is a purely infinite-dimensional phenomenon, quite unexpected. In the classical case, i.e., if you take a split semisimple Lie algebra for the symmetrizable Kac-Moody algebra, it coincides with a split semisimple simply connected algebraic group. Put in another way, there seem to exist fundamentally different infinite-dimensional generalizations of a split semisimple simply connected algebraic group.

The results obtained in [M1], [M2], [M3], and [M4] show that the face monoid \widehat{G} has similar structural and algebraic geometric properties as a reductive algebraic monoid, e.g., the monoid of $(n \times n)$ -matrices. The face monoid \widehat{G} is the first example of an infinite-dimensional reductive algebraic monoid. Actually, it is particular. The investigation of the conjugacy classes in [M6] will show, that the relation between the face monoid \widehat{G} and its unit group G , the Kac-Moody group, is much closer than for a general reductive algebraic monoid.

Obviously, there is the following question: Does the face monoid \widehat{G} fit in some way to the building theory of the Kac-Moody group G ? In [M5] we investigated how to extend the natural action of the Kac-Moody group G on its building Ω to actions of the face monoid \widehat{G} on Ω . To explain the results obtained in [M5] note the following facts:

For the face monoid \widehat{G} an infinite Renner monoid \widehat{W} plays the same role as the Weyl group W does for the Kac-Moody group G . For example, there are Bruhat and Birkhoff decompositions of \widehat{G} , similar as for G , but the Weyl group W replaced by the monoid \widehat{W} . The monoid \widehat{W} can be constructed from the Weyl group W and the face lattice of the Tits cone X , where the term "face" means a face of the convex cone X in the sense of convex geometry. The Weyl group W is the unit group of \widehat{W} .

The building Ω is covered by certain subcomplexes, the apartments, which are isomorphic to the Coxeter complex \mathcal{C} associated to the Weyl group W . This connects the action of G on Ω and the action of W on \mathcal{C} .

Now let \mathcal{A} be the standard apartment associated to a fixed BN-pair (B, N) of the Kac-Moody group G . The group N may be obtained as normalizer $N = N_G(T)$ of the standard torus $T = B \cap N$, and the Weyl group \mathcal{W} identifies with N/T . If we define similarly $\widehat{N} := N_{\widehat{G}}(T)$, then the monoid $\widehat{\mathcal{W}}$ identifies with \widehat{N}/T . Consider the following diagram:

$$\begin{array}{ccccc}
\Omega & \supset & \mathcal{A} & \cong & \mathcal{C} \\
\circlearrowleft & & \circlearrowleft & & \circlearrowleft \\
G & \supset & N & \twoheadrightarrow & \mathcal{W} \\
\cap & & \cap & & \cap \\
\widehat{G} & \supset & \widehat{N} & \twoheadrightarrow & \widehat{\mathcal{W}}
\end{array} \tag{1}$$

Suppose it is given an action of $\widehat{\mathcal{W}}$ on \mathcal{C} , extending the natural action of \mathcal{W} on \mathcal{C} . Then this action induces uniquely an action of \widehat{N} on \mathcal{A} , extending the natural action of N on \mathcal{A} . There are the following questions: Does there exist an action of \widehat{G} on Ω , extending the natural action of G on Ω , such that diagram (1) commutes? If such an action exists, is it uniquely determined? Which actions can be obtained in this way?

In Theorem 45 and Corollary 48 of [M5] we obtained the following beautiful result: There is a bijective correspondence between:

- (i) The actions of $\widehat{\mathcal{W}}$ on \mathcal{C} , extending the natural action of \mathcal{W} on \mathcal{C} , satisfying certain conditions which we do not state here.
- (ii) The actions of \widehat{G} on Ω , extending the natural action of G on Ω .

Furthermore, the action of \widehat{G} on Ω is obtained by an explicit formula from the action of $\widehat{\mathcal{W}}$ on \mathcal{C} .

There exist quite general actions of \widehat{G} on Ω , extending the natural action of G on Ω . Compare for example the action of Remark 46 in [M5]. Is it possible to single out some actions of \widehat{G} on Ω , which extend the natural action of G on Ω and also keep some of its properties? The natural action of G on the building Ω satisfies:

- (a) It preserves the natural order on Ω .
- (b) $\text{Stab}_G(Q) = Q$ for all standard parabolic subgroups $Q \in \Omega$.

Now every standard parabolic subgroup Q of G extends naturally to a standard parabolic submonoid \widehat{Q} of \widehat{G} . An action of \widehat{G} on Ω , which extends the natural action of G on Ω , is called good if it satisfies:

- (a) It preserves the natural order on Ω .
- (b) $\text{Stab}_{\widehat{G}}(Q) = \widehat{Q}$ for all standard parabolic subgroups $Q \in \Omega$.

It is open to determine all good actions. In [M5] we found two good actions of \widehat{G} , good action 1 and good action 2. Both have been obtained by the correspondence mentioned above from actions of $\widehat{\mathcal{W}}$ on \mathcal{C} , which in turn have been found by an action of $\widehat{\mathcal{W}}$ on the facets of the Tits cone and by an action of $\widehat{\mathcal{W}}$ on Looijenga's modified Tits cone. It worked perfectly. But if you look at the formulas of these actions in Corollary 50 and Corollary 51 of [M5], you would like to have some natural models of these actions, if it is possible.

In this paper we give a natural algebraic geometric model of good action 1 of \widehat{G} . It is obtained as follows: Let P^+ be the set of dominant weights of the weight lattice P . Let $L(\Lambda)$ be the irreducible highest weight module of highest weight Λ , let $L(\Lambda)^{(*)}$ be its restricted dual. The Cartan algebra

$$CA = \bigoplus_{\Lambda \in P^+} L(\Lambda)^{(*)}$$

is a P^+ -graded algebra over the field \mathbb{F} of characteristic zero, whose multiplication on the graded parts

$$L(\Lambda_1)^{(*)} \otimes L(\Lambda_2)^{(*)} \rightarrow L(\Lambda_1 + \Lambda_2)^{(*)}$$

is obtained dually to G -equivariant embeddings $L(\Lambda_1 + \Lambda_2) \rightarrow L(\Lambda_1) \otimes L(\Lambda_2)$, $\Lambda_1, \Lambda_2 \in P^+$.

The spectrum $\text{Proj}(CA)$ of all P^+ -graded prime ideals, no irrelevant ideals excluded, is a locally ringed space. It is no scheme, but it is stratified by schemes.

The building Ω , which is as a set the union

$$\Omega = \bigcup_{Q \text{ standard parabolic}} G/Q$$

of all thin flag spaces, identifies with a dense part of the \mathbb{F} -valued points $\text{Proj}(CA)(\mathbb{F})$ of this spectrum, i.e., there exists a natural injection

$$\Omega \hookrightarrow \text{Proj}(CA)(\mathbb{F})$$

with dense image. A natural completion Ω_{fn} of the building Ω , which is as a set the union of all thick flag spaces, identifies with the \mathbb{F} -valued points $\text{Proj}(CA)(\mathbb{F})$ itself.

Now the face monoid \widehat{G} acts on every module $L(\Lambda)$, $\Lambda \in P^+$. Dually we get actions of the opposite monoid \widehat{G}^{op} on the restricted duals $L(\Lambda)^{(*)}$, $\Lambda \in P^+$, which fit together to an action of \widehat{G}^{op} on the Cartan algebra CA by morphisms of graded algebras. This in turn induces actions of \widehat{G} on $\text{Proj}(CA)$ and $\text{Proj}(CA)(\mathbb{F})$ by morphisms. The image of Ω in $\text{Proj}(CA)(\mathbb{F})$ is invariant under the action of \widehat{G} . Pulling back the action of the face monoid \widehat{G} to the building Ω gives good action 1.

This algebraic geometric model of an action of the face monoid \widehat{G} on the building Ω has been found as follows. V. Kac reviews in Section 2.6 of [K2] the construction of the flag space G/B from [KP1]. It uses the Kostant cone $\mathcal{V}_\Lambda := G(L(\Lambda)_\Lambda)$, the G -orbit of the highest weight space $L(\Lambda)_\Lambda$ of $L(\Lambda)$, where Λ is chosen arbitrarily from the set of strongly dominant weights P^{++} . V. Kac suggests on page 199, line 3 and 4, to take $\text{Proj}(CA)$ as a definition of the flag space G/B , independent of the choice of $\Lambda \in P^{++}$. Now the author had obtained different algebraic actions of the face monoid \widehat{G} on the building Ω , and was looking for everything linked to algebraic geometric actions of the face monoid on flag spaces. As shown in [M1], Proposition 5.4, the face monoid acts naturally on the Kostant cones \mathcal{V}_Λ , $\Lambda \in P^+$. These actions are related to an action of the face monoid \widehat{G} on the building Ω , which does not preserve its natural order, the bad action of [M5]. The author tried $\text{Proj}(CA)$, which led to this article.

A Cartan algebra can be obtained for every normal reductive algebraic monoid. It should be possible to investigate its spectrum of homogeneous prime ideals and the action of the reductive algebraic monoid on this spectrum similarly as in this article. Maybe also results of [M5] generalize in some way. As far as the author knows, actions of reductive algebraic monoids on the buildings of their unit groups or on building-like objects associated to their unit groups have not been investigated up to now. It may be interesting to do it.

1 Preliminaries

In this section we collect some facts about Kac-Moody algebras, minimal Kac-Moody groups, its associated face monoids, and formal Kac-Moody groups. It recalls the representation theoretic constructions of these groups and monoids. It is a reference for the notation used in this article.

The facts on Kac-Moody algebras, which we state in this section, can be found in the books [K1], [Ku], and [Mo,Pi], most results of [K1] and [Ku] also valid for a field of characteristic zero with the same proofs. The facts on minimal Kac-Moody groups in [KP1], [KP2], [KP3], [Ré], and [Mo,Pi]. The facts on formal Kac-Moody groups in [Ku], [Ré], and [Sl], most results of [Sl] and [Ku] also valid for a field of characteristic zero with the same proofs. The facts on face monoids in [M1] and [M5].

We denote by $\mathbb{N} = \mathbb{Z}^+$, \mathbb{Q}^+ , resp. \mathbb{R}^+ the sets of strictly positive numbers of \mathbb{Z} , \mathbb{Q} , resp. \mathbb{R} , and the sets $\mathbb{N}_0 = \mathbb{Z}_0^+$, \mathbb{Q}_0^+ , \mathbb{R}_0^+ contain, in addition, the zero. \mathbb{F} is a field of characteristic 0 and \mathbb{F}^\times its group of units.

The starting data: The starting data for the representation theoretic construction of the simply connected minimal and formal Kac-Moody groups and its face monoids are:

- A symmetrizable generalized Cartan matrix $A = (a_{ij})_{i,j \in I}$ with finite index set $I := \{1, 2, \dots, n\}$. We denote by l the rank of A .
- A *simply connected minimal free root base* for A consisting of:
 - Dual free \mathbb{Z} -modules H, P of rank $2n - l$.
 - Linear independent sets $\{h_1, \dots, h_n\} \subseteq H$, $\{\alpha_1, \dots, \alpha_n\} \subseteq P$, such that $\alpha_i(h_j) = a_{ji}$, $i, j = 1, \dots, n$.
 - Furthermore, there exist $\Lambda_1, \dots, \Lambda_n \in P$ such that $\Lambda_i(h_j) = \delta_{ij}$, $i, j = 1, \dots, n$.

P is called the *weight lattice*, $P^+ := \{\Lambda \in P \mid \Lambda(h_i) \geq 0 \text{ for all } i \in I\}$ the set of *dominant weights* of P . $Q := \mathbb{Z}\text{-span}\{\alpha_i \mid i \in I\}$ is called the *root lattice*. Set $Q_0^\pm := \mathbb{Z}_0^\pm\text{-span}\{\alpha_i \mid i \in I\}$ and $Q^\pm := Q_0^\pm \setminus \{0\}$. The height function $ht : Q \rightarrow \mathbb{Z}$ is defined by $ht(\sum_{i \in I} n_i \alpha_i) := \sum_{i \in I} n_i$.

$\{\alpha_i \mid i \in I\}$ is called the set of *simple roots*. Note: To cut short many formulas we often identify this set with the index set I .

We fix a system of elements $\Lambda_1, \dots, \Lambda_n \in P$ such that $\Lambda_i(h_j) = \delta_{ji}$, $i, j = 1, \dots, n$. We extend to dual bases $h_1, \dots, h_{2n-l} \in H$ and $\Lambda_1, \dots, \Lambda_{2n-l} \in P$. We call $\Lambda_1, \dots, \Lambda_{2n-l}$ a system of *fundamental dominant weights* of P .

Let $J \subseteq I$. Set $A_J := (a_{ij})_{i,j \in J}$, which is a symmetrizable generalized Cartan matrix if J is nonempty. Set $P_J := \mathbb{Z}\text{-span}\{\Lambda_i \mid i \in J\}$ and $P_J^+ := \mathbb{Z}_0^+\text{-span}\{\Lambda_i \mid i \in J\}$. Set $Q_J := \mathbb{Z}\text{-span}\{\alpha_i \mid i \in J\}$ and $(Q_J)_0^\pm := \mathbb{Z}_0^\pm\text{-span}\{\alpha_i \mid i \in J\}$ and $Q_J^\pm := (Q_J)_0^\pm \setminus \{0\}$. Set $P_{rest} := \mathbb{Z}\text{-span}\{\Lambda_i \mid i = n+1, \dots, 2n-l\}$. As always, a span of the empty set is defined to be $\{0\}$.

The linear spaces \mathbf{h} and \mathbf{h}^* : Define the \mathbb{F} -linear spaces

$$\mathbf{h} := \mathbf{h}_{\mathbb{F}} := H \otimes_{\mathbb{Z}} \mathbb{F} \quad \text{and} \quad \mathbf{h}^* = \mathbf{h}_{\mathbb{F}}^* := P \otimes_{\mathbb{Z}} \mathbb{F}.$$

Identify H and P with the corresponding sublattices of \mathbf{h} and \mathbf{h}^* . Interpret \mathbf{h}^* as the dual of \mathbf{h} . Order the elements of \mathbf{h}^* by $\lambda \leq \lambda'$ if and only if $\lambda' - \lambda \in Q_0^+$.

For $J \subseteq I$ set $\mathbf{h}_J := \text{span}\{h_i \mid i \in J\}$. Set $\mathbf{h}_{rest} := \text{span}\{h_i \mid i = n+1, \dots, 2n-l\}$. In the same way as in §2.1 of [K1] equip \mathbf{h} with a non-degenerate symmetric bilinear form (\mid) adapted to the decomposition $\mathbf{h} = \mathbf{h}_J \oplus \mathbf{h}_{rest}$. Denote by $\nu : \mathbf{h} \rightarrow \mathbf{h}^*$ be the induced linear isomorphism. Denote also by (\mid) the induced non-degenerate symmetric bilinear form on \mathbf{h}^* .

The Weyl group \mathcal{W} : The *Weyl group* $\mathcal{W} = \mathcal{W}(A)$ is the Coxeter group with generators σ_i , $i \in I$, and relations

$$\sigma_i^2 = 1 \quad (i \in I) \quad , \quad (\sigma_i \sigma_j)^{m_{ij}} = 1 \quad (i, j \in I, i \neq j),$$

where the m_{ij} are given by:

$a_{ij}a_{ji}$	0	1	2	3	≥ 4
m_{ij}	2	3	4	6	no relation between σ_i and σ_j

Denote by $l : \mathcal{W} \rightarrow \mathbb{N}_0$ the *length function*. Denote by $\mathcal{W}_J \cong \mathcal{W}(A_J)$, $J \subseteq I$, the *standard parabolic subgroups* of \mathcal{W} . For $J, K \subseteq I$ denote by \mathcal{W}^J the set of minimal coset representatives of $\mathcal{W}/\mathcal{W}_J$, by ${}^K\mathcal{W}$ the set of minimal coset representatives of $\mathcal{W}_K \backslash \mathcal{W}$, and by ${}^K\mathcal{W}^J$ the set of minimal double coset representatives of $\mathcal{W}_K \backslash \mathcal{W}/\mathcal{W}_J$.

The Tits cone X : The Weyl group \mathcal{W} acts faithfully \mathbf{h}^* by

$$\sigma_i \lambda := \lambda - \lambda(h_i) \alpha_i \quad i \in I, \quad \lambda \in \mathbf{h}^*.$$

The lattices Q and P are left invariant by this action.

The *Tits cone* X is a convex \mathcal{W} -invariant cone in $\mathbf{h}_{\mathbb{R}}^*$ obtained by

$$X := \mathcal{W}\overline{C} \quad \text{with} \quad \overline{C} := \{\lambda \in \mathbf{h}_{\mathbb{R}}^* \mid \lambda(h_i) \geq 0 \text{ for all } i \in I\}.$$

Every \mathcal{W} -orbit of X contains exactly one point of the (closed) *standard fundamental chamber* \overline{C} . Now we recall shortly:

1. A \mathcal{W} -invariant partition of X given by the interiors of polyhedral cones, which we call the facets of X .
2. The set $\text{Fa}(X)$ of faces of the convex cone X in the sense of convex geometry. It is a \mathcal{W} -space, partially ordered by the inclusion of sets, even a complete lattice.

To 1. For $J \subseteq I$ set

$$\begin{aligned} F_J &:= \{ \lambda \in \mathbf{h}_{\mathbb{R}}^* \mid \lambda(h_i) = 0 \text{ for } i \in J, \lambda(h_i) > 0 \text{ for } i \in I \setminus J \}, \\ \overline{F_J} &:= \{ \lambda \in \mathbf{h}_{\mathbb{R}}^* \mid \lambda(h_i) = 0 \text{ for } i \in J, \lambda(h_i) \geq 0 \text{ for } i \in I \setminus J \} = \bigcup_{K \supseteq J} F_K. \end{aligned}$$

In particular, $\overline{C} = \overline{F_{\emptyset}}$. Here $\overline{F_J}$ is a polyhedral cone with relative interior F_J . Its faces in the sense of convex geometry are $\overline{F_K}$, $K \supseteq J$. We call F_J resp. $\overline{F_J}$ an open resp. closed *standard facet* of type J .

The parabolic subgroup \mathcal{W}_J of \mathcal{W} is the stabilizer of any element $\lambda \in F_J$. It is the stabilizer of F_J as well as of $\overline{F_J}$ as a whole.

The set $\{\sigma F_J \mid \sigma \in \mathcal{W}, J \subseteq I\}$ is a \mathcal{W} -invariant partition of X . Partially ordered by the reverse closure relation it gives the Coxeter complex of \mathcal{W} . We call σF_J resp. $\sigma \overline{F_J}$ an open resp. closed *facet* of type J .

To 2. For $\emptyset \neq \Theta \subseteq I$ we denote by Θ^0 , resp. Θ^∞ the set of indices which correspond to the sum of the components of the generalized Cartan submatrix A_Θ of finite, resp. non-finite type. We set $\emptyset^0 := \emptyset^\infty := \emptyset$. For $J \subseteq I$ set $J^\perp := \{i \in I \mid a_{ij} = 0 \text{ for all } j \in J\}$.

A set $\Theta \subseteq I$ such that $\Theta = \Theta^\infty$ is called *special*. If Θ is a special set then

$$R(\Theta) := \mathcal{W}_{\Theta^\perp} \overline{F_\Theta} = \{ \lambda \in X \mid \lambda(h_i) = 0 \text{ for all } i \in \Theta \}$$

is a face of X with relative interior

$$ri(R(\Theta)) = \mathcal{W}_{\Theta^\perp} \bigcup_{J \subseteq \Theta^\perp, J=J^0} F_{\Theta \cup J}.$$

The \mathcal{W} -stabilizers of $R(\Theta)$, pointwise and as a whole, are

$$Z_{\mathcal{W}}(R(\Theta)) = \mathcal{W}_\Theta \quad \text{and} \quad N_{\mathcal{W}}(R(\Theta)) = \mathcal{W}_{\Theta \cup \Theta^\perp}.$$

It holds $\{R(\Theta) \mid \Theta \text{ special}\} = \{R \in \text{Fa}(X) \mid ri(R) \cap \overline{C} \neq \emptyset\}$. Furthermore, it holds

$$\text{Fa}(X) = \bigcup_{\Theta \text{ special}} \{ \sigma R(\Theta) \mid \sigma \in \mathcal{W} \}.$$

The special set Θ is called the *type* of the face $\sigma R(\Theta)$, $\sigma \in \mathcal{W}$.

Let $\sigma, \sigma' \in \mathcal{W}$, Θ, Θ' be special. It holds

$$\sigma' R(\Theta') \subseteq \sigma R(\Theta) \iff \Theta' \supseteq \Theta \text{ and } \sigma^{-1} \sigma' \in \mathcal{W}_{\Theta^\perp} \mathcal{W}_{\Theta'}.$$

Different faces of X of the same type are not comparable by \subseteq .

A well defined function $red : \mathcal{W} \rightarrow I$ is obtained as follows: Set $red(1) := \emptyset$. If $\sigma \in \mathcal{W} \setminus \{1\}$ and $\sigma_{i_1} \cdots \sigma_{i_k}$ is a reduced expression for σ set $red(\sigma) := \{i_1, \dots, i_k\} \subseteq I$.

The lattice intersection, which coincides with the set theoretical intersection, and the lattice join of two arbitrary faces can be reduced easily by the formulas for the stabilizers to the following formulas: Let $\tau \in {}_{\Theta_1 \cup \Theta_1^\perp} \mathcal{W}^{\Theta_2 \cup \Theta_2^\perp}$, Θ_1, Θ_2 be special. It holds

$$\begin{aligned} R(\Theta_1) \cap \tau R(\Theta_2) &= R(\Theta_1 \cup \Theta_2 \cup red(\tau)), \\ R(\Theta_1) \vee \tau R(\Theta_2) &= R((\Theta_1 \cap \tau \Theta_2)^\infty). \end{aligned}$$

The Kac-Moody algebra \mathfrak{g} : The *Kac-Moody algebra* $\mathfrak{g} = \mathfrak{g}(A)$ is the Lie algebra over \mathbb{F} generated by the abelian Lie algebra \mathfrak{h} , and the elements $e_i, f_i, (i \in I)$, with the following relations, which hold for any $i, j \in I, h \in \mathfrak{h}$:

$$\begin{aligned} [e_i, f_j] &= \delta_{ij} h_i, \quad [h, e_i] = \alpha_i(h) e_i, \quad [h, f_i] = -\alpha_i(h) f_i, \\ (ad e_i)^{1-\alpha_{ij}} e_j &= (ad f_i)^{1-\alpha_{ij}} f_j = 0 \quad (i \neq j). \end{aligned}$$

The abelian Lie algebra \mathfrak{h} and the elements $e_i, f_i, (i \in I)$, identify with their images in \mathfrak{g} .

There is the *root space decomposition*

$$\mathfrak{g} = \bigoplus_{\alpha \in \mathfrak{h}^*} \mathfrak{g}_\alpha \quad \text{where} \quad \mathfrak{g}_\alpha := \{x \in \mathfrak{g} \mid [h, x] = \alpha(h)x \text{ for all } h \in \mathfrak{h}\}.$$

In particular, it holds $\mathfrak{g}_0 = \mathfrak{h}$, $\mathfrak{g}_{\alpha_i} = \mathbb{F}e_i$, and $\mathfrak{g}_{-\alpha_i} = \mathbb{F}f_i, i \in I$. The set of *roots* $\Delta := \{\alpha \in \mathfrak{h}^* \setminus \{0\} \mid \mathfrak{g}_\alpha \neq 0\}$ satisfies $\Delta \subseteq Q, \Delta = -\Delta$, and it is invariant under the Weyl group. $\Delta_{re} := \mathcal{W}\{\alpha_i \mid i \in I\}$ is the set of *real roots*. $\Delta_{im} := \Delta \setminus \Delta_{re}$ is the set of *imaginary roots*. Every root space $\mathfrak{g}_\alpha, \alpha \in \Delta$, is finite dimensional.

Δ, Δ_{re} , and Δ_{im} decompose into the disjoint union of the sets of *positive* and *negative roots* $\Delta^\pm := \Delta \cap Q^\pm, \Delta_{re}^\pm := \Delta_{re} \cap Q^\pm, \Delta_{im}^\pm := \Delta_{im} \cap Q^\pm$. There is the corresponding *triangular decomposition*

$$\mathfrak{g} = \mathfrak{n}^- \oplus \mathfrak{h} \oplus \mathfrak{n}^+ \quad \text{where} \quad \mathfrak{n}^\pm := \bigoplus_{\alpha \in \Delta^\pm} \mathfrak{g}_\alpha.$$

For every $\alpha \in \Delta_{re}$ the subalgebra $\mathfrak{g}_\alpha \oplus [\mathfrak{g}_\alpha, \mathfrak{g}_{-\alpha}] \oplus \mathfrak{g}_{-\alpha}$ of \mathfrak{g} is isomorphic to $sl(2, \mathbb{F})$.

The non-degenerate symmetric bilinear form (\mid) on \mathfrak{h} extends uniquely to a non-degenerate symmetric invariant bilinear form (\mid) on \mathfrak{g} . For $\alpha, \beta \in \Delta \cup \{0\}$ such that $\alpha + \beta \neq 0$ it holds $(\mathfrak{g}_\alpha \mid \mathfrak{g}_\beta) = 0$. The restriction $(\mid) : \mathfrak{g}_\alpha \times \mathfrak{g}_{-\alpha} \rightarrow \mathbb{F}$ is nondegenerate, and it holds $[x, y] = (x \mid y)\nu^{-1}(\alpha)$ for all $x \in \mathfrak{g}_\alpha, y \in \mathfrak{g}_{-\alpha}, \alpha \in \Delta$.

Let $J \subseteq I$. Set

$$\mathfrak{g}_J := \mathfrak{h}_J \oplus \bigoplus_{\alpha \in \Delta_J} \mathfrak{g}_\alpha = \mathfrak{n}_J^- \oplus \mathfrak{h}_J \oplus \mathfrak{n}_J^+ \quad \text{where} \quad \mathfrak{n}_J^\pm := \bigoplus_{\alpha \in \Delta_J^\pm} \mathfrak{g}_\alpha$$

and $\Delta_J := \Delta \cap Q_J, \Delta_J^\pm := \Delta^\pm \cap Q_J$. Then $\mathfrak{g}_J \cong \mathfrak{g}(A_J)'$ for $J \neq \emptyset$. In particular, \mathfrak{g}_I is the derived Lie algebra of \mathfrak{g} .

An ideal of \mathfrak{n}^\pm is obtained by

$$(\mathfrak{n}^\pm)^J := \bigoplus_{\alpha \in (\Delta^\pm)^J} \mathfrak{g}_\alpha \quad \text{with} \quad (\Delta^\pm)^J := \Delta^\pm \setminus \Delta_J^\pm.$$

The (minimal) Kac-Moody group G : A \mathfrak{g} -module V is called *integrable* if V is \mathfrak{h} -diagonalizable with set of weights $P(V) \subseteq P$ and the elements of \mathfrak{g}_α act locally nilpotent on V for all $\alpha \in \Delta_{re}$. Examples of integrable representations are the adjoint representation \mathfrak{g} , the irreducible highest weight representations $L(\Lambda)$ with highest weights $\Lambda \in P^+$, and the irreducible lowest weight representations $L(\Lambda)^{(*)}$ with lowest weights $-\Lambda \in -P^+$.

Let \mathcal{I} be the category whose objects are the integrable \mathfrak{g} -modules and whose morphisms are the morphisms of \mathfrak{g} -modules. This category generalizes the category of locally finite dimensional representations of a semisimple Lie algebra.

Let $Nat(\mathcal{I})$ be the set of natural transformations of the forgetful functor from the category \mathcal{I} into the category of \mathbb{F} -linear spaces. Explicitly, $Nat(\mathcal{I})$ consists of the families of linear maps

$$m = (m_V \in End(V))_{V \text{ obj. of } \mathcal{I}},$$

such that for all objects V, W , and all morphisms $\phi : V \rightarrow W$ of \mathcal{I} the diagram

$$\begin{array}{ccc} V & \xrightarrow{m_V} & V \\ \phi \downarrow & & \downarrow \phi \\ W & \xrightarrow{m_W} & W \end{array}$$

commutes. $Nat(\mathcal{I})$ gets in the obvious way the structure of an associative \mathbb{F} -algebra with unit. In particular, there are the following elements of $Nat(\mathcal{I})$:

- (1) For every $h \in H$, $s \in \mathbb{F}^\times$ there exists $t_h(s) \in Nat(\mathcal{I})$, such that for every object V of \mathcal{I} it holds

$$t_h(s)v_\lambda = s^{\lambda(h)}v_\lambda \quad , \quad v_\lambda \in V_\lambda \quad , \quad \lambda \in P(V).$$

- (2) For every $x \in \mathfrak{g}_\alpha$, $\alpha \in \Delta_{re}$, there exists $exp(x) \in Nat(\mathcal{I})$, such that for every object V of \mathcal{I} it holds

$$exp(x)v = \sum_{k \in \mathbb{N}_0} \frac{x^k v}{k!} \quad , \quad v \in V.$$

The *minimal Kac-Moody group* G , which we call *Kac-Moody group* for short, is the submonoid of $Nat(\mathcal{I})$ generated by the elements of (1) and (2). Its elements are compatible with \oplus , \otimes , act as identity on the trivial representation, and induce dual maps on the integrable duals, i.e., on the biggest integrable submodules of the full duals. The Kac-Moody group G has the following important structural properties:

A root group data system $(T, (U_\alpha)_{\alpha \in \Delta_{re}})$ of G is obtained as follows: The elements of (1) induce an embedding of the torus $H \otimes_{\mathbb{Z}} \mathbb{F}^\times = Hom((P, +), (\mathbb{F}^\times, \cdot))$ into G , whose image is T . For every $\alpha \in \Delta_{re}$ the elements of (2) induce an embedding of $(\mathfrak{g}_\alpha, +)$ into G , whose image is the *root group* U_α .

The normalizer $N := N_G(T)$ is generated by T and the elements $n_i := exp(e_i)exp(-f_i)exp(e_i)$, $i \in I$. An isomorphism $\kappa : \mathcal{W} \rightarrow N/T$ is induced by $\kappa(\sigma_i) := n_i T$, $i \in I$.

We denote by $\widetilde{\cdot} : \mathcal{W} \rightarrow N$ the cross section of the canonical map from N to \mathcal{W} defined by $\widetilde{1} = 1$, $\widetilde{\sigma}_i = n_i$, $i \in I$, and $\widetilde{\sigma\tau} = \widetilde{\sigma}\widetilde{\tau}$ if $l(\sigma\tau) = l(\sigma) + l(\tau)$, $\sigma, \tau \in \mathcal{W}$.

We denote an arbitrary element $n \in N$ with $\kappa^{-1}(nT) = \sigma \in \mathcal{W}$ by n_σ . The set of weights $P(V)$ of an integrable \mathfrak{g} -module V is \mathcal{W} -invariant, and it holds $n_\sigma V_\lambda = V_{\sigma\lambda}$, $\lambda \in P(V)$, $\sigma \in \mathcal{W}$.

Let U^\pm be the subgroups of G generated by U_α , $\alpha \in \Delta_{re}^\pm$. Then U^\pm are normalized by T . The pairs $(B^\pm := T \ltimes U^\pm, N)$ are twin BN-pairs of G with the property $B^+ \cap B^- = B^\pm \cap N = T$. There are the *Bruhat* and *Birkhoff decompositions*

$$G = \dot{\bigcup}_{\sigma \in N} U^\epsilon n U^\delta = \dot{\bigcup}_{\sigma \in \mathcal{W}} B^\epsilon \sigma B^\delta \quad , \quad \epsilon, \delta \in \{+, -\}.$$

There are also *Levi decompositions* of the standard parabolic subgroups P_J^\pm , $J \subseteq I$, and of U^\pm :

$$P_J^\pm = L_J^\pm \ltimes (U^\pm)^J \quad \text{and} \quad U^\pm = U_J^\pm \ltimes (U^\pm)^J.$$

Here U_J^\pm is the group generated by U_α , $\alpha \in (\Delta_J^\pm)_{re} := \Delta_J^\pm \cap \Delta_{re}$. $(U^\pm)^J$ is the smallest normal subgroup of U^\pm containing U_α , $\alpha \in (\Delta^\pm)^J_{re} := (\Delta^\pm)^J \cap \Delta_{re}$. This group equals $\bigcap_{\sigma \in \mathcal{W}_J} \sigma U^\pm \sigma^{-1}$. Furthermore, $L_J^\pm := U_J^\pm (T \mathcal{W}_J) U_J^\pm$.

For $J \subseteq I$ let G_J be the subgroup of G generated by U_α , $\alpha \in (\Delta_J)_{re} := \Delta_J \cap \Delta_{re}$. It holds $G_\emptyset = \{1\}$ and $G_J \cong G(A_J)'$ for $J \neq \emptyset$. The torus $T_J := G_J \cap T$ is generated by the subtori $t_{h_i}(\mathbb{F}^\times)$, $i \in J$. The group $N_J := G_J \cap N$ is generated by T_J and the elements n_i , $i \in J$.

In particular, G_I is the derived group of G , which identifies with the Kac-Moody group as defined in [KP1]. It holds $G = G_I \rtimes T_{rest}$, where T_{rest} is the subtorus of T generated by the subtori $t_{h_i}(\mathbb{F}^\times)$, $i = n+1, \dots, 2n-l$.

The adjoint action ad of the Kac-Moody algebra \mathfrak{g} on itself is integrable. Therefore, the Kac-Moody group G acts on \mathfrak{g} . This action is called the *adjoint action* Ad of G .

Some stabilizers which are often used: Let $J \subseteq I$ and $\Lambda \in P^+ \cap F_J$. Let T^J be the subtorus of T generated by $t_{h_i}(\mathbb{F}^\times)$, $i \in \{1, 2, \dots, 2n-l\} \setminus J$. Set

$$Y_\Lambda := \left\{ \prod_{i=1, \dots, 2n-l, i \notin J} t_i(s_i) \in T \mid \prod_{i=1, \dots, 2n-l, i \notin J} (s_i)^{\Lambda(h_i)} = 1 \right\} \subseteq T^J.$$

It holds

$$\begin{aligned} N_G(L(\Lambda)_\Lambda) &:= \{g \in G \mid gL(\Lambda)_\Lambda = L(\Lambda)_\Lambda\} = P_J = U^J G_J T^J = U N_J T^J U, \\ Z_G(L(\Lambda)_\Lambda) &:= \{g \in G \mid gv = v \text{ for all } v \in L(\Lambda)_\Lambda\} = U^J G_J Y_\Lambda = U N_J Y_\Lambda U. \end{aligned}$$

The face monoid \widehat{G} associated to the Kac-Moody group G : The category \mathcal{O}^p is defined as follows: Its objects are the \mathfrak{g} -modules V , which have the properties:

1. V is \mathbf{h} -diagonalizable with finite dimensional weight spaces.
2. There exist finitely many elements $\lambda_1, \dots, \lambda_m \in \mathbf{h}^*$, such that the set of weights $P(V)$ of V is contained in the union $\bigcup_{i=1}^m \{\lambda \in \mathbf{h}^* \mid \lambda \leq \lambda_i\}$.

The morphisms of \mathcal{O}^p are the morphisms of \mathfrak{g} -modules.

Let \mathcal{O}_{int}^p the full subcategory of the category \mathcal{O}^p , whose objects are integrable modules. This category generalizes the category of finite dimensional representations of a semisimple Lie algebra, keeping the complete reducibility theorem: Every object of \mathcal{O}_{int}^p is isomorphic to a direct sum of the integrable irreducible highest weight modules $L(\Lambda)$, $\Lambda \in P^+$. Denote the set of weights of $L(\Lambda)$ by $P(\Lambda)$, $\Lambda \in P^+$. The set of weights of every object of \mathcal{O}_{int}^p is contained in $X \cap P$ because of $\bigcup_{\Lambda \in P^+} P(\Lambda) = X \cap P$.

Let V be an object of \mathcal{O}_{int}^p and $V = \bigoplus_{\lambda \in P(V)} V_\lambda$ its weight space decomposition. We may associate two \mathbb{F} -linear duals, the restricted dual $V^{(*)}$ and the full dual V^* as follows

$$V^{(*)} := \bigoplus_{\lambda \in P(V)} V_\lambda^* \subseteq V^* = \prod_{\lambda \in P(V)} V_\lambda^*.$$

Let $Nat(\mathcal{O}_{int}^p)$ be the set of natural transformations of the forgetful functor from the category \mathcal{O}_{int}^p into the category of \mathbb{F} -linear spaces. In particular, there are the following elements of the \mathbb{F} -algebra $Nat(\mathcal{O}_{int}^p)$:

(1) For every $h \in H$, $s \in \mathbb{F}^\times$ there exists $t_h(s) \in Nat(\mathcal{O}_{int}^p)$, such that for every object V of \mathcal{O}_{int}^p it holds

$$t_h(s)v_\lambda = s^{\lambda(h)}v_\lambda \quad , \quad v_\lambda \in V_\lambda \quad , \quad \lambda \in P(V).$$

(2) For every $x \in \mathfrak{g}_\alpha$, $\alpha \in \Delta_{re}$, there exists $exp(x) \in Nat(\mathcal{O}_{int}^p)$, such that for every object V of \mathcal{O}_{int}^p it holds

$$exp(x)v = \sum_{k \in \mathbb{N}_0} \frac{x^k v}{k!} \quad , \quad v \in V.$$

(3) For every $R \in Fa(X)$ there exists $e(R) \in Nat(\mathcal{O}_{int}^p)$, such that for every object V of \mathcal{O}_{int}^p it holds

$$e(R)v_\lambda = \begin{cases} v_\lambda & \lambda \in R \\ 0 & \lambda \in X \setminus R \end{cases} \quad , \quad v_\lambda \in V_\lambda \quad , \quad \lambda \in P(V).$$

The *face monoid* \widehat{G} is the submonoid of $Nat(\mathcal{O}_{int}^p)$ generated by the elements of (1), (2), and (3). It is a Tannaka monoid: It is the biggest part of $Nat(\mathcal{O}_{int}^p)$, whose elements are compatible with \oplus , \otimes , act as identity on the trivial representation, and induce dual maps on the restricted duals.

The unit group of \widehat{G} is generated by the elements of (1) and (2). It is isomorphic to the Kac-Moody group G in the obvious way, and for simplicity of notation we identify it with G .

Let Θ be special. There are the following decompositions of certain parabolic subgroups of G :

$$\begin{aligned} P_{\Theta \cup \Theta^\perp} &= \left(G_{\Theta^\perp} \rtimes T^{\Theta \cup \Theta^\perp} \right) \ltimes \left(G_\Theta \rtimes U^{\Theta \cup \Theta^\perp} \right). \\ P_{\Theta \cup \Theta^\perp}^- &= \left((U^-)^{\Theta \cup \Theta^\perp} \rtimes G_\Theta \right) \ltimes \left(G_{\Theta^\perp} \rtimes T^{\Theta \cup \Theta^\perp} \right). \end{aligned}$$

The projections belonging to these semidirect products are denoted by $p_{\Theta}^{\pm} : P_{\Theta \cup \Theta^{\perp}}^{\pm} \rightarrow G_{\Theta^{\perp}} \rtimes T^{\Theta \cup \Theta^{\perp}}$. Now the monoid \widehat{G} may be described algebraically as follows: It holds

$$\widehat{G} = \bigcup_{\Theta \text{ special}} Ge(R(\Theta))G.$$

Let $g_1, g_2, h_1, h_2 \in G$, let Θ_1, Θ_2 be special. It holds $g_1 e(R(\Theta_1))h_1 = g_2 e(R(\Theta_2))h_2$ if and only if $\Theta_2 = \Theta_1$ and

$$(g_2)^{-1}g_1 \in P_{\Theta_1 \cup \Theta_1^{\perp}}, \quad h_2(h_1)^{-1} \in P_{\Theta_1 \cup \Theta_1^{\perp}}^{-} \quad \text{with} \quad p_{\Theta_1}((g_2)^{-1}g_1) = p_{\Theta_1}^{-}(h_2(h_1)^{-1}).$$

To describe the multiplication of $g_1 e(R(\Theta_1))h_1$ and $g_2 e(R(\Theta_2))h_2$ write $h_1 g_2 = a_1 \tilde{\sigma} a_2$ with $a_1 \in P_{\Theta_1 \cup \Theta_1^{\perp}}^{-}$, $a_2 \in P_{\Theta_2 \cup \Theta_2^{\perp}}$, and $\sigma \in {}_{\Theta_1 \cup \Theta_1^{\perp}}\mathcal{W}^{\Theta_2 \cup \Theta_2^{\perp}}$. It holds

$$g_1 e(R(\Theta_1))h_1 g_2 e(R(\Theta_2))h_2 = g_1 p_{\Theta_1}^{-}(a_1) e(R(\Theta_1 \cup \Theta_2 \cup \text{red}(\sigma))) p_{\Theta_2}(a_2) h_2.$$

The monoid $\widehat{N} := N_{\widehat{G}}(T) := \{\hat{g} \in \widehat{G} \mid \hat{g}T = T\hat{g}\}$ is generated by N and $\{e(R) \mid R \in \text{Fa}(X)\}$. There are the Bruhat and Birkhoff decompositions

$$\widehat{G} = \bigcup_{\hat{n} \in \widehat{N}} U^{\epsilon} \hat{n} U^{\delta} = \bigcup_{\hat{w} \in \widehat{N}/T} B^{\epsilon} \hat{w} B^{\delta} \quad , \quad \epsilon, \delta \in \{+, -\}.$$

The Weyl group \mathcal{W} acts on the monoid $(\text{Fa}(X), \cap)$. The semidirect product $\mathcal{W} \ltimes \text{Fa}(X)$ consists of the set $\mathcal{W} \times \text{Fa}(X)$ on which the structure of a monoid is given by

$$(\sigma, R) \cdot (\tau, S) := (\sigma\tau, \tau^{-1}R \cap S).$$

A congruence relation on $\mathcal{W} \ltimes \text{Fa}(X)$ is obtained by

$$(\sigma, R) \sim (\sigma', R') \quad : \Longleftrightarrow \quad R = R' \quad \text{and} \quad \sigma^{-1}\sigma' \in Z_{\mathcal{W}}(R).$$

We denote the congruence class of (σ, R) by $\sigma\epsilon(R)$. Now the monoid \widehat{N}/T is isomorphic to the monoid $\widehat{\mathcal{W}} := (\mathcal{W} \ltimes \text{Fa}(X))/\sim$, which we call the *face monoid* associated to \mathcal{W} , by

$$\begin{array}{ccc} \kappa : & \widehat{\mathcal{W}} & \rightarrow \widehat{N}/T \\ & \sigma\epsilon(R) & \mapsto \tilde{\sigma}e(R)T \end{array}$$

The Weyl group \mathcal{W} is the unit group of $\widehat{\mathcal{W}}$. There are standard parabolic submonoids $\widehat{\mathcal{W}}_J \cong \widehat{\mathcal{W}}(A_J)$, $J \subseteq I$, of $\widehat{\mathcal{W}}$. These give the standard parabolic submonoids $\widehat{P}_J := B\widehat{\mathcal{W}}_J B$, $J \subseteq I$, of \widehat{G} .

The formal Kac-Moody group G_{fn} : The Lie bracket of \mathfrak{g} extends in the obvious way to a Lie bracket of

$$\mathfrak{g}_{fn} := \mathfrak{n}_f^{-} \oplus \mathfrak{h} \oplus \mathfrak{n} \quad \text{with} \quad \mathfrak{n}_f^{-} := \prod_{\alpha \in \Delta^{-}} \mathfrak{g}_{\alpha}.$$

This Lie algebra is called the *formal Kac-Moody algebra*. The Lie subalgebra \mathfrak{n}_f^{-} is a pronilpotent Lie algebra with (non-complete) defining system of ideals $\mathfrak{n}_f^{-}(k) := \prod_{\alpha \in \Delta^{-}, -ht(\alpha) \geq k+1} \mathfrak{g}_{\alpha}$, $k \in \mathbb{N}_0$.

Define the category \mathcal{O}^n similarly as the category \mathcal{O}^p , but replace \leq by \geq in the second condition. Let \mathcal{O}_{int}^n the full subcategory of the category \mathcal{O}^n , whose objects are integrable modules. This category generalizes the category of finite dimensional representations of a semisimple Lie algebra, keeping the complete reducibility theorem: Every object of \mathcal{O}_{int}^n is isomorphic to a direct sum of the integrable irreducible lowest weight modules $L(\Lambda)^{(*)}$ of lowest weights $-\Lambda \in -P^{+}$.

For every object V of \mathcal{O}_{int}^n the action of the Kac-Moody algebra \mathfrak{g} on V extends in the obvious way to an action of the formal Kac-Moody algebra \mathfrak{g}_{fn} on V .

Let $Nat(\mathcal{O}_{int}^n)$ be the set of natural transformations of the forgetful functor from the category \mathcal{O}_{int}^n into the category of \mathbb{F} -linear spaces. In particular, there are the following elements of the \mathbb{F} -algebra $Nat(\mathcal{O}_{int}^n)$:

(1) For every $h \in H$, $s \in \mathbb{F}^\times$ there exists $t_h(s) \in Nat(\mathcal{O}_{int}^n)$, such that for every object V of \mathcal{O}_{int}^n it holds

$$t_h(s)v_\lambda = s^{\lambda(h)}v_\lambda \quad , \quad v_\lambda \in V_\lambda \quad , \quad \lambda \in P(V).$$

(2) For every $x \in \mathbf{n}_f^- \cup \bigcup_{\alpha \in \Delta_{re}^+} \mathbf{g}_\alpha$ there exists $exp(x) \in Nat(\mathcal{O}_{int}^n)$, such that for every object V of \mathcal{O}_{int}^n it holds

$$exp(x)v = \sum_{k \in \mathbb{N}_0} \frac{x^k v}{k!} \quad , \quad v \in V.$$

The *formal Kac-Moody group* G_{fn} is the submonoid of $Nat(\mathcal{O}_{int}^n)$ generated by the elements of (1) and (2). Its elements are compatible with \oplus , \otimes , act as identity on the trivial representation, and induce dual maps now on the full duals. The formal Kac-Moody group G_{fn} has the following important structural properties:

$U_f^- := exp(\mathbf{n}_f^-)$ is a subgroup of G_{fn} . As a group U_f^- is the pronipotent group with pronilpotent Lie algebra \mathbf{n}_f^- and exponential map $exp : \mathbf{n}_f^- \rightarrow U_f^-$.

Now certain subsets of Δ^- induce pronipotent subgroups and pronipotent normal subgroups of U_f^- . Certain decompositions induce multiplicative decompositions of U_f^- and of pronipotent subgroups of U_f^- . In this article we use many times: For $J \subseteq I$ the decomposition $\Delta^- = \Delta_J^- \dot{\cup} (\Delta^-)^J$ induces the semidirect decomposition

$$\mathbf{n}_f^- = (\mathbf{n}_f^-)_J \ltimes (\mathbf{n}_f^-)^J \quad \text{with} \quad (\mathbf{n}_f^-)_J := \prod_{\alpha \in \Delta_J^-} \mathbf{g}_\alpha \quad \text{and} \quad (\mathbf{n}_f^-)^J := \prod_{\alpha \in (\Delta^-)^J} \mathbf{g}_\alpha ,$$

which in turn induces the semidirect decomposition

$$U_f^- = (U_f^-)_J \ltimes (U_f^-)^J \quad \text{with} \quad (U_f^-)_J := exp((\mathbf{n}_f^-)_J) \quad \text{and} \quad (U_f^-)^J := exp((\mathbf{n}_f^-)^J).$$

The formal Kac-Moody group G_{fn} contains in the obvious way a subgroup isomorphic the Kac-Moody group G , and for simplicity of notation we identify it with G .

It holds $G_{fn} = U_f^- G = G U_f^-$. Furthermore, $G_{fn} = G$ if and only if $U_f^- = U^-$ if and only if the generalized Cartan matrix A contains only components of finite type.

U_f^- is normalized by T and $(B_f^- := T U_f^-, N)$ is a BN-pair of G_{fn} with $B_f^- \cap N = T$. More generally, G_{fn} , U_f^- , U , N , T , $\{\sigma_i \in \mathcal{W} \cong N/T \mid i \in I\}$ is a refined Tits system. (To adapt the notation used for refined Tits systems in [KP3], Section 2, or in [Ku], Section 5.2 to our situation, the groups U^+ and U^- in the axioms have to be interchanged.) In particular, there are the *Bruhat* and *Birkhoff decompositions*

$$G_{fn} = \bigcup_{\sigma \in \mathcal{W}} B_f^- \sigma B_f^- = \bigcup_{\sigma \in \mathcal{W}} B_f^\epsilon \sigma B_f^- = \bigcup_{\sigma \in \mathcal{W}} B_f^- \sigma B_f^\epsilon \quad , \quad \epsilon \in \{+, -\}.$$

There are *Levi decompositions* of the standard parabolic subgroups

$$(P_{fn}^-)_J = (L_{fn})_J \ltimes (U_f^-)^J$$

with $(L_{fn})_J = (U_f^-)_J (\mathcal{W}_J T) (U_f^-)_J = (U_f^-)_J L_J = L_J (U_f^-)_J$, $J \subseteq I$.

For $J \subseteq I$ let $(G_{fn})_J$ be the subgroup of G_{fn} generated by $(U_f^-)_J$ and U_α , $\alpha \in (\Delta_J^+)_{re}$. It holds $(G_{fn})_\emptyset = \{1\}$ and $(G_{fn})_J \cong G_{fn}(A_J)'$ for $J \neq \emptyset$.

The formal Kac-Moody group G_{fn} acts on the formal Kac-Moody algebra \mathbf{g}_{fn} by the *adjoint action* Ad . On the generators (1) and (2) of G_{fn} it is given as follows. For $h \in H$ and $s \in \mathbb{F}^\times$ it is

$$Ad(t_h(s))y = \sum_{\alpha \in \Delta \cup \{0\}} s^{\alpha(h)} y_\alpha \quad \text{where} \quad y = \sum_{\alpha \in \Delta \cup \{0\}} y_\alpha \in \mathbf{g}_{fn}.$$

For $x \in \mathfrak{n}_f^- \cup \bigcup_{\alpha \in \Delta_{re}^+} \mathfrak{g}_\alpha$ it is

$$Ad(exp(x))y = exp(ad(x))y = \sum_{k \in \mathbb{N}_0} \frac{1}{k!} (ad(x))^k y \quad \text{where } y \in \mathfrak{g}_{fn}.$$

This action extends the adjoint action Ad of G on \mathfrak{g} .

It is important to note: If V is a \mathfrak{g} -module contained in \mathcal{O}_{int}^p , then the restricted dual \mathfrak{g} -module $V^{(*)}$ is contained in \mathcal{O}_{int}^n . The Kac-Moody group G acts on V and $V^{(*)}$ dually. The formal Kac-Moody group G_{fn} only acts on

$$V_f := \prod_{\lambda \in P(V)} V_\lambda = \prod_{\lambda \in P(V)} (V_\lambda^*)^* = (V^{(*)})^*$$

and $V^{(*)}$ dually, where V_λ has been identified canonically with $(V_\lambda^*)^*$, $\lambda \in P(V)$. The G_{fn} -module V_f extends the G -module V . The linear subspace V of V_f is G -invariant, but in general not G_{fn} -invariant.

Note also: In the literature it is more common to consider the formal Kac-Moody algebra

$$\mathfrak{g}_{fp} := \mathfrak{n}^- \oplus \mathfrak{h} \oplus \mathfrak{n}_f \quad \text{with} \quad \mathfrak{n}_f := \prod_{\alpha \in \Delta^+} \mathfrak{g}_\alpha.$$

For every object V of \mathcal{O}_{int}^p the action of the Kac-Moody algebra \mathfrak{g} on V extends to an action of the formal Kac-Moody algebra \mathfrak{g}_{fp} on V . Now the *formal Kac-Moody group* G_{fp} is the submonoid of $Nat(\mathcal{O}_{int}^p)$ generated by the elements given in (1) and (2):

(1) For every $h \in H$, $s \in \mathbb{F}^\times$ there exists $t_h(s) \in Nat(\mathcal{O}_{int}^p)$, such that for every object V of \mathcal{O}_{int}^p it holds

$$t_h(s)v_\lambda = s^{\lambda(h)}v_\lambda \quad , \quad v_\lambda \in V_\lambda \quad , \quad \lambda \in P(V).$$

(2) For every $x \in \mathfrak{n}_f \cup \bigcup_{\alpha \in \Delta_{re}^-} \mathfrak{g}_\alpha$ there exists $exp(x) \in Nat(\mathcal{O}_{int}^p)$, such that for every object V of \mathcal{O}_{int}^p it holds

$$exp(x)v = \sum_{k \in \mathbb{N}_0} \frac{x^k v}{k!} \quad , \quad v \in V.$$

The formal Kac-Moody group G_{fp} now contains the prounipotent group $U_f = exp(\mathfrak{n}_f)$.

The group G_{fp} identifies with the Kac-Moody group of Kapitel 5 in [Sl] for a simply connected minimal free realization. It identifies with the Kac-Moody group of Section 6 in [Ku]. In [Sl], [Ku] the field is assumed to be \mathbb{C} , but it works in the same way over a field \mathbb{F} of characteristic 0. The results for G_{fn} stated above can be proved similarly as the corresponding results for G_{fp} .

The coordinate ring of the face monoid \widehat{G} and its restriction to the (minimal) Kac-Moody group G : The Kac-Moody group G acts on every \mathfrak{g} -module contained in \mathcal{O}_{int}^p and on its restricted dual $V^{(*)}$. We now change this action of G on $V^{(*)}$ to an action of G^{op} on $V^{(*)}$ by concatenation the corresponding representation of G with the inverse map. For a \mathfrak{g} -module V contained in \mathcal{O}_{int}^p , for $v \in V$ and $\phi \in V^{(*)}$ the map

$$\begin{aligned} \phi(\cdot v) : G &\rightarrow \mathbb{F} \\ g &\mapsto \phi(gv) \end{aligned}$$

is called the matrix coefficient of ϕ and v on G .

The coordinate ring $\mathbb{F}[G]$ of the Kac-Moody group G is the set of matrix coefficients obtained in this way by the modules contained in \mathcal{O}_{int}^p and their restricted duals. It is a \mathbb{F} -algebra on which $G^{op} \times G$ acts.

As explained above, it holds $G = G_I \rtimes T_{rest}$. Here, the derived group G_I of G identifies with the Kac-Moody group of [KP1] and T_{rest} is a certain subtorus of T . The multiplication map $m : G_I \times T_{rest} \rightarrow G$ induces an isomorphism $m^* : \mathbb{F}[G] \rightarrow \mathbb{F}[G_I] \otimes \mathbb{F}[T_{rest}]$. The restriction $\mathbb{F}[G_I] := \mathbb{F}[G]|_{G_I}$

coincides with the algebra of strongly regular functions of [KP2]. The restriction $\mathbb{F}[T_{rest}] := \mathbb{F}[G]|_{T_{rest}}$ is the classical coordinate ring of the torus T_{rest} , which identifies with the group algebra $\mathbb{F}[P_{rest}]$ of the lattice P_{rest} .

V. Kac and D. Peterson showed in Theorem 1 of [KP2] a Peter-Weyl theorem for $\mathbb{F}[G_I]$. They showed in Corollary 2.2 and Remark 2.1 of [KP2] a Borel-Weil theorem for $\mathbb{F}[G_I]^U$. It is easy to transcribe these theorems to theorems for $\mathbb{F}[G]$ and $\mathbb{F}[G]^U$. (Use in addition: P_I identifies with the weight lattice of [KP2]. It holds $P = P_I \oplus P_{rest}$ and $P^+ = P_I^+ \oplus P_{rest}$. The modules $L(N)$, $N \in P_{rest}$, are one-dimensional. In particular, it holds $L(\Lambda) \otimes L(N) = L(\Lambda + N)$ for all $\Lambda \in P^+$ and $N \in P_{rest}$.) Also the proofs of these theorems could be easily adapted to obtain the following theorems.

The Peter-Weyl theorem: A $G^{op} \times G$ -equivariant bijective linear map

$$PW : \bigoplus_{\Lambda \in P^+} L(\Lambda)^{(*)} \otimes L(\Lambda) \rightarrow \mathbb{F}[G]$$

is given by $PW(\phi \otimes v) := \phi(\cdot v)$ for all $\phi \in L(\Lambda)^{(*)}$, $v \in L(\Lambda)$, $\Lambda \in P^+$.

The Borel-Weil theorem: Denote by $\mathbb{F}[G]^U$ the functions of $\mathbb{F}[G]$ which take constant values on the elements of every coset gU , $g \in G$. Choose a highest weight vector $v_\Lambda \in L(\Lambda)_\Lambda$ for every $\Lambda \in P^+$. A G^{op} -equivariant linear bijective map

$$BW : \bigoplus_{\Lambda \in P^+} L(\Lambda)^{(*)} \rightarrow \mathbb{F}[G]^U$$

is given by $BW(\phi) := \phi(\cdot v_\Lambda)$ for all $\phi \in L(\Lambda)^{(*)}$, $\Lambda \in P^+$. Pulling back the algebra structure of $\mathbb{F}[G]^U$ by BW gives the Cartan algebra structure on $\bigoplus_{\Lambda \in P^+} L(\Lambda)^{(*)}$, which will be explained in subsection 3.1.

Note also that the coordinate ring $\mathbb{F}[G]$, and therefore also $\mathbb{F}[G]^U$, is an integral domain. This follows with Theorem 2.1 of [KP2], or could be shown directly.

Similarly as above we may define matrix coefficients for the face monoid \widehat{G} , and the algebra of matrix coefficients $\mathbb{F}[\widehat{G}]$, on which $G^{op} \times \widehat{G}$ acts. By definition, it holds $\mathbb{F}[G] = \mathbb{F}[\widehat{G}]|_G$. It has been shown in [M1] that the Kac-Moody group G is the Zariski open dense unit group of the face monoid \widehat{G} . In particular, the coordinate ring $\mathbb{F}[G]$ is isomorphic to the coordinate ring $\mathbb{F}[\widehat{G}]$ by the restriction map.

The Peter-Weyl theorem and Borel-Weil theorem of above give corresponding theorems for $\mathbb{F}[\widehat{G}]$ and $\mathbb{F}[\widehat{G}]^U$, the $G^{op} \times G$ and G^{op} actions now extended to $\widehat{G}^{op} \times \widehat{G}$ and \widehat{G}^{op} actions, but we do not need it in the article.

Note that this is a reinterpretation of the coordinate rings used in [KP2], [Kas], and in subsequent articles of other authors. These coordinate rings should be looked at as the coordinate rings of face monoids \widehat{G}_I , \widehat{G} . Classically, i.e., if the generalized Cartan matrix has only components of finite type, it holds $\widehat{G} = \widehat{G}_I = G$. This explains why the face monoid \widehat{G} may turn up in algebraic geometric constructions, where classically only the group G is involved.

2 Projective spectra of graded algebras

In this section we briefly work out the theory of spectra of homogeneous prime ideals of graded algebras as far as we will need it in this article. We call such spectra projective spectra for short. Proofs which are only variants of the nongraded or \mathbb{N}_0 -graded case are omitted.

To explain the difference to the projective spectra considered in algebraic geometry let $A = \bigoplus_{n \in \mathbb{N}_0} A_n$ be a \mathbb{N}_0 -graded \mathbb{F} -algebra such that $\mathbb{F}1 \subseteq A_0$. For simplicity assume equality. Denote by $\text{Proj}(A)$ all \mathbb{N}_0 -homogeneous prime ideals of A and by $m := \bigoplus_{n \in \mathbb{N}} A_n$ the irrelevant ideal. Then

$$\text{Proj}(A) = \text{Proj}(A) \setminus \{m\} \dot{\cup} \{m\}.$$

Suppose that $\text{Proj}(A) \setminus \{m\} \neq \emptyset$. Then $\text{Proj}(A) \setminus \{m\}$ can be made into a projective scheme as for example explained in [Ha], Chapter II, Section 2. But it is only an open dense part of $\text{Proj}(A)$. Its boundary is given by the closed point m .

The example investigated in the following sections 3 and 4 is the projective spectrum $\text{Proj}(CA)$ of the P^+ -graded Cartan algebra CA associated to a Kac-Moody group G . The projective spectrum $\text{Proj}(CA)$ is a locally ringed space. There is an open dense part of $\text{Proj}(CA)$ which is a scheme, but now also its boundary is interesting. Actually, $\text{Proj}(CA)$ is stratified by schemes. The algebraic geometric action of the face monoid \widehat{G} on $\text{Proj}(CA)$, which we will obtain naturally, does not respect the strata but the closures of the strata. The existence of such an action is not at all obvious if we would only consider the union of these strata.

2.1 Projective spectra and their \mathbb{F} -valued points as topological spaces

In this article a \mathcal{M} -graded \mathbb{F} -algebra A , or graded algebra A for short, consists of the following data:

- (a) A commutative monoid $(\mathcal{M}, +)$ with the cancellation property, i.e., for all $\Lambda_1, \Lambda_2, N \in \mathcal{M}$ it holds:

$$\Lambda_1 + N = \Lambda_2 + N \quad \Rightarrow \quad \Lambda_1 = \Lambda_2.$$

- (b) A commutative, associative, \mathbb{F} -linear algebra (A, \bullet) with unit 1. Furthermore, a \mathbb{F} -linear decomposition

$$A = \bigoplus_{\Lambda \in \mathcal{M}} A_\Lambda \quad \text{such that} \quad \mathbb{F}1 \subseteq A_0 \quad \text{and} \quad A_\Lambda \bullet A_N \subseteq A_{\Lambda+N} \text{ for all } \Lambda, N \in \mathcal{M}.$$

Since the cancelation property holds, the commutative monoid \mathcal{M} embeds into a commutative group $\mathcal{M} - \mathcal{M}$, minimal over \mathcal{M} .

Let A be a \mathcal{M} -graded algebra. The projective spectrum of A (as a topological space) consists of the set

$$\text{Proj}(A) := \{\mathcal{M}\text{-homogeneous prime ideals of } A\}$$

equipped with the Zariski topology, whose closed sets are obtained by

$$\mathcal{V}(I) := \{Q \in \text{Proj}(A) \mid Q \supseteq I\}, \quad I \text{ a } \mathcal{M}\text{-homogeneous ideal of } A.$$

Fix a point $Q \in \text{Proj}(A)$. Since Q is \mathcal{M} -homogeneous it has the decomposition $Q = \bigoplus_{\Lambda \in \mathcal{M}} Q_\Lambda$ with $Q_\Lambda := Q \cap A_\Lambda$. Since Q is prime the complement $A \setminus Q$ is a multiplicatively closed subset of A . Also the set $\bigcup_{\Lambda \in \mathcal{M}} A_\Lambda \setminus Q_\Lambda$ of homogeneous elements of $A \setminus Q$ is a multiplicatively closed subset of A . Now

$$A_{(Q)} := \left\{ \frac{a}{b} \mid a \in A_\Lambda, b \in A_\Lambda \setminus Q_\Lambda, \Lambda \in \mathcal{M} \text{ such that } A_\Lambda \setminus Q_\Lambda \neq \emptyset \right\} \subseteq \left(\bigcup_{\Lambda \in \mathcal{M}} A_\Lambda \setminus Q_\Lambda \right)^{-1} A$$

is a commutative, associative, unital, local, \mathbb{F} -linear algebra with maximal ideal

$$m_{(Q)} := \left\{ \frac{a}{b} \mid a \in Q_\Lambda, b \in A_\Lambda \setminus Q_\Lambda, \Lambda \in \mathcal{M} \text{ such that } A_\Lambda \setminus Q_\Lambda \neq \emptyset \right\}.$$

We define the residue field of Q to be

$$\mathbb{F}_{(Q)} := A_{(Q)} / m_{(Q)}.$$

We identify the field \mathbb{F} with the corresponding subfield of $\mathbb{F}_{(Q)}$.

We call $Q \in \text{Proj}(A)$ an \mathbb{F} -valued point of $\text{Proj}(A)$ if $\mathbb{F}_{(Q)} = \mathbb{F}$. We set

$$\text{Proj}(A)(\mathbb{F}) := \{Q \in \text{Proj}(A) \mid \mathbb{F}_{(Q)} = \mathbb{F}\}.$$

A useful characterization of the \mathbb{F} -valued points of the projective spectrum is given by

Theorem 2.1 *Let $Q \in \text{Proj}(A)$. It is equivalent:*

(i) $Q \in \text{Proj}(A)(\mathbb{F})$.

(ii) $\dim(A_\Lambda/Q_\Lambda) \leq 1$ for all $\Lambda \in \mathcal{M}$.

Proof: For every $\Lambda \in \mathcal{M}$ we choose a linear complement R_Λ of Q_Λ in A_Λ , i.e., $A_\Lambda = Q_\Lambda \oplus R_\Lambda$. We first show

$$\mathbb{F}_{(Q)} = \left\{ \frac{r}{s} + m_{(Q)} \mid r \in R_\Lambda, s \in R_\Lambda \setminus \{0\}, \Lambda \in \mathcal{M} \text{ such that } R_\Lambda \neq \{0\} \right\}. \quad (2)$$

Obviously, the inclusion " \supseteq " holds. Now let $\frac{a}{b} + m_{(Q)} \in \mathbb{F}_{(Q)}$ where $a \in A_\Lambda$, $b \in A_\Lambda \setminus Q_\Lambda$ and $\Lambda \in \mathcal{M}$ such that $A_\Lambda \setminus Q_\Lambda \neq \emptyset$. Decompose

$$\begin{aligned} a &= q_a + r & \text{with} & & q_a \in Q_\Lambda, r \in R_\Lambda, \\ b &= q_b + s & \text{with} & & q_b \in Q_\Lambda, s \in R_\Lambda \setminus \{0\}. \end{aligned}$$

It holds

$$\frac{a}{b} - \frac{r}{s} = \frac{a \bullet s - r \bullet b}{b \bullet s} = \frac{q_a \bullet s - r \bullet q_b}{b \bullet s} \in m_{(Q)},$$

because of $q_a \bullet s, r \bullet q_b \in Q_\Lambda \bullet R_\Lambda \subseteq Q_{2\Lambda}$ and $b \bullet s \in (A_\Lambda \setminus Q_\Lambda) \bullet (A_\Lambda \setminus Q_\Lambda) \subseteq A_{2\Lambda} \setminus Q_{2\Lambda}$.

Now suppose that (ii) holds. Let $\Lambda \in \mathcal{M}$ such that $R_\Lambda \neq \{0\}$. If $r \in R_\Lambda$, $s \in R_\Lambda \setminus \{0\}$ then $r = \mu s$ there for some $\mu \in \mathbb{F}$. Therefore, $\frac{r}{s} = \mu \frac{1}{1}$. Now (i) follows by (2).

Suppose that (i) holds. Let $\Lambda \in \mathcal{M}$ such that $R_\Lambda \neq \{0\}$. Choose $s \in R_\Lambda \setminus \{0\}$. By (2) it follows that for every $r \in R_\Lambda$ there exists some $\mu \in \mathbb{F}$ such that

$$\frac{r}{s} - \mu \frac{1}{1} = \frac{r - \mu s}{s} \in m_{(Q)}.$$

Therefore, there exist $a \in Q_N$, $b \in A_N \setminus Q_N$, $N \in \mathcal{M}$ such that

$$\frac{r - \mu s}{s} = \frac{a}{b}.$$

It follows that there exists $c \in A_M \setminus Q_M$, $M \in \mathcal{M}$ such that

$$(r - \mu s) \bullet b \bullet c = a \bullet s \bullet c. \quad (3)$$

Since $a \in Q$ the right hand side of (3) is contained in Q . Assume that $r - \mu s \neq 0$. Since $r - \mu s \in R_\Lambda \setminus \{0\} \subseteq A \setminus Q$ and $b, c \in A \setminus Q$, the left hand side of (3) is contained in $A \setminus Q$, which is not possible. It follows $r = \mu s$. Therefore, (ii) holds. \square

The subset $\text{Proj}(A)(\mathbb{F})$ of $\text{Proj}(A)$ is equipped with its relative topology. The relative closure of $Z \subseteq \text{Proj}(A)(\mathbb{F})$ is denoted by \overline{Z}^{pts} .

For a subset $S \subseteq \text{Proj}(A)$ we define

$$S(\mathbb{F}) := S \cap \text{Proj}(A)(\mathbb{F}).$$

For the \mathcal{M} -graded algebras considered in the following sections 3 and 4 the commutative group $\mathcal{M} - \mathcal{M}$ is isomorphic to some \mathbb{Z}^m , $m \in \mathbb{N}_0$. It applies:

Proposition 2.2 *Let A be a \mathcal{M} -graded algebra such that $\mathcal{M} - \mathcal{M}$ is a free commutative group. Then it is $\text{Proj}(A) \neq \emptyset$.*

Remark 2.3 *A similar statement does not hold for the \mathbb{F} -valued points, as it does not hold in the nongraded case. Take for example the \mathbb{R} -algebra \mathbb{C} , then $\text{Spec}(\mathbb{C})(\mathbb{R}) = \emptyset$.*

Proof: It is $\text{Spec}(A) \neq \emptyset$, compare for example Proposition 3.9 and Remark 3.25 (i) of [Sh]. Therefore, the Proposition holds for $\mathcal{M} = \{0\}$. Now let $\mathcal{M} \neq \{0\}$ and choose $I \in \text{Spec}(A)$. It is easy to check that

$$I^h := \bigoplus_{\Lambda \in \mathcal{M}} I \cap A_\Lambda.$$

is a homogeneous ideal. We show that I^h is prime.

For $x \in A$ with homogeneous components $x_\Lambda \in A_\Lambda$, $\Lambda \in \mathcal{M}$, set

$$cs(x) := \{\Lambda \in \mathcal{M} \mid x_\Lambda \notin I\}.$$

Let $x, \tilde{x} \in A \setminus I^h$. Then $cs(x)$ and $cs(\tilde{x})$ are nonempty finite subsets of \mathcal{M} . Choose a base of the nonzero free commutative group $\mathcal{M} - \mathcal{M}$. There exists a finite subset $\{N_1, N_2, \dots, N_k\}$ of this base such that $cs(x)$ and $cs(\tilde{x})$ are contained in the span $\mathbb{Z}N_1 \oplus \mathbb{Z}N_2 \oplus \dots \oplus \mathbb{Z}N_k$. Equip the elements of this span with the lexicographical order induced by the standard order on \mathbb{Z} . This is a total order compatible with the addition.

Let M be the biggest element of $cs(x)$ and \tilde{M} the biggest element of $cs(\tilde{x})$. Then $M + \tilde{M}$ is the biggest element of

$$\{\Lambda + \tilde{\Lambda} \mid \Lambda \in cs(x), \tilde{\Lambda} \in cs(\tilde{x})\}$$

Therefore, the $M + \tilde{M}$ -homogeneous component $(x\tilde{x})_{M+\tilde{M}}$ of $x\tilde{x}$ is given modulo the linear space $I \cap A_{M+\tilde{M}}$ by

$$(x\tilde{x})_{M+\tilde{M}} = \sum_{\Lambda, \tilde{\Lambda} \in \mathcal{M}, \Lambda+\tilde{\Lambda}=M+\tilde{M}} x_\Lambda \tilde{x}_{\tilde{\Lambda}} \equiv \sum_{\Lambda \in cs(x), \tilde{\Lambda} \in cs(\tilde{x}), \Lambda+\tilde{\Lambda}=M+\tilde{M}} x_\Lambda \tilde{x}_{\tilde{\Lambda}} = x_M \tilde{x}_{\tilde{M}} \pmod{I \cap A_{M+\tilde{M}}}.$$

Because I is prime and $x_M, \tilde{x}_{\tilde{M}} \notin I$ it is $x_M \tilde{x}_{\tilde{M}} \notin I \cap A_{M+\tilde{M}}$. It follows $x\tilde{x} \in A \setminus I^h$. \square

Let A be a \mathcal{M}_A -graded algebra and B be a \mathcal{M}_B -graded algebra. A morphism $f^* : A \rightarrow B$ of graded algebras is a unital morphism of algebras $f^* : A \rightarrow B$ for which there exists a map $F^* : \mathcal{M}_A \rightarrow \mathcal{M}_B$ such that it holds:

- (a) $f^*(A_\Lambda) \subseteq B_{F^*(\Lambda)}$ for all $\Lambda \in \mathcal{M}_A$.
- (b) F^* is injective on $\{\Lambda \in \mathcal{M}_A \mid f^*(A_\Lambda) \neq \{0\}\}$.

Note that the map F^* is uniquely determined only on $\{\Lambda \in \mathcal{M}_A \mid f^*(A_\Lambda) \neq \{0\}\}$. Outside this set the map F^* is not relevant and can be chosen arbitrarily. The definition has been formulated in this way to avoid working with partial maps.

Condition (b) guaranties the following indispensable property: If J is a homogeneous ideal of B then the inverse image $(f^*)^{-1}(J)$ is a homogeneous ideal of A . Now it is not difficult to check

Theorem 2.4 *Let $f^* : A \rightarrow B$ be a morphism of graded algebras.*

- (a) *A continuous map is given by*

$$\begin{aligned} f : \text{Proj}(B) &\rightarrow \text{Proj}(A) \\ Q &\rightarrow (f^*)^{-1}(Q) \end{aligned}.$$

It maps $\text{Proj}(B)(\mathbb{F})$ into $\text{Proj}(A)(\mathbb{F})$.

- (b) *For every $Q \in \text{Proj}(B)$ the morphism $f^* : A \rightarrow B$ induces a morphism of algebras*

$$\begin{aligned} f_Q^* : A_{(f(Q))} &\rightarrow B_{(Q)} \\ \frac{a}{b} &\rightarrow \frac{f^*(a)}{f^*(b)}, \end{aligned}$$

which is local, i.e., $f_Q^(m_{(f(Q))}) \subseteq m_{(Q)}$. This morphism induces a injective morphism $f_Q^* : \mathbb{F}_{(f(Q))} \rightarrow \mathbb{F}_{(Q)}$ of fields over \mathbb{F} .*

The graded algebras and its morphisms form a category, in which the concatenation of morphisms is the concatenation of maps.

Remark 2.5 *A morphism $f^* : A \rightarrow B$ of graded algebras is an isomorphism of graded algebras if and only if it is bijective.*

If $f^ : A \rightarrow B$ is an isomorphism of graded algebras then a map F^* as in the definition maps the set $\{\Lambda \in \mathcal{M}_A \mid A_\Lambda \neq \{0\}\}$ bijectively to $\{N \in \mathcal{M}_B \mid B_N \neq \{0\}\}$. But the monoids \mathcal{M}_A and \mathcal{M}_B may be non-isomorphic, even non-bijective.*

Let A be a \mathcal{M} -graded algebra. Let $S \subseteq A \setminus \{0\}$ be a multiplicatively closed subset of homogeneous elements. For $\Lambda \in \mathcal{M}$ denote by $S_\Lambda := S \cap A_\Lambda$ the Λ -homogeneous elements of S . It is $\mathcal{M}_S := \{\Lambda \in \mathcal{M} \mid S_\Lambda \neq \emptyset\}$ a submonoid of \mathcal{M} and $\mathcal{M} - \mathcal{M}_S := \{\Lambda - N \in \mathcal{M} - \mathcal{M} \mid \Lambda \in \mathcal{M}, N \in \mathcal{M}_S\}$ a submonoid of the group $\mathcal{M} - \mathcal{M}$. Set

$$D(S) := \{Q \in \text{Proj}(A) \mid S \subseteq A \setminus Q\}.$$

The localization $S^{-1}A$ is a commutative associative unital $(\mathcal{M} - \mathcal{M}_S)$ -graded \mathbb{F} -algebra. The canonical morphism $i^* : A \rightarrow S^{-1}A$ is a morphism of graded algebras. (The cancellation property for \mathcal{M} in our definition of a \mathcal{M} -graded algebra A guaranties this property, which was the reason to demand it.) It is not difficult to check

Theorem 2.6 (a) *The continuous map*

$$\begin{array}{ccc} i : \text{Proj}(S^{-1}A) & \rightarrow & \text{Proj}(A) \\ Q & \mapsto & (i^*)^{-1}(Q) \end{array}$$

maps $\text{Proj}(S^{-1}A)$ homeomorphically to $D(S)$. It also maps $\text{Proj}(S^{-1}A)(\mathbb{F})$ homeomorphically to $D(S)(\mathbb{F})$.

(b) *For every $Q \in \text{Proj}(S^{-1}A)$ the local morphism $i_Q^* : A_{(i(Q))} \rightarrow (S^{-1}A)_{(Q)}$ is an isomorphism.*

In particular, the theorem applies to the principal open sets

$$D(a) := \{Q \in \text{Proj}(A) \mid a \notin Q\}$$

with $a \in A$ homogeneous and not nilpotent, because it holds $D(a) = D(\{a^n \mid n \in \mathbb{N}_0\})$. These sets form a base of the open sets of the Zariski topology of $\text{Proj}(A)$.

Let A be a \mathcal{M} -graded algebra. Let $S \subseteq A \setminus \{0\}$ be a multiplicatively closed subset of homogeneous elements of A . We call $s_0 \in S$ a *principal element* of S if the following property holds: For every $s \in S$ there exist $\tilde{s} \in S$ and $n \in \mathbb{N}_0$ such that $s\tilde{s} = s_0^n$.

A subset $I \subseteq S$ is called a semigroup ideal of S if $S \bullet I \subseteq I$. Note that we also allow I to be the empty set. A submonoid $F \subseteq S$, for which $S \setminus F$ is a semigroup ideal of S , is called a face of S . The relative interior $ri(S)$ of S is the semigroup ideal of S defined by

$$ri(S) := S \setminus \bigcup_{F \text{ a face of } S, F \neq S} F.$$

It is not difficult to check

Theorem 2.7 *Suppose there exists a principal element $s_0 \in S$. Then it holds:*

(a) $D(s_0) = D(S)$.

(b) *An isomorphism of graded algebras is given by*

$$\begin{array}{ccc} \{s_0^n \mid n \in \mathbb{N}_0\}^{-1} A & \rightarrow & S^{-1}A \\ \frac{a}{s_0^n} & \mapsto & \frac{a}{s_0^n} \end{array}.$$

Furthermore, the set of principal elements of S is a semigroup ideal of S contained in $\text{ri}(S)$.

If there exists a principal element of S we also call $D(S)$ a *principal open set*.

Remark 2.8 If S is generated by finitely many elements s_1, s_2, \dots, s_k then $s_0 := s_1 s_2 \cdots s_k$ is a principal element of S . It is easy to find examples where S is not finitely generated and principal elements exist.

Let A be a \mathcal{M} -graded algebra. Let I be a homogeneous ideal of A , $I \neq A$. The quotient algebra A/I is \mathcal{M} -graded. The canonical projection $\pi^* : A \rightarrow A/I$ is a morphism of graded algebras. It is not difficult to check

Theorem 2.9 (a) *The continuous map*

$$\begin{array}{ccc} \pi : \text{Proj}(A/I) & \rightarrow & \text{Proj}(A) \\ Q & \mapsto & (\pi^*)^{-1}(Q) \end{array}$$

maps $\text{Proj}(A/I)$ homeomorphically to $\mathcal{V}(I)$. It also maps $\text{Proj}(A/I)(\mathbb{F})$ homeomorphically to $\mathcal{V}(I)(\mathbb{F})$.

(b) *For every $Q \in \text{Proj}(A/I)$ the local morphism $\pi_Q^* : A_{(\pi(Q))} \rightarrow (A/I)_{(Q)}$ is surjective.*

If A is a \mathcal{M} -graded algebra and B is a \mathcal{N} -graded algebra then the tensor product $A \otimes B$ of the algebras A and B over \mathbb{F} is a $\mathcal{M} \oplus \mathcal{N}$ -graded algebra. The canonical injections

$$\begin{array}{ccc} i_A^* : A & \rightarrow & A \otimes B \\ a & \mapsto & a \otimes 1 \end{array} \quad \text{and} \quad \begin{array}{ccc} i_B^* : B & \rightarrow & A \otimes B \\ b & \mapsto & 1 \otimes b \end{array}$$

are morphisms of graded algebras. But note that in general $(A \otimes B, i_A^*, i_B^*)$ is no coproduct of A and B in the category of graded algebras. It holds

Theorem 2.10 (a) *The map*

$$\begin{array}{ccc} i_A : \text{Proj}(A \otimes B) & \rightarrow & \text{Proj}(A) \\ Q & \mapsto & (i_A^*)^{-1}(Q) \end{array}$$

is a continuous map, mapping $\text{Proj}(A \otimes B)(\mathbb{F})$ into $\text{Proj}(A)(\mathbb{F})$.

(b) *For every $Q \in \text{Proj}(A \otimes B)$ the local morphism $(i_A^*)_Q : A_{(i_A(Q))} \rightarrow (A \otimes B)_{(Q)}$ is injective.*

We now consider particular graded algebras to obtain some results on the projective spectra of tensor products, which will be used later. Let \mathcal{N} be a commutative monoid with cancellation property. Its monoid algebra

$$\mathbb{F}[\mathcal{N}] = \bigoplus_{N \in \mathcal{N}} \mathbb{F}e_N, \quad e_N \bullet e_{N'} = e_{N+N'} \text{ for all } N, N' \in \mathcal{N},$$

over the field \mathbb{F} is a \mathcal{N} -graded algebra. The following proposition should be known. We give a proof because we will need soon a variant of it.

Proposition 2.11 *Let C be a commutative, associative, \mathbb{F} -linear algebra without zero divisors. Let \mathcal{N} be a free commutative group, and let $\mathbb{F}[\mathcal{N}]$ be its group algebra. Then the algebra $C \otimes \mathbb{F}[\mathcal{N}]$ has no zero divisors.*

Remark 2.12 *For $C = \mathbb{F}$ the proposition specializes to: The group algebra $\mathbb{F}[\mathcal{N}]$ of a free commutative group \mathcal{N} has no zero divisors.*

Proof: The proposition is trivial for $\mathcal{N} = \{0\}$. Let $\mathcal{N} \neq \{0\}$. An element $x \in C \otimes \mathbb{F}[\mathcal{N}]$ can be uniquely written in the form $x = \sum_{N \in \mathcal{N}} c_N \otimes e_N$ with $c_N \in C$, $N \in \mathcal{N}$, and $\text{supp}(x) := \{N \in \mathcal{N} \mid c_N \neq 0\}$ is finite.

Suppose there exist nonzero elements $x = \sum_{N \in \mathcal{N}} c_N \otimes e_N$, $\tilde{x} = \sum_{N \in \mathcal{N}} \tilde{c}_N \otimes e_N$ of $C \otimes \mathbb{F}[\mathcal{N}]$ such that $x\tilde{x} = 0$.

Choose a base of the free commutative group \mathcal{N} . There exists a finite subset $\{N_1, N_2, \dots, N_k\}$ of this base such that $\text{supp}(x)$ and $\text{supp}(\tilde{x})$ are contained in the span $\mathbb{Z}N_1 \oplus \mathbb{Z}N_2 \oplus \dots \oplus \mathbb{Z}N_k$. Equip the elements of this span with the lexicographical order induced by the standard order on \mathbb{Z} . This is a total order compatible with the addition.

Let M be the maximal element of $\text{supp}(x)$ and \tilde{M} the maximal element of $\text{supp}(\tilde{x})$. Then $M + \tilde{M}$ is the maximal element of $\text{supp}(x) + \text{supp}(\tilde{x})$. Therefore, $x\tilde{x}$ is of the form

$$x\tilde{x} = c_M \tilde{c}_{\tilde{M}} \otimes e_{M+\tilde{M}} + \sum_{N < M+\tilde{M}} d_N \otimes e_N.$$

Since C has no zero divisors it holds $c_M \tilde{c}_{\tilde{M}} \neq 0$. This contradicts $x\tilde{x} = 0$. \square

The last Proposition is used to show

Theorem 2.13 *Let A be a \mathcal{M} -graded algebra. Let \mathcal{N} be a free commutative group, and let $\mathbb{F}[\mathcal{N}]$ be its \mathcal{N} -graded group algebra. The map*

$$\begin{array}{ccc} i_A : \text{Proj}(A \otimes \mathbb{F}[\mathcal{N}]) & \rightarrow & \text{Proj}(A) \\ Q & \mapsto & (i_A^*)^{-1}(Q) \end{array}$$

is a homeomorphism, mapping $\text{Proj}(A \otimes \mathbb{F}[\mathcal{N}])(\mathbb{F})$ bijectively to $\text{Proj}(A)(\mathbb{F})$. Its inverse is

$$(i_A)^{-1} : \begin{array}{ccc} \text{Proj}(A) & \rightarrow & \text{Proj}(A \otimes \mathbb{F}[\mathcal{N}]) \\ Q & \mapsto & Q \otimes \mathbb{F}[\mathcal{N}] \end{array}.$$

For every $Q \in \text{Proj}(A)$ the local morphism $(i_A^)_Q : A_{(Q)} \rightarrow (A \otimes \mathbb{F}[\mathcal{N}]_{(Q \otimes \mathbb{F}[\mathcal{N}]})$ is an isomorphism.*

Remark 2.14 *For $A = \mathbb{F}$ the theorem specializes to: $\text{Proj}(\mathbb{F}[\mathcal{N}]) = \text{Proj}(\mathbb{F}[\mathcal{N}])(\mathbb{F}) = \{\{0\}\}$ for the group algebra $\mathbb{F}[\mathcal{N}]$ of a free commutative group \mathcal{N} .*

Proof: It is i_A a continuous map by Theorem 2.10 (a). If $Q \in \text{Proj}(A)$ then $Q \otimes \mathbb{F}[\mathcal{N}]$ is a homogeneous ideal of $A \otimes \mathbb{F}[\mathcal{N}]$ and it holds

$$(i_A^*)^{-1}(Q \otimes \mathbb{F}[\mathcal{N}]) = \{a \in A \mid a \otimes 1 \in Q \otimes \mathbb{F}[\mathcal{N}]\} = Q.$$

The ideal $Q \otimes \mathbb{F}[\mathcal{N}]$ is also prime because $(A \otimes \mathbb{F}[\mathcal{N}]) / (Q \otimes \mathbb{F}[\mathcal{N}]) \cong (A/Q) \otimes \mathbb{F}[\mathcal{N}]$ has no zero divisors by Proposition 2.11. This shows that the map $(i_A)^{-1}$ given in the theorem is well defined and it holds $i_A \circ (i_A)^{-1} = \text{id}$.

Let J be a homogeneous ideal of $A \otimes \mathbb{F}[\mathcal{N}]$. Then $(i_A^*)^{-1}(J)$ is a homogeneous ideal of A because i_A^* is a morphism of graded algebras. We show that it holds

$$J = (i_A^*)^{-1}(J) \otimes \mathbb{F}[\mathcal{N}]. \quad (4)$$

Since J is an ideal, the inclusion \supseteq holds. Now let $x \in J$. Write x in the form

$$x = \sum_{N \in \mathcal{N}} a_N \otimes e_N \quad \text{with} \quad a_N \in A, \quad a_N \neq 0 \text{ for only finitely many } N.$$

Since J is homogeneous it follows $a_N \otimes e_N \in J$ for all $N \in \mathcal{N}$. Since J is an ideal it follows

$$a_N \otimes 1 = (a_N \otimes e_N) \bullet (1 \otimes e_{-N}) \in J \quad \text{for all} \quad N \in \mathcal{N},$$

which is equivalent to $a_N \in (i_A^*)^{-1}(J)$ for all $N \in \mathcal{N}$, which in turn is equivalent to $x \in (i_A^*)^{-1}(J) \otimes \mathbb{F}[\mathcal{N}]$.

From (4) it follows $(i_A)^{-1} \circ i_A = id$. Furthermore, it follows $i_A(\mathcal{V}(J)) = \mathcal{V}((i_A^*)^{-1}(J))$ for every homogeneous ideal J of $A \otimes \mathbb{F}[\mathcal{N}]$, which shows that $(i_A)^{-1}$ is continuous.

Let $Q \in \text{Proj}(A)$. By Theorem 2.10 (b) the morphism $(i_A^*)_Q : A_{(Q)} \rightarrow (A \otimes \mathbb{F}[\mathcal{N}])_{(Q \otimes \mathbb{F}[\mathcal{N}]})$ is injective. Let $M \in \mathcal{M}$ and $N \in \mathcal{N}$. For $a \otimes e_N \in A_M \otimes e_N$ and $b \in A_M \otimes e_N \setminus Q_M \otimes e_N$ it holds

$$(i_A^*)_Q\left(\frac{a}{b}\right) = \frac{a \otimes 1}{b \otimes 1} = \frac{a \otimes e_N}{b \otimes e_N}.$$

Therefore, $(i_A^*)_Q$ is also surjective. It follows that $\text{Proj}(A \otimes \mathbb{F}[\mathcal{N}])(\mathbb{F})$ is mapped bijectively to $\text{Proj}(A)(\mathbb{F})$. \square

The following theorem is instructive. We don't need to make use of it due to the quite explicit descriptions of the spectra of sections 3 and 4.

Theorem 2.15 *Let A be a \mathcal{M} -graded algebra. Let B a \mathcal{N} -graded algebra such that $\mathcal{N} - \mathcal{N}$ is a free commutative group.*

(a) *It holds: $\text{Proj}(A \otimes B)(\mathbb{F}) \neq \emptyset \iff \text{Proj}(A)(\mathbb{F}) \neq \emptyset$ and $\text{Proj}(B)(\mathbb{F}) \neq \emptyset$.*

(b) *The map*

$$\begin{aligned} (i_A, i_B) : \text{Proj}(A \otimes B)(\mathbb{F}) &\rightarrow \text{Proj}(A)(\mathbb{F}) \times \text{Proj}(B)(\mathbb{F}) \\ Q &\mapsto ((i_A^*)^{-1}(Q), (i_B^*)^{-1}(Q)) \end{aligned}$$

is a bijective continuous map. Its inverse is given by

$$\begin{aligned} (i_A, i_B)^{-1} : \text{Proj}(A)(\mathbb{F}) \times \text{Proj}(B)(\mathbb{F}) &\rightarrow \text{Proj}(A \otimes B)(\mathbb{F}) \\ (R, S) &\mapsto R \otimes B + A \otimes S \end{aligned}$$

Proof: Let C be a commutative, associative, \mathbb{F} -linear algebra without zero divisors. Let $D = \bigoplus_{N \in \mathcal{N}} D_N$ be a \mathcal{N} -graded algebra without zero divisors such that $\dim(D_N) \leq 1$ for all $N \in \mathcal{N}$. By adapting the proof of Proposition 2.11 it is easy to show that $C \otimes D$ has no zero divisors.

To (a): If $Q \in \text{Proj}(A \otimes B)(\mathbb{F})$ then by Theorem 2.10 (a) it follows $(i_A^*)^{-1}(Q) \in \text{Proj}(A)(\mathbb{F})$ and $(i_B^*)^{-1}(Q) \in \text{Proj}(B)(\mathbb{F})$.

Now let $R \in \text{Proj}(A)(\mathbb{F})$ and $S \in \text{Proj}(B)(\mathbb{F})$. Obviously, $R \otimes B + A \otimes S$ is a homogeneous ideal of $A \otimes B$ with homogeneous parts

$$(R \otimes B + A \otimes S)_{(M, N)} = R_M \otimes B_N + A_M \otimes S_N \quad \text{for } M \in \mathcal{M}, N \in \mathcal{N}.$$

It holds

$$\begin{aligned} A \otimes B / (R \otimes B + A \otimes S) &\cong A/R \otimes B/S, \\ (A \otimes B)_{(M, N)} / (R \otimes B + A \otimes S)_{(M, N)} &\cong A_M/R_M \otimes B_N/S_N \quad \text{for all } M \in \mathcal{M}, N \in \mathcal{N}. \end{aligned}$$

The algebras A/R and B/S have no zero divisors. Furthermore, it holds $B/S = \bigoplus_{N \in \mathcal{N}} B_N/S_N$ with $\dim(B_N/S_N) \leq 1$ for all $N \in \mathcal{N}$ by Theorem 2.1. It follows that $A/R \otimes B/S$ has no zero divisors. Therefore, $R \otimes B + A \otimes S \in \text{Proj}(A \otimes B)$. By Theorem 2.1 it holds $\dim(A_M/R_M \otimes B_N/S_N) = \dim(A_M/R_M)\dim(B_N/S_N) \leq 1$ for all $M \in \mathcal{M}$ and $N \in \mathcal{N}$. By the same theorem it follows $R \otimes B + A \otimes S \in \text{Proj}(A \otimes B)(\mathbb{F})$.

To (b): By Theorem 2.10 (a) the map (i_A, i_B) is continuous. It remains to show that the maps of the theorem are inverse maps. For $R \in \text{Proj}(A)(\mathbb{F})$ and $S \in \text{Proj}(B)(\mathbb{F})$ it holds

$$(i_A^*)^{-1}(R \otimes B + A \otimes S) = \{a \in A \mid a \otimes 1 \in R \otimes B + A \otimes S\} = R,$$

because of $1 \notin S$. Similarly, it holds $(i_B^*)^{-1}(R \otimes B + A \otimes S) = S$. Therefore, $(i_A, i_B) \circ (i_A, i_B)^{-1} = id$.

Now we show $(i_A, i_B)^{-1} \circ (i_A, i_B) = id$, which is equivalent to

$$i_A(Q) \otimes B + A \otimes i_B(Q) = Q \quad \text{for all } Q \in \text{Proj}(A \otimes B)(\mathbb{F}).$$

For all $M \in \mathcal{M}$ and $N \in \mathcal{N}$ it holds

$$i_A(Q)_M \otimes B_N + A_M \otimes i_B(Q)_N \subseteq Q_{M+N} \subseteq A_M \otimes B_N. \quad (5)$$

By Theorem 2.1 it holds $\dim(A_M/i_A(Q)_M) \leq 1$ and $\dim(B_N/i_B(Q)_N) \leq 1$.

If $i_A(Q)_M = A_M$ or $i_B(Q)_N = B_N$ then in (5) both inclusions are equalities. Now suppose that $i_A(Q)_M \neq A_M$, $i_B(Q)_N \neq B_N$, and $i_A(Q)_M \otimes B_N + A_M \otimes i_B(Q)_N \subsetneq Q_{M+N}$. Then $Q_{M+N} = A_M \otimes B_N$. Let $a \in A_M \setminus i_A(Q)_M$, $b \in B_N \setminus i_B(Q)_N$. Then it holds

$$\underbrace{(a \otimes 1)}_{\notin Q} \bullet \underbrace{(1 \otimes b)}_{\notin Q} = a \otimes b \in A_M \otimes B_N = Q_{M+N},$$

which contradicts that Q is prime. \square

2.2 The structure sheaves of projective spectra

Let A be a \mathcal{M} -graded algebra. We now introduce a locally ringed space $(\text{Proj}(A), \mathcal{O}_{\text{Proj}(A)})$. We call $\mathcal{O}_{\text{Proj}(A)}$ the structure sheaf on $\text{Proj}(A)$.

A presheaf $\tilde{\mathcal{O}}_{\text{Proj}(A)}$ of \mathbb{F} -algebras on $\text{Proj}(A)$ is obtained as follows: For $\emptyset \neq U \subseteq \text{Proj}(A)$, U open, let $\tilde{\mathcal{O}}_{\text{Proj}(A)}(U)$ be the commutative, associative, unital, \mathbb{F} -linear algebra

$$\tilde{\mathcal{O}}_{\text{Proj}(A)}(U) := \left\{ \frac{a}{b} \mid \begin{array}{l} a \in A_\Lambda, b \in \bigcap_{Q \in U} A_\Lambda \setminus Q_\Lambda, \\ \Lambda \in \mathcal{M} \text{ such that } \bigcap_{Q \in U} A_\Lambda \setminus Q_\Lambda \neq \emptyset \end{array} \right\} \subseteq \left(\bigcup_{\Lambda \in \mathcal{M}} \bigcap_{Q \in U} A_\Lambda \setminus Q_\Lambda \right)^{-1} A.$$

For $\emptyset \neq U_1 \subseteq U_2 \subseteq \text{Proj}(A)$, U_1 and U_2 open, let

$$\widetilde{res}_{U_1}^{U_2} : \tilde{\mathcal{O}}_{\text{Proj}(A)}(U_2) \xrightarrow{\frac{a}{b}} \tilde{\mathcal{O}}_{\text{Proj}(A)}(U_1)$$

be the restriction morphism. It is easy to check that the stalk in $Q \in \text{Proj}(A)$ and its restriction morphisms are given by the local \mathbb{F} -algebra $A_{(Q)}$ and the morphisms

$$\widetilde{res}_Q^U : \tilde{\mathcal{O}}_{\text{Proj}(A)}(U) \xrightarrow{\frac{a}{b}} A_{(Q)}$$

where U is open in $\text{Proj}(A)$ with $Q \in U$.

The structure sheaf $\mathcal{O}_{\text{Proj}(A)}$ on $\text{Proj}(A)$ is the sheaf associated to the presheaf $\tilde{\mathcal{O}}_{\text{Proj}(A)}$ on $\text{Proj}(A)$. The stalk in $Q \in \text{Proj}(A)$ identifies with the local \mathbb{F} -algebra $A_{(Q)}$. (For the sheaf associated to a presheaf compare for example with Proposition-Definition 1.2 and its proof in Chapter II of [Ha].)

Let $f^* : A \rightarrow B$ be a morphism of graded algebras. We now construct a morphism (f, f^*) of the locally ringed spaces $(\text{Proj}(B), \mathcal{O}_{\text{Proj}(B)})$ and $(\text{Proj}(A), \mathcal{O}_{\text{Proj}(A)})$, which we call the morphism induced by $f^* : A \rightarrow B$.

For f we take the continuous map

$$f : \begin{array}{ccc} \text{Proj}(B) & \rightarrow & \text{Proj}(A) \\ Q & \rightarrow & (f^*)^{-1}(Q) \end{array}.$$

of Theorem 2.4.

A morphism of presheaves $\tilde{f}^* : \tilde{\mathcal{O}}_{\text{Proj}(A)} \rightarrow f_*(\tilde{\mathcal{O}}_{\text{Proj}(B)})$ is obtained as follows: For $\emptyset \neq U \subseteq \text{Proj}(A)$, U open, set

$$\tilde{f}^*(U) : \begin{array}{ccc} \tilde{\mathcal{O}}_{\text{Proj}(A)}(U) & \rightarrow & \tilde{\mathcal{O}}_{\text{Proj}(B)}(f^{-1}(U)) \\ \frac{a}{b} & \rightarrow & \frac{f^*(a)}{f^*(b)} \end{array}.$$

It is easy to check that for $Q \in \text{Proj}(B)$ the morphism induced on the stalks of the structure presheaves is given by the local morphism

$$\begin{aligned} f_Q^* : A_{(f(Q))} &\rightarrow B_{(Q)} \\ \frac{a}{b} &\rightarrow \frac{f^*(a)}{f^*(b)}. \end{aligned}$$

of Theorem 2.4.

This morphism of presheaves induces a morphism of sheaves $f^* : \mathcal{O}_{\text{Proj}(A)} \rightarrow f_*(\mathcal{O}_{\text{Proj}(B)})$ such that the local morphisms $f_Q^* : A_{(f(Q))} \rightarrow B_{(Q)}$, $Q \in \text{Proj}(B)$, are the induced morphisms on the stalks of the structure sheaves. (Construct the associated sheaves as in the proof of Proposition-Definition 1.2 in Chapter II of [Ha]. Let U be a nonempty open set in $\text{Proj}(A)$. Then $f^*(U)$ maps the section $s : U \rightarrow \prod_{P \in U} A_{(P)}$ in $\mathcal{O}_{\text{Proj}(A)}(U)$ to the section $f^*(U)s : f^{-1}(U) \rightarrow \prod_{Q \in f^{-1}(U)} B_{(Q)}$ in $\mathcal{O}_{\text{Proj}(B)}(f^{-1}(U))$ obtained by $(f^*(U)s)(Q) := f_Q^*(s(f(Q)))$, $Q \in f^{-1}(U)$.)

From Theorem 2.6 it follows immediately

Corollary 2.16 *Let A be a graded algebra and $a \in A$ be homogeneous, not nilpotent. The canonical morphism $i^* : A \rightarrow \{a^n | n \in \mathbb{N}_0\}^{-1}A$ induces an isomorphism of the restriction $(D(a), \mathcal{O}_{\text{Proj}(A)}|_{D(a)})$ and $(\text{Proj}(\{a^n | n \in \mathbb{N}_0\}^{-1}A), \mathcal{O}_{\text{Proj}(\{a^n | n \in \mathbb{N}_0\}^{-1}A)})$.*

From Theorem 2.13 it follows immediately

Corollary 2.17 *Let A be a \mathcal{M} -graded algebra. Let \mathcal{N} be a free commutative group, and let $\mathbb{F}[\mathcal{N}]$ be its \mathcal{N} -graded group algebra. The canonical map $i_A^* : A \rightarrow A \otimes \mathbb{F}[\mathcal{N}]$ induces an isomorphism of $(\text{Proj}(A), \mathcal{O}_{\text{Proj}(A)})$ and $(\text{Proj}(A \otimes \mathbb{F}[\mathcal{N}]), \mathcal{O}_{\text{Proj}(A \otimes \mathbb{F}[\mathcal{N}]})$.*

3 The projective spectrum of the Cartan algebra and its \mathbb{F} -valued points

In this section we describe the projective spectrum $\text{Proj}(CA)$ of the P^+ -graded Cartan algebra CA associated to a Kac-Moody group G . We show that its \mathbb{F} -valued points $\text{Proj}(CA)(\mathbb{F})$ identify with a completion Ω_{f_n} of the building Ω associated to G .

3.1 The Cartan algebra

The Cartan algebra CA associated to the Kac-Moody group G is a commutative associative unital P^+ -graded algebra over the field \mathbb{F} without zero-divisors. It is obtained by the following construction: Take as P^+ -graded \mathbb{F} -linear space

$$CA := \bigoplus_{\Lambda \in P^+} L(\Lambda)^{(*)},$$

where $L(\Lambda)^{(*)} := \bigoplus_{\lambda \in P(\Lambda)} L(\Lambda)_\lambda^* \subseteq L(\Lambda)^*$ is the restricted \mathbb{F} -linear dual of $L(\Lambda)$. To define the product of CA fix for every $\Lambda \in P^+$ a highest weight vector $v_\Lambda \in L(\Lambda)_\Lambda \setminus \{0\}$. For $\Lambda, N \in P^+$ the G -module $L(\Lambda) \otimes L(N)$ decomposes in the form

$$L(\Lambda) \otimes L(N) = \underbrace{L_{high}}_{\cong L(\Lambda+N)} \oplus \underbrace{L_{low}}_{\text{only isotypical components of type } L(M), M < \Lambda+N}.$$

Since $v_\Lambda \otimes v_N$ is a highest weight vector of L_{high} , it is possible to define a G -equivariant linear map

$$\Phi : L(\Lambda + N) \rightarrow L(\Lambda) \otimes L(N) \quad \text{by} \quad \Phi(v_{\Lambda+N}) := v_\Lambda \otimes v_N.$$

The product \bullet of the Cartan algebra CA between the parts $L(\Lambda)^{(*)}$, $L(N)^{(*)}$ is now obtained dually to Φ :

$$\bullet : L(\Lambda)^{(*)} \times L(N)^{(*)} \rightarrow L(\Lambda)^{(*)} \otimes L(N)^{(*)} \rightarrow (L(\Lambda) \otimes L(N))^{(*)} \xrightarrow{\Phi^{(*)}} L(\Lambda + N)^{(*)}.$$

The unit of the Cartan algebra CA is given by the element $\delta_0 \in L(0)_0^* \subseteq CA$ which satisfies $\delta_0(v_0) = 1$.

Algebras obtained in this way by different choices of highest weight vectors in $L(\Lambda)_\Lambda \setminus \{0\}$, $\Lambda \in P^+$, are isomorphic.

The Borel-Weil map $BW : CA \rightarrow \mathbb{F}[G]^U$ of Section 1 is a G -equivariant isomorphism of algebras.

3.2 Some actions on the projective spectrum of the Cartan algebra

Fix $\Lambda \in P^+$. The face monoid \widehat{G} acts on the highest weight module $L(\Lambda)$. This action induces an action of the opposite monoid \widehat{G}^{op} on $L(\Lambda)^{(*)}$.

The \mathfrak{g} -module $L(\Lambda)^{(*)}$ is a lowest weight module of lowest weight $-\Lambda$. The formal Kac-Moody group G_{fn} acts on $L(\Lambda)^{(*)}$. Dually, we get an action of G_{fn} on

$$L(\Lambda)_f := \prod_{\lambda \in P(\Lambda)} L(\Lambda)_\lambda = \prod_{\lambda \in P(\Lambda)} (L(\Lambda)_\lambda^*)^* = (L(\Lambda)^{(*)})^*,$$

where $L(\Lambda)_\lambda$ has been identified canonically with $(L(\Lambda)_\lambda^*)^*$, $\lambda \in P(\Lambda)$.

The Kac-Moody group G sits inside \widehat{G} and G_{fn} . The actions introduced before restrict to the same action of G on $L(\Lambda)$. They restrict to opposite actions of G on $L(\Lambda)^{(*)}$, which is cumbersome. Therefore, in this article we work with the action $(G_{fn})^{op}$ on $L(\Lambda)^{(*)}$ obtained from the action of G_{fn} on $L(\Lambda)^{(*)}$ by concatenation the corresponding representation with the inverse map. Similarly, we work with the action of $(\mathfrak{g}_{fn})^{op} \supseteq \mathfrak{g}^{op}$ on $L(\Lambda)^{(*)}$ obtained from the action of $\mathfrak{g}_{fn} \supseteq \mathfrak{g}$ on $L(\Lambda)^{(*)}$ by concatenation the corresponding representation with the multiplication by -1 .

The actions of \widehat{G}^{op} and $(G_{fn})^{op}$ on all $L(\Lambda)^{(*)}$, $\Lambda \in P^+$, combine to actions of \widehat{G}^{op} and $(G_{fn})^{op}$ on the Cartan algebra CA by morphisms of graded algebras.

To keep our notation clear and simple we denote the representations of the opposite monoid \widehat{G}^{op} , the opposite group $(G_{fn})^{op}$, the opposite Lie algebra $(\mathfrak{g}_{fn})^{op}$ by the same symbol π .

Now, for $x \in \widehat{G}$ resp. $x \in G_{fn}$ we get a continuous map on the spectrum $\text{Proj}(CA)$ of all P^+ -homogeneous prime ideals of CA by

$$xQ := \pi(x)^{-1}(Q) = \bigoplus_{\Lambda \in P^+} \pi(x)^{-1}(Q_\Lambda) \quad \text{where} \quad Q = \bigoplus_{\Lambda \in P^+} Q_\Lambda \in \text{Proj}(CA).$$

It leaves the spectrum of \mathbb{F} -valued points $\text{Proj}(CA)(\mathbb{F})$ invariant. Therefore, we get an action of \widehat{G} as well as an action of G_{fn} on the spectra $\text{Proj}(CA)$ and $\text{Proj}(CA)(\mathbb{F})$. (We could have combined the actions of \widehat{G} and G_{fn} to the action of a bigger monoid. For the aims of this article it is not advantageous.)

3.3 The completed building

The Kac-Moody group G has the opposite BN-pairs (B^\pm, N) with corresponding standard parabolic subgroups P_J^\pm , $J \subseteq I$.

The formal Kac-Moody group G_{fn} has the BN-pair (B_f^-, N) with corresponding standard parabolic subgroups $(P_{fn}^-)_J$, $J \subseteq I$. It holds $(P_{fn}^-)_J \cap G = P_J^-$. Furthermore, for every $\Lambda \in P^+ \cap F_J$ it holds

$$N_{G_{fn}}(L(\Lambda)_\Lambda^*) := \{g \in G_{fn} \mid \pi(g)L(\Lambda)_\Lambda^* = L(\Lambda)_\Lambda^*\} = (P_{fn}^-)_J.$$

Now we similarly extend the parabolic subgroups P_J , $J \subseteq I$, of G to subgroups of G_{fn} . For $J \subseteq I$ set

$$(P_{fn})_J := (U_f^-)_J P_J = (U_f^-)_J (W_J T) U.$$

In particular, $(P_{fn})_\emptyset = B$ and $(P_{fn})_I = G_{fn}$.

Theorem 3.1 *Let $J \subseteq I$.*

(a) $(P_{fn})_J$ is a subgroup of G_{fn} for which it holds $(P_{fn})_J \cap G = P_J$.

(b) For every $\Lambda \in P^+ \cap F_J$ it holds

$$N_{G_{fn}}(L(\Lambda)_\Lambda) := \{g \in G_{fn} \mid gL(\Lambda)_\Lambda = L(\Lambda)_\Lambda\} = (P_{fn})_J.$$

Proof: We first show (b). It is well-known that

$$N_G(L(\Lambda)_\Lambda) = P_J \quad \text{and} \quad N_{\mathbf{g}}(L(\Lambda)_\Lambda) := \{x \in \mathbf{g} \mid xL(\Lambda)_\Lambda \subseteq L(\Lambda)_\Lambda\} = \mathbf{p}_J := \mathbf{n}_J^- \oplus \mathbf{h} \oplus \mathbf{n}.$$

In particular, it holds $\mathbf{n}_J^- L(\Lambda)_\Lambda \subseteq L(\Lambda)_\Lambda$, from which it follows $(\mathbf{n}_J^-)_J L(\Lambda)_\Lambda \subseteq L(\Lambda)_\Lambda$, from which in turn it follows $(U_f^-)_J L(\Lambda)_\Lambda = L(\Lambda)_\Lambda$. Therefore, $(P_{fn})_J = (U_f^-)_J P_J \subseteq N_{G_{fn}}(L(\Lambda)_\Lambda)$. Now let $g \in N_{G_{fn}}(L(\Lambda)_\Lambda)$. Decompose g in the form $g = u^J u_J n_\sigma v$ with $u^J \in (U_f^-)^J$, $u_J \in (U_f^-)_J$, $n_\sigma \in N$ projecting to $\sigma \in \mathcal{W}$, and $v \in U^+$. From

$$L(\Lambda)_\Lambda = gL(\Lambda)_\Lambda = u^J u_J n_\sigma L(\Lambda)_\Lambda = u^J u_J L(\Lambda)_{\sigma\Lambda}$$

it follows $\sigma\Lambda = \Lambda$, which is equivalent to $\sigma \in \mathcal{W}_J$, and $u^J L(\Lambda)_\Lambda = L(\Lambda)_\Lambda$. Now u^J acts as

$$u^J = \exp(x) \quad \text{for some} \quad x = \sum_{\alpha \in (\Delta^-)^J} x_\alpha \in \prod_{\alpha \in (\Delta^-)^J} \mathbf{g}_\alpha.$$

Assume that $u^J \neq 1$. Choose a nonzero homogeneous component x_β with β of maximal height. Then $x_\beta v_\Lambda$ is the $(\Lambda + \beta)$ -homogeneous part of $u^J v_\Lambda$. It is nonzero because of $x_\beta \notin \mathbf{p}_J = N_{\mathbf{g}}(L(\Lambda)_\Lambda)$. This contradicts $u^J L(\Lambda)_\Lambda = L(\Lambda)_\Lambda$. Therefore, $u^J = 1$ and $g = u_J n_\sigma v \in (U_f^-)_J (\mathcal{W}_J T) U = (P_{fn})_J$.

Now (a) follows from (b) and $N_{G_{fn}}(L(\Lambda)_\Lambda) \cap G = N_G(L(\Lambda)_\Lambda) = P_J$. \square

The following easy Lemma will be used several times:

Lemma 3.2 *Let C be a subgroup of U_f^- , let D be a subgroup of N , and let E be a subgroup of U . It holds:*

$$CDE \cap U_f^- = C \quad \text{and} \quad CDE \cap N = D \quad \text{and} \quad CDE \cap U = E.$$

Proof: The inclusions ' \supseteq ' hold, because the unit 1 is contained in C , D , and E . The inclusion ' \subseteq ' of the second equality follows directly from the Birkhoff decomposition of G_{fn} . The inclusion ' \subseteq ' of the first and third equality are shown similarly, for the first as follows: Let $c \in C$, $d \in D$, $e \in E$, and $u \in U_f^-$ such that $cde = u$. Then it holds $(u^{-1}c)de = 1$, from which it follows $u^{-1}c = d = e = 1$. In particular, $u = c \in C$. \square

Proposition 3.3 *Let $J \subseteq I$. It holds $(P_{fn})_J = P_J$ if and only if $J = J^0$.*

Proof: It holds $(U_f^-)_J = U_J^-$ if and only if $J = J^0$. Now the Proposition follows from the definition of $(P_{fn})_J$ and Lemma 3.2. \square

Let $J \subseteq I$. The parabolic subgroup $(P_{fn}^-)_J$ has the Levi decomposition

$$(P_{fn}^-)_J = (U_f^-)^J \rtimes (L_{fn})_J$$

with $(L_{fn})_J := (U_f^-)_J (\mathcal{W}_J T) (U_f^-)_J = (U_f^-)_J (\mathcal{W}_J T) U_J^- = (U_f^-)_J (\mathcal{W}_J T) U_J$. Similarly, $(P_{fn})_J$ has a Levi decomposition:

Theorem 3.4 *For $J \subseteq I$ it holds $(P_{fn})_J = (L_{fn})_J \rtimes U^J$.*

Proof: This holds trivially for $J = I$. Let $J \neq I$. $(L_{fn})_J$ and U^J are subgroups of G_{fn} such that it holds

$$(P_{fn})_J = (U_f^-)_J(\mathcal{W}_J T)U = (U_f^-)_J(\mathcal{W}_J T)U_J U^J = (L_{fn})_J U^J.$$

With Lemma 3.2 it follows $(L_{fn})_J \cap U^J = (L_{fn})_J \cap U \cap U^J = U_J \cap U^J = \{1\}$.

It remains to show that $(L_{fn})_J = (U_f^-)_J(\mathcal{W}_J T)(U_f^-)_J$ normalizes U^J . The group $\mathcal{W}_J T$ normalizes U^J because it holds $U^J = \bigcap_{\sigma \in \mathcal{W}_J} \sigma U \sigma^{-1}$. To show that $(U_f^-)_J$ normalizes U^J we need some preparations before:

(a) We first show

$$U = \{g \in G_{fn} \mid gv_\Lambda = v_\Lambda \text{ for all } \Lambda \in P^+ \cap C\}. \quad (6)$$

Trivially, the inclusion ' \subseteq ' holds. To show ' \supseteq ' fix $\Lambda \in P^+ \cap C$ and let $g \in G_{fn}$ such that $gv_\Lambda = v_\Lambda$. Then also $gL(\Lambda)_\Lambda = L(\Lambda)_\Lambda$. By Theorem 3.1 (b) it follows $g \in B = UT$. Write g in the form

$$g = u \prod_{i=1}^{2n-l} t_i(s_i) \quad \text{with} \quad u \in U \text{ and } s_1, s_2, \dots, s_{2n-l} \in \mathbb{F}^\times.$$

From $gv_\Lambda = v_\Lambda$ it follows $\prod_{i=1}^{2n-l} (s_i)^{\Lambda(h_i)} = 1$. For $k \in \{1, 2, \dots, 2n-l\}$ it holds $\Lambda + \Lambda_k \in P^+ \cap C$. Suppose that in addition $gv_{\Lambda+\Lambda_k} = v_{\Lambda+\Lambda_k}$. Then also $\prod_{i=1}^{2n-l} (s_i)^{(\Lambda+\Lambda_k)(h_i)} = s_k \prod_{i=1}^{2n-l} (s_i)^{\Lambda(h_i)} = 1$. It follows $s_k = 1$.

For every $\sigma \in \mathcal{W}_J$ choose $n_\sigma \in N$ projecting to σ . By (6) we find

$$U^J = \bigcap_{\sigma \in \mathcal{W}_J} \sigma U \sigma^{-1} = \{g \in G_{fn} \mid g n_\sigma v_\Lambda = n_\sigma v_\Lambda \text{ for all } \sigma \in \mathcal{W}_J \text{ and all } \Lambda \in P^+ \cap C\}. \quad (7)$$

(b) It holds $(U_f^-)_J = \exp((\mathbf{n}_f^-)_J)$ with $(\mathbf{n}_f^-)_J = \bigoplus_{\alpha \in \Delta_J^-} \mathbf{g}_\alpha$. If $v_\lambda \in L(\Lambda)_\lambda$, $\lambda \in P$, it follows

$$(U_f^-)_J v_\lambda \subseteq \prod_{\mu \in \lambda - (Q_0^+)_J} L(\Lambda)_\mu.$$

The group U^J is also a subgroup of G_{fp} , which acts on $L(\Lambda)$. In particular, it holds $U^J \subseteq (U_f^+)^J = \exp((\mathbf{n}_f^+)^J)$ with $(\mathbf{n}_f^+)^J = \bigoplus_{\alpha \in \Delta^+ \setminus \Delta_J^+} \mathbf{g}_\alpha$. If $v_\lambda \in L(\Lambda)_\lambda$, $\lambda \in P$, it follows

$$U^J v_\lambda \subseteq v_\lambda + \bigoplus_{\mu \in \lambda + Q_0^+ \setminus (Q_0^+)_J} L(\Lambda)_\mu.$$

(c) Let $u \in U^J$ and $u^- \in (U_f^-)_J$. We use (7) to show $u^- u (u^-)^{-1} \in U^J$. Let $\sigma \in \mathcal{W}_J$ and $\Lambda \in P^+ \cap C$. By (b) it follows

$$u(u^-)^{-1} n_\sigma v_\Lambda \in (u^-)^{-1} n_\sigma v_\Lambda + \prod_{\mu \in \sigma \Lambda - (Q_0^+)_J + Q_0^+ \setminus (Q_0^+)_J} L(\Lambda)_\mu.$$

By Theorem 3.12 d) of [K1] applied to the fundamental chamber in $\mathbf{h}_\mathbb{R}^*$ it holds $\Lambda - \sigma \Lambda \in Q_0^+$. Because of $\sigma \in \mathcal{W}_J$ we find $\Lambda - \sigma \Lambda \in (Q_0^+)_J$. Therefore, we get

$$\sigma \Lambda - (Q_0^+)_J + Q_0^+ \setminus (Q_0^+)_J \subseteq \Lambda - (Q_0^+)_J + Q_0^+ \setminus (Q_0^+)_J.$$

An element μ of the set on the right is of the form

$$\mu = \Lambda - \sum_{i \in J} n_i \alpha_i + \sum_{i \in I} m_i \alpha_i = \Lambda - \sum_{i \in J} (n_i - m_i) \alpha_i + \sum_{i \in I \setminus J} m_i \alpha_i$$

with $n_i \in \mathbb{N}_0$, $i \in J$, with $m_i \in \mathbb{N}_0$, $i \in I$, and $m_i \neq 0$ for at least one $i \in I \setminus J$. Therefore, $\mu \notin \Lambda - Q_0^+ \supseteq P(\Lambda)$. It follows $L(\Lambda)_\mu = \{0\}$.

We have shown $u^- u (u^-)^{-1} n_\sigma v_\Lambda = u^- (u^-)^{-1} n_\sigma v_\Lambda = n_\sigma v_\Lambda$. \square

There are the following modified generalized Birkhoff decompositions:

Theorem 3.5 *Let $J \subseteq I$. It holds*

$$G_{fn} = \bigcup_{\sigma \in \mathcal{W}^J} U_f^- \sigma (P_{fn})_J = \bigcup_{\sigma \in \mathcal{W}^J} (U_f^- \cap \sigma(U_f^-)^J \sigma^{-1}) \sigma (P_{fn})_J.$$

Furthermore, for $\sigma \in \mathcal{W}^J$ and $u_1, u_2 \in U_f^- \cap \sigma(U_f^-)^J \sigma^{-1}$ it holds

$$u_1 \sigma (P_{fn})_J = u_2 \sigma (P_{fn})_J \iff u_1 = u_2.$$

Proof: (a) It holds $G_{fn} = G_{fn}(P_{fn})_J = U_f^- G(P_{fn})_J$. From the generalized Birkhoff decomposition of G it follows

$$G_{fn} = U_f^- \left(\bigcup_{\sigma \in \mathcal{W}^J} U^- \sigma P_J \right) (P_{fn})_J = \bigcup_{\sigma \in \mathcal{W}^J} U_f^- \sigma (P_{fn})_J.$$

This union is also disjoint. Suppose that $U_f^- \sigma (P_{fn})_J \cap U_f^- \sigma' (P_{fn})_J \neq \emptyset$. Then there exist $u, u' \in U_f^-$, $p, p' \in (P_{fn})_J$, and $n_\sigma, n_{\sigma'} \in N$ projecting to σ, σ' such that $un_\sigma p = u'n_{\sigma'} p'$. Apply this element to $L(\Lambda)_\Lambda$, where $\Lambda \in P^+ \cap F_J$ is chosen arbitrarily. With Theorem 3.1 (b) it follows $uL(\Lambda)_{\sigma\Lambda} = u'L(\Lambda)_{\sigma'\Lambda}$, from which in turn it follows $\sigma\Lambda = \sigma'\Lambda$, which is equivalent to $\sigma\mathcal{W}_J = \sigma'\mathcal{W}_J$. Because σ, σ' are minimal coset representatives, we find $\sigma = \sigma'$.

(b) Let $\sigma \in \mathcal{W}^J$. It holds $U_f^- = (U_f^- \cap \sigma(U_f^-) \sigma^{-1})(U_f^- \cap \sigma U \sigma^{-1})$, which is a property of refined Tits systems. Therefore, we find

$$\begin{aligned} U_f^- \sigma (P_{fn})_J &= (U_f^- \cap \sigma(U_f^-) \sigma^{-1})(U_f^- \cap \sigma U \sigma^{-1}) \sigma (P_{fn})_J \\ &= (U_f^- \cap \sigma(U_f^-) \sigma^{-1}) \sigma (\sigma^{-1} U_f^- \sigma \cap U) (P_{fn})_J = (U_f^- \cap \sigma(U_f^-) \sigma^{-1}) \sigma (P_{fn})_J. \end{aligned} \quad (8)$$

Next we show

$$U_f^- \cap \sigma(U_f^-) \sigma^{-1} = (U_f^- \cap \sigma(U_f^-)^J \sigma^{-1}) \sigma (U_f^-)_J \sigma^{-1}. \quad (9)$$

Because σ has minimal length in $\sigma\mathcal{W}_J$ it holds $l(\sigma\sigma_j) > l(\sigma)$ for all $j \in J$, from which it follows $\sigma\Delta_J^- \subseteq \Delta^-$ by Lemma 3.11 (a) in [K1]. Choose $n_\sigma \in N$ projecting to σ . Then it holds

$$Ad(n_\sigma)(\mathbf{n}_f^-)_J = Ad(n_\sigma) \prod_{\alpha \in \Delta_J^-} \mathbf{g}_\alpha = \prod_{\alpha \in \Delta_J^-} \mathbf{g}_{\sigma\alpha} \subseteq \mathbf{n}_f^- ,$$

from which we find

$$\sigma(U_f^-)_J \sigma^{-1} = n_\sigma \exp((\mathbf{n}_f^-)_J) n_\sigma^{-1} = \exp(Ad(n_\sigma)(\mathbf{n}_f^-)_J) \subseteq U_f^- . \quad (10)$$

It follows the inclusion ' \supseteq ' of (9). To show the reverse inclusion ' \subseteq ' let $a = bc$ with $a \in U_f^-$ and $b \in \sigma(U_f^-)^J \sigma^{-1}$, $c \in \sigma(U_f^-)_J \sigma^{-1}$. With (10) it follows $b = ac^{-1} \in U_f^- \cap \sigma(U_f^-)^J \sigma^{-1}$.

Inserting (9) in (8) we find

$$U_f^- \sigma (P_{fn})_J = (U_f^- \cap \sigma(U_f^-)^J \sigma^{-1}) \sigma (U_f^-)_J (P_{fn})_J = (U_f^- \cap \sigma(U_f^-)^J \sigma^{-1}) \sigma (P_{fn})_J.$$

(c) Let $\sigma \in \mathcal{W}^J$ and $u_1, u_2 \in U_f^- \cap \sigma(U_f^-)^J \sigma^{-1}$ that $u_1 \sigma (P_{fn})_J = u_2 \sigma (P_{fn})_J$. With Lemma 3.2 it follows

$$u_1^{-1} u_2 \in \sigma (P_{fn})_J \sigma^{-1} \cap (U_f^- \cap \sigma(U_f^-)^J \sigma^{-1}) \subseteq \sigma((P_{fn})_J \cap (U_f^-)^J) \sigma^{-1} = \sigma((U_f^-)_J \cap (U_f^-)^J) \sigma^{-1} = \{1\}.$$

□

The next two theorems show: The system of nonparabolic subgroups $(P_J)_{fn}$, $J \subseteq I$, of G_{fn} has many properties in common with systems of parabolic subgroups.

Theorem 3.6 *Let $J, K \subseteq I$. It holds:*

(a) $(P_{fn})_J \subseteq (P_{fn})_K$ if and only if $J \subseteq K$.

(b) $(P_{fn})_J \cap (P_{fn})_K = (P_{fn})_{J \cap K}$.

(c) In general, $(P_{fn})_{J \cup K}$ is not generated by $(P_{fn})_J$ and $(P_{fn})_K$ as a group.

Proof: (a) If $J \subseteq K$ it holds $(U_f^-)_J \subseteq (U_f^-)_K$, from which it follows $(P_{fn})_J \subseteq (P_{fn})_K$. If $(P_{fn})_J \subseteq (P_{fn})_K$ we find $P_J \subseteq P_K$ by Theorem 3.1 (a). It follows $J \subseteq K$.

The inclusion " \supseteq " of part (b) follows from part (a). Now let $g \in (P_{fn})_J \cap (P_{fn})_K$. By the Birkhoff decomposition of G_{fn} , and by the definitions of $(P_{fn})_J$, $(P_{fn})_K$, the element g can be written in the forms

$$g = u_J n_\sigma u = \tilde{u}_K n_\sigma \tilde{u}$$

with $u, \tilde{u} \in U$, $n_\sigma \in N$ projecting to $\sigma \in \mathcal{W}_J \cap \mathcal{W}_K = \mathcal{W}_{J \cap K}$ and $u_J \in (U_f^-)_J$, $\tilde{u}_K \in (U_f^-)_K$. It follows

$$(u_J)^{-1} \tilde{u}_K \in U_f^- \cap \sigma U \sigma^{-1} = U^- \cap \sigma U \sigma^{-1} = \prod_{\alpha \in \Delta_{re}^- \cap \sigma \Delta_{re}^+} U_\alpha.$$

Since $\sigma \in \mathcal{W}_{J \cap K}$ it holds $\Delta_{re}^- \cap \sigma \Delta_{re}^+ \subseteq (\Delta_{J \cap K})_{re}^-$. Therefore, we get

$$\tilde{u}_K \in (U_f^-)_K \cap (U_f^-)_J U_{J \cap K}^- = (U_f^-)_K \cap (U_f^-)_J = \exp((\mathbf{n}_f^-)_K) \cap \exp((\mathbf{n}_f^-)_J).$$

Because the exponential function $\exp : \mathbf{n}_f^- \rightarrow U_f^-$ is bijective it holds

$$\exp((\mathbf{n}_f^-)_K) \cap \exp((\mathbf{n}_f^-)_J) = \exp((\mathbf{n}_f^-)_K \cap (\mathbf{n}_f^-)_J) = \exp((\mathbf{n}_f^-)_{J \cap K}) = (U_f^-)_{J \cap K}.$$

It follows $g = \tilde{u}_K n_\sigma \tilde{u} \in (U_f^-)_{J \cap K} (\mathcal{W}_{J \cap K} T) U = (P_{fn})_{J \cap K}$.

To show (c) suppose that for all $J, K \subseteq I$ the group $(P_{fn})_{J \cup K}$ is generated by the groups $(P_{fn})_J$ and $(P_{fn})_K$. Then $G_{fn} = (P_{fn})_I$ is generated by the groups $(P_{fn})_{\{i\}} = P_{\{i\}}$, $i \in I$. It follows $G_{fn} = G$, which is only possible if all components of the generalized Cartan matrix A are of finite type. \square

Theorem 3.7 *Let $J, K \subseteq I$ and $g \in G_{fn}$. It is equivalent:*

(i) $g(P_{fn})_J g^{-1} \subseteq (P_{fn})_K$.

(ii) $J \subseteq K$ and $g \in (P_{fn})_K$.

In particular: The subgroups $(P_{fn})_J$, $(P_{fn})_K$ of G_{fn} are conjugated if and only if $J = K$. The subgroup $(P_{fn})_J$ of G_{fn} coincides with its normalizer.

Proof: It only remains to show that (i) implies (ii). Here a well-known argument to compute representation theoretically the normalizers of parabolics also applies:

(a) Fix $\Lambda \in P^+$. Suppose that V is a one-dimensional subspace of $L(\Lambda)_f$ such that $BV \subseteq V$. We show $V = L(\Lambda)_\Lambda$:

Let $v \in V \setminus \{0\}$. Decompose

$$v = \sum_{\lambda \in \text{supp}(v)} v_\lambda \quad \text{with} \quad \text{supp}(v) := \{\lambda \in P(\Lambda) \mid v_\lambda \neq 0\} \neq \emptyset$$

according to the decomposition $L(\Lambda)_f = \prod_{\lambda \in P(\Lambda)} L(\Lambda)_\lambda$.

We first show $|\text{supp}(v)| = 1$. Suppose there exist $\lambda_1, \lambda_2 \in \text{supp}(v)$, $\lambda_1 \neq \lambda_2$. Choose $h \in H$ such that $\lambda_1(h) \neq \lambda_2(h)$. Choose $s \in \mathbb{N} \subseteq \mathbb{F}^\times$ such that $s^{\lambda_1(h)} \neq s^{\lambda_2(h)}$. Then it follows

$$t_h(s)v = \sum_{\lambda \in \text{supp}(v)} s^{\lambda(h)} v_\lambda \notin \mathbb{F}v,$$

which contradicts $TV \subseteq V$.

Now let $v = v_\lambda$. Suppose that $\mathbf{n}v_\lambda \neq \{0\}$. Since \mathbf{n} is generated by e_i , $i \in I$, there exists an index i such that $e_i v_\lambda \neq 0$. It follows

$$\exp(e_i)v_\lambda = \underbrace{v_\lambda}_{\in L(\Lambda)_\lambda \setminus \{0\}} + \underbrace{e_i v_\lambda}_{\in L(\Lambda)_{\lambda+\alpha_i} \setminus \{0\}} + \sum_{k \in \mathbb{N}_0, k \geq 2} \underbrace{\frac{1}{k!}(e_i)^k v_\lambda}_{\in L(\Lambda)_{\lambda+k\alpha_i}} \notin \mathbb{F}v_\lambda,$$

which contradicts $UV \subseteq V$.

We have shown $v \in L(\Lambda)_\lambda$ for some $\lambda \in P(\Lambda)$ and $\mathbf{n}v = \{0\}$. It follows $v \in L(\Lambda)_\Lambda$ by Proposition 9.3 b) in [K1].

(b) Now suppose that (i) holds. Choose $\Lambda \in P^+ \cap F_K$. By (i) and Theorem 3.1 (b) it follows

$$Bg^{-1}L(\Lambda)_\Lambda \subseteq (P_{fn})_J g^{-1}L(\Lambda)_\Lambda \subseteq g^{-1}(P_{fn})_K L(\Lambda)_\Lambda = g^{-1}L(\Lambda)_\Lambda.$$

Now $g^{-1}L(\Lambda)_\Lambda$ is a one-dimensional subspace of $L(\Lambda)_f$. By (a) it follows $g^{-1}L(\Lambda)_\Lambda = L(\Lambda)_\Lambda$. Again by Theorem 3.1 (b) we find $g \in (P_{fn})_K$. With (i) we get $(P_{fn})_J \subseteq (P_{fn})_K$. By Theorem (3.6) (a) it follows $J \subseteq K$. \square

We take as *completed building* Ω_{fn} associated to the formal Kac-Moody group G_{fn} the set

$$\Omega_{fn} := \bigcup_{J \subseteq I} G_{fn}/(P_{fn})_J = \{g(P_{fn})_J \mid g \in G_{fn}, J \subseteq I\}$$

partially ordered by the reverse inclusion, i.e., for $g, g' \in G_{fn}$ and $J, J' \subseteq I$,

$$g(P_{fn})_J \leq g'(P_{fn})_{J'} : \iff g(P_{fn})_J \supseteq g'(P_{fn})_{J'}.$$

The group G_{fn} acts order preservingly on Ω_{fn} by multiplication from the left. We denote by

$$\mathcal{A}_{fn} := \{n(P_{fn})_J \mid n \in N, J \subseteq I\}$$

the *standard apartment* of Ω_{fn} . The completed building Ω_{fn} is covered by the apartments $g\mathcal{A}_{fn}$, $g \in G_{fn}$.

Remark 3.8 (a) At first, the construction of Ω_{fn} by the nonparabolic subgroups $(P_J)_{fn}$, $J \subseteq I$, of G_{fn} may look strange. We give a hand-waving motivation, which indicates that it is natural:

Let $J \subseteq I$. Consider first the classical case, i.e., all components of the generalized Cartan matrix A are of finite type, and take $\mathbb{F} = \mathbb{C}$. Here G/P_J is a well-known manifold, a partial flag manifold. It is covered by big cells, which are charts of the manifold. The standard big cell $BC(J)$ of G/P_J is obtained as the image of U^- in G/P_J . As a set it is given by

$$BC(J) \cong U^- / (U^- \cap P_J) = U^- / U_J^- \cong (U^-)^J.$$

Now consider the case of an arbitrary generalized Cartan matrix. By analogy, the standard big cell of $G_{fn}/(P_{fn})_J$ is obtained as the image of U_f^- in $G_{fn}/(P_{fn})_J$. As a set it is given by

$$BC(J) \cong U_f^- / (U_f^- \cap (P_{fn})_J) = U_f^- / (U_f^-)_J \cong (U_f^-)^J.$$

(b) The completed building Ω_{fn} should be looked at as an equivariant completion of the building Ω associated to G as described in Section 4. It does not satisfy all properties of a building.

3.4 The projective spectrum of the Cartan algebra and its \mathbb{F} -valued points

We first show that the completed building Ω_{fn} embeds G_f -equivariantly into the \mathbb{F} -valued points $\text{Proj}(CA)(\mathbb{F})$ of the spectrum $\text{Proj}(CA)$.

For $\Lambda \in P^+$ let $\delta_\Lambda \in L(\Lambda)_\Lambda^* \subseteq CA$ be defined by $\delta_\Lambda(v_\Lambda) := 1$. It is easy to check that for all $\Lambda, N \in P^+$ it holds $\delta_\Lambda \bullet \delta_N = \delta_{\Lambda+N}$. Therefore,

$$\widetilde{P^+} := \{ \delta_\Lambda \mid \Lambda \in P^+ \}$$

is a multiplicatively closed subset of the Cartan algebra CA , isomorphic to $(P^+, +)$. For $M \subseteq P^+$ we set

$$\widetilde{M} := \{ \delta_\Lambda \mid \Lambda \in M \} \subseteq \widetilde{P^+}.$$

For $\Lambda \in P^+$ we set

$$L(\Lambda)_{\neq \Lambda}^{(*)} := \left\{ \phi \in L(\Lambda)^{(*)} \mid \phi(v_\Lambda) = 0 \right\} = \bigoplus_{\lambda \in P(\Lambda) \setminus \{\Lambda\}} L(\Lambda)_\lambda^*.$$

Here, as always, a sum over the empty set is defined to be $\{0\}$.

Theorem 3.9 *For $J \subseteq I$ define*

$$P(J) := \bigoplus_{\Lambda \in P^+ \cap \overline{F_J}} L(\Lambda)_{\neq \Lambda}^{(*)} \oplus \bigoplus_{\Lambda \in P^+ \setminus \overline{F_J}} L(\Lambda)^{(*)} \subseteq CA.$$

A G_{fn} -equivariant embedding of the G_{fn} -set Ω_{fn} into the G_{fn} -set $\text{Proj}(CA)(\mathbb{F})$ is given by

$$\begin{array}{ccc} \omega : & \Omega_{fn} & \rightarrow \text{Proj}(CA)(\mathbb{F}) \\ & g(P_{fn})_J & \mapsto gP(J) \end{array}.$$

Furthermore, for $g(P_{fn})_J, h(P_{fn})_K \in \Omega_{fn}$ it holds

$$g(P_{fn})_J \leq h(P_{fn})_K \iff \overline{\{gP(J)\}}^{pts} \subseteq \overline{\{hP(K)\}}^{pts}.$$

Proof: Obviously, $P(J)$ is a P^+ -homogeneous linear space different from CA . We first show that $P(J)$ is an ideal. Let $N \in P^+$. If $\Lambda \in P^+ \setminus \overline{F_J}$ then also $N + \Lambda \in P^+ \setminus \overline{F_J}$, because $P^+ \cap \overline{F_J}$ is a face of P^+ . It follows

$$L(N)^{(*)} \bullet P(J)_\Lambda = L(N)^{(*)} \bullet L(\Lambda)^{(*)} \subseteq L(N + \Lambda)^{(*)} = P(J)_{N+\Lambda}.$$

Now let $\Lambda \in P^+ \cap \overline{F_J}$. For every $\phi \in L(N)^{(*)}$ and every $\psi \in L(\Lambda)_{\neq \Lambda}^{(*)}$ it holds

$$(\phi \bullet \psi)(v_{N+\Lambda}) = \phi(v_N) \underbrace{\psi(v_\Lambda)}_{=0} = 0.$$

It follows

$$L(N)^{(*)} \bullet P(J)_\Lambda = L(N)^{(*)} \bullet L(\Lambda)_{\neq \Lambda}^{(*)} \subseteq L(N + \Lambda)_{\neq \Lambda}^{(*)} \subseteq P(J)_{N+\Lambda}.$$

Next we show that $P(J)$ is prime. $CA/P(J)$ is isomorphic in the obvious way to the monoid algebra $\mathbb{F}[\widetilde{P^+ \cap \overline{F_J}}]$, which identifies with the monoid algebra $\mathbb{F}[P^+ \cap \overline{F_J}]$, which is a subalgebra of the group algebra $\mathbb{F}[P]$ of the lattice P . By Remark 2.12 the algebra $\mathbb{F}[P]$ has no zero divisors. Therefore, $CA/P(J)$ has no zero divisors.

It holds $\dim(L(\Lambda)^{(*)}/P(J)_\Lambda) \leq 1$ for all $\Lambda \in P^+$. By Theorem 2.1 it follows $P(J) \in \text{Proj}(CA)(\mathbb{F})$. ω is a well-defined G_{fn} -equivariant embedding if and only if for all $J \subseteq I$ it holds

$$\text{Stab}_{G_{fn}}(P(J)) = (P_{fn})_J.$$

Let $g \in G_{fn}$. It holds $P(J) = gP(J) = \pi(g)^{-1}(P(J))$ if and only if the following equations are satisfied:

$$\pi(g)^{-1}L(\Lambda)^{(*)} = L(\Lambda)^{(*)} \quad \text{for all } \Lambda \in P^+ \setminus \overline{F_J}, \quad (11)$$

$$\pi(g)^{-1}L(\Lambda)_{\neq \Lambda}^{(*)} = L(\Lambda)_{\neq \Lambda}^{(*)} \quad \text{for all } \Lambda \in P^+ \cap \overline{F_J}. \quad (12)$$

Equations (11) always hold. Written differently, equations (12) are

$$\left\{ \phi \in L(\Lambda)^{(*)} \mid \phi(gv_\Lambda) = 0 \right\} = \left\{ \phi \in L(\Lambda)^{(*)} \mid \phi(v_\Lambda) = 0 \right\} \quad \text{for all } \Lambda \in P^+ \cap \overline{F_J}.$$

Here $\mathbb{F}gv_\Lambda$ and $\mathbb{F}v_\Lambda$ are finite dimensional subspaces of $L(\Lambda)_f$, which is paired nondegenerately with $L(\Lambda)^{(*)}$. Therefore, these equations are equivalent to $\mathbb{F}gv_\Lambda = \mathbb{F}v_\Lambda$ for all $\Lambda \in P^+ \cap \overline{F_J}$. (To check this directly use a base adapted to the decomposition $L(\Lambda)^{(*)} = \bigoplus_{\lambda \in P(\Lambda)} L(\Lambda)_\lambda^*$, $\Lambda \in P^+ \cap \overline{F_J}$.) By Theorem 3.1 (b) and Theorem 3.6 (a) this is in turn equivalent to $g \in \bigcap_{K \supseteq J} (P_{fn})_K = (P_{fn})_J$.

Now, $g(P_{fn})_J \leq h(P_{fn})_K$ is by definition equivalent to $g(P_{fn})_J \supseteq h(P_{fn})_K$, which is equivalent $(P_{fn})_J \supseteq g^{-1}h(P_{fn})_K$, which in turn is equivalent to $J \supseteq K$ and $g^{-1}h \in (P_{fn})_J$. On the other hand, it holds $\overline{\{gP(J)\}}^{pts} \subseteq \overline{\{hP(K)\}}^{pts}$ if and only if $gP(J) \in \overline{\{hP(K)\}}^{pts} = \overline{\{hP(K)\}} \cap \text{Proj}(CA)(\mathbb{F})$ if and only if $gP(J) \supseteq hP(K)$ if and only if $P(J) \supseteq g^{-1}hP(K) = \pi(g^{-1}h)^{-1}(P(K))$, which is equivalent to the inclusions

$$\bigoplus_{\Lambda \in P^+ \setminus \overline{F_J}} L(\Lambda)^{(*)} \supseteq \bigoplus_{\Lambda \in P^+ \setminus \overline{F_K}} L(\Lambda)^{(*)}, \quad (13)$$

$$\bigoplus_{\Lambda \in P^+ \cap \overline{F_J}} L(\Lambda)_{\neq \Lambda}^{(*)} \supseteq \bigoplus_{\Lambda \in P^+ \cap \overline{F_K} \cap \overline{F_J}} \pi(g^{-1}h)^{-1}L(\Lambda)_{\neq \Lambda}^{(*)}. \quad (14)$$

Inclusion (13) is equivalent to $P^+ \setminus \overline{F_J} \supseteq P^+ \setminus \overline{F_K}$, which is equivalent to $\overline{F_J} \subseteq \overline{F_K}$, which in turn is equivalent to $J \supseteq K$. Therefore, inclusion (14) is equivalent to

$$L(\Lambda)_{\neq \Lambda}^{(*)} \supseteq \pi(g^{-1}h)^{-1}L(\Lambda)_{\neq \Lambda}^{(*)} \quad \text{for all } \Lambda \in P^+ \cap \overline{F_J}.$$

Here it holds always equality, because $L(\Lambda)_{\neq \Lambda}^{(*)}$ and $\pi(g^{-1}h)^{-1}L(\Lambda)_{\neq \Lambda}^{(*)} = \pi(h^{-1}g)L(\Lambda)_{\neq \Lambda}^{(*)}$ are 1-codimensional subspaces of $L(\Lambda)^{(*)}$. As we have seen by the calculation of the stabilizers, these equations are equivalent to $g^{-1}h \in (P_{fn})_J$. \square

It remains to show that the map ω of Theorem 3.9 is surjective, which is not straightforward. It is reached in Corollary 3.38 as a consequence of our description of the full projective spectrum $\text{Proj}(CA)$, which we investigate next. For that we first introduce a G_{fn} -invariant stratification of $\text{Proj}(CA)$, whose strata extend the orbits $G_{fn}P(J)$, $J \subseteq I$.

The Cartan algebra CA is a P^+ -graded algebra without zero divisors. The faces of P^+ are $P^+ \cap \overline{F_J}$, $J \subseteq I$. It follows that for every $J \subseteq I$ there is the semidirect decomposition

$$CA = \bigoplus_{\Lambda \in P^+ \cap \overline{F_J}} L(\Lambda)^{(*)} \oplus \bigoplus_{\Lambda \in P^+ \setminus \overline{F_J}} L(\Lambda)^{(*)} \quad (15)$$

with graded subalgebra $\bigoplus_{\Lambda \in P^+ \cap \overline{F_J}} L(\Lambda)^{(*)}$ and graded prime ideal $\bigoplus_{\Lambda \in P^+ \setminus \overline{F_J}} L(\Lambda)^{(*)}$, a sum over the empty set defined to be $\{0\}$.

For every $J \subseteq I$ set

$$\overline{Or(J)} := \mathcal{V}\left(\bigoplus_{\Lambda \in P^+ \setminus \overline{F_J}} L(\Lambda)^{(*)}\right) = \overline{\left\{ \bigoplus_{\Lambda \in P^+ \setminus \overline{F_J}} L(\Lambda)^{(*)} \right\}},$$

which is a G_{fn} -invariant closed subset of $\text{Proj}(CA)$. In particular, it holds $\overline{Or(\emptyset)} = \text{Proj}(CA)$ and $\overline{Or(I)} = \left\{ \bigoplus_{\Lambda \in P^+ \setminus \overline{F_I}} L(\Lambda)^{(*)} \right\}$. The following properties are trivial to check:

Proposition 3.10 *Let $K, J \subseteq I$. It holds:*

- (a) $\overline{Or(K)} \subseteq \overline{Or(J)}$ if and only if $K \supseteq J$.
- (b) $\overline{Or(J)} \cap \overline{Or(K)} = \overline{Or(J \cup K)}$.

As a closed set $\overline{Or(J)}$ can be described as a projective spectrum:

Proposition 3.11 *Let $J \subseteq I$. The map*

$$\begin{aligned} Proj\left(\bigoplus_{\Lambda \in P^+ \cap \overline{F_J}} L(\Lambda)^{(*)}\right) &\rightarrow \overline{Or(J)} \\ Q &\mapsto Q \oplus \bigoplus_{\Lambda \in P^+ \setminus \overline{F_J}} L(\Lambda)^{(*)} \end{aligned}$$

is a homeomorphism, mapping $Proj(\bigoplus_{\Lambda \in P^+ \cap \overline{F_J}} L(\Lambda)^{()})(\mathbb{F})$ bijectively to $\overline{Or(J)}(\mathbb{F})$.*

Proof: The restriction of the canonical map $CA \rightarrow CA / \bigoplus_{\Lambda \in P^+ \setminus \overline{F_J}} L(\Lambda)^{(*)}$ to the graded subalgebra $\bigoplus_{\Lambda \in P^+ \cap \overline{F_J}} L(\Lambda)^{(*)}$ is an isomorphism of graded algebras. The proposition now follows by Theorem 2.9. \square

For every $J \subseteq I$ set

$$Or(J) := \overline{Or(J)} \setminus \bigcup_{I \supseteq K \supsetneq J} \overline{Or(K)} = \overline{Or(J)} \setminus \bigcup_{i \in I \setminus J} \overline{Or(J \cup \{i\})}.$$

Proposition 3.12 *Let $J \subseteq I$.*

- (a) $Or(J)$ is G_{fn} -invariant and $P(J) \in Or(J)$.
- (b) $Or(J)$ is open and dense in $\overline{Or(J)}$. In particular, $Or(\emptyset)$ is open and dense in $Proj(CA)$.
- (c) It holds

$$\overline{Or(J)} = \bigcup_{I \supseteq K \supseteq J} Or(K).$$

$$\text{In particular, } Proj(CA) = \bigcup_{K \subseteq I} Or(K).$$

Proof: To (a) and (b): By its definition, $Or(J)$ is open in $\overline{Or(J)}$. It is a G_{fn} -invariant set because the sets $\overline{Or(K)}$, $K \supseteq J$, are G_{fn} -invariant. It is trivial to check that $P(J)$ as well as $\bigoplus_{\Lambda \in P^+ \setminus \overline{F_J}} L(\Lambda)^{(*)}$ are contained in $Or(J)$. It follows that $Or(J)$ is dense in $\overline{Or(J)}$.

To (c): By Proposition 3.10 it holds

$$\overline{Or(J)} = \bigcup_{I \supseteq K \supseteq J} \overline{Or(K)} \supseteq \bigcup_{I \supseteq K \supseteq J} Or(K).$$

Let $Q \in \overline{Or(J)}$. Choose a set $I \supseteq K_{max} \supseteq J$, maximal with respect to the inclusion, such that $Q \in Or(K_{max})$. Then $Q \in Or(K_{max})$.

Let $K_1, K_2 \subseteq I$ and $K_1 \neq K_2$. By Proposition 3.10 it follows

$$Or(K_1) \cap Or(K_2) \subseteq \overline{Or(K_1)} \cap \overline{Or(K_2)} \subseteq \overline{Or(K_1 \cup K_2)}. \quad (16)$$

At least one of the sets K_1, K_2 is nonempty. Assume K_2 is nonempty. Then it holds $K_1 \cup K_2 \supsetneq K_1$. By the definition of $Or(K_1)$ it holds

$$Or(K_1) \cap \overline{Or(K_1 \cup K_2)} = \emptyset. \quad (17)$$

From (16) and (17) it follows

$$Or(K_1) \cap Or(K_2) = Or(K_1) \cap Or(K_2) \cap \overline{Or(K_1 \cup K_2)} = \emptyset.$$

□

Let $J \subseteq I$. The Or-stratification at $P(J)$ will be investigated by restricting to a principal open subset at $P(J)$. Define

$$D(J) := \{ Q \in \text{Proj}(CA) \mid \delta_N \notin Q \text{ for all } N \in P^+ \cap \overline{F_J} \}.$$

In particular, $D(I) = \text{Proj}(CA)$. This follows because δ_N is a unit of CA for every $N \in P^+ \cap \overline{F_I} = \sum_{i=n+1}^{2n-l} \mathbb{Z}\Lambda_i$.

To describe $D(J)$ as a projective spectrum we identify CA with its image in $(\widetilde{P^+ \cap \overline{F_J}})^{-1}CA$, which is possible because CA has no zero divisors. By Theorem 2.6 it holds:

Proposition 3.13 *Let $J \subseteq I$. The map*

$$\begin{array}{ccc} \text{Proj}((\widetilde{P^+ \cap \overline{F_J}})^{-1}CA) & \rightarrow & D(J) \\ Q & \mapsto & Q \cap CA \end{array} \quad (18)$$

is a homeomorphism, mapping $\text{Proj}((\widetilde{P^+ \cap \overline{F_J}})^{-1}CA)(\mathbb{F})$ bijectively to $D(J)(\mathbb{F})$.

Remark 3.14 *It is easy to check that $ri(P^+ \cap \overline{F_J}) = P^+ \cap F_J$ is the set of principal elements of $P^+ \cap \overline{F_J}$. By Theorem 2.7 the set $D(J)$ is principal open: For every $\Lambda \in P^+ \cap F_J$ it holds*

$$\begin{array}{ccc} D(J) & = & D(\delta_\Lambda), \\ (\widetilde{P^+ \cap \overline{F_J}})^{-1}CA & \cong & \{ \delta_\Lambda^n \mid n \in \mathbb{N}_0 \}^{-1}CA. \end{array}$$

We will use the description which is independent of the choice of $\Lambda \in P^+ \cap F_J$ in the formulation of theorems and propositions. In proofs we will make use of both descriptions.

$D(J)$ decomposes by the Or-stratification as follows:

Proposition 3.15 *Let $J \subseteq I$. It holds*

$$D(J) = \bigcup_{K \subseteq J} (D(J) \cap Or(K)) \quad \text{and} \quad P(K) \in D(J) \cap Or(K) \text{ for all } K \subseteq J.$$

Proof: It holds $P(K) \in Or(K)$ for all $K \subseteq I$ by Proposition 3.12 (a). It is trivial to check that also $P(K) \in D(J)$ for all $K \subseteq J$.

Let $K \subseteq I$. If $\emptyset \neq D(J) \cap Or(K) \subseteq D(J) \cap \overline{Or(K)}$ there exists $Q \in \text{Proj}(CA)$ such that

$$Q \supseteq \bigoplus_{\Lambda \in P^+ \setminus \overline{F_K}} L(\Lambda)^{(*)} \quad \text{and} \quad \delta_\Lambda \notin Q \text{ for all } \Lambda \in P^+ \cap \overline{F_J}.$$

It follows $\overline{F_J} \subseteq \overline{F_K}$, which is equivalent to $J \supseteq K$. Now use Proposition 3.12 (c). □

Next we describe the open set obtained as the G_{fn} -orbit of $D(J)$. For the theorem we use the definition $\mathcal{V}(S) := \{ Q \in \text{Proj}(CA) \mid Q \supseteq S \}$ for subsets $S \subseteq CA$. It holds $\mathcal{V}(S) = \mathcal{V}((S_{hom}))$, where (S_{hom}) is the ideal generated by the set S_{hom} of homogeneous components of the elements of S .

Theorem 3.16 *Let $J \subseteq I$.*

(a) It holds

$$\bigcup_{g \in G_{fn}} gD(J) = \text{Proj}(CA) \setminus \mathcal{V}\left(\bigoplus_{\Lambda \in P^+ \cap F_J} L(\Lambda)^{(*)}\right).$$

(b) It holds $(P_{fn}^-)_J D(J) = D(J)$. In particular, the union in (a) can be taken over sets of coset representatives of $G_{fn}/(P_{fn}^-)_J$.

Proof: We first show (a). Let $Q \in \text{Proj}(CA)$. Then

$$Q \notin \bigcup_{g \in G_{fn}} gD(J)$$

is equivalent to $\pi(g)Q \notin D(J)$ for all $g \in G_{fn}$. By Remark 3.14 this is equivalent to

$$\delta_\Lambda \in \pi(g)Q \quad \text{for all } g \in G_{fn} \text{ and } \Lambda \in P^+ \cap F_J. \quad (19)$$

The irreducible G_{fn}^{op} -module $L(\Lambda)^{(*)}$ is spanned by $\pi(G_{fn})\delta_\Lambda$. Therefore, since Q is homogeneous, (19) is equivalent to

$$Q \supseteq \bigoplus_{\Lambda \in P^+ \cap F_J} L(\Lambda)^{(*)}.$$

Let $\Lambda \in P^+ \cap F_J$. By Remark 3.14 it holds $D(J) = \{Q \in \text{Proj}(CA) \mid \delta_\Lambda \notin Q\}$. Now (b) follows from $\pi((P_{fn}^-)_J)\delta_\Lambda \in \mathbb{F}\delta_\Lambda \setminus \{0\}$. \square

Let $J \subseteq I$. From the description of $\overline{Or(J)}$ in Proposition 3.12 (c) and from Proposition 3.15 it follows

$$BC(J) := \overline{Or(J)} \cap D(J) = Or(J) \cap D(J) \subseteq Or(J).$$

Furthermore, $P(J) \in BC(J)$. We call $BC(J)$ the *standard big cell* of $Or(J)$. We call every $gBC(J)$, $g \in G_{fn}$, a *big cell* of $Or(J)$.

Theorem 3.17 *For $J \subseteq I$ it holds:*

- (a) *Every big cell $gBC(J)$, $g \in G_{fn}$, is principal open in $\overline{Or(J)}$ and dense in $\overline{Or(J)}$.*
- (b) *$Or(J)$ can be covered by big cells:*

$$Or(J) = \bigcup_{g \in G_{fn}} gBC(J).$$

It holds $(P_{fn})_J^- BC(J) = BC(J)$. In particular, the union can be taken over sets of coset representatives of $G_{fn}/(P_{fn})_J^-$.

Proof: It is sufficient to show part (a) for $BC(J)$ because $\overline{Or(J)}$ is G_{fn} -invariant. $BC(J)$ is principal open in $\overline{Or(J)}$ because $D(J)$ is principal open in $\text{Proj}(CA)$ by Remark 3.14. By the definition of $Or(J)$ and $BC(J)$ it follows

$$\bigoplus_{\Lambda \in P^+ \setminus \overline{F_J}} L(\Lambda)^{(*)} \in BC(J) = \overline{Or(J)} \cap D(J) \subseteq \overline{Or(J)} = \overline{\bigoplus_{\Lambda \in P^+ \setminus \overline{F_J}} L(\Lambda)^{(*)}}.$$

Therefore, $BC(J)$ is dense in $\overline{Or(J)}$.

For part (b) it is sufficient to show

$$Or(J) = \overline{Or(J)} \cap \text{Proj}(CA) \setminus \mathcal{V}\left(\bigoplus_{\Lambda \in P^+ \cap F_J} L(\Lambda)^{(*)}\right). \quad (20)$$

Then (b) follows from Theorem 3.16 (a), (b) and the G_{fn} -invariance of $\overline{Or(J)}$. Taking the complement in $\overline{Or(J)}$, the equation (20) is equivalent to

$$\bigcup_{K \supsetneq J} \overline{Or(K)} = \overline{Or(J)} \cap \mathcal{V}\left(\bigoplus_{\Lambda \in P^+ \cap F_J} L(\Lambda)^{(*)}\right).$$

Inserting the definition of $\overline{Or(K)}$, $K \supseteq J$, this is equivalent to

$$\bigcup_{K \supsetneq J} \mathcal{V} \left(\bigoplus_{\Lambda \in P^+ \setminus \overline{F_K}} L(\Lambda)^{(*)} \right) = \mathcal{V} \left(\bigoplus_{\Lambda \in (P^+ \cap F_J) \cup P^+ \setminus \overline{F_J}} L(\Lambda)^{(*)} \right). \quad (21)$$

The inclusion " \subseteq " is valid, because for all $K \supsetneq J$ it holds

$$P^+ \setminus \overline{F_K} = P^+ \setminus \overline{F_J} \cup (P^+ \cap (\overline{F_J} \setminus \overline{F_K})) \supseteq P^+ \setminus \overline{F_J} \cup (P^+ \cap F_J).$$

Now we show " \supseteq ". Let Q be in the right hand side of (21). Suppose that Q is not contained in the left hand side of (21). Then for every $K \supsetneq J$ there exists a weight $\Lambda_K \in P^+ \setminus \overline{F_K}$ and an element $\phi_K \in L(\Lambda_K)^{(*)}$ such that $\phi_K \notin Q$. Since Q is prime it follows

$$Q \not\supseteq \prod_{K \supsetneq J} \phi_K \in L \left(\sum_{K \supsetneq J} \Lambda_K \right)^{(*)}. \quad (22)$$

Since $P^+ \setminus \overline{F_K}$ is a semigroup ideal of P^+ it holds

$$\begin{aligned} \sum_{K \supsetneq J} \Lambda_K &\in \bigcap_{K \supsetneq J} P^+ \setminus \overline{F_K} = \bigcap_{K \supsetneq J} (P^+ \setminus \overline{F_J} \cup (P^+ \cap (\overline{F_J} \setminus \overline{F_K}))) \\ &= P^+ \setminus \overline{F_J} \cup \bigcap_{K \supsetneq J} (P^+ \cap (\overline{F_J} \setminus \overline{F_K})) = P^+ \setminus \overline{F_J} \cup (P^+ \cap F_J). \end{aligned}$$

Therefore, (22) contradicts that Q is contained in the right hand side of (21). \square

As a principal open set of a closed set the standard big cell $BC(J)$ can be described as a projective spectrum:

Proposition 3.18 *Let $J \subseteq I$. Then*

$$\begin{aligned} Proj((P^+ \cap \overline{F_J})^{-1} \bigoplus_{\Lambda \in P^+ \cap \overline{F_J}} L(\Lambda)^{(*)}) &\rightarrow BC(J) \\ Q &\mapsto (Q \cap \bigoplus_{\Lambda \in P^+ \cap \overline{F_J}} L(\Lambda)^{(*)}) \oplus \bigoplus_{\Lambda \in P^+ \setminus \overline{F_J}} L(\Lambda)^{(*)} \end{aligned}$$

is a homeomorphism, mapping $Proj((P^+ \cap \overline{F_J})^{-1} \bigoplus_{\Lambda \in P^+ \cap \overline{F_J}} L(\Lambda)^{()})(\mathbb{F})$ bijectively to $BC(J)(\mathbb{F})$.*

Proof: The projective spectrum $Proj(\bigoplus_{\Lambda \in P^+ \cap \overline{F_J}} L(\Lambda)^{(*)})$ maps homeomorphically onto $\overline{Or(J)}$ by the map given in Proposition 3.11. The preimage of $BC(J) = \overline{Or(J)} \cap D(J)$ is the set

$$\left\{ Q \in Proj \left(\bigoplus_{\Lambda \in P^+ \cap \overline{F_J}} L(\Lambda)^{(*)} \right) \mid \delta_\Lambda \notin Q \text{ for all } \Lambda \in P^+ \cap \overline{F_J} \right\}.$$

Now the proposition follows with Theorem 2.6. \square

Next we introduce for $J \subseteq I$ a closed subset of $D(J)$, which is a transversal slice to $Or(J)$ at the point $P(J)$ as we will see later. To give its definition we need some easy preparations before.

For $\emptyset \neq J \subseteq I$ the Lie subalgebra \mathfrak{g}_J of \mathfrak{g} identifies with the derived Kac-Moody algebra $\mathfrak{g}(A_J)'$ to the generalized Cartan submatrix A_J of A . The sublattice Q_J of Q identifies with the root lattice, the sublattice P_J of P identifies with the weight lattice of $\mathfrak{g}(A_J)'$. The following theorem is well-known. Since the author has not found a reference where the proof is given, some details of the proof are added.

Theorem 3.19 *Let $\emptyset \neq J \subseteq I$ and $\Lambda \in P^+$. Decompose $\Lambda = \Lambda_J + \Lambda^J$ with $\Lambda_J \in P_J^+$ and $\Lambda^J \in P^+ \cap \overline{F_J}$. Regard $L(\Lambda)$ as a \mathfrak{g}_J -module.*

(a) Define an equivalence relation on $P(\Lambda)$ by

$$\lambda \sim \mu \quad : \Longleftrightarrow \quad \lambda - \mu \in Q_J.$$

If C is an equivalence class, then $L(\Lambda)_C := \bigoplus_{\lambda \in C} L(\Lambda)_\lambda$ is a \mathfrak{g}_J -module of $L(\Lambda)$. It decomposes into a direct sum of irreducible highest weight modules of \mathfrak{g}_J with highest weights in P_J^+ .

(b) The equivalence class $[\Lambda]$ of Λ is given by $[\Lambda] = P(\Lambda) \cap (\Lambda - (Q_J)_0^+)$. The \mathfrak{g}_J -module $L(\Lambda)_{[\Lambda]}$ is an irreducible highest weight module of \mathfrak{g}_J of highest weight Λ_J and highest weight space $L(\Lambda)_\Lambda$.

Proof: Part (a) follows with Theorem 10.7 a) of [K1]. For (b) only the nontrivial equality $U(\mathfrak{n}_J^-)L(\Lambda)_\Lambda = \bigoplus_{\lambda \in [\Lambda]} L(\Lambda)_\lambda$ is shown. It holds

$$L(\Lambda) = U(\mathfrak{n}^-)L(\Lambda)_\Lambda = U((\mathfrak{n}^-)^J)U(\mathfrak{n}_J^-)L(\Lambda)_\Lambda = U(\mathfrak{n}_J^-)L(\Lambda)_\Lambda + \sum_{\beta \in Q^+ \setminus Q_J^+} U((\mathfrak{n}^-)^J)_{-\beta} U(\mathfrak{n}_J^-)L(\Lambda)_\Lambda.$$

Here

$$U(\mathfrak{n}_J^-)L(\Lambda)_\Lambda \subseteq \bigoplus_{\lambda \in P(\Lambda) \cap (\Lambda - (Q_J)_0^+)} L(\Lambda)_\lambda, \quad (23)$$

$$\sum_{\beta \in Q^+ \setminus Q_J^+} U((\mathfrak{n}^-)^J)_{-\beta} U(\mathfrak{n}_J^-)L(\Lambda)_\Lambda \subseteq \bigoplus_{\lambda \in P(\Lambda) \cap (\Lambda - (Q_J)_0^+ - Q^+ \setminus Q_J^+)} L(\Lambda)_\lambda. \quad (24)$$

Now $L(\Lambda)$ is the direct sum of the sums in (23) and (24) on the right. It follows that in (23) and (24) it holds equality. \square

For $J \subseteq I$ and $\Lambda \in P^+$ set

$$P_J(\Lambda) := P(\Lambda) \cap (\Lambda - (Q_J)_0^+).$$

Remark 3.20 In particular, it holds:

(a) $P_I(\Lambda) = P(\Lambda)$ for all $\Lambda \in P^+$.

(b) $P_J(\Lambda) = \{\Lambda\}$ for all $\Lambda \in P^+ \cap \overline{F_J}$ and $J \subseteq I$.

Here (a) and the case $J = \emptyset$ in (b) follow from the definition. (b) holds also for $J \neq \emptyset$, because the \mathfrak{g}_J -module $L(\Lambda)_{[\Lambda]}$ is trivial.

For $J \subseteq I$ and $\Lambda \in P^+$ set

$$\begin{aligned} L_J(\Lambda) &:= \bigoplus_{\lambda \in P_J(\Lambda)} L(\Lambda)_\lambda \subseteq L(\Lambda) \quad \text{and} \quad L_J(\Lambda)_f := \prod_{\lambda \in P_J(\Lambda)} L(\Lambda)_\lambda \subseteq L(\Lambda)_f, \\ L_J(\Lambda)^{(*)} &:= \bigoplus_{\lambda \in P_J(\Lambda)} L(\Lambda)_\lambda^* \subseteq L(\Lambda)^{(*)}. \end{aligned}$$

Here $L_J(\Lambda)^{(*)}$ is the restricted dual of the \mathfrak{g}_J -module $L_J(\Lambda)$. Note that it holds

$$P(\Lambda) \setminus P_J(\Lambda) = P(\Lambda) \cap (\Lambda - (Q_0^+ \setminus (Q_J)_0^+)), \quad (25)$$

where we adopted the convention $B \pm \emptyset := \pm \emptyset + B := \emptyset$ for $B \subseteq P$. Now set

$$\begin{aligned} R_J(\Lambda) &:= \bigoplus_{\lambda \in P(\Lambda) \setminus P_J(\Lambda)} L(\Lambda)_\lambda \subseteq L(\Lambda) \quad \text{and} \quad R_J(\Lambda)_f := \prod_{\lambda \in P(\Lambda) \setminus P_J(\Lambda)} L(\Lambda)_\lambda \subseteq L(\Lambda)_f, \\ R_J(\Lambda)^{(*)} &:= \bigoplus_{\lambda \in P(\Lambda) \setminus P_J(\Lambda)} L(\Lambda)_\lambda^* \subseteq L(\Lambda)^{(*)}, \end{aligned}$$

a sum or product over the empty set defined to be $\{0\}$.

$L_J(\Lambda)^{(*)}$, $R_J(\Lambda)^{(*)}$ are $(G_{fn})_J^{op}$ -invariant subspaces of $L(\Lambda)^{(*)}$, and $L_J(\Lambda)_f$, $R_J(\Lambda)_f$ are $(G_{fn})_J$ -invariant subspaces of $L(\Lambda)_f$. Furthermore, $L_J(\Lambda)$, $R_J(\Lambda)$ are G_J -invariant subspaces of $L(\Lambda)$.

Proposition 3.21 *Let $J \subseteq I$. In CA it holds for all $\Lambda, N \in P^+$:*

- (a) $L_J(\Lambda)^{(*)} \bullet L_J(N)^{(*)} \subseteq L_J(\Lambda + N)^{(*)}$.
- (b) $R_J(\Lambda)^{(*)} \bullet L(N)^{(*)} \subseteq R_J(\Lambda + N)^{(*)}$ and $L(\Lambda)^{(*)} \bullet R_J(N)^{(*)} \subseteq R_J(\Lambda + N)^{(*)}$.

Proof: We first show

$$L(\Lambda)_\lambda^* \bullet L(N)_\mu^* \subseteq L(\Lambda + N)_{\lambda+\mu}^* \quad \text{for all } \lambda, \mu \in P. \quad (26)$$

Let $\phi_\lambda \in L(\Lambda)_\lambda^*$, $\psi_\mu \in L(N)_\mu^*$, $\lambda, \mu \in P$. With the notation of Subsection 3.1 the Cartan product $\phi_\lambda \bullet \psi_\mu \in L(\Lambda + N)^{(*)}$ is given by

$$(\phi_\lambda \bullet \psi_\mu)(w) := (\phi_\lambda \otimes \psi_\mu)(\Phi(w)) \quad \text{where } w \in L(\Lambda + N).$$

The G -equivariant linear map Φ preserves the weights. It follows immediately $\phi_\lambda \bullet \psi_\mu \in L(\Lambda + N)_{\lambda+\mu}^*$.

As an example we show $R_J(\Lambda)^{(*)} \bullet L(N)^{(*)} \subseteq R_J(\Lambda + N)^{(*)}$. The other inclusions can be treated similarly. This inclusion holds trivially for $J = I$, because $R_I(\Lambda)^{(*)} = \{0\}$ and $R_I(\Lambda + N)^{(*)} = \{0\}$. For $J \neq I$ use (25) to rewrite the definitions of $R_J(\Lambda)^{(*)}$ and $R_J(\Lambda + N)^{(*)}$. Then the inclusion follows from

$$(\Lambda - (Q_0^+ \setminus (Q_J)_0^+)) + (N - Q_0^+) \subseteq (\Lambda + N) - (Q_0^+ \setminus (Q_J)_0^+).$$

with (26). □

Let $J \subseteq I$. The Cartan algebra CA has no zero divisors. With Proposition 3.21 it follows the semidirect decomposition

$$CA = \bigoplus_{\Lambda \in P^+} L_J(\Lambda)^{(*)} \oplus \bigoplus_{\Lambda \in P^+} R_J(\Lambda)^{(*)} \quad (27)$$

with graded subalgebra $\bigoplus_{\Lambda \in P^+} L_J(\Lambda)^{(*)}$ and graded prime ideal $\bigoplus_{\Lambda \in P^+} R_J(\Lambda)^{(*)}$. We call

$$S(J) := \mathcal{V}(\bigoplus_{\Lambda \in P^+} R_J(\Lambda)^{(*)}) \cap D(J)$$

the standard transversal slice to $Or(J)$ at the point $P(J) \in Or(J)$. This name will be justified later. We now only show:

Theorem 3.22 *Let $J \subseteq I$. It holds*

$$Or(J) \cap S(J) = BC(J) \cap S(J) = \{P(J)\}.$$

Proof: Because of $S(J) \subseteq D(J)$ it follows

$$BC(J) \cap S(J) \subseteq Or(J) \cap S(J) \subseteq \overline{Or(J)} \cap S(J) = \overline{Or(J)} \cap D(J) \cap S(J) = BC(J) \cap S(J).$$

By the definition of $BC(J)$, $S(J)$, and by Remark 3.20 (b) we get

$$\begin{aligned} BC(J) \cap S(J) &= \mathcal{V}(\bigoplus_{\Lambda \in P^+} R_J(\Lambda)^{(*)}) \cap \mathcal{V}(\bigoplus_{\Lambda \in P^+ \setminus \overline{F_J}} L(\Lambda)^{(*)}) \cap D(J) \\ &= \mathcal{V}(\bigoplus_{\Lambda \in P^+} R_J(\Lambda)^{(*)} \oplus \bigoplus_{\Lambda \in P^+ \setminus \overline{F_J}} L(\Lambda)^{(*)}) \cap D(J) \\ &= \mathcal{V}(\bigoplus_{\Lambda \in P^+ \cap \overline{F_J}} L(\Lambda)^{(*)}_{\neq \Lambda} \oplus \bigoplus_{\Lambda \in P^+ \setminus \overline{F_J}} L(\Lambda)^{(*)}) \cap D(J) = \mathcal{V}(P(J)) \cap D(J). \end{aligned}$$

With Proposition 3.15 it follows $P(J) \in \mathcal{V}(P(J)) \cap D(J)$. Now let $Q \in \mathcal{V}(P(J)) \cap D(J)$. By definition it holds $Q \supseteq P(J)$ and $\delta_\Lambda \notin Q$ for all $\Lambda \in P^+ \cap \overline{F_J}$. We show $Q \subseteq P(J)$. Every $\phi \in Q$ is of the form

$$\phi = \sum_{\Lambda \in P^+ \cap \overline{F_J}} c_\Lambda \delta_\Lambda + \psi$$

with $\psi \in P(J)$ and $c_\Lambda \in \mathbb{F}$, $c_\Lambda \neq 0$ for at most finitely many $\Lambda \in P^+ \cap \overline{F_J}$. It follows $\sum_{\Lambda \in P^+ \cap \overline{F_J}} c_\Lambda \delta_\Lambda \in Q$. Since Q is graded we get $c_\Lambda \delta_\Lambda \in Q$, which implies $c_\Lambda = 0$, $\Lambda \in P^+ \cap \overline{F_J}$. We found $\phi = \psi \in P(J)$. \square

$S(J)$ decomposes with respect to the Or-stratification as follows:

Proposition 3.23 *Let $J \subseteq I$. It holds*

$$S(J) = \bigcup_{K \subseteq J} \left(S(J) \cap Or(K) \right) \quad \text{and} \quad P(K) \in S(J) \cap Or(K) \text{ for all } K \subseteq J.$$

Proof: It is trivial to check that $P(K) \supseteq \bigoplus_{\Lambda \in P^+} R_J(\Lambda)^{(*)}$ for all $K \subseteq I$. Now the proposition follows from the definition of $S(J)$ and Proposition 3.15. \square

As a principal open set of a closed set the standard transversal slice $S(J)$ can be described as a projective spectrum. The proof is completely similar as for $BC(J)$, the semidirect decomposition (15) replaced by the semidirect decomposition (27):

Proposition 3.24 *Let $J \subseteq I$. The map*

$$\begin{aligned} Proj(\widetilde{(P^+ \cap \overline{F_J})}^{-1} \bigoplus_{\Lambda \in P^+} L_J(\Lambda)^{(*)}) &\rightarrow S(J) \\ Q &\mapsto (Q \cap \bigoplus_{\Lambda \in P^+} L_J(\Lambda)^{(*)}) \oplus \bigoplus_{\Lambda \in P^+} R_J(\Lambda)^{(*)} \end{aligned}$$

is a homeomorphism, mapping $Proj(\widetilde{(P^+ \cap \overline{F_J})}^{-1} \bigoplus_{\Lambda \in P^+} L_J(\Lambda)^{()})(\mathbb{F})$ bijectively to $S(J)(\mathbb{F})$.*

The projective spectrum of the graded algebra $\widetilde{(P^+ \cap \overline{F_J})}^{-1} CA$ gives the principal open set $D(J)$ by Proposition 3.13. Projective spectra of graded subalgebras of $\widetilde{(P^+ \cap \overline{F_J})}^{-1} CA$ give the standard big cell $BC(J)$ and the standard transversal slice $S(J)$ by Propositions 3.18 and 3.24. The next aim is to describe the graded algebra $\widetilde{(P^+ \cap \overline{F_J})}^{-1} CA$ explicitly as a tensor product of three graded algebras. It will be reached in Theorem 3.34. We need some preparations before.

Let $J \subseteq I$. We equip $(\mathbf{n}_f^-)^J$ with the unital coordinate ring of functions $\mathbb{F}[(\mathbf{n}_f^-)^J]$, which is generated by the linear functions contained in the subspace

$$((\mathbf{n}^-)^J)^{(*)} := \bigoplus_{\alpha \in (\Delta^-)^J} \mathbf{g}_\alpha^* \subseteq ((\mathbf{n}_f^-)^J)^*.$$

It is easy to check that this coordinate ring is the symmetric algebra

$$\mathbb{F}[(\mathbf{n}_f^-)^J] = Sym\left(((\mathbf{n}^-)^J)^{(*)}\right).$$

We push this coordinate ring forward by the bijective exponential map $exp : (\mathbf{n}_f^-)^J \rightarrow (U_f^-)^J$ and obtain a coordinate ring $\mathbb{F}[(U_f^-)^J]$ on $(U_f^-)^J$ such that the comorphism

$$exp^* : \mathbb{F}[(U_f^-)^J] \rightarrow \mathbb{F}[(\mathbf{n}_f^-)^J]$$

exists and is an isomorphism of algebras. We equip $\mathbb{F}[(U_f^-)^J]$ with the trivial graduation $\{0\}$.

In particular, there is the easy case $(\mathbf{n}_f^-)^I = \{0\}$, $\mathbb{F}[(\mathbf{n}_f^-)^I] \cong \mathbb{F}$ and $(U_f^-)^I = \{1\}$, $\mathbb{F}[(U_f^-)^I] \cong \mathbb{F}$ included.

It is also worth to note the following remark, which will not be used in this article:

Remark 3.25 $\mathbb{F}[(\mathbf{n}_f^-)^J]$ is the coordinate ring of the pronilpotent Lie algebra $(\mathbf{n}_f^-)^J$ as follows: For $k \in \mathbb{N}_0$ it is

$$(\mathbf{n}_f^-)^J(k) := \prod_{\alpha \in (\Delta^-)^J, -ht(\alpha) \geq k+1} \mathbf{g}_\alpha.$$

an ideal of $(\mathbf{n}_f^-)^J$, such that the quotient $(\mathbf{n}_f^-)^J / (\mathbf{n}_f^-)^J(k)$ is a finite dimensional nilpotent Lie algebra. There is the system of projections

$$p_{kl} : (\mathbf{n}_f^-)^J / (\mathbf{n}_f^-)^J(k) \rightarrow (\mathbf{n}_f^-)^J / (\mathbf{n}_f^-)^J(l) \quad \text{where} \quad k, l \in \mathbb{N}_0, k \geq l.$$

It is easy to check that a projective limit is given by the projections

$$p_k : (\mathbf{n}_f^-)^J \rightarrow (\mathbf{n}_f^-)^J / (\mathbf{n}_f^-)^J(k) \quad \text{where} \quad k \in \mathbb{N}_0.$$

Equip $(\mathbf{n}_f^-)^J / (\mathbf{n}_f^-)^J(k)$ with its coordinate ring $\mathbb{F}[(\mathbf{n}_f^-)^J / (\mathbf{n}_f^-)^J(k)]$ as a finite dimensional \mathbb{F} -linear space. There is the system of comorphisms

$$p_{kl}^* : \mathbb{F}[(\mathbf{n}_f^-)^J / (\mathbf{n}_f^-)^J(l)] \rightarrow \mathbb{F}[(\mathbf{n}_f^-)^J / (\mathbf{n}_f^-)^J(k)] \quad \text{where} \quad l, k \in \mathbb{N}_0, l \leq k.$$

It is easy to check that an inductive limit is given by the comorphisms

$$p_k^* : \mathbb{F}[(\mathbf{n}_f^-)^J / (\mathbf{n}_f^-)^J(k)] \rightarrow \mathbb{F}[(\mathbf{n}_f^-)^J] \quad \text{where} \quad k \in \mathbb{N}_0.$$

Similarly, $\mathbb{F}[(U_f^-)^J]$ is the coordinate ring of the prounipotent group $(U_f^-)^J$.

Proposition 3.26 Let $J \subseteq I$. It holds

$$\text{Spec} \left(\mathbb{F}[(U_f^-)^J] \right) (\mathbb{F}) = \left\{ I(u) \mid u \in (U_f^-)^J \right\},$$

where $I(u)$ denotes the vanishing ideal in $u \in (U_f^-)^J$.

Proof: By the linear isomorphisms

$$((\mathbf{n}^-)^J)^{(*)} = \left(\bigoplus_{\alpha \in (\Delta^-)^J} \mathbf{g}_\alpha^* \right)^* \cong \prod_{\alpha \in (\Delta^-)^J} (\mathbf{g}_\alpha^*)^* \cong \prod_{\alpha \in (\Delta^-)^J} \mathbf{g}_\alpha = (\mathbf{n}_f^-)^J$$

the elements of the dual of $((\mathbf{n}^-)^J)^{(*)}$ are given by the evaluations in the elements of $(\mathbf{n}_f^-)^J$. By the universal property of the symmetric algebra it follows $\text{Spec}(\mathbb{F}[(\mathbf{n}_f^-)^J])(\mathbb{F}) = \{I(x) \mid x \in (\mathbf{n}_f^-)^J\}$, where $I(x)$ is the vanishing ideal in $x \in (\mathbf{n}_f^-)^J$. Since the comorphism $\exp^* : \mathbb{F}[(U_f^-)^J] \rightarrow \mathbb{F}[(\mathbf{n}_f^-)^J]$ is an isomorphism of coordinate rings the Corollary follows. \square

For a \mathbf{g} -module V contained in the category \mathcal{O}_{int}^p , for $v \in V_f$ and $\phi \in V^{(*)}$ the map

$$\begin{aligned} \phi(\cdot v) : U_f^- &\rightarrow \mathbb{F} \\ u &\mapsto (\pi(u)\phi)(v) = \phi(uv) \end{aligned}$$

is called the matrix coefficient of ϕ and v on U_f^- . To keep our notation easy we often denote its restriction to nonempty subsets of U_f^- by the same symbol.

Theorem 3.27 Let $J \subseteq I$. For every \mathbf{g} -module V from the category \mathcal{O}_{int}^p , for every $v \in V_f$ and $\phi \in V^{(*)}$ it is

$$\phi(\cdot v) \in \mathbb{F}[(U_f^-)^J]. \quad (28)$$

Furthermore, it holds:

$$\mathbb{F}[(U_f^-)^J] = \text{span} \left\{ \phi(\cdot v_N) \mid \phi \in L(N)^{(*)}, N \in P^+ \cap \overline{F_J} \right\}. \quad (29)$$

For every $\Lambda \in P^+$ with $\Lambda(h_i) > 0$ for all $i \in I \setminus J$ the linear map

$$\begin{aligned} \mathbf{n}^J &\rightarrow \mathbb{F}[(U_f^-)^J] \\ x &\mapsto (\pi(x)\delta_\Lambda)(\cdot v_\Lambda) \end{aligned}$$

is injective and it holds

$$\mathbb{F}[(U_f^-)^J] = \text{Sym}(\{(\pi(x)\delta_\Lambda)(\cdot v_\Lambda) \mid x \in \mathbf{n}^J\}). \quad (30)$$

Proof: This holds trivially for $J = I$. Let $J \neq I$. For every $\alpha \in (\Delta^-)^J$ choose a base $y_{\alpha i} \in \mathbf{g}_\alpha$, $i = 1, \dots, m_\alpha$. Let $\tilde{\psi}_{\alpha i} \in \mathbf{g}_\alpha^*$, $i = 1, \dots, m_\alpha$, be the dual base. Interpret these elements as elements of $((\mathbf{n}^-)^J)^{(*)} \subseteq ((\mathbf{n}_f^-)^J)^*$ and set

$$\psi_{\alpha i} := (\exp^*)^{-1}(\tilde{\psi}_{\alpha i}) \in \mathbb{F}[(U_f^-)^J], \quad i = 1, \dots, m_\alpha.$$

Note that it holds

$$\tilde{\psi}_{\alpha i}(y_{\beta j}) = \delta_{\alpha\beta}\delta_{ij} \quad \text{for} \quad \alpha, \beta \in (\Delta^-)^J, \quad i = 1, \dots, m_\alpha, \quad j = 1, \dots, m_\beta.$$

For every $\alpha \in (\Delta^+)^J$ the pairing $(\mid) : \mathbf{g}_\alpha \times \mathbf{g}_{-\alpha} \rightarrow \mathbb{F}$ is nondegenerate. Let $z_{\alpha i} \in \mathbf{g}_\alpha$, $\alpha \in (\Delta^+)^J$, $i = 1, \dots, m_\alpha$, be the dual base of $y_{-\alpha i} \in \mathbf{g}_{-\alpha}$, $i = 1, \dots, m_{-\alpha}$. Because of $(\mathbf{g}_\alpha \mid \mathbf{g}_{-\beta}) = \{0\}$ for $\alpha, \beta \in (\Delta^+)^J$, $\alpha \neq \beta$, it holds

$$(z_{\alpha i} \mid y_{-\beta j}) = \delta_{\alpha\beta}\delta_{ij} \quad \text{for} \quad \alpha, \beta \in (\Delta^+)^J, \quad i = 1, \dots, m_\alpha, \quad j = 1, \dots, m_{-\beta}.$$

(a) Let $v = \sum_{\lambda \in P(V)} v_\lambda \in V_f$ with $v_\lambda \in V_\lambda$ for all $\lambda \in P(V)$. Let $\phi_\mu \in V_\mu^*$, $\mu \in P(V)$. It holds

$$\phi_\mu(\exp(x)v) = \sum_{\lambda \in P(V)} \phi_\mu(\exp(x)v_\lambda) = \sum_{\lambda \in P(V), \lambda \geq \mu} \phi_\mu(\exp(x)v_\lambda) = \left(\sum_{\lambda \in P(V), \lambda \geq \mu} \phi_\mu(\cdot v_\lambda) \right) (\exp(x))$$

for all $x \in (\mathbf{n}_f^-)^J$, a sum over the empty set defined to be zero. The sum on the right is finite, because V is contained in \mathcal{O}_{int}^p .

Therefore, it is sufficient to show (28) for $v_\lambda \in V_\lambda$, $\phi_\mu \in V_\mu^*$, $\lambda, \mu \in P(V)$, $\lambda \geq \mu$. For every $x = \sum_{\alpha \in (\Delta^-)^J} x_\alpha \in (\mathbf{n}_f^-)^J$ it holds:

$$\phi_\mu(\exp(x)v_\lambda) = \sum_{k=0}^{ht(\lambda-\mu)} \frac{1}{k!} \phi_\mu(x^k v_\lambda) = \phi_\mu(v_\lambda) + \sum_{k=1}^{ht(\lambda-\mu)} \frac{1}{k!} \sum_{\substack{\beta_1, \dots, \beta_k \in (\Delta^-)^J \\ \beta_1 + \dots + \beta_k = \mu - \lambda}} \phi_\mu(x_{\beta_1} \cdots x_{\beta_k} v_\lambda). \quad (31)$$

For every $\beta \in (\Delta^-)^J$ we develop x_β in the basis $y_{\beta i} \in \mathbf{g}_\beta$, $i = 1, \dots, m_\beta$, as follows:

$$x_\beta = \sum_{i=1}^{m_\beta} \tilde{\psi}_{\beta i}(x_\beta) y_{\beta i} = \sum_{i=1}^{m_\beta} \tilde{\psi}_{\beta i}(x) y_{\beta i} = \sum_{i=1}^{m_\beta} \psi_{\beta i}(\exp(x)) y_{\beta i}.$$

Inserting in the inner summand of (31) we get:

$$\phi_\mu(x_{\beta_1} \cdots x_{\beta_k} v_\lambda) = \sum_{i_1, \dots, i_k} \psi_{\beta_1 i_1}(\exp(x)) \cdots \psi_{\beta_k i_k}(\exp(x)) \phi_\mu(y_{\beta_1 i_1} \cdots y_{\beta_k i_k} v_\lambda).$$

Inserting back in (31) we find

$$\phi_\mu(\cdot v_\lambda) = \phi_\mu(v_\lambda)1 + \sum_{k=1}^{ht(\lambda-\mu)} \frac{1}{k!} \sum_{\substack{\beta_1, \dots, \beta_k \in (\Delta^-)^J, i_1, \dots, i_k \\ \beta_1 + \dots + \beta_k = \mu - \lambda}} \phi_\mu(y_{\beta_1 i_1} \cdots y_{\beta_k i_k} v_\lambda) \psi_{\beta_1 i_1} \cdots \psi_{\beta_k i_k} \in \mathbb{F}[(U_f^-)^J]. \quad (32)$$

(b) It holds

$$\{N \in P^+ \mid N(h_i) > 0 \text{ for all } i \in I \setminus J\} = P^+ \cap \bigcup_{K \subseteq J} F_K.$$

Fix $K \subseteq J$ and $\Lambda \in P^+ \cap F_K$. It follows

$$\text{span} \left\{ \phi(\cdot v_N) \mid \phi \in L(N)^{(*)}, N \in P^+ \cap \overline{F_K} \right\} \subseteq \mathbb{F}[(U_f^-)^J]$$

from (28). This span is even a subalgebra of $\mathbb{F}[(U_f^-)^J]$. It contains $\delta_0(\cdot v_0)$, which is the unit of $\mathbb{F}[(U_f^-)^J]$. Now let $\phi \in L(N)^{(*)}$, $N \in P^+ \cap \overline{F_K}$ and $\psi \in L(M)^{(*)}$, $M \in P^+ \cap \overline{F_K}$. It holds

$$\phi(\cdot v_N)\psi(\cdot v_M) = (\phi \otimes \psi)(\cdot(v_N \otimes v_M)) = (\phi \bullet \psi)(\cdot v_{N+M})$$

with $\phi \bullet \psi \in L(N+M)^{(*)}$ and $N+M \in P^+ \cap \overline{F_K}$.

Let $\text{Alg}(\{(\pi(x)\delta_\Lambda)(\cdot v_\Lambda) \mid x \in \mathbf{n}^J\})$ be the subalgebra of $\mathbb{F}[(U_f^-)^J]$ generated by $\{(\pi(x)\delta_\Lambda)(\cdot v_\Lambda) \mid x \in \mathbf{n}^J\}$. We have shown the inclusions

$$\text{Alg}(\{(\pi(x)\delta_\Lambda)(\cdot v_\Lambda) \mid x \in \mathbf{n}^J\}) \subseteq \text{span} \left\{ \phi(\cdot v_N) \mid \phi \in L(N)^{(*)}, N \in P^+ \cap \overline{F_K} \right\} \subseteq \mathbb{F}[(U_f^-)^J].$$

(c) Now we show

$$\psi_{-\alpha i} \in \text{Alg}(\{(\pi(x)\delta_\Lambda)(\cdot v_\Lambda) \mid x \in \mathbf{n}^J\}) \quad \text{for all } \alpha \in (\Delta^+)^J, \quad i = 1, \dots, m_{-\alpha},$$

which implies $\mathbb{F}[(U_f^-)^J] \subseteq \text{Alg}(\{(\pi(x)\delta_\Lambda)(\cdot v_\Lambda) \mid x \in \mathbf{n}^J\})$.

Specialize formula (32) to the matrix coefficient $(\pi(z_{\alpha i})\delta_\Lambda)(\cdot v_\Lambda)$: Because $z_{\alpha i}$ kills v_Λ it holds

$$(\pi(z_{\alpha i})\delta_\Lambda)(v_\Lambda) = \delta_\Lambda(z_{\alpha i}v_\Lambda) = 0.$$

Transform the summand for $k = 1$ of the sum in (32) as follows:

$$\sum_j \delta_\Lambda(z_{\alpha i} y_{-\alpha j} v_\Lambda) \psi_{-\alpha j} = \sum_j \delta_\Lambda([z_{\alpha i}, y_{-\alpha j}] v_\Lambda) \psi_{-\alpha j} = \sum_j \delta_\Lambda((z_{\alpha i} \mid y_{-\alpha j}) \nu^{-1}(\alpha) v_\Lambda) \psi_{-\alpha j} = (\Lambda \mid \alpha) \psi_{-\alpha i}.$$

We have found:

$$\begin{aligned} & (\pi(z_{\alpha i})\delta_\Lambda)(\cdot v_\Lambda) \\ &= (\Lambda \mid \alpha) \psi_{-\alpha i} + \sum_{k=2}^{ht(\alpha)} \frac{1}{k!} \sum_{\substack{\beta_1, \dots, \beta_k \in (\Delta^-)^J, i_1, \dots, i_k \\ \beta_1 + \dots + \beta_k = -\alpha}} \delta_\Lambda(z_{\alpha i} y_{\beta_1 i_1} \cdots y_{\beta_k i_k} v_\Lambda) \psi_{\beta_1 i_1} \cdots \psi_{\beta_k i_k}. \end{aligned} \quad (33)$$

Here α is of the form $\alpha = \sum_{i \in I} n_i \alpha_i$ with $n_i \in \mathbb{N}_0$ and $n_{j_0} \neq 0$ for at least one $j_0 \in J$. With $\nu^{-1}(\alpha_i) = \frac{1}{2}(\alpha_i \mid \alpha_i) h_i$ and $(\alpha_i \mid \alpha_i) > 0$, $i \in I$, it follows

$$(\Lambda \mid \alpha) = \Lambda(\nu^{-1}(\alpha)) = \sum_{i \in I, i \neq j_0} \underbrace{n_i}_{\geq 0} \frac{(\alpha_i \mid \alpha_i)}{2} \underbrace{\Lambda(h_i)}_{\geq 0} + \underbrace{n_{j_0}}_{> 0} \frac{(\alpha_{j_0} \mid \alpha_{j_0})}{2} \underbrace{\Lambda(h_{j_0})}_{> 0} > 0. \quad (34)$$

Let $\alpha \in (\Delta^+)^J$ with $ht(\alpha) = 1$. Then $m_\alpha = m_{-\alpha} = 1$ and from (33) and (34) it follows

$$\psi_{-\alpha 1} = \frac{1}{(\Lambda \mid \alpha)} (\pi(z_{\alpha 1})\delta_\Lambda)(\cdot v_\Lambda) \in \text{Alg}(\{(\pi(x)\delta_\Lambda)(\cdot v_\Lambda) \mid x \in \mathbf{n}^J\}).$$

Suppose $\psi_{-\beta i} \in \text{Alg}(\{(\pi(x)\delta_\Lambda)(\cdot v_\Lambda) \mid x \in \mathbf{n}^J\})$ for all $\beta \in (\Delta^+)^J$ with $ht(\beta) < p \in \mathbb{N}$ and $i = 1, \dots, m_{-\beta}$. Let $\alpha \in (\Delta^+)^J$ with $ht(\alpha) = p$. Then from (33) and (34) it follows

$$\psi_{-\alpha i} \in \text{Alg}(\{(\pi(x)\delta_\Lambda)(\cdot v_\Lambda) \mid x \in \mathbf{n}^J\}) \quad \text{for all } i = 1, \dots, m_\alpha.$$

(d) At last we show that the linear map $\mathbf{n}^J \rightarrow \mathbb{F}[(U_f^-)^J]$ of the theorem is injective and that $\mathbb{F}[(U_f^-)^J] = \text{Alg}(\{(\pi(x)\delta_\Lambda)(\cdot v_\Lambda) \mid x \in \mathbf{n}^J\})$ is a symmetric algebra in $\{(\pi(x)\delta_\Lambda)(\cdot v_\Lambda) \mid x \in \mathbf{n}^J\}$.

Choose a counting (β_m, i_m) , $m \in \mathbb{N}$, of the set

$$\{(\beta, i) \mid \beta \in (\Delta^+)^J, i = 1, \dots, m_\beta\}.$$

Since $z_{\beta_m i_m}$, $m \in \mathbb{N}$, is a base of \mathbf{n}^J it is sufficient to show that the monomials in

$$f_m := (\pi(z_{\beta_m i_m})\delta_\Lambda)(\cdot v_\Lambda), \quad m \in \mathbb{N},$$

are linearly independent.

Note that by (33) and (34) for all $m \in \mathbb{N}$ it holds

$$f_m = \underbrace{(\Lambda \mid \beta_m)}_{\neq 0} \psi_{-\beta_m i_m} + r_m \quad (35)$$

where r_m is a sum of homogeneous components of degree ≥ 2 .

Now let $r \in \mathbb{N}$ and $L \in \mathbb{N}_0$ and suppose

$$0 = \sum_{\substack{k_1, \dots, k_r \in \mathbb{N}_0 \\ k_1 + \dots + k_r \leq L}} c_{k_1 \dots k_r} f_1^{k_1} \dots f_r^{k_r} \quad (36)$$

with coefficients $c_{k_1 \dots k_r} \in \mathbb{F}$.

First we compare the 0-homogeneous components of both sides of equation (36). Because of (35) we get $c_{0 \dots 0} = 0$. Let $l \in \mathbb{N}$, $l \leq L$, and suppose we have shown

$$c_{k_1 \dots k_r} = 0 \quad \text{for all } k_1, \dots, k_r \in \mathbb{N}_0 \text{ with } k_1 + \dots + k_r \leq l - 1.$$

Now we compare the l -homogeneous components of both sides of equation (36). Because of (35) we get

$$0 = \sum_{\substack{k_1, \dots, k_r \in \mathbb{N}_0 \\ k_1 + \dots + k_r = l}} c_{k_1 \dots k_r} (\Lambda \mid \beta_1)^{k_1} \dots (\Lambda \mid \beta_r)^{k_r} \psi_{-\beta_1 i_1}^{k_1} \dots \psi_{-\beta_r i_r}^{k_r}.$$

Since the monomials in $\psi_{-\beta_m i_m}$, $m \in \mathbb{N}$, are linearly independent and $(\Lambda \mid \beta_m) \neq 0$ for all $m \in \mathbb{N}$ it follows

$$c_{k_1 \dots k_r} = 0 \quad \text{for all } k_1, \dots, k_r \in \mathbb{N}_0 \text{ with } k_1 + \dots + k_r = l.$$

□

Next we draw two Corollaries for the groups $(U^-)^J$ and its coordinate rings, $J \subseteq I$.

Corollary 3.28 *Let $J \subseteq I$. The subgroup $(U^-)^J$ of $(U_f^-)^J$ is dense. In particular, its coordinate ring*

$$\mathbb{F}[(U^-)^J] := \mathbb{F}[(U_f^-)^J]|_{(U^-)^J}$$

is isomorphic to $\mathbb{F}[(U_f^-)^J]$ by the restriction map.

Proof: This holds trivially for $J = I$. Let $J \neq I$. Let $f \in \mathbb{F}[(U_f^-)^J]$ such that $f|_{(U^-)^J} = 0$. By the last theorem we can write f in the form

$$f = \sum_{N \in P^+ \cap \overline{E_J}} \phi_N(\cdot v_N)$$

with $\phi_N \in L(N)^{(*)}$, $\phi_N \neq 0$ for at most finitely many $N \in P^+ \cap \overline{F_J}$. Because U_J^- stabilizes $L(N)_N$ pointwise for all $N \in P^+ \cap \overline{F_J}$ and because of $U^- = (U^-)^J U_J$ we find

$$0 = \sum_{N \in P^+ \cap \overline{F_J}} \phi_N(uv_N) \quad \text{for all } u \in U^-.$$

For $p \in \mathbb{N}$, $\beta_1, \dots, \beta_p \in \Delta_{re}^-$, and $x_{\beta_1} \in \mathfrak{g}_{\beta_1}, \dots, x_{\beta_p} \in \mathfrak{g}_{\beta_p}$ it follows

$$0 = \sum_{N \in P^+ \cap \overline{F_J}} \phi_N(\exp(t_1 x_{\beta_1}) \cdots \exp(t_p x_{\beta_p}) v_N) = \sum_{k_1, \dots, k_p \in \mathbb{N}_0} \frac{t_1^{k_1} \cdots t_p^{k_p}}{k_1! \cdots k_p!} \sum_{N \in P^+ \cap \overline{F_J}} \phi_N(x_{\beta_1}^{k_1} \cdots x_{\beta_p}^{k_p} v_N).$$

for all $t_1, \dots, t_p \in \mathbb{F}$. The last expression is well defined and polynomial in t_1, \dots, t_p because $x_{\beta_1}, \dots, x_{\beta_p}$ act locally nilpotent. The coefficients of the polynomial vanish. In particular, we find

$$0 = \sum_{N \in P^+ \cap \overline{F_J}} \phi_N(v_N) \quad \text{and} \quad 0 = \sum_{N \in P^+ \cap \overline{F_J}} \phi_N(x_{\beta_1} \cdots x_{\beta_p} v_N).$$

By §1.3 of [K1] the Lie algebra \mathfrak{n}^- is generated by the elements $f_i \in \mathfrak{g}_{-\alpha_i}$, $i \in I$. Therefore, also the universal enveloping algebra $U(\mathfrak{n}^-)$ is generated by $f_i \in \mathfrak{g}_{-\alpha_i}$, $i \in I$. Because of $-\alpha_i \in \Delta_{re}^-$, $i \in I$, we have found

$$0 = \sum_{N \in P^+ \cap \overline{F_J}} \phi_N(xv_N) \quad \text{for all } x \in U(\mathfrak{n}^-). \quad (37)$$

Now let $y = \sum_{\beta \in (\Delta^-)^J} y_\beta \in (\mathfrak{n}_f^-)^J$. With (37) it follows

$$\begin{aligned} f(\exp(y)) &= \sum_{N \in P^+ \cap \overline{F_J}} \phi_N(\exp(y)v_N) \\ &= \sum_{N \in P^+ \cap \overline{F_J}} \phi_N(v_N) + \sum_{\gamma \in Q^-} \frac{1}{k!} \sum_{k=1}^{-ht(\gamma)} \sum_{\substack{\beta_1, \dots, \beta_k \in (\Delta^-)^J \\ \beta_1 + \dots + \beta_k = \gamma}} \sum_{N \in P^+ \cap \overline{F_J}} \phi_N(y_{\beta_1} \cdots y_{\beta_k} v_N) = 0. \end{aligned}$$

We have shown $f = 0$. □

Recall from Section 1 that the Kac-Moody group G is equipped with a coordinate ring of matrix coefficients such that there holds a Peter-Weyl Theorem: A $G^{op} \times G$ -equivariant bijective linear map

$$PW : \bigoplus_{\Lambda \in P^+} L(\Lambda)^{(*)} \otimes L(\Lambda) \rightarrow \mathbb{F}[G]$$

is given by $PW(\phi \otimes v) := \phi(\cdot v)$ for all $\phi \in L(\Lambda)^{(*)}$, $v \in L(\Lambda)$, $\Lambda \in P^+$. Here the matrix coefficient $\phi(\cdot v)$ denotes the function which assigns to $g \in G$ the value $\phi(gv) \in \mathbb{F}$. To keep our notation easy we often denote its restriction to nonempty subsets of G by the same symbol.

Now $(U^-)^J$ is also contained in G . It holds:

Corollary 3.29 *Let $J \subseteq I$. The restriction of the coordinate ring $\mathbb{F}[G]$ to $(U^-)^J$ coincides with the restriction of the coordinate ring $\mathbb{F}[(U_f^-)^J]$ to $(U^-)^J$, which we denoted by $\mathbb{F}[(U^-)^J]$.*

Proof: By the Peter-Weyl theorem it holds

$$\mathbb{F}[G]|_{(U^-)^J} = \text{span} \left\{ \phi(\cdot v)|_{(U^-)^J} \mid \phi \in L(\Lambda)^{(*)}, v \in L(\Lambda), \Lambda \in P^+ \right\}.$$

Now the Corollary follows from Theorem 3.27. □

To avoid trivialities let $J \subsetneq I$. The pairing $(\mid) : \mathbf{n}^J \times (\mathbf{n}^-)^J \rightarrow \mathbb{F}$ is nondegenerate. Therefore, we get a linear bijective map

$$\begin{aligned} \tilde{\psi} : \mathbf{n}^J &\rightarrow ((\mathbf{n}^-)^J)^{(*)} \\ x &\mapsto \tilde{\psi}_x \end{aligned}$$

where $\tilde{\psi}_x \in ((\mathbf{n}^-)^J)^{(*)} \subseteq ((\mathbf{n}_f^-)^J)^*$ is defined by

$$\tilde{\psi}_x(y) := \sum_{\alpha \in (\Delta^-)^J} (x \mid y_\alpha) \quad \text{for} \quad y = \sum_{\alpha \in (\Delta^-)^J} y_\alpha \in (\mathbf{n}_f^-)^J.$$

Set $\psi_x := (\exp^*)^{-1}(\tilde{\psi}_x) \in \mathbb{F}[(U_f^-)^J]$ for every $x \in \mathbf{n}^J$. By definition it is

$$\mathbb{F}[(U_f^-)^J] = \text{Sym}(\{ \psi_x \mid x \in \mathbf{n}^J \}).$$

Fix $\Lambda \in P^+$ with $\Lambda(h_i) > 0$ for all $i \in I \setminus J$. By Theorem 3.27 it holds

$$\mathbb{F}[(U_f^-)^J] = \text{Sym}(\{ (\pi(x)\delta_\Lambda)(\cdot v_\Lambda) \mid x \in \mathbf{n}^J \}).$$

The linear map which maps ψ_x to $(\pi(x)\delta_\Lambda)(\cdot v_\Lambda)$, $x \in \mathbf{n}^J$, induces an automorphism of the algebra $\mathbb{F}[(U_f^-)^J]$. The next theorem describes this automorphism as a comorphism. It will not be used in this article, but it rounds off the description of $\mathbb{F}[(U_f^-)^J]$ nicely.

Theorem 3.30 *Let $J \subsetneq I$ and $\Lambda \in P^+$ with $\Lambda(h_i) > 0$ for all $i \in I \setminus J$. Set $h := \nu^{-1}(\Lambda) \in \mathbf{h}$.*

(a) *We get a bijective map $d\Xi_h : (\mathbf{n}_f^-)^J \rightarrow (\mathbf{n}_f^-)^J$ by*

$$(d\Xi_h)(x) := \text{Ad}(\exp(-x))h - h = \sum_{k \in \mathbb{N}} \frac{1}{k!} (-\text{ad}(x))^k h, \quad x \in (\mathbf{n}_f^-)^J.$$

(b) *The map*

$$\begin{aligned} \Xi_h : (U_f^-)^J &\rightarrow (U_f^-)^J \\ u &\mapsto \exp(\text{Ad}(u^{-1})h - h) \end{aligned}$$

is bijective. Its comorphism $\Xi_h^ : \mathbb{F}[(U_f^-)^J] \rightarrow \mathbb{F}[(U_f^-)^J]$ exists. It is an isomorphism of algebras for which it holds*

$$\Xi_h^*(\psi_x) = (\pi(x)\delta_\Lambda)(\cdot v_\Lambda), \quad x \in \mathbf{n}^J.$$

Proof: To (a): Let $x = \sum_{\beta \in (\Delta^+)^J} x_{-\beta} \in (\mathbf{n}_f^-)^J$. For $\gamma \in \Delta \cup \{0\}$ the $(-\gamma)$ -homogeneous component of the defining sum of $(d\Xi_h)(x)$ is given by

$$\sum_{k \in \mathbb{N}} \frac{1}{k!} \sum_{\substack{\beta_1, \beta_2, \dots, \beta_k \in (\Delta^+)^J \\ \beta_1 + \beta_2 + \dots + \beta_k = \gamma}} (-\text{ad } x_{-\beta_1})(-\text{ad } x_{-\beta_2}) \cdots (-\text{ad } x_{-\beta_k})h,$$

a sum over the empty set defined to be zero. For $\gamma \notin (\Delta^+)^J$ this expression is zero. For $\gamma \in (\Delta^+)^J$ it is a well defined expression: The roots in $(\Delta^+)^J$ have at least height 1. Therefore, it is sufficient to sum from $k = 1$ to $k = ht \gamma$ in the first sum.

Let $\gamma \in (\Delta^+)^J$ and write this component as follows:

$$(d\Xi_h)(x)_{-\gamma} = -\gamma(h)x_{-\gamma} + \sum_{k \in \mathbb{N}, k \geq 2} \frac{(-1)^k}{k!} \sum_{\substack{\beta_1, \beta_2, \dots, \beta_k \in (\Delta^+)^J \\ \beta_1 + \beta_2 + \dots + \beta_k = \gamma}} [x_{-\beta_1}, [x_{-\beta_2}, [\cdots [x_{-\beta_k}, h] \cdots]]]. \quad (38)$$

Here γ is of the form $\gamma = \sum_{i \in I} n_i \alpha_i$ with $n_i \in \mathbb{N}_0$ and $n_{j_0} \neq 0$ for at least one $j_0 \in J$. With $\nu^{-1}(\alpha_i) = \frac{1}{2}(\alpha_i \mid \alpha_i) h_i$ and $(\alpha_i \mid \alpha_i) > 0$, $i \in I$, it follows

$$\gamma(h) = \gamma(\nu^{-1}(\Lambda)) = (\gamma \mid \Lambda) = \Lambda(\nu^{-1}(\gamma)) = \sum_{i \in I, i \neq j_0} \underbrace{n_i}_{\geq 0} \underbrace{\frac{(\alpha_i \mid \alpha_i)}{2}}_{\geq 0} \underbrace{\Lambda(h_i)}_{\geq 0} + \underbrace{n_{j_0}}_{> 0} \underbrace{\frac{(\alpha_{j_0} \mid \alpha_{j_0})}{2}}_{> 0} \underbrace{\Lambda(h_{j_0})}_{> 0} > 0. \quad (39)$$

Let $y = \sum_{\beta \in (\Delta^+)^J} y_{-\beta} \in (\mathbf{n}_f^-)^J$. We show that there exists a unique $x = \sum_{\beta \in (\Delta^+)^J} x_{-\beta} \in (\mathbf{n}_f^-)^J$ such that $(d\Xi_h)(x) = y$ or equivalently

$$(d\Xi_h)(x)_{-\gamma} = y_{-\gamma} \quad \text{for all } \gamma \in (\Delta^+)^J.$$

We do this by induction over $ht \gamma$. If $ht \gamma = 1$, then from (38) and (39) it follows $x_{-\gamma} = \frac{1}{-\gamma(h)} y_{-\gamma}$ uniquely. Now let $ht \gamma = m + 1$, $m \in \mathbb{N}$, and suppose that $x_{-\gamma'}$ has been determined uniquely for all $\gamma' \in (\Delta^+)^J$ with $ht \gamma' \leq m$. Then from (38) and (39) it follows

$$x_{-\gamma} = \frac{1}{-\gamma(h)} y_{-\gamma} - \frac{1}{-\gamma(h)} \sum_{k \in \mathbb{N}, k \geq 2} \frac{(-1)^k}{k!} \sum_{\substack{\beta_1, \beta_2, \dots, \beta_k \in (\Delta^+)^J \\ \beta_1 + \beta_2 + \dots + \beta_k = \gamma}} [x_{-\beta_1}, [x_{-\beta_2}, [\dots [x_{-\beta_k}, h] \dots]]]$$

uniquely.

To (b): The map $d\Xi_h : (\mathbf{n}_f^-)^J \rightarrow (\mathbf{n}_f^-)^J$ and the map $exp : (\mathbf{n}_f^-)^J \rightarrow (U_f^-)^J$ are bijective. Therefore, by its definition also the map $\Xi_h : (U_f^-)^J \rightarrow (U_f^-)^J$ is bijective. Since ψ_x , $x \in \mathbf{n}^J$, as well as $(\pi(x)\delta_\Lambda)(\cdot v_\Lambda)$, $x \in \mathbf{n}^J$, generate the algebra $\mathbb{F}[U_f^-]$ it only remains to show $\psi_x \circ \Xi_h = (\pi(x)\delta_\Lambda)(\cdot v_\Lambda)$ for $x \in \mathbf{n}^J$. For all $y \in (\mathbf{n}_f^-)^J$ it holds:

$$\begin{aligned} \psi_x(\Xi_h(exp(y))) &= ((exp^*)^{-1} \tilde{\psi}_x) \left(exp((d\Xi_h)(y)) \right) = \tilde{\psi}_x((d\Xi_h)(y)) = \tilde{\psi}_x \left(\sum_{k \in \mathbb{N}} \frac{1}{k!} (-ad(y))^k h \right) \\ &= \sum_{\gamma \in (\Delta^+)^J} \sum_{k \in \mathbb{N}} \frac{(-1)^k}{k!} \sum_{\substack{\beta_1, \beta_2, \dots, \beta_k \in (\Delta^+)^J \\ \beta_1 + \beta_2 + \dots + \beta_k = \gamma}} \tilde{\psi}_x([y_{-\beta_1}, [y_{-\beta_2}, [\dots [y_{-\beta_k}, h] \dots]]]) \end{aligned} \quad (40)$$

We transform the inner summand of this expression:

$$\begin{aligned} \tilde{\psi}_x([y_{-\beta_1}, [\dots [y_{-\beta_k}, h] \dots]]) &= (x \mid [y_{-\beta_1}, [\dots [y_{-\beta_k}, h] \dots]]) = ([y_{-\beta_k}, [\dots [y_{-\beta_1}, x] \dots]] \mid h) \\ &= ([y_{-\beta_k}, [\dots [y_{-\beta_1}, x] \dots]]_0 \mid h) \end{aligned}$$

Here we used $(\mathbf{g}_\alpha \mid \mathbf{h}) = \{0\}$ for all $\alpha \in \Delta$ and $[y_{-\beta_k}, [\dots [y_{-\beta_1}, x] \dots]]_0$ denotes the 0-homogeneous component of $[y_{-\beta_k}, [\dots [y_{-\beta_1}, x] \dots]]$. Now we insert $h = \nu^{-1}(\Lambda)$ and transform further:

$$\begin{aligned} \Lambda([y_{-\beta_k}, [\dots [y_{-\beta_1}, x] \dots]]_0) &= \delta_\Lambda([y_{-\beta_k}, [\dots [y_{-\beta_1}, x] \dots]]_0 v_\Lambda) = \delta_\Lambda([y_{-\beta_k}, [\dots [y_{-\beta_1}, x] \dots]] v_\Lambda) \\ &= (\pi([y_{-\beta_k}, [\dots [y_{-\beta_1}, x] \dots]]) \delta_\Lambda)(v_\Lambda) = (-1)^k (\pi(y_{-\beta_k}), [\dots [\pi(y_{-\beta_1}), \pi(x)] \dots]) \delta_\Lambda(v_\Lambda) \end{aligned}$$

Since $\pi(y_{-\beta_1}), \dots, \pi(y_{-\beta_k})$ kills δ_Λ it only remains

$$(-1)^k (\pi(y_{-\beta_k}) \dots \pi(y_{-\beta_1}) \pi(x) \delta_\Lambda)(v_\Lambda) = (-1)^k (\pi(x) \delta_\Lambda)(y_{-\beta_1} \dots y_{-\beta_k} v_\Lambda).$$

Inserting in (40) we get

$$\psi_x(\Xi_h(exp(y))) = (\pi(x) \delta_\Lambda) \left(\sum_{\gamma \in (\Delta^+)^J} \sum_{k \in \mathbb{N}} \frac{1}{k!} \sum_{\substack{\beta_1, \beta_2, \dots, \beta_k \in (\Delta^+)^J \\ \beta_1 + \beta_2 + \dots + \beta_k = \gamma}} y_{-\beta_1} y_{-\beta_2} \dots y_{-\beta_k} v_\Lambda \right) \quad (41)$$

Here the sum over $\gamma \in (\Delta^+)^J$ can be replaced by a sum over $\gamma \in Q^+$, because of $\pi(x) \delta_\Lambda \in \bigoplus_{\lambda \in \Lambda - (\Delta^+)^J} L(\Lambda)_\lambda^*$. If we use in addition $(\pi(x) \delta_\Lambda)(v_\Lambda) = \delta_\Lambda(x v_\Lambda) = 0$ we find

$$\psi_x(\Xi_h(exp(y))) = (\pi(x) \delta_\Lambda) \left(\sum_{k \in \mathbb{N}} \frac{1}{k!} y^k v_\Lambda \right) = (\pi(x) \delta_\Lambda)(exp(y) v_\Lambda).$$

□

Remark 3.31 (a) V. Kac and D. Peterson described in Lemma 4.3 of [KP2] the coordinate ring $\mathbb{F}[G]_{|U}$ as a symmetric algebra in the matrix coefficients $\delta_\Lambda(\cdot xv_\Lambda)$, $x \in \mathfrak{n}^-$, where $\Lambda \in P^+ \cap C$ is arbitrary, the key steps as follows: Equip \mathfrak{n} with the coordinate ring $\mathbb{F}[\mathfrak{n}] = \text{Sym}(\bigoplus_{\alpha \in \Delta^+} \mathfrak{g}_\alpha^*)$. Choose $h \in \mathfrak{h}$ such that $\alpha(h) \neq 0$ for all $\alpha \in \Delta$. V. Kac and D. Peterson showed that the map $\psi : U \rightarrow \mathfrak{n}$ given by $\psi(u) = \text{Ad}(u)h - h$, $u \in U$, induces a bijective comorphism $\psi^* : \mathbb{F}[\mathfrak{n}] \rightarrow \mathbb{F}[U]$. This is tricky in the context of the minimal Kac-Moody group, in particular: The group U is only defined by generators. The map ψ is in general not surjective, and it is not at all obvious that its image is dense. Look at §4 of [KP2] and check the details.

The author tried in [M1], Theorem 5.6, to adapt the proof of V. Kac and D. Peterson to describe for $J \subseteq I$ the coordinate ring $\mathbb{F}[G]_{|U^J}$, but only reached that $\mathbb{F}[G]_{|U^J}$ is generated by the matrix coefficients $\delta_\Lambda(\cdot xv_\Lambda)$, $x \in (\mathfrak{n}^-)^J$, where $\Lambda \in P^+ \cap F_J$ is arbitrary.

As shown above, the descriptions of these coordinate rings can be reached more directly in the context of formal Kac-Moody groups.

(b) The proofs of Theorem 3.27 and Theorem 3.30 work quite generally: It is easy to modify Theorem 3.27 such that it holds in the case where $(\Delta^-)^J$ is replaced by a subset $\Omega \subseteq \Delta^-$. It is easy to modify Theorem 3.30 such that it holds in the case where $(\Delta^-)^J$ is replaced by a bracket closed subset $\Omega \subseteq \Delta^-$. (Bracket closed means: If $\alpha, \beta \in \Omega$ and $\alpha + \beta \in \Delta^-$, then also $\alpha + \beta \in \Omega$.)

We will also use

Proposition 3.32 Let $J \subseteq I$. Let $\phi \in L_J(\Lambda)^{(*)}$, $v \in L_J(\Lambda)_f$, $\Lambda \in P^+$. It holds

$$\phi(uv) = \phi(v) \quad \text{for all } u \in (U_f^-)^J.$$

Proof: This holds trivially for $J = I$. Let $J \neq I$. It holds $\phi \in \bigoplus_{\lambda \in \Lambda - (Q_J)_0^+} L(\Lambda)_\lambda^*$ and $v \in \prod_{\lambda \in \Lambda - (Q_J)_0^+} L(\Lambda)_\lambda$. Furthermore, $u \in (U_f^-)^J$ acts on v as $\exp(x)$ for some $x \in \prod_{\alpha \in (\Delta^-)^J} \mathfrak{g}_\alpha$. It follows

$$uv - v \in \prod_{\lambda \in \Lambda - (Q_J)_0^+ - Q_0^+ \setminus (Q_J)_0^+} L(\Lambda)_\lambda.$$

Therefore, $\phi(uv - v) = 0$. □

For $J \subseteq I$ set

$$CA_J := \bigoplus_{\Lambda \in P_J^+} L_J(\Lambda)^{(*)},$$

which is a P_J^+ -graded subalgebra of CA by Proposition 3.21 (a). It holds $CA_\emptyset = \mathbb{F}1$ and:

Proposition 3.33 For $\emptyset \neq J \subseteq I$ the algebra CA_J is the Cartan algebra associated to G_J , the highest weight vectors chosen as $v_\Lambda \in L_J(\Lambda)_\Lambda$, $\Lambda \in P_J^+$.

Proof: For every $\Lambda \in P^+$ the G -orbit Gv_Λ spans $L(\Lambda)$, because $L(\Lambda)$ is an irreducible G -module. Now let $\Lambda, N \in P^+$ and $\phi \in L(\Lambda)^{(*)}$, $\psi \in L(N)^{(*)}$. Then the Cartan product $\phi \bullet \psi \in L(\Lambda + N)^{(*)}$ is determined by

$$(\phi \bullet \psi)(gv_{\Lambda+N}) = (\phi \otimes \psi)(g(v_\Lambda \otimes v_N)) = (\phi \otimes \psi)(gv_\Lambda \otimes gv_N) = \phi(gv_\Lambda)\psi(gv_N), \quad g \in G. \quad (42)$$

The subgroup G_J of G identifies with the derived Kac-Moody group $G(A_J)'$ to the generalized Cartan submatrix A_J of A . The sublattice P_J of P identifies with the weight lattice, and P_J^+ identifies with the set of dominant weights of the weight lattice of $G(A_J)'$. For $\Lambda \in P_J^+$ the G_J -module $L_J(\Lambda)$ identifies by Theorem 3.19 with the irreducible highest weight module of $G(A_J)'$, whose highest weight corresponds to Λ .

Let $\Lambda, N \in P_J^+$ and $\phi \in L_J(\Lambda)^{(*)}$, $\psi \in L_J(N)^{(*)}$. Then $\phi \bullet \psi \in L_J(\Lambda + N)^{(*)}$ by Proposition 3.21 (a). Restricting (42) to $g \in G_J$ shows: The product \bullet is also the Cartan product of CA_J , where the

highest weight vectors have been choosen as $v_\Lambda \in L_J(\Lambda)_\Lambda$, $\Lambda \in P_J^+$. \square

For $J \subseteq I$ let

$$\mathbb{F}[P \cap (\overline{F_J} - \overline{F_J})] = \bigoplus_{\Lambda \in P \cap (\overline{F_J} - \overline{F_J})} \mathbb{F}e_\Lambda$$

be the group algebra of the lattice $P \cap (\overline{F_J} - \overline{F_J})$, equipped with its natural $P \cap (\overline{F_J} - \overline{F_J})$ -graduation.

Theorem 3.34 *Let $J \subseteq I$. It holds:*

(a) *There exists an isomorphism of graded algebras*

$$\Gamma : (\widetilde{P^+ \cap \overline{F_J}})^{-1} CA \rightarrow \mathbb{F}[(U_f^-)^J] \otimes CA_J \otimes \mathbb{F}[P \cap (\overline{F_J} - \overline{F_J})]$$

such that

$$\Gamma\left(\frac{\delta_N}{\delta_\Lambda}\right) = 1 \otimes 1 \otimes e_{N-\Lambda} \quad \text{for all } N, \Lambda \in P^+ \cap \overline{F_J}, \quad (43)$$

$$\Gamma(\phi) = 1 \otimes \phi \otimes 1 \quad \text{for all } \phi \in L_J(N)^{(*)}, N \in P_J^+, \quad (44)$$

$$\Gamma\left(\frac{\phi}{\delta_N}\right) = \phi(\cdot v_N) \otimes 1 \otimes 1 \quad \text{for all } \phi \in L(N)^{(*)}, N \in P^+ \cap \overline{F_J}. \quad (45)$$

(b) *The map*

$$\begin{aligned} \alpha : \text{Proj}(\mathbb{F}[(U_f^-)^J] \otimes CA_J) &\rightarrow D(J) \\ Q &\mapsto \Gamma^{-1}(Q \otimes \mathbb{F}[P \cap (\overline{F_J} - \overline{F_J})]) \cap CA \end{aligned}$$

is a homeomorphism, mapping $\text{Proj}(\mathbb{F}[(U_f^-)^J] \otimes CA_J)(\mathbb{F})$ bijectively to $D(J)(\mathbb{F})$.

Proof: Part (b) of the theorem follows with part (a) from Proposition 3.13 and Proposition 2.13. To show part (a) we first interpret and describe in (α) , (β) , (γ) , (δ) the algebras involved as coordinate rings of functions. In (ϵ) we realize Γ as a comorphism. By this approach it is automatic, that Γ is a well defined map. It also shows how Γ can be found.

(α) The functions of $\mathbb{F}[G]^U$ take constant values on the elements of every coset gU , $g \in G$. Therefore, they induce a coordinate ring of functions on G/U isomorphic to $\mathbb{F}[G]^U$. We denote this coordinate ring on G/U also by $\mathbb{F}[G]^U$. For $\Lambda \in P^+$ set

$$\theta_\Lambda := \delta_\Lambda(\cdot v_\Lambda) \in \mathbb{F}[G]^U.$$

Define

$$D_{G/U}(J) := \{gU \in G/U \mid \theta_\Lambda(gU) \neq 0 \text{ for all } \Lambda \in P^+ \cap \overline{F_J}\},$$

and equip this set with the coordinate ring of functions

$$\mathbb{F}[D_{G/U}(J)] := \left\{ \frac{f}{\theta_\Lambda} \Big|_{D_{G/U}(J)} \mid f \in \mathbb{F}[G]^U, \Lambda \in P^+ \cap \overline{F_J} \right\}.$$

The Borel-Weil isomorphism of algebras

$$BW : CA = \bigoplus_{\Lambda \in P^+} L(\Lambda)^{(*)} \rightarrow \mathbb{F}[G]^U$$

is given by $BW(\phi) := \phi(\cdot v_\Lambda)$ for all $\phi \in L(\Lambda)^{(*)}$, $\Lambda \in P^+$, compare Section 1 and Subsection 3.1. It is easy to check that it induces a well defined surjective homomorphism of algebras

$$BW^J : (\widetilde{P^+ \cap \overline{F_J}})^{-1} CA \rightarrow \mathbb{F}[D_{G/U}(J)] \quad (46)$$

by $BW^J(\frac{\phi}{\theta_\Lambda}) := \frac{BW(\phi)}{\theta_\Lambda}|_{D_{G/U}(J)}$ for all $\phi \in CA$, $\Lambda \in P^+ \cap \overline{F_J}$. This map is also injective, because $D_{G/U}(J)$ is dense in G/U , which can be seen as follows: Let $N \in P \cap C$. By the Birkhoff decomposition of G it holds

$$\{gU \in G/U \mid \theta_N(gU) \neq 0\} = U^-TU/U \subseteq D_{G/U}(J).$$

Now $\mathbb{F}[G]^U$ is an integral domain, compare Section 1. Therefore, the principal open set on the left and also $D_{G/U}(J)$ is dense in G/U .

(β) We denote by $\mathbb{F}[G_J]$ the restriction of $\mathbb{F}[G]$ to G_J . It holds $G_\emptyset = \{1\}$ by definition, and $\mathbb{F}[G_\emptyset] = \mathbb{F}1$. For $J \neq \emptyset$ the subgroup G_J of G identifies with the derived Kac-Moody group $G(A_J)'$ to the generalized Cartan submatrix A_J of A . It is well-known that $\mathbb{F}[G_J]$ identifies with the algebra of strongly regular functions on $G(A_J)'$ of [KP2]. (To check this use Theorem 3.19.)

We denote by $\mathbb{F}[G_J]^{U_J}$ the functions of $\mathbb{F}[G_J]$ which take constant values on the elements of every coset gU_J , $g \in G_J$. If $J \neq \emptyset$ then by Corollary 2.2 and Remark 2.1 of [KP2] there holds a Borel-Weil Theorem : An isomorphism of algebras

$$BW_J : CA_J = \bigoplus_{\Lambda \in P_J^+} L_J(\Lambda)^{(*)} \rightarrow \mathbb{F}[G_J]^{U_J} \quad (47)$$

is given by $BW_J(\phi) := \phi(\cdot v_\Lambda)$ for all $\phi \in L_J(\Lambda)^{(*)}$, $\Lambda \in P_J^+$. By our definitions this also holds for $J = \emptyset$. We identify $\mathbb{F}[G_J]^{U_J}$ with its corresponding coordinate ring of functions on G_J/U_J .

(γ) It holds

$$P \cap (\overline{F_J} - \overline{F_J}) = \{\lambda \in P \mid \lambda(h_i) = 0 \text{ for all } i \in J\} = \bigoplus_{i \in \{1, \dots, 2n-l\} \setminus J} \mathbb{Z}\Lambda_i.$$

The torus T^J has been defined as the subtorus of T generated by $t_{h_i}(\mathbb{F}^\times)$, $i \in \{1, 2, \dots, 2n-l\} \setminus J$. Therefore, the group algebra $\mathbb{F}[P \cap (\overline{F_J} - \overline{F_J})]$ identifies with the classical coordinate ring of the torus T^J . Explicitly, the generator $e_\lambda \in \mathbb{F}[P \cap (\overline{F_J} - \overline{F_J})]$, $\lambda \in P \cap (\overline{F_J} - \overline{F_J})$, identifies with the function

$$e_\lambda \left(\prod_{i \in \{1, \dots, 2n-l\} \setminus J} h_i(s_i) \right) = \prod_{i \in \{1, \dots, 2n-l\} \setminus J} (s_i)^{\lambda(h_i)}, \text{ where } s_i \in \mathbb{F}^\times, i \in \{1, \dots, 2n-l\} \setminus J.$$

(δ) By Corollary 3.28 the restriction map $\mathbb{F}[(U_f^-)^J] \rightarrow \mathbb{F}[(U^-)^J]$ is an isomorphism of algebras. By Theorem 3.27 it follows

$$\mathbb{F}[(U^-)^J] = \text{span} \left\{ \phi(\cdot v_N) \mid \phi \in L(N)^{(*)}, N \in P^+ \cap \overline{F_J} \right\}.$$

By Corollary 3.29 it holds $\mathbb{F}[G]|_{(U^-)^J} = \mathbb{F}[(U^-)^J]$.

(ϵ) Now we show: The map

$$\begin{aligned} m : (U^-)^J \times G_J/U_J \times T^J &\rightarrow D_{G/U}(J) \\ (u^-, gU_J, t) &\mapsto u^-gtU \end{aligned}$$

is well defined and induces a comorphism

$$m^* : \mathbb{F}[D_{G/U}(J)] \rightarrow \mathbb{F}[(U^-)^J] \otimes \mathbb{F}[G_J]^{U_J} \otimes \mathbb{F}[P \cap (\overline{F_J} - \overline{F_J})],$$

which is an isomorphism of algebras. Furthermore, it holds:

$$m^* \left(\frac{\theta_N}{\theta_\Lambda} |_{D_{G/U}(J)} \right) = 1 \otimes 1 \otimes e_{N-\Lambda} \quad \text{for } N, \Lambda \in P^+ \cap \overline{F_J}, \quad (48)$$

$$m^* \left(\phi(\cdot v_N) |_{D_{G/U}(J)} \right) = 1 \otimes \phi(\cdot v_N) |_{G_J/U_J} \otimes 1 \quad \text{for } \phi \in L_J(N)^{(*)}, N \in P_J^+. \quad (49)$$

$$m^* \left(\frac{\phi(\cdot v_N)}{\theta_N} |_{D_{G/U}(J)} \right) = \phi(\cdot v_N) |_{(U^-)^J} \otimes 1 \otimes 1 \quad \text{for } \phi \in L(N)^{(*)}, N \in P^+ \cap \overline{F_J}. \quad (50)$$

We first show that m is well defined as a map to G/U . Let $u^- \in (U^-)^J$, $gU_J = g'U_J \in G_J/U_J$, and $t \in T^J$. Then it holds $u^-gtU = u^-g'tU$ if and only if $gtUt^{-1} = g'tUt^{-1}$ if and only if $gU = g'U$, which follows from $gU_J = g'U_J$.

Next we show that m is a surjective map onto $D_{G/U}(J)$. Let $gU \in G/U$. Write gU in the form $gU = u^-n_\sigma U$ with $u^- \in U^-$, $n_\sigma \in N$ projecting to $\sigma \in \mathcal{W}$. Then it holds $gU \in D_{G/U}(J)$ if and only if

$$\delta_\Lambda(u^-n_\sigma v_\Lambda) \neq 0 \quad \text{for all} \quad \Lambda \in P^+ \cap \overline{F_J},$$

if and only if $\sigma\Lambda = \Lambda$ for all $\Lambda \in P^+ \cap \overline{F_J}$, if and only if $\sigma \in \mathcal{W}_J$. Therefore,

$$D_{G/U}(J) = U^-N_JT^J/U = (U^-)^JU_J^-N_JT^J/U = m((U^-)^J, G_J/U_J, T^J).$$

Now let $\phi \in L(N)_\lambda^*$, $\lambda \in P(N)$, and $N \in P^+$. Choose dual bases $\phi_{\mu j} \in L(N)_\mu^*$, $v_{\mu j} \in L(N)_\mu$, $\mu \in P(N)$, $j = 1, \dots, \dim(L(N)_\mu)$. Decompose $N = N_J + N^J$ with $N_J \in P_J^+$ and $N^J \in P^+ \cap \overline{F_J}$. Let $\Lambda \in P^+ \cap \overline{F_J}$. Note that $G_J \subseteq Z_G(L(\Lambda)_\Lambda)$. For all $u^- \in (U^-)^J$, $g \in G_J$, and $t \in T^J$ it holds

$$m^* \left(\frac{\phi(\cdot v_N)}{\theta_\Lambda} \Big|_{D_{G/U}(J)} \right) (u^-, gU, t) = \frac{\phi(u^-gtv_N)}{\delta_\Lambda(u^-gtv_\Lambda)} = \phi(u^-gv_N)e_{N^J-\Lambda}(t) \quad (51)$$

$$\begin{aligned} &= \sum_{\mu, j} \phi(u^-v_{\mu j})\phi_{\mu j}(gv_N)e_{N^J-\Lambda}(t) \\ &= \left(\sum_{\substack{\mu, j \\ \mu \geq \lambda}} \phi(\cdot v_{\mu j}) \otimes \phi_{\mu j}(\cdot v_N) \otimes e_{N^J-\Lambda} \right) (u^-, gU, t). \end{aligned} \quad (52)$$

The sum in (52) is finite. It follows that the comorphism m^* exists. Again by $G_J \subseteq Z_G(L(N)_N)$ for all $N \in P^+ \cap \overline{F_J}$ it follows (48) and (50) from (51). By Proposition 3.32 it follows (49) from (51). The elements in (48) on the right generate $\mathbb{F}1 \otimes \mathbb{F}1 \otimes \mathbb{F}[P \cap (\overline{F_J} - \overline{F_J})]$, in (49) they generate $\mathbb{F}1 \otimes \mathbb{F}[G_J]^{U_J} \otimes \mathbb{F}1$, and in (50) they generate $\mathbb{F}[(U^-)^J] \otimes \mathbb{F}1 \otimes \mathbb{F}1$. It follows that the morphism of algebras m^* is surjective. Furthermore, the comorphism m^* is injective, because m is surjective.

Now, by the isomorphisms and identifications of (α) , (β) , (γ) , (δ) the isomorphism m^* coincides with Γ .

The algebra $(\widetilde{P^+ \cap \overline{F_J}})^{-1} CA$ is graded by

$$P^+ - (P^+ \cap \overline{F_J}).$$

The algebra $\mathbb{F}[(U_f^-)^J] \otimes CA_J \otimes \mathbb{F}[P \cap (\overline{F_J} - \overline{F_J})]$ is graded by

$$\{0\} \oplus P_J^+ \oplus (P \cap (\overline{F_J} - \overline{F_J})).$$

Both gradings identify with the submonoid $P \cap (\overline{C} - \overline{F_J})$ of P . The isomorphism Γ preserves the degree of the generators in (43), (44), (45). It follows that Γ is an isomorphism of graded algebras, the induced map on the gradings $P \cap (\overline{C} - \overline{F_J})$ given by the identity map. \square

The first conclusion of the last theorem is the following description of the standard big cells:

Corollary 3.35 *Let $J \subseteq I$. It holds:*

(a) *There exists an isomorphism of graded algebras*

$$\Gamma : (\widetilde{P^+ \cap \overline{F_J}})^{-1} \bigoplus_{\Lambda \in P^+ \cap \overline{F_J}} L(\Lambda)^{(*)} \rightarrow \mathbb{F}[(U_f^-)^J] \otimes \mathbb{F}[P \cap (\overline{F_J} - \overline{F_J})]$$

such that

$$\begin{aligned} \Gamma\left(\frac{\delta_N}{\delta_\Lambda}\right) &= 1 \otimes e_{N-\Lambda} & \text{for all} & \quad N, \Lambda \in P^+ \cap \overline{F_J}, \\ \Gamma\left(\frac{\phi}{\delta_N}\right) &= \phi(\cdot v_N) \otimes 1 & \text{for all} & \quad \phi \in L(N)^{(*)}, N \in P^+ \cap \overline{F_J}. \end{aligned}$$

(b) The map

$$\begin{aligned} \beta : \text{Spec}(\mathbb{F}[(U_f^-)^J]) &\rightarrow BC(J) \\ Q &\mapsto \left(\Gamma^{-1}(Q \otimes \mathbb{F}[P \cap (\overline{F_J} - \overline{F_J})]) \cap \bigoplus_{\Lambda \in P^+ \cap \overline{F_J}} L(\Lambda)^{(*)} \right) \oplus \bigoplus_{\Lambda \in P^+ \setminus \overline{F_J}} L(\Lambda)^{(*)} \end{aligned}$$

is a homeomorphism, mapping $\text{Spec}(\mathbb{F}[(U_f^-)^J])(\mathbb{F})$ bijectively to $BC(J)(\mathbb{F})$.

Proof: The isomorphism of graded algebras of part (a) is obtained by restricting the graded isomorphism of Theorem 3.34. Part (b) follows with (a) from Proposition 3.18 and Proposition 2.13. \square

For $J \subseteq I$ identify the topological space $\overline{Or(J)}$ with $Proj(\bigoplus_{\Lambda \in P^+ \cap \overline{F_J}} L(\Lambda)^{(*)})$ by Proposition 3.11. By this identification $\overline{Or(J)}$ gets the structure of a locally ringed space of \mathbb{F} -algebras. The covering of $Or(J)$ by big cells leads to:

Corollary 3.36 *Let $J \subseteq I$. The open set $Or(J)$ of $\overline{Or(J)}$ equipped with its locally ringed substructure is a scheme.*

Proof: Follows from Theorem 3.17, from Corollary 3.18, Corollary 2.16, and from Corollary 3.35, Corollary 2.17. \square

With Corollary 3.35 we can also determine the \mathbb{F} -valued points contained in a standard big cell:

Theorem 3.37 *For $J \subseteq I$ it holds $BC(J)(\mathbb{F}) = (U_f^-)^J P(J)$.*

Proof: By Proposition 3.26 the vanishing ideals $I(u)$, $u \in (U_f^-)^J$, give the \mathbb{F} -valued points of $\mathbb{F}[(U_f^-)^J]$. By Corollary 3.35 (b) its images under the map β give the \mathbb{F} -valued points of $BC(J)$. For $u \in (U_f^-)^J$ we now compute $\beta(I(u))$ explicitly. Let $\phi \in L(\Lambda)^{(*)}$, $\Lambda \in P^+ \cap \overline{F_J}$. Then it holds

$$\frac{\phi}{\delta_0} \in \Gamma^{-1}(I(u) \otimes \mathbb{F}[P \cap (\overline{F_J} - \overline{F_J})])$$

if and only if

$$\Gamma\left(\frac{\phi}{\delta_0}\right) = \phi(\cdot v_\Lambda) \otimes e_\Lambda \in I(u) \otimes \mathbb{F}[P \cap (\overline{F_J} - \overline{F_J})]$$

if and only if

$$0 = \phi(uv_\Lambda) = (\pi(u)\phi)(v_\Lambda)$$

if and only if $\pi(u)\phi \in L(\Lambda)^{(*)}_{\neq \Lambda}$. Since $\Gamma^{-1}(I(u) \otimes \mathbb{F}[P \cap (\overline{F_J} - \overline{F_J})])$ is a homogeneous ideal we have shown

$$\beta(I(u)) = \bigoplus_{\Lambda \in P^+ \cap \overline{F_J}} \pi(u)^{-1} L(\Lambda)^{(*)}_{\neq \Lambda} \oplus \bigoplus_{\Lambda \in P^+ \setminus \overline{F_J}} L(\Lambda)^{(*)} = uP(J).$$

\square

Now we use again the covering of the Or -strata by big cells and finally reach:

Corollary 3.38 *It holds*

$$Proj(CA)(\mathbb{F}) = \bigcup_{J \subseteq I} Or(J)(\mathbb{F}) \quad \text{with} \quad Or(J)(\mathbb{F}) = G_{fn} P(J) \quad \text{for all } J \subseteq I.$$

In particular, the map ω of Theorem 3.9 is also surjective.

Proof: The decomposition of $\text{Proj}(CA)(\mathbb{F})$ is given by Proposition 3.12 (c). From the last theorem and Theorem 3.17 (b) it follows

$$\text{Or}(J)(\mathbb{F}) = \bigcup_{g \in G_{fn}} gBC(J)(\mathbb{F}) = G_{fn}(U_f^-)^J P(J) = G_{fn} P(J).$$

□

At last we investigate the transversal behaviour of the Or-stratification of $\text{Proj}(CA)$. For $J \subseteq I$ we get a stratification of $\text{Proj}(CA_J)$, similarly as for $\text{Proj}(CA)$, as follows: For every $K \subseteq J$ it is $\bigoplus_{\Lambda \in P_J^+ \setminus P_{J \setminus K}^+} L_J(\Lambda)^{(*)}$ a homogeneous prime ideal of CA_J . Set

$$\overline{\text{Or}_J(K)} := \mathcal{V}\left(\bigoplus_{\Lambda \in P_J^+ \setminus P_{J \setminus K}^+} L_J(\Lambda)^{(*)}\right) \subseteq \text{Proj}(CA_J).$$

For $K \subseteq J$ define

$$\text{Or}_J(K) := \overline{\text{Or}_J(K)} \setminus \bigcup_{K \subsetneq T \subseteq J} \overline{\text{Or}_J(T)}.$$

Then $\text{Or}_J(K) \neq \emptyset$, $\text{Or}_J(K)$ is open and dense in $\overline{\text{Or}_J(K)}$, and $\text{Proj}(CA_J) = \bigcup_{K \subseteq J} \text{Or}_J(K)$.

As a second consequence of Theorem 3.34 we obtain:

Corollary 3.39 *Let $J \subseteq I$. It holds:*

(a) *There exists an isomorphism of graded algebras*

$$\Gamma : (\widetilde{P^+ \cap \overline{F_J}})^{-1} \bigoplus_{\Lambda \in P^+} L_J(\Lambda)^{(*)} \rightarrow CA_J \otimes \mathbb{F}[P \cap (\overline{F_J} - \overline{F_J})]$$

such that

$$\begin{aligned} \Gamma\left(\frac{\delta_N}{\delta_\Lambda}\right) &= 1 \otimes e_{N-\Lambda} & \text{for all } N, \Lambda \in P^+ \cap \overline{F_J}, \\ \Gamma(\phi) &= \phi \otimes 1 & \text{for all } \phi \in L_J(N)^{(*)}, N \in P_J^+. \end{aligned}$$

(b) *The map*

$$\begin{aligned} \gamma : \text{Proj}(CA_J) &\rightarrow S(J) \\ Q &\mapsto \left(\Gamma^{-1}(Q \otimes \mathbb{F}[P \cap (\overline{F_J} - \overline{F_J})]) \cap \bigoplus_{\Lambda \in P^+} L_J(\Lambda)^{(*)} \right) \oplus \bigoplus_{\Lambda \in P^+} R_J(\Lambda)^{(*)} \end{aligned}$$

is a homeomorphism, mapping $\text{Proj}(CA_J)(\mathbb{F})$ bijectively to $S(J)(\mathbb{F})$. Furthermore, it holds

$$\gamma(\text{Or}_J(K)) = S(J) \cap \overline{\text{Or}_J(K)} \quad \text{for all } K \subseteq J. \quad (53)$$

Proof: The isomorphism of graded algebras of part (a) is obtained by restricting the graded isomorphism of Theorem 3.34. Part (b), except (53), follows with (a) from Proposition 3.24 and Proposition 2.13. To prove (53) we first show

$$\gamma(\overline{\text{Or}_J(K)}) = S(J) \cap \overline{\text{Or}_J(K)} \quad \text{for all } K \subseteq J. \quad (54)$$

Because it holds

$$\gamma(\overline{\text{Or}_J(K)}) = \gamma\left(\mathcal{V}\left(\bigoplus_{\Lambda \in P_J^+ \setminus P_{J \setminus K}^+} L_J(\Lambda)^{(*)}\right)\right) = \mathcal{V}\left(\gamma\left(\bigoplus_{\Lambda \in P_J^+ \setminus P_{J \setminus K}^+} L_J(\Lambda)^{(*)}\right) \cap S(J)\right), \quad (55)$$

we first determine $\gamma(\bigoplus_{\Lambda \in P_J^+ \setminus P_{J \setminus K}^+} L_J(\Lambda)^{(*)})$. It holds $P^+ = P_J^+ \oplus (P^+ \cap \overline{F_J})$. Let $\Lambda_1 \in P_J^+$ and $\Lambda_2 \in P^+ \cap \overline{F_J}$. From the definition of the map γ it follows easily

$$\gamma\left(\bigoplus_{\Lambda \in P_J^+ \setminus P_{J \setminus K}^+} L_J(\Lambda)^{(*)}\right)_{\Lambda_1 + \Lambda_2} = \begin{cases} L_J(\Lambda_1)^{(*)} \bullet \delta_{\Lambda_2} + R_J(\Lambda_1 + \Lambda_2)^{(*)} & \text{if } \Lambda_1 \in P_J^+ \setminus P_{J \setminus K}^+ \\ R_J(\Lambda_1 + \Lambda_2)^{(*)} & \text{if } \Lambda_1 \in P_{J \setminus K}^+ \end{cases}.$$

Here, it holds $L_J(\Lambda_1)^{(*)} \bullet \delta_{\Lambda_2} \subseteq L_J(\Lambda_1 + \Lambda_2)^{(*)}$ by Proposition 3.21 (a). To show equality note that the \mathfrak{g}_J -module $L_J(\Lambda_2)$ is trivial. It follows that G_J^{op} acts trivially on $L_J(\Lambda_2)^{(*)} = \mathbb{F}\delta_{\Lambda_2}$. It holds

$$\pi(g)\delta_{\Lambda_1 + \Lambda_2} = \pi(g)(\delta_{\Lambda_1} \bullet \delta_{\Lambda_2}) = (\pi(g)\delta_{\Lambda_1}) \bullet (\pi(g)\delta_{\Lambda_2}) = (\pi(g)\delta_{\Lambda_1}) \bullet \delta_{\Lambda_2} \in L_J(\Lambda_1)^{(*)} \bullet \delta_{\Lambda_2}$$

for all $g \in G_J$. Since the G_J^{op} -orbit of $\delta_{\Lambda_1 + \Lambda_2}$ spans the G_J^{op} -module $L_J(\Lambda_1 + \Lambda_2)^{(*)}$ it follows $L_J(\Lambda_1)^{(*)} \bullet \delta_{\Lambda_2} = L_J(\Lambda_1 + \Lambda_2)^{(*)}$. We have shown

$$\gamma\left(\bigoplus_{\Lambda \in P_J^+ \setminus P_{J \setminus K}^+} L_J(\Lambda)^{(*)}\right)_{\Lambda_1 + \Lambda_2} = \begin{cases} L(\Lambda_1 + \Lambda_2)^{(*)} & \text{if } \Lambda_1 \in P_J^+ \setminus P_{J \setminus K}^+ \\ R_J(\Lambda_1 + \Lambda_2)^{(*)} & \text{if } \Lambda_1 \in P_{J \setminus K}^+ \end{cases}.$$

With $P_{J \setminus K}^+ + (P^+ \cap \overline{F_J}) = P^+ \cap \overline{F_K}$ we get

$$\gamma\left(\bigoplus_{\Lambda \in P_J^+ \setminus P_{J \setminus K}^+} L_J(\Lambda)^{(*)}\right) = \bigoplus_{\Lambda \in P^+ \cap \overline{F_K}} R_J(\Lambda)^{(*)} \oplus \bigoplus_{\Lambda \in P^+ \setminus \overline{F_K}} L(\Lambda)^{(*)}.$$

Now insert in (55) and use that it holds $S(J) \subseteq \mathcal{V}(\bigoplus_{\Lambda \in P^+} R_J(\Lambda)^{(*)})$ by definition:

$$\begin{aligned} \gamma(\overline{Or_J(K)}) &= \left\{ Q \in S(J) \mid Q \supseteq \bigoplus_{\Lambda \in P^+ \cap \overline{F_K}} R_J(\Lambda)^{(*)} \oplus \bigoplus_{\Lambda \in P^+ \setminus \overline{F_K}} L(\Lambda)^{(*)} \right\} \\ &= \left\{ Q \in S(J) \mid Q \supseteq \bigoplus_{\Lambda \in P^+ \setminus \overline{F_K}} L(\Lambda)^{(*)} \right\} = S(J) \cap \overline{Or(K)}. \end{aligned}$$

From the definition of $Or_J(K)$ and (54) it follows

$$\gamma(Or_J(K)) = (S(J) \cap \overline{Or(K)}) \setminus \bigcup_{K \subsetneq T \subseteq J} (S(J) \cap \overline{Or(T)}). \quad (56)$$

Suppose that $T \subseteq I$, $T \not\subseteq J$. Then by Proposition 3.12 (c) and Proposition 3.23 we find

$$S(J) \cap \overline{Or(T)} = \bigcup_{I \supseteq L \supseteq T} (S(J) \cap Or(L)) = \emptyset.$$

Therefore, the union in (56) can be taken over all T with $K \subsetneq T \subseteq I$. It follows (53). \square

Next we determine the \mathbb{F} -valued points of the standard transversal slices $S(J)$, $J \subseteq I$. To do this we don't work with the last Corollary but with the description of the \mathbb{F} -valued points of $\text{Proj}(CA)$ instead. This way we also find the \mathbb{F} -valued points of the principal open sets $D(J)$, $J \subseteq I$.

Theorem 3.40 *For $J \subseteq I$ it holds*

$$D(J)(\mathbb{F}) = \bigcup_{K \subseteq J} (U_f^-)^J (G_{fn})_J P(K) = \bigcup_{K \subseteq J} (G_{fn})_J (U_f^-)^J P(K) \quad \text{and} \quad S(J)(\mathbb{F}) = \bigcup_{K \subseteq J} (G_{fn})_J P(K).$$

Proof: (a) By Corollary 3.38, by the Birkhoff decomposition of G_{fn} , and by $U \subseteq (P_{fn})_K = \text{Stab}_{G_{fn}}(P(K))$ for all $K \subseteq I$, compare Theorem 3.9, we find

$$\text{Proj}(CA)(\mathbb{F}) = \bigcup_{K \subseteq I} G_{fn} P(K) = \bigcup_{K \subseteq I} U_f^- N P(K).$$

We use Remark 3.14 to describe $D(J)(\mathbb{F})$. Fix an arbitrary $\Lambda \in P^+ \cap F_J$. For $K \subseteq I$, $u \in U_f^-$, and $n_\sigma \in N$ projecting to $\sigma \in \mathcal{W}$ we find

$$\begin{aligned} \delta_\Lambda \notin un_\sigma P(K) &= \bigoplus_{N \in P^+ \cap \overline{F_K}} \pi(un_\sigma)^{-1} L(N)_{\neq \Lambda}^{(*)} \oplus \bigoplus_{N \in P^+ \setminus \overline{F_K}} L(N)^{(*)} \\ \iff \Lambda \in P^+ \cap \overline{F_K} \text{ and } \delta_\Lambda \notin \pi(un_\sigma)^{-1} L(\Lambda)_{\neq \Lambda}^{(*)} &\iff J \supseteq K \text{ and } \delta_\Lambda(un_\sigma v_\Lambda) \neq 0 \\ \iff J \supseteq K \text{ and } \sigma\Lambda = \Lambda &\iff J \supseteq K \text{ and } \sigma \in \mathcal{W}_J. \end{aligned}$$

Because of T , $U_J \subseteq (P_{fn})_K = \text{Stab}_{G_{fn}}(P(K))$, $K \subseteq I$, it follows

$$U_f^-(\mathcal{W}_J T)P(K) = (U_f^-)^J (U_f^-)_J N_J U_J P(K) = (U_f^-)^J (G_{fn})_J P(K).$$

From the Levi decomposition of $(P_{fn})_J$ it follows that $(G_{fn})_J \subseteq (L_{fn})_J$ normalizes $(U_f^-)^J$. Therefore, we get $(U_f^-)^J (G_{fn})_J = (G_{fn})_J (U_f^-)^J$.

(b) It holds $S(J) = \mathcal{V}(\bigoplus_{\Lambda \in P^+} R_J(\Lambda)^{(*)}) \cap D(J)$ by definition. We have seen in (a) that the elements of $D(J)(\mathbb{F})$ are of the form $guP(K)$ with $g \in (G_{fn})_J$, $u \in (U_f^-)^J$, and $K \subseteq J$. Because $R_J(\Lambda)^{(*)}$ is invariant under the action of $(G_{fn})_J^{op}$ for all $\Lambda \in P^+$ it holds:

$$\begin{aligned} guP(K) &\supseteq \bigoplus_{\Lambda \in P^+} R_J(\Lambda)^{(*)} \\ \iff uP(K) &= \bigoplus_{\Lambda \in P^+ \cap \overline{F_K}} \pi(u)^{-1} L(\Lambda)_{\neq \Lambda}^{(*)} \oplus \bigoplus_{\Lambda \in P^+ \setminus \overline{F_K}} L(\Lambda)^{(*)} \supseteq \bigoplus_{\Lambda \in P^+} R_J(\Lambda)^{(*)} \\ \iff \pi(u)^{-1} L(\Lambda)_{\neq \Lambda}^{(*)} &= \left\{ \phi \in L(\Lambda)^{(*)} \mid \phi(uv_\Lambda) = 0 \right\} \supseteq R_J(\Lambda)^{(*)} \text{ for all } \Lambda \in P^+ \cap \overline{F_K} \\ \iff uv_\Lambda \in L_J(\Lambda)_f &= \prod_{\lambda \in \Lambda - (Q_0^+)_J} L(\Lambda)_\lambda \text{ for all } \Lambda \in P^+ \cap \overline{F_K}. \end{aligned}$$

Because of $u \in (U_f^-)^J = \exp((\mathbf{n}_f^-)^J)$ with $(\mathbf{n}_f^-)^J = \bigoplus_{\alpha \in (\Delta^-)_J} \mathbf{g}_\alpha$ this is equivalent to

$$uL(\Lambda)_\Lambda \subseteq \prod_{\lambda \in \Lambda - (\{0\} \cup Q_0^+ \setminus (Q_0^+)_J)} L(\Lambda)_\lambda \cap \prod_{\lambda \in \Lambda - (Q_0^+)_J} L(\Lambda)_\lambda = L(\Lambda)_\Lambda$$

for all $\Lambda \in P^+ \cap \overline{F_K}$. These inclusions are equalities, because the highest weight spaces are one-dimensional. By Theorem 3.1 (b), Theorem 3.6 (b), the definition of $(P_{fn})_K$, and Lemma 3.2 this in turn is equivalent to

$$u \in (U_f^-)^J \cap \bigcap_{T \supseteq K} (P_{fn})_T = (U_f^-)^J \cap (P_{fn})_K = (U_f^-)^J \cap (U_f^-)_K (\mathcal{W}_K T) U = (U_f^-)^J \cap (U_f^-)_K.$$

Because of $K \subseteq J$ it holds $(U_f^-)_K \subseteq (U_f^-)_J$, from which it follows $(U_f^-)^J \cap (U_f^-)_K = \{1\}$. \square

Let $J \subseteq I$. The canonical injections

$$\mathbb{F}[(U_f^-)^J] \rightarrow \mathbb{F}[(U_f^-)^J] \otimes CA_J \quad \text{and} \quad CA_J \rightarrow \mathbb{F}[(U_f^-)^J] \otimes CA_J$$

induce a pair of continuous maps

$$\text{Proj}(\mathbb{F}[(U_f^-)^J] \otimes CA_J) \rightarrow \text{Spec}(\mathbb{F}[(U_f^-)^J]) \times \text{Proj}(CA_J).$$

The projective spectrum $\text{Proj}(\mathbb{F}[(U_f^-)^J] \otimes CA_J)$ identifies by Theorem 3.34 with the principal open set $D(J)$. The spectrum $\text{Spec}(\mathbb{F}[(U_f^-)^J])$ identifies by Corollary 3.35 with the standard big cell $BC(J)$. The projective spectrum $\text{Proj}(CA_J)$ identifies by Corollary 3.39 with the standard transversal slice $S(J)$. Therefore, we get a pair of continuous maps

$$p = (p_1, p_2) : D(J) \rightarrow BC(J) \times S(J),$$

whose properties are investigated next. Recall also: $BC(J)$ and $S(J)$ are closed subsets of $D(J)$ by definition. It holds $BC(J) \cap S(J) = \{P(J)\}$ by Theorem 3.22.

We first describe the maps p_1 and p_2 explicitly:

Proposition 3.41 *Let $J \subseteq I$. The map $p_1 : D(J) \rightarrow BC(J)$ is given by*

$$p_1(Q) = \bigoplus_{\Lambda \in P^+ \cap \overline{F_J}} (Q \cap L(\Lambda)^{(*)}) \oplus \bigoplus_{\Lambda \in P^+ \setminus \overline{F_J}} L(\Lambda)^{(*)}, \quad Q \in D(J).$$

The map $p_2 : D(J) \rightarrow S(J)$ is given by

$$p_2(Q) = \bigoplus_{\Lambda \in P^+} \left((Q \cap L_J(\Lambda)^{(*)}) \oplus R_J(\Lambda)^{(*)} \right), \quad Q \in D(J).$$

Proof: We only show the statement for p_1 , the statement for p_2 follows similarly. Consider the following commutative diagram:

$$\begin{array}{ccccc} \mathbb{F}[(U_f^-)^J] \otimes CA_J & \rightarrow & \mathbb{F}[(U_f^-)^J] \otimes CA_J \otimes \mathbb{F}[P \cap (\overline{F_J} - \overline{F_J})] & \rightarrow & (\widetilde{P^+ \cap \overline{F_J}})^{-1} CA \\ \uparrow & & \uparrow & & \uparrow \\ \mathbb{F}[(U_f^-)^J] & \rightarrow & \mathbb{F}[(U_f^-)^J] \otimes \mathbb{F}[P \cap (\overline{F_J} - \overline{F_J})] & \rightarrow & (\widetilde{P^+ \cap \overline{F_J}})^{-1} \bigoplus_{\Lambda \in P^+ \cap \overline{F_J}} L(\Lambda)^{(*)} \end{array}$$

Here the maps of the square on the left are the canonical injections of the tensor products. The remaining horizontal maps on the right are the inverse isomorphisms of graded algebras of Theorem 3.34 (a) and Corollary 3.35 (a). The remaining vertical map on the right is the inclusion map, which is a morphism of graded algebras.

This diagram induces a commutative diagram of continous maps between the corresponding spectra. The two horizontal maps on the left induce homeomorphisms between the corresponding projective spectra by Proposition 2.13. Trivially, the two horizontal maps on the right induce homeomorphisms between the corresponding projective spectra.

By these homeomorphisms, the map induced by $\mathbb{F}[(U_f^-)^J] \rightarrow \mathbb{F}[(U_f^-)^J] \otimes CA_J$ identifies with the map

$$\begin{aligned} \text{Proj}((\widetilde{P^+ \cap \overline{F_J}})^{-1} CA) &\rightarrow \text{Proj}((\widetilde{P^+ \cap \overline{F_J}})^{-1} \bigoplus_{\Lambda \in P^+ \cap \overline{F_J}} L(\Lambda)^{(*)}) \\ \tilde{Q} &\mapsto \tilde{Q} \cap (\widetilde{P^+ \cap \overline{F_J}})^{-1} \bigoplus_{\Lambda \in P^+ \cap \overline{F_J}} L(\Lambda)^{(*)} \end{aligned}$$

which in turn identifies by Propositions 3.13 and 3.18 with the map $p_1 : D(J) \rightarrow BC(J)$. For $Q \in D(J)$ denote by Q^e the ideal in $(\widetilde{P^+ \cap \overline{F_J}})^{-1} CA$ generated by Q . It holds $Q^e \cap CA = Q$ since Q is prime. We find

$$\begin{aligned} p_1(Q) &= \left((Q^e \cap (\widetilde{P^+ \cap \overline{F_J}})^{-1} \bigoplus_{\Lambda \in P^+ \cap \overline{F_J}} L(\Lambda)^{(*)}) \cap \bigoplus_{\Lambda \in P^+ \cap \overline{F_J}} L(\Lambda)^{(*)} \right) \oplus \bigoplus_{\Lambda \in P^+ \setminus \overline{F_J}} L(\Lambda)^{(*)} \\ &= \left(Q \cap \bigoplus_{\Lambda \in P^+ \cap \overline{F_J}} L(\Lambda)^{(*)} \right) \oplus \bigoplus_{\Lambda \in P^+ \setminus \overline{F_J}} L(\Lambda)^{(*)} = \bigoplus_{\Lambda \in P^+ \cap \overline{F_J}} (Q \cap L(\Lambda)^{(*)}) \oplus \bigoplus_{\Lambda \in P^+ \setminus \overline{F_J}} L(\Lambda)^{(*)}. \end{aligned}$$

□

The map $p = (p_1, p_2) : D(J) \rightarrow BC(J) \times S(J)$ has the following properties:

Corollary 3.42 *Let $J \subseteq I$. It holds:*

(a) $p(Q) = (Q, P(J))$ for all $Q \in BC(J)$.

$p(Q) = (P(J), Q)$ for all $Q \in S(J)$.

(b) $p(D(J) \cap \overline{Or(K)}) \subseteq BC(J) \times (S(J) \cap \overline{Or(K)})$ for all $K \subseteq J$.

Proof: To (a): We only show the second equation, the first follows similarly. Let $Q \in S(J)$. The definition $S(J) = \mathcal{V}(\bigoplus_{\Lambda \in P^+} R_J(\Lambda)^{(*)}) \cap D(J)$ implies

$$Q \supseteq \bigoplus_{\Lambda \in P^+} R_J(\Lambda)^{(*)}. \quad (57)$$

From (57) and the description of p_1 in Proposition 3.41 it follows

$$p_1(Q) \supseteq \bigoplus_{\Lambda \in P^+} R_J(\Lambda)^{(*)}.$$

By $BC(J) \subseteq D(J)$, by the definition of $S(J)$, and by Theorem 3.22 we find

$$p_1(Q) \in BC(J) \cap \mathcal{V}(\bigoplus_{\Lambda \in P^+} R_J(\Lambda)^{(*)}) = BC(J) \cap S(J) = \{P(J)\}.$$

From (57) and the description of p_2 in Proposition 3.41 it follows

$$Q = \bigoplus_{\Lambda \in P^+} \left((Q \cap L_J(\Lambda)^{(*)}) \oplus R_J(\Lambda)^{(*)} \right) = p_2(Q).$$

To (b): Let $K \subseteq J$. Let $Q \in D(J)$ with $Q \supseteq \bigoplus_{\Lambda \in P^+ \setminus \overline{F_K}} L(\Lambda)^{(*)}$. Then from the description of p_2 in Proposition 3.41 it follows $p_2(Q) \supseteq \bigoplus_{\Lambda \in P^+ \setminus \overline{F_K}} L(\Lambda)^{(*)}$. \square

On the \mathbb{F} -valued points there holds the following description:

Theorem 3.43 *Let $J \subseteq I$. The map $p : D(J) \rightarrow BC(J) \times S(J)$ restricts to a bijective map $p : D(J)(\mathbb{F}) \rightarrow BC(J)(\mathbb{F}) \times S(J)(\mathbb{F})$, which is given by*

$$p(ugP(K)) = (uP(J), gP(K)) \quad \text{with} \quad u \in (U_f^-)^J, g \in (G_{fn})_J, K \subseteq J.$$

In particular, it holds $p((D(J) \cap Or(K))(\mathbb{F})) = BC(J)(\mathbb{F}) \times (S(J) \cap Or(K))(\mathbb{F})$ for all $K \subseteq J$.

Proof: Let $u \in (U_f^-)^J$, $g \in (G_{fn})_J$, and $K \subseteq J$. If we apply p_1 to

$$ugP(K) = \bigoplus_{\Lambda \in P^+ \cap \overline{F_K}} \pi(ug)^{-1} L(\Lambda)^{(*)}_{\neq \Lambda} \oplus \bigoplus_{\Lambda \in P^+ \setminus \overline{F_K}} L(\Lambda)^{(*)}$$

and use $\overline{F_K} \supseteq \overline{F_J}$ we find

$$p_1(ugP(K)) = \bigoplus_{\Lambda \in P^+ \cap \overline{F_J}} \pi(ug)^{-1} L(\Lambda)^{(*)}_{\neq \Lambda} \oplus \bigoplus_{\Lambda \in P^+ \setminus \overline{F_J}} L(\Lambda)^{(*)} = ugP(J).$$

By Theorem 3.9 it holds $Stab_{G_{fn}}(P(J)) = (P_{fn})_J \supseteq (G_{fn})_J$. Therefore, we get $p_1(ugP(K)) = ugP(J) = uP(J)$.

If we apply p_2 to $ugP(K)$ we get

$$p_2(ugP(K)) = \bigoplus_{\Lambda \in P^+ \cap \overline{F_K}} \left((\pi(ug)^{-1} L(\Lambda)^{(*)}_{\neq \Lambda} \cap L_J(\Lambda)^{(*)}) \oplus R_J(\Lambda)^{(*)} \right) \oplus \bigoplus_{\Lambda \in P^+ \setminus \overline{F_K}} L(\Lambda)^{(*)}.$$

Let $\Lambda \in P^+ \cap \overline{F_K}$. Because of $gv_\Lambda \in L_J(\Lambda)_f$ and Proposition 3.32 it follows

$$\begin{aligned} & (\pi(ug)^{-1} L(\Lambda)^{(*)}_{\neq \Lambda} \cap L_J(\Lambda)^{(*)}) \oplus R_J(\Lambda)^{(*)} = \left\{ \phi \in L_J(\Lambda)^{(*)} \mid \phi(ugv_\Lambda) = 0 \right\} \oplus R_J(\Lambda)^{(*)} \\ (G_{fn})_J & = \left\{ \phi \in L_J(\Lambda)^{(*)} \mid \phi(gv_\Lambda) = 0 \right\} \oplus R_J(\Lambda)^{(*)} = \left\{ \phi \in L(\Lambda)^{(*)} \mid \phi(gv_\Lambda) = 0 \right\} = \pi(g)^{-1} L(\Lambda)^{(*)}_{\neq \Lambda} \end{aligned}$$

Therefore, we get $p_2(ugP(K)) = gP(K)$.

The bijectivity can be concluded from Theorem 2.15, but it is also easy to check it directly: The surjectivity follows by Theorem 3.37 and Theorem 3.40. To show the injectivity let $p(ugP(K)) = p(u'g'P(K'))$ with $u, u' \in (U_f^-)^J$ and $g, g' \in (G_{fn})_J$, $K, K' \subseteq I$. From $uP(J) = u'P(J)$ it follows with Theorem 3.9 and Lemma 3.2:

$$u^{-1}u' \in \text{Stab}_{G_{fn}}(P(J)) \cap (U_f^-)^J = (P_{fn})_J \cap (U_f^-)^J = (U_f^-)_J \cap (U_f^-)^J = \{1\}.$$

With $gP(K) = g'P(K')$ we find $ugP(K) = u'g'P(K')$. □

4 An action of the face monoid on its building

Now we use the results of the last section to obtain an algebraic geometric model of an action of the face monoid \widehat{G} on the building of its unit group, the Kac-Moody group G .

We take as building the set

$$\Omega := \bigcup_{J \subseteq I} G/P_J = \{gP_J \mid g \in G, J \subseteq I\}$$

partially ordered by the reverse inclusion, i.e., for $g, g' \in G$ and $J, J' \subseteq I$,

$$gP_J \leq g'P_{J'} : \iff gP_J \supseteq g'P_{J'}.$$

The Kac-Moody group G acts order preservingly on Ω by multiplication from the left. We denote by

$$\mathcal{A} := \{nP_J \mid n \in N, J \subseteq I\}$$

the standard apartment of Ω . The building Ω is covered by the apartments $g\mathcal{A}$, $g \in G$.

The building Ω is a substructure of the completed building Ω_{fn} :

Proposition 4.1 *We get an order preserving, G -equivariant embedding by*

$$\begin{aligned} j : \quad \Omega &\rightarrow \Omega_{fn} \\ gP_J &\mapsto g(P_{fn})_J \end{aligned}$$

The standard apartment \mathcal{A} is mapped bijectively to the standard apartment \mathcal{A}_{fn} .

Proof: This follows immediately from Theorem 3.1 (a) and Theorem 3.6 (a). □

Recall that for $J \subseteq I$ we set

$$P(J) := \bigoplus_{\Lambda \in P^+ \cap \overline{F_J}} L(\Lambda)_{\neq \Lambda}^{(*)} \oplus \bigoplus_{\Lambda \in P^+ \setminus \overline{F_J}} L(\Lambda)^{(*)} \subseteq CA$$

with $L(\Lambda)_{\neq \Lambda}^{(*)} := \bigoplus_{\lambda \in P(\Lambda) \setminus \{\Lambda\}} L(\Lambda)_{\lambda}^*$, a sum over the empty set defined to be $\{0\}$.

Corollary 4.2 *We get a G -equivariant embedding of the G -set Ω into the G -set $\text{Proj}(CA)(\mathbb{F})$ by*

$$\begin{aligned} \omega : \quad \Omega &\rightarrow \text{Proj}(CA)(\mathbb{F}) \\ gP_J &\mapsto gP(J) \end{aligned}$$

Its image is dense in $\text{Proj}(CA)(\mathbb{F})$. Furthermore, for $gP_J, hP_K \in \Omega$ it holds

$$gP_J \leq hP_K \iff \overline{\{gP(J)\}}^{pts} \cap \omega(\Omega) \subseteq \overline{\{hP(K)\}}^{pts} \cap \omega(\Omega).$$

Proof: The existence of the G -equivariant embedding $\omega : \Omega \rightarrow \text{Proj}(CA)(\mathbb{F})$ follows from Theorem 3.9 and Proposition 4.1.

For $\omega(\Omega)$ to be dense in $\text{Proj}(CA)(\mathbb{F})$ it is sufficient that the G -orbit of $P(\emptyset)$ is dense in $\text{Proj}(CA)$. It holds

$$\begin{aligned} \bigcap_{g \in G} gP(\emptyset) &= \bigcap_{g \in G} \left(\bigoplus_{\Lambda \in P^+} \pi(g)^{-1} L(\Lambda)^{(*)}_{\neq \Lambda} \right) = \bigoplus_{\Lambda \in P^+} \left(\bigcap_{g \in G} \pi(g)^{-1} L(\Lambda)^{(*)}_{\neq \Lambda} \right) \\ &= \bigoplus_{\Lambda \in P^+} \left\{ \phi \in L(\Lambda)^{(*)} \mid \pi(g)\phi \in L(\Lambda)^{(*)}_{\neq \Lambda} \text{ for all } g \in G \right\} \\ &= \bigoplus_{\Lambda \in P^+} \left\{ \phi \in L(\Lambda)^{(*)} \mid \phi(gv_\Lambda) = 0 \text{ for all } g \in G \right\}. \end{aligned}$$

For every $\Lambda \in P^+$ the orbit Gv_Λ spans $L(\Lambda)$, because $L(\Lambda)$ is an irreducible G -module. It follows $\bigcap_{g \in G} gP(\emptyset) = \{0\}$. Therefore, $\overline{GP(\emptyset)} = \mathcal{V}(\{0\}) = \text{Proj}(CA)$.

By Theorem 3.9 and Proposition 4.1 it holds $gP_J \leq hP_K$ if and only if

$$\overline{\{gP(J)\}}^{pts} \subseteq \overline{\{hP(K)\}}^{pts}, \quad (58)$$

from which it follows

$$\overline{\{gP(J)\}}^{pts} \cap \omega(\Omega) \subseteq \overline{\{hP(K)\}}^{pts} \cap \omega(\Omega). \quad (59)$$

Now suppose that (59) holds. Because of

$$gP(J) \in \overline{\{gP(J)\}}^{pts} \cap \omega(\Omega) \subseteq \overline{\{hP(K)\}}^{pts} \cap \omega(\Omega) \subseteq \overline{\{hP(K)\}}^{pts}$$

it follows (58). \square

The face monoid \widehat{G} acts on $\text{Proj}(CA)(\mathbb{F})$. It holds

Theorem 4.3 *The image $\omega(\Omega)$ is a \widehat{G} -invariant subset of $\text{Proj}(CA)(\mathbb{F})$, the action of \widehat{G} on $\omega(\Omega)$ obtained as follows: Let $g_1 e(R(\Theta)) g_2 \in \widehat{G}$ and $hP(J) \in \omega(\Omega)$. Decompose $g_2 h$ in the form*

$$g_2 h = a n_y c$$

with $a \in P_{\Theta \cup \Theta^\perp}^-$, $c \in P_J$, and $n_y \in N$ projecting to $y \in \Theta \cup \Theta^\perp \mathcal{W}^J$. Then it holds

$$(g_1 e(R(\Theta)) g_2) hP(J) = g_1 p_{\Theta}^-(a) P(\Theta \cup J \cup \text{red}(y)). \quad (60)$$

If we identify the building Ω with its image $\omega(\Omega)$, then this action coincides with good action 1 of \widehat{G} on Ω given in Corollary 50 of [M5].

Proof: It remains to show formula (60). By Theorem 3.9 and by Theorem 9 (b) of [M5] it holds

$$(g_1 e(R(\Theta)) g_2) hP(J) = g_1 e(R(\Theta)) a n_y c P(J) = g_1 p_{\Theta}^-(a) e(R(\Theta)) n_y P(J).$$

Therefore, we only have to show

$$e(R(\Theta)) n_y P(J) = P(\Theta \cup J \cup \text{red}(y)). \quad (61)$$

Let $\Lambda \in P^+ \setminus \overline{F_J}$. Trivially, it holds

$$\pi(e(R(\Theta)) n_y)^{-1} P(J)_\Lambda = \pi(e(R(\Theta)) n_y)^{-1} L(\Lambda)^{(*)} = L(\Lambda)^{(*)}. \quad (62)$$

Let $\Lambda \in P^+ \cap \overline{F_J}$. Then it holds

$$\begin{aligned}
\pi(e(R(\Theta))n_y)^{-1}P(J)_\Lambda &= \pi(e(R(\Theta))n_y)^{-1}L(\Lambda)_{\neq \Lambda}^{(*)} \\
&= \left\{ \phi \in L(\Lambda)^{(*)} \mid \pi(e(R(\Theta))n_y)\phi \in L(\Lambda)_{\neq \Lambda}^{(*)} \right\} \\
&= \left\{ \phi \in L(\Lambda)^{(*)} \mid \phi(e(R(\Theta))n_y v_\Lambda) = 0 \right\} \\
&= \begin{cases} \left\{ \phi \in L(\Lambda)^{(*)} \mid \phi(n_y v_\Lambda) = 0 \right\} & \text{if } y\Lambda \in R(\Theta) \\ \left\{ \phi \in L(\Lambda)^{(*)} \mid \phi(0) = 0 \right\} & \text{if } y\Lambda \notin R(\Theta) \end{cases} \\
&= \begin{cases} \left\{ \phi \in L(\Lambda)^{(*)} \mid \phi(n_y v_\Lambda) = 0 \right\} & \text{if } y\Lambda \in R(\Theta) \\ L(\Lambda)^{(*)} & \text{if } y\Lambda \notin R(\Theta) \end{cases}.
\end{aligned}$$

By Theorem 7 of [M5] it holds $R(\Theta) \cap y\overline{F_J} = \overline{F_{\Theta \cup J \cup \text{red}(y)}}$. Note also that y fixes the points of $\overline{F_{\Theta \cup J \cup \text{red}(y)}}$. It follows

$$y\Lambda \in R(\Theta) \iff y\Lambda \in \overline{F_{\Theta \cup J \cup \text{red}(y)}} \iff \Lambda \in \overline{F_{\Theta \cup J \cup \text{red}(y)}}.$$

Furthermore, for $\Lambda \in \overline{F_{\Theta \cup J \cup \text{red}(y)}}$ we get $0 \neq n_y v_\Lambda \in L(\Lambda)_{y\Lambda} = L(\Lambda)_\Lambda$. Therefore, for $\Lambda \in P^+ \cap \overline{F_J}$, we have shown

$$\pi(e(R(\Theta))n_y)^{-1}P(J)_\Lambda = \begin{cases} L(\Lambda)_{\neq \Lambda}^{(*)} & \text{if } \Lambda \in \overline{F_{\Theta \cup J \cup \text{red}(y)}} \\ L(\Lambda)^{(*)} & \text{if } \Lambda \notin \overline{F_{\Theta \cup J \cup \text{red}(y)}} \end{cases}. \quad (63)$$

Since $\overline{F_{\Theta \cup J \cup \text{red}(y)}} \subseteq \overline{F_J}$, from (62) and (63) it follows (61). \square

5 Some open problems

- (a) $\text{Proj}(CA)$ is stratified by schemes for the P^+ -graded Cartan algebra CA . $\text{Proj}(A)$ is also stratified by schemes for a \mathbb{N}_0 -graded algebra A . For which \mathcal{M} -graded algebras A exists a similar stratification of $\text{Proj}(A)$.
- (b) In the literature, there are many constructions of flag schemes, flag varieties, and flag manifolds for Kac-Moody groups. Compare the flag schemes $Or(J)$, $J \subseteq I$, with these constructions. In particular:
 - Compare the scheme $Or(\emptyset)$ with the thick flag scheme of M. Kashiwara in [Kas].
 - Let $\Lambda \in P^+ \cap F_J$, $J \subseteq I$. V. Kac and D. Peterson constructed in [KP1], [KP2] a thin flag variety as follows. They showed that the Kostant cone $\mathcal{V}_\Lambda := G(L(\Lambda)_\Lambda)$ is a closed subvariety of the linear space $L(\Lambda)$, its \mathbb{N}_0 -graded coordinate ring $\mathbb{C}[\mathcal{V}_\Lambda]$ isomorphic to $\bigoplus_{k \in \mathbb{N}_0} L(\Lambda)^{(*)}_k \subseteq CA$. This realizes the thin flag variety G/P_J as a closed subvariety of the projective space $\mathbb{P}(L(\Lambda))$. Compare the scheme $\text{Proj}(\mathbb{C}[\mathcal{V}_\Lambda]) \setminus \{ \text{the irrelevant ideal} \}$ with the scheme $Or(J)$.
- (c) Many structures and maps linked with buildings haven't been considered in the article. Do some of them fit into the algebraic geometric context? Do they make sense for the whole projective spectrum $\text{Proj}(CA)$?
- (d) To define the Cartan algebra CA the restricted duals $L(\Lambda)^{(*)}$, $\Lambda \in P^+$, have been used. Similarly, it is possible to obtain a Cartan algebra CA_f , the restricted duals replaced by the full duals $L(\Lambda)^*$, $\Lambda \in P^+$. Investigate $\text{Proj}(CA_f)$. Does it hold $\text{Proj}(CA_f)(\mathbb{F}) \cong \bigcup_{J \subseteq I} G_{fp}/(P_{fp})_J$ where now the parabolic subgroups $(P_{fp})_J$, $J \subseteq I$, of G_{fp} enter?
- (e) A Cartan algebra CA_M can be obtained for every normal reductive algebraic monoid M . Investigate $\text{Proj}(CA_M)$ and the action of M on $\text{Proj}(CA_M)$.

References

- [Ha] R. Hartshorne, Algebraic geometry, Springer-Verlag, sixth printing, 1993.
- [K1] V. G. Kac, Infinite dimensional Lie algebras, Cambridge Univ. Press, 1990.
- [K2] V. G. Kac, Constructing groups associated to infinite-dimensional Lie algebras, in Infinite Dimensional Groups with Applications, MSRI Publ. **4** (1985), 167-216.
- [KP1] V. G. Kac, D. H. Peterson, Infinite flag varieties and conjugacy theorems, Proc. Natl. Acad. Sci. U.S.A. **80** (1983), 1778-1782.
- [KP2] V. G. Kac, D. H. Peterson, Regular functions on certain infinite-dimensional groups, in: Arithmetic and Geometry, Progr. in Math. **36**, Birkhäuser, Boston, 1983, 141-166.
- [KP3] V. G. Kac, D. H. Peterson, Defining relations on certain infinite-dimensional groups, Proceedings of the Cartan conference, Lyon 1984, Astérisque, 1985, Numero hors serie, 165-208.
- [Kas] M. Kashiwara, The Flag Manifold of Kac-Moody Lie Algebra, Amer. J. of Math. **111** (1989), Supplement Volume: Algebraic Analysis, Geometry and Number Theory, Ed. J.-i. Igusa, John Hopkins Univ., Baltimore, 1989, 161-190.
- [Ku] S. Kumar, Kac-Moody Groups, their Flag Varieties and Representation Theory, Progress in Mathematics **204**, Birkhäuser, 2002.
- [M1] C. Mokler, An analogue of a reductive algebraic monoid whose unit group is a Kac-Moody group, Memoirs of the AMS **823** (2005).
- [M2] C. Mokler, The \mathbb{F} -valued points of the algebra of strongly regular functions of a Kac-Moody group, Transformation Groups **7** (2002), 343-378.
- [M3] C. Mokler, Extending the Bruhat order and the length function from the Weyl group to the Weyl monoid, J. of Algebra **275** (2004), 815-855.
- [M4] C. Mokler, The maximal chains of the extended Bruhat orders on the $\mathcal{W} \times \mathcal{W}$ -orbits of an infinite Renner monoid, Communications in Algebra **35** (2007), 2298-2323.
- [M5] C. Mokler, Actions of the face monoid associated to a Kac-Moody group on its building, Journal of Algebra **321** (2009), 2384-2421.
- [M6] C. Mokler, The full adjoint quotient of a Kac-Moody group, in preparation.
- [Mo,Pi] R. V. Moody, A. Pianzola, Lie Algebras With Triangular Decompositions, Canadian Mathematical Society Series of Monographs and Advanced Texts, John Wiley & Sons, 1995.
- [Ré] Bertrand Rémy, Groupes de Kac-Moody déployés et presque déployés, Astérisque 277, Société Mathématique de France, 2002.
- [Sh] R. Y. Sharp, Steps in Commutative Algebra, Cambridge Univ. Press, 1990.
- [Sl] P. Slodowy, Singularitäten, Kac-Moody-Liealgebren und assoziierte Gruppen, Habilitationsschrift, Bonn 1984.