

# On the existence of Stone-Čech compactification

Giovanni Curi

Dipartimento di Informatica - Università di Verona  
 Strada le Grazie 15 - 37134 Verona, Italy.  
 e-mail: giovanni.curi@univr.it

January 9, 2010.

**Introduction.** In 1937 E. Čech and M.H. Stone, independently, introduced the maximal compactification of a completely regular topological space, thereafter called Stone-Čech compactification [8, 19]. In the introduction of [8] the non-constructive character of this result is so described: “It must be emphasized that  $\beta(S)$  [the Stone-Čech compactification of  $S$ ] may be defined only formally (not constructively) since it exists only in virtue of Zermelo’s theorem”.<sup>1</sup>

By replacing topological spaces with locales, Banaschewski and Mulvey [4, 5, 6], and Johnstone [14] obtained choice-free intuitionistic proofs of Stone-Čech compactification. Although valid in any topos, these localic constructions rely - essentially, as is to be demonstrated - on highly impredicative principles, and thus cannot be considered as constructive in the sense of the main systems for constructive mathematics, such as Martin-Löf’s constructive type theory and Aczel’s constructive set theory.

In [10] I characterized the locales of which the Stone-Čech compactification can be defined in constructive type theory CTT, and in the formal system CZF+uREA+DC, a natural extension of Aczel’s system for constructive set theory CZF by a strengthening of the Regular Extension Axiom REA and the principle of Dependent Choice.

In this paper I show that this characterization continues to hold over the standard system CZF<sup>+</sup>, i.e., CZF plus REA, thus removing in particular any dependency from a choice principle. This will follow by a result of

---

<sup>1</sup>Over ZF, the existence of Stone-Čech compactification has later been proved equivalent to the Prime Ideal Theorem [4].

independent interest, namely the proof that the class of continuous mappings from a compact regular locale  $X$  to a regular a set-presented locale  $Y$  is a set in CZF (even without REA).

By exploiting the consistency of CZF+REA with a general form of Troelstra's *principle of uniformity* [24], I then prove that the existence of Stone-Čech compactification of a non-degenerate Boolean locale is independent of the axioms of CZF (+REA), so that the aforementioned characterization characterizes a proper subcollection of the collection of all locales. The same also holds for several, even impredicative, extensions of CZF+REA, as well as for CTT. This is in strong contrast with what happens in the context of Higher-order Heyting arithmetic HHA - and thus in any topos-theoretic universe: within HHA the constructions in [4, 5, 6, 14] of Stone-Čech compactification can be carried out for every locale.

**1 Constructive Set Theory and the Principle of Uniformity.** The reader is invited to consult [3, 17] for background on Aczel's constructive set theory, CZF, and Martin-Löf's type theory, CTT. Here I shall be working in constructive set theory, but the arguments to be presented may be adapted also to the setting of CTT.

The system CZF is a subsystem of ZF that uses intuitionistic rather than classical logic, has only a restricted form of the Separation Scheme, and does not have the Powerset Axiom. More specifically, CZF is based on first-order intuitionistic logic with equality, has  $\in$  as the only non-logical symbol, and has Extensionality, Pair, Union, Infinity, Set Induction, Restricted Separation (i.e., Separation for bounded formulae), Strong Collection and Subset Collection as non-logical axioms and schemes. Strong Collection is the following scheme:

**Strong Collection** For every set  $a$ , if  $(\forall x \in a)(\exists y) \phi(x, y)$ , then there is a set  $b$  such that  $(\forall x \in a)(\exists y \in b) \phi(x, y)$  and  $(\forall y \in b)(\exists x \in a) \phi(x, y)$ .

A purely logical consequence of Strong Collection, to be often applied in the following (sometimes tacitly), is the usual Replacement Scheme.

**Replacement** For every set  $a$ , if  $(\forall x \in a)(\exists!y) \phi(x, y)$ , then there is a set  $b$  such that  $\forall y(y \in b \leftrightarrow (\exists x \in a) \phi(x, y))$ .

The Subset Collection Scheme is a strengthening of Myhill's Exponentiation Axiom, asserting that the class of functions between any two sets is a set. Subset Collection is often presented in the equivalent form (over the

remaining axioms of CZF) of the Fullness Axiom. For sets  $a, b$ , let  $\mathbf{mv}(b^a)$  be the class of subsets  $r$  of  $a \times b$  such that  $(\forall x \in a)(\exists y \in b) (x, y) \in r$ .

**Fullness** Given sets  $a, b$  there is a subset  $c$  of  $\mathbf{mv}(a^b)$  such that for every  $r \in \mathbf{mv}(a^b)$  there is  $r_0 \in c$  with  $r_0 \subseteq r$ .

This principle may in some cases be used to replace applications of the fully impredicative Powerset Axiom. The system one obtains replacing Subset Collection with the Exponentiation Axiom is usually denoted by  $\text{CZF}_{exp}$ .

The Regular Extension Axiom REA, stating that every set is a subset of a regular set, is often added to the axioms of CZF; the resulting system is denoted by  $\text{CZF}^+$ . REA is needed in order to prove that certain inductively defined classes are sets (see [3] for more information).

Both  $\text{CZF}+\text{REA}$  and  $\text{CTT}$  can consistently be extended with a general form of Troelstra's principle of uniformity (van den Berg & Moerdijk [7], Coquand, cf. [12]). In constructive set theory, this is formulated as the following schema:

**GUP** For every set  $a$ , if  $(\forall x)(\exists y \in a)\phi(x, y)$ , then  $(\exists y \in a)(\forall x)\phi(x, y)$ .

This principle may be seen as resulting from the particular case in which  $a$  is the set  $\omega$  of natural numbers, i.e., from the uniformity principle in its standard form, and the principle that every set is subcountable, also consistent with  $\text{CZF}+\text{REA}$  ([7]; see also [15, 20, 21, 22, 23]). In fact, several extensions of CZF, as  $\text{CZF}+\text{REA}+\text{PA}+\text{Sep}$ , where PA is the Presentation Axiom (implying the Dependent and Countable Choice principles), and Sep is the impredicative full Separation Scheme, have been proved consistent with GUP [7].

As is customary in classical set theory, class notation and terminology can be exploited in this context [3]. Recall also that a set is *finite* in this setting if it (is empty, or) can be finitely enumerated (possibly with repetitions), and that the class  $\text{Pow}_{fin}(S)$  of finite subsets of a set  $S$  is a set in CZF (and  $\text{CZF}_{exp}$ ).

We conclude this introductory section by proving a first important general consequence of the consistency of the systems we are considering with GUP. A (large)  $\vee$ -*semilattice* is a partially ordered class (i.e., a class together with a class relation on it satisfying the usual axioms for a partial order) that has suprema for arbitrary *subsets*. A  $\vee$ -semilattice need not be a (large) *complete lattice*, i.e., need not have also infima of arbitrary subsets. A *class-frame*, or *class-locale*,  $X$  is a  $\vee$ -semilattice that has a top element

$\top$ , binary meets, and that is such that meets distribute over suprema of arbitrary sets of elements of  $X$  [1]. A *class-preframe* is defined in the same way as a class-frame, but suprema have to exist only of directed subsets, and meets are required to distribute only over these suprema. Note that a preframe need not have a smallest element. A partially ordered class is *degenerate* if it consists of a single element. Then, a  $\bigvee$ -semilattice  $L$  is degenerate iff  $L = \{\perp\}$ , with  $\perp = \bigvee \emptyset$ , while a class-preframe  $P$  is degenerate iff  $P = \{\top\}$ .

Lattices of the above kinds arise everywhere in mathematics; the fact that they are carried by sets is often an essential tacit assumption in the theory and applications of these structures, e.g. when classes of ideals are considered. A consequence of the consistency of CZF with the generalized uniformity principle is that, constructively, in no non-trivial case this assumption is legitimate.

**Lemma 1.1 (The Main Lemma)** *No non-degenerate  $\bigvee$ -semilattice, or class-preframe, and hence no non-degenerate class-frame, can be proved to have a set of elements in CZF (+REA+PA+Sep).*

**Proof.** Let  $L$  be a non-degenerate  $\bigvee$ -semilattice, and assume  $L$  is carried by a set. Then the class  $\{x \in L : \emptyset \in y\}$  is a set for every set  $y$ . Therefore,  $(\forall y)(\exists a \in L)a = \bigvee\{x \in L : \emptyset \in y\}$ . In CZF (+REA+PA+Sep)+GUP one then gets  $(\exists a \in L)(\forall y)a = \bigvee\{x \in L : \emptyset \in y\}$ , so that  $L$  must be degenerate, as follows by first taking  $y = \emptyset$ , then  $y = \{\emptyset\}$ . So  $L$  is not a set in CZF (+REA+PA+Sep)+GUP, and thus cannot be proved to be a set in CZF (+REA+PA+Sep).

The proof for  $P$  a non-degenerate preframe (and in fact for any non-degenerate partially ordered class with a greatest element and joins of directed subsets) is similar, but one considers instead the set  $\{x \in P : \emptyset \in y\} \cup \{b\}$ , for  $b \in P$ . It is an easy exercise in intuitionistic logic to check that, for every  $y$ , this set is directed, so that it has a join in  $P$  for every  $y$ . Reasoning as in the previous case, one gets that  $b = \top$ . As this holds for every  $b \in P$ , it follows that  $P$  is degenerate, against the hypothesis.

Clearly, given any set  $X$ , its powerclass  $\text{Pow}(X)$  with intersection as meet and union of arbitrary set-indexed families of subsets as join is a frame. Therefore, no non-trivial instance of the Powerset Axiom is constructively derivable.

**Corollary 1.2** *For no non-empty set  $X$  the powerclass  $\text{Pow}(X)$  can be proved to form a set in CZF(+REA+PA+Sep).*

**2 Small homsets of continuous maps in CZF.** Locales, or formal spaces, provide a suitable substitute to the concept of topological space in choice-free and/or intuitionistic settings [14, 13]. In CZF, or CTT, due to the absence of powersets, the concept of locale needs to be formulated with special care [1, 9] (recall also the Main Lemma). In CZF, a class-locale is said to be *set-generated* by a subclass  $B$  if:

- i.  $B$  is a set,
- ii. the class  $\{b \in B : b \leq x\}$  is a set and  $x = \bigvee\{b \in B : b \leq x\}$ , for all  $x \in X$ .

In a fully impredicative context as intuitionistic set theory IZF, set-generated class-locales and ordinary locales come to the same thing. Here a set-generated class-locale  $(X, B)$  will simply be referred to as a *locale*  $X$ , omitting the explicit mention of the base  $B$ . A *continuous map* of locales  $f : X \rightarrow Y$  is a class function  $f^- : B_Y \rightarrow X$  (note the reverse direction) satisfying:

- 1.  $\bigvee_{a \in B_Y} f^-(a) = \top$ ,
- 2.  $f^-(a) \wedge f^-(b) = \bigvee\{f^-(c) : c \in B_Y, c \leq a, c \leq b\}$ , for all  $a, b \in B_Y$ ,
- 3.  $f^-(a) \leq \bigvee_{b \in U} f^-(b)$ , for all  $a \in B_Y, U \in \mathbf{Pow}(B_Y)$  with  $a \leq \bigvee U$ .

By Replacement, one may assume that any such function is a set. The (in general proper) class of these maps is denoted by  $Hom(X, Y)$ . Observe that  $Hom(X, Y)$  is in a one-to-one correspondence with the collection of frame homomorphisms from  $Y$  to  $X$  (i.e., class functions from  $Y$  to  $X$  preserving the frame structure).

A locale  $X$  is *compact* iff every covering of  $\top$  by basic elements (i.e., every  $U \in \mathbf{Pow}(B)$  such that  $\top = \bigvee U$ ) has a finite subcover.  $X$  is *regular* if, for all  $a \in B$ ,  $a = \bigvee\{b \in B : b \prec a\}$ , where, for  $x, y \in X$ ,  $y \prec x \iff \top = x \vee y^*$ , with  $y^* = \bigvee\{c \in B : c \wedge y = \perp\}$  the *pseudocomplement* of  $y$ .

Let  $rc : B \rightarrow \mathbf{Pow}(B)$  be a function with the property that for all  $b \in rc(a)$  a scale exists from  $b$  to  $a$ , i.e., a family  $\{c_p\}_{p \in \mathbb{I}}$  of elements of  $X$ , indexed on the rational unit interval  $\mathbb{I}$ , satisfying:  $c_0 = b, c_1 = a$  and, for  $p < q$ ,  $c_p \prec c_q$ .  $X$  is *completely regular* if a function  $rc : B \rightarrow \mathbf{Pow}(B)$  of this kind is given with the property that for all  $a \in B$ ,  $a = \bigvee rc(a)$ . Over CZF (or HHA) plus the principle of dependent choice, a compact regular locale is completely regular. In ZFC, compact (completely) regular locales and compact Hausdorff spaces define equivalent categories [14].

A locale  $X$  is *set-presented* [1, 9] iff there is a function  $C : B \rightarrow \text{Pow}(\text{Pow}(B))$ , with  $\text{Pow}(\text{Pow}(B))$  the class of subsets of the class  $\text{Pow}(B)$ , such that

$$a \leq \bigvee U \iff \exists V \in C(a) V \subseteq U.$$

Observe that this implies  $a \leq \bigvee V$ , for all  $a \in B, V \in C(a)$ .

For  $x, y \in X$ , one says that  $y$  is *way-below*  $x$  iff for all  $U \in \text{Pow}(B)$ ,  $x \leq \bigvee U$  implies  $y \leq \bigvee u$  for  $u$  a finite subset of  $U$ . Due to the unbounded quantification over  $\text{Pow}(B)$ , the way-below relation is a class in CZF even when restricted to basic elements. However, if  $X$  is set-presented, for  $a, b \in B$ , one has  $b$  way-below  $a$  iff  $b \ll a \equiv (\forall V \in C(a))(\exists v \in \text{Pow}_{fin}(V))b \leq \bigvee v$ . Since in CZF (or  $\text{CZF}_{exp}$ ), the class  $\text{Pow}_{fin}(A)$  of finite subsets of a given set  $A$  is a set, the last formula is easily seen to be equivalent to a bounded formula, and thus defines a subset of  $B \times B$ . A *locally compact* locale is a set-presented locale  $X$  satisfying  $a = \bigvee \{b \in B : b \ll a\}$ , for all  $a \in B$ . A key fact for what follows is that, in a locally compact locale, for  $x, y \in X$ ,  $y$  way-below  $x$  implies  $y$  way-below  $\bigvee u$  and  $\bigvee u$  way-below  $x$ , for  $u$  a finite subset of  $B$  [10].

Finally, every compact regular locale is locally compact, with  $\ll = \prec$ , and is therefore set-presented in CZF [14, 1].

The reader may consult [14] for the definition of the locale of the real numbers  $R$  and of its sublocale  $[0, 1]$ , the localic real unit interval. Both locales are completely regular,  $R$  is locally compact,  $[0, 1]$  is compact (hence both are set-presented in CZF).

The following result has an independent interest, in particular in connection with the theory of (rings of) continuous real-valued functions.

**Theorem 2.1 (CZF)** *If  $X$  is a compact regular locale and  $Y$  is a regular and set-presented locale, the class  $\text{Hom}(X, Y)$  is a set.*

**Proof.** Note first that, for  $X$  any locale and  $Y$  a set-presented locale, a mapping  $f^- : B_Y \rightarrow X$  satisfies conditions from 1 to 3 on continuous mappings iff it satisfies 1, 2 and

$$3'. f^-(a) \leq \bigvee_{b \in V} f^-(b), \text{ for all } a \in B_Y, V \in C(a).$$

Given any continuous map  $f \in \text{Hom}(X, Y)$ , the associated morphism  $f^- : B_Y \rightarrow X$  is easily seen to preserve the well-inside relation  $\prec$ , in the sense that  $f^-(b) \prec f^-(a)$ , for all  $a, b \in B_Y$  with  $b \prec a$ . As we assumed here  $Y$  to be regular, we also have  $f^-(a) = \bigvee_{b \prec a} f^-(b)$ . Since in a compact regular locale the way-below relation coincides with the well-inside relation,

for all  $a, b \in B_Y$  with  $b \prec a$ , there is a finite subset  $u$  of  $B_X$  such that  $f^-(b) \ll \vee u \ll f^-(a)$ . In particular,  $f^-(b) \leq \vee u \leq f^-(a)$ .

Now let  $W$  be the set  $\{(a, b) \in B_Y \times B_Y : b \prec a\}$ , and consider the class  $mv(\mathbf{Pow}_{fin}(B_X)^W)$  of multivalued functions from  $W$  to the set of finite subsets of  $B_X$ . By Fullness there is a set

$$K \subseteq mv(\mathbf{Pow}_{fin}(B_X)^W)$$

such that for all  $R \in mv(\mathbf{Pow}_{fin}(B_X)^W)$  there is  $R^0 \in K$  with  $R^0 \subseteq R$ . There is then a class function from  $K$  to the class  $X^{B_Y}$  of mappings from  $B_Y$  to  $X$ , defined by letting, for  $F \in K$  and  $a \in B_Y$ ,  $f_F^-(a) = \bigvee(\bigcup_{b \prec a} \{\vee u : F((a, b), u)\})$ . By Replacement, the range  $K' \subseteq X^{B_Y}$  of this function is a set. We show that  $Hom(X, Y)$  is a subset of this set. Indeed, let  $D$  be the subclass of the set  $K'$  defined by  $f^- \in D$  iff  $f^-$  satisfies conditions 1, 2 and 3' on continuous mappings. As  $f^-$  is restricted to belong to the set  $K'$ , using Replacement these conditions can equivalently be expressed by a bounded formula, so that by Restricted Separation  $D$  is a set.

Clearly,  $D \subseteq Hom(X, Y)$ . Conversely, given  $f \in Hom(X, Y)$ , let the set  $R_f \subseteq W \times \mathbf{Pow}_{fin}(B_X)$  be defined by:

$$R_f((a, b), u) \iff f^-(b) \leq \vee u \leq f^-(a).$$

We have shown that for all  $(a, b) \in W$  there is  $u \in \mathbf{Pow}_{fin}(B_X)$  such that  $R_f((a, b), u)$ , so that  $R_f \in mv(\mathbf{Pow}_{fin}(B_X)^W)$ . Then there is  $R_f^0 \in K \subseteq mv(\mathbf{Pow}_{fin}(B_X)^W)$  such that  $R_f^0 \subseteq R_f$ . Since it also holds  $(\forall (a, b) \in W)(\exists u \in \mathbf{Pow}_{fin}(B_X))R_f^0((a, b), u)$ , by  $f^-(a) = \bigvee_{b \prec a} f^-(b)$ , one obtains

$$f^-(a) = \bigvee(\bigcup_{b \prec a} \{\vee u : R_f^0((a, b), u)\}),$$

for every  $a \in B_Y$ . Thus,  $f^- = f_{R_f^0}^- \in K'$ , and since  $f^-$  satisfies conditions 1, 2 and 3',  $f^- \in D$ .

Recall that compact (completely) regular locales have the role in intuitionistic settings that compact Hausdorff spaces play in ordinary classical topology [14]. Let  $\mathbf{KRLoc}$  (resp.  $\mathbf{KCRLoc}$ ) denote the full subcategory of the category  $\mathbf{Loc}$  of locales whose objects are the compact regular (resp. compact completely regular) locales. By the above theorem we directly have:

**Corollary 2.2 (CZF)**  *$\mathbf{KRLoc}$  and (hence)  $\mathbf{KCRLoc}$  are locally small categories.*

**Corollary 2.3 (CZF)** *If  $X$  is a compact regular locale, then the classes  $\text{Hom}(X, R)$  and  $\text{Hom}(X, [0, 1])$  are sets.*

**Remarks.** *i.* Note that the last corollary does not hold in  $\text{CZF}_{exp}$ : for  $X = \text{Pow}(\{1\})$ ,  $\text{Hom}(X, R)$  is isomorphic with the class of Dedekind reals which form a proper class in  $\text{CZF}_{exp}$  [16]. However, Corollary 2.3 can be proved in  $\text{CZF}_{exp}$  plus the principle of countable choice  $\text{AC}_\omega$ .

*ii.* In [10] I proved that in stronger systems, as  $\text{CZF} + \text{uREA} + \text{DC}$ , or  $\text{CTT}$ , the class  $\text{Hom}(X, Y)$  is a set whenever  $X$  is locally compact and  $Y$  is set-presented and regular. Using the type-theoretic axiom of choice and regular universes, Palmgren [18] generalized this result in  $\text{CTT}$  by weakening  $X$  locally compact to  $X$  set-presented.

*iii.* Over  $\text{CZF} + \text{REA}$ , the concept of set-presented locale and of *inductively generated* formal space [9] are equivalent, cf. [1]. Although this is no more the case over  $\text{CZF}$ , one may prove with essentially the same argument that Theorem 2.1 also holds if one replaces (locales with formal spaces and) ‘set-presented’ with ‘inductively generated’.

*iv.* Finally, we note that the proof of Theorem 2.1 can easily be seen to apply in the more general case where  $X$  is any locally compact locale and  $\text{Hom}(X, Y)$  is replaced by the class of mapping  $f$  in  $\text{Hom}(X, Y)$  such that  $f^-$  sends the well-inside relation on a base of  $Y$  in the way-below relation (i.e.,  $f^-(b) \ll f^-(a)$  whenever  $b \prec a$ ,  $a, b \in B_Y$ ). A further generalization of a more formalistic nature is implicit in the proof of Theorem 2.1, but will not be spelt out here.

**3 Existence of Stone-Čech compactification.** The (generalized) Stone-Čech compactification of a space or locale  $X$  is its compact completely regular reflection, i.e., it is a continuous map

$$\eta : X \rightarrow \beta X,$$

with  $\beta X$  compact and completely regular, which satisfies the following universal property: for every continuous map  $f : X \rightarrow Y$  to a compact completely regular codomain  $Y$ , a unique map  $f^\beta : \beta X \rightarrow Y$  exists such that  $f^\beta \circ \eta = f$ . Moreover,  $\eta$  is a (dense) embedding precisely when  $X$  is completely regular.

In [10] one finds the proof in  $\text{CTT}$  and  $\text{CZF} + \text{REA}$  that, on the assumption that for every compact completely regular locale  $Y$  the class  $\text{Hom}(Y, [0, 1])$  is a set, if  $X$  is such that  $\text{Hom}(X, [0, 1])$  is a set, the Stone-Čech compactification of  $X$  exists [10, Corollary 6.2] (see also [11]).

By Theorem 2.1 we may then conclude that if  $\text{Hom}(X, [0, 1])$  is a set then  $\beta X$  exists in  $\text{CZF} + \text{REA}$ , in particular without any intervention of a choice principle. As, moreover, the universal property of  $\beta$  directly yields a bijection  $\text{Hom}(X, [0, 1]) \cong \text{Hom}(\beta X, [0, 1])$ , and since the latter is a set by Theorem 2.1 when  $\beta X$  exists, in summary we have:

**Theorem 3.1 (CZF+REA)** *The Stone-Čech compactification  $\beta X$  of a locale  $X$  exists if, and only if,  $\text{Hom}(X, [0, 1])$  is a set.*

The universal property of  $\beta$  yields more generally a bijection  $\text{Hom}(X, Y) \cong \text{Hom}(\beta X, Y)$ , for every compact completely regular  $Y$ , and, again by Theorem 2.1, one has that  $\text{Hom}(\beta X, Y)$  is a set. Thus:

**Corollary 3.2 (CZF+REA)** *Given any locale  $X$ , if  $\beta X$  exists,  $\text{Hom}(X, Y)$  is a set for every compact completely regular locale  $Y$ . Therefore, for every locale  $X$ ,  $\text{Hom}(X, Y)$  is a set for every compact completely regular locale  $Y$  if and only if  $\text{Hom}(X, [0, 1])$  is a set.*

In [10], in the context of  $\text{CZF} + \text{uREA} + \text{DC}$ , these results similarly followed by the results recalled in Remark *ii.* of the previous section.

We now turn to the proof that there are locales  $X$  of which  $\beta X$  does not exist constructively. A well-known classical identification is that of the subsets of a set  $S$  with the mappings of  $S$  in the two-element set  $\{0, 1\}$ . We may regard this identification, that fails intuitionistically, as a special case of a bijection that exists between the frame of opens  $\Omega(X)$  of a Boolean space  $X$  (i.e., a space  $X$  such that  $\Omega(X)$  is Boolean), and the continuous functions from  $X$  to  $\{0, 1\}$ . The next lemma shows that this classically more general fact does constructively carry over to the localic setting. Recall that a locale is *Boolean* if it is a (complete) Boolean algebra.

**Lemma 3.3 (CZF<sub>exp</sub>)** *The compact completely regular locale of opens of the discrete space  $\{0, 1\}$  classifies the opens of any Boolean locale  $X$ . I.e., a bijection exists between the class of elements of  $X$  and  $\text{Hom}(X, \text{Pow}(\{0, 1\}))$ .*

**Proof.** To  $a \in X$  one associates the map  $f_a^- : \{\{0\}, \{1\}\} \rightarrow X$ , defined by  $f_a^-(\{0\}) = a$ ,  $f_a^-(\{1\}) = a^*$ . We leave to the reader to check that this defines a continuous map. Conversely,  $f : X \rightarrow \text{Pow}(\{0, 1\})$  defines the open  $a_f = f^-(\{0\})$ .

Note that  $\text{Pow}(\{0, 1\})$  is not Boolean intuitionistically. To prove that there are locales of which Stone-Čech compactification does not exist, we shall

need the result that the class of continuous mappings from a compact completely regular locale to  $\mathbf{Pow}(\{0, 1\})$  is a set. In CZF, this follows by Theorem 2.1, as  $\mathbf{Pow}(\{0, 1\})$  is set-presented and regular. However, this also holds in  $\mathbf{CZF}_{exp}$ .

**Lemma 3.4 (CZF<sub>exp</sub>)** *Let  $X$  be any compact locale. Then the class  $\mathit{Hom}(X, \mathbf{Pow}(\{0, 1\}))$  is a set.*

**Proof.** Note first that, since  $\mathbf{Pow}(\{0, 1\})$  is set-presented,  $f \in \mathit{Hom}(X, \mathbf{Pow}(\{0, 1\}))$  iff  $f^- : \{\{0\}, \{1\}\} \rightarrow X$  satisfies conditions 1, 2 and 3' on continuous mappings (cf. the proof of Theorem 2.1). By Replacement, the class  $\bar{B}_X = \mathit{range}(\vee), \vee : \mathbf{Pow}_{fin}(B_X) \rightarrow X$ , is a set, so that by Exponentiation also the class  $\bar{B}_X^{\{\{0\}, \{1\}\}}$  is a set. We show that  $\mathit{Hom}(X, \mathbf{Pow}(\{0, 1\}))$  coincides with the subclass  $D$  of  $\bar{B}_X^{\{\{0\}, \{1\}\}}$  given by the maps in  $\bar{B}_X^{\{\{0\}, \{1\}\}}$  satisfying conditions 1, 2 and 3'. As  $\bar{B}_X^{\{\{0\}, \{1\}\}}$  is a set, exploiting the Replacement Scheme these conditions can equivalently be expressed by a bounded formula, so that  $D = \mathit{Hom}(X, \mathbf{Pow}(\{0, 1\}))$  is a set by Restricted Separation.

Let then  $f \in \mathit{Hom}(X, \mathbf{Pow}(\{0, 1\}))$  be a continuous map. One must have  $\top = f^-(\{0\}) \vee f^-(\{1\})$ , and  $\perp = f^-(\{0\}) \wedge f^-(\{1\})$ . As  $X$  is compact, there is a finite subset  $v$  of  $\{b \in B_X : b \leq f^-(\{0\})\} \cup \{b \in B_X : b \leq f^-(\{1\})\}$  such that  $\top = \vee v$ . There are thus finite subsets  $v_0, v_1$  such that  $v_0 \cup v_1 = v$ ,  $v_0 \subseteq \{b \in B_X : b \leq f^-(\{0\})\}$ , and  $v_1 \subseteq \{b \in B_X : b \leq f^-(\{1\})\}$ . Hence,  $\top = \vee(v_0 \cup v_1)$ , and  $\perp = (\vee v_0) \wedge (\vee v_1)$ . This gives  $f^-(\{0\}) = \vee v_0$  and  $f^-(\{1\}) = \vee v_1$ . Thus,  $f^-$  is in fact a map from  $\{\{0\}, \{1\}\}$  to  $\bar{B}_X$  satisfying conditions 1, 2 and 3', i.e.,  $f^- \in D$ . The converse inclusion is trivial.

These two lemmas, together with Lemma 1.1, give us:

**Theorem 3.5** *The Stone-Ćech compactification of a non-degenerate Boolean locale  $X$  cannot be defined in CZF (+REA+PA+Sep). Moreover,  $\mathit{Hom}(X, \mathbf{Pow}(\{0, 1\}))$ , and  $\mathit{Hom}(X, [0, 1])$ ,  $\mathit{Hom}(X, \mathbb{R})$ , are proper classes in this setting.*

**Proof.** By Lemma 3.4 (or Theorem 2.1), if  $\beta X$  exists,  $\mathit{Hom}(\beta X, \mathbf{Pow}(\{0, 1\}))$  is a set in CZF (and a fortiori in every stronger system). Moreover, by the universal property of  $\beta$ ,  $\mathit{Hom}(X, \mathbf{Pow}(\{0, 1\})) \cong \mathit{Hom}(\beta X, \mathbf{Pow}(\{0, 1\}))$ . Thus  $\mathit{Hom}(X, \mathbf{Pow}(\{0, 1\}))$  is a set, so that, by Lemma 3.3,  $X$  is a set in  $\mathbf{CZF}_{exp}$ . However, by Lemma 1.1, no non-degenerate locale can be proved to have a set of elements in CZF (+REA+PA+Sep).

Boolean locales abound in nature also constructively: given any (non-degenerate) locale  $X$ , the set-generated class-frame of its regular elements [13] is a (non-degenerate) Boolean locale. Note also that a Boolean locale is (completely) regular.

**Remarks.** *i.* As, e.g. in IZF, which is obtained from CZF by replacing Restricted Separation with Separation and adding Powerset, Stone-Čech compactification of every locale can be defined, Theorem 3.5 is an independence result.

*ii.* In the statements of the Main Lemma and of Theorem 3.5 one may of course replace CZF+REA+PA+Sep with any extension of CZF (or even of CZF<sub>exp</sub>) that is compatible with GUP. Similarly, the above results can be shown to hold (mutatis mutandis) also with respect to every extension of CTT compatible with (the type-theoretic version of) GUP.

*iii.* The given proof of Theorem 3.5 is entirely self-contained (it does not depend on results in [10]); a corresponding result is analogously seen to hold also for the compact zero-dimensional reflection of a Boolean locale, as a zero-dimensional locale is regular, and as  $\text{Pow}(\{0, 1\})$  is (compact and) zero-dimensional.

*iv.* By contrast with Theorem 3.5, the ‘approximation’ to Stone-Čech compactification introduced in [10] exists for every locale  $L$  (and every given set-indexed family of continuous maps of the appropriate type, see [10]).

**4 Conclusion.** If one agrees in considering a necessary condition in order for an argument to be defined constructive that it may be formulated within an extension of CZF or CTT compatible with the form of the uniformity principle we are considering, Theorem 3.5 is read as saying that the Stone-Čech compactification of a non-degenerate Boolean locale  $X$  does not exist constructively. This goes very much against what holds in the topos-theoretic context: in any topos, the Stone-Čech compactification of a Boolean locale  $X$  is simply given by the lattice of ideals on  $X$ .

Note that the given necessary condition for constructivity is by no means sufficient: the theory CZF+Sep, where Sep is impredicative unbounded separation, has the same proof-theoretic strength of second-order Heyting arithmetic [15], and is however consistent with the generalized uniformity principle. We find it remarkable, and somewhat surprising, that, due to this fact, Lemma 1.1 and Theorem 3.5 also hold with respect to this theory.

## References

- [1] P. Aczel, “Aspects of general topology in constructive set theory”. *Ann. Pure Appl. Logic* 137 (2006), 1-3, 3–29.
- [2] P. Aczel, G. Curi, “On the  $T_1$  axiom and other separation properties in constructive topology”. *Ann. Pure Appl. Logic*, to appear. doi:10.1016/j.apal.2009.03.005
- [3] P. Aczel, M. Rathjen, “Notes on Constructive Set Theory”, Mittag-Leffler Technical Report No.40, 2000/2001.
- [4] B. Banaschewski, C.J. Mulvey “Stone-Čech compactification of locales I”. *Houston J. Math.* 6 (1980), 301-312.
- [5] B. Banaschewski, C. J. Mulvey, “Stone-Čech compactification of locales. II”. *J. Pure Appl. Algebra* 33 (1984), 107–122.
- [6] B. Banaschewski, C. J. Mulvey, “Stone-Čech compactification of locales. III”. *J. Pure Appl. Algebra* 185 (2003), 25–33.
- [7] B. van den Berg, I. Moerdijk, “Aspects of predicative algebraic set theory II: realizability”. *Theor. Comp. Science*. To appear. [Available from: <http://arxiv.org/abs/0801.2305>].
- [8] E. Čech, “On bicomact spaces”. *Annals of Mathematics* (2) 38 (1937), 823-844.
- [9] T. Coquand, G. Sambin, J. Smith, S. Valentini, “Inductively generated formal topologies”. *Ann. Pure Appl. Logic* 124, 1-3 (2003), 71–106.
- [10] G. Curi, “Exact approximations to Stone-Čech compactification”. *Ann. Pure Appl. Logic*, 146, 2-3 (2007), 103-123.
- [11] G. Curi, “Remarks on the Stone-Čech and Alexandroff compactifications of locales”. *J. Pure Appl. Algebra* 212, 5, (2008), 1134-1144.
- [12] G. Curi, “On some peculiar aspects of the constructive theory of point-free spaces”. *MLQ*, to appear.
- [13] M. Fourman, D.S. Scott, “Sheaves and logic”. In *Applications of sheaves*, M. Fourman et al. eds., Springer *LNM* 753, Springer-Verlag, 1979, 302-401.
- [14] P. T. Johnstone, *Stone Spaces*, Cambridge University Press, 1982.
- [15] R.S. Lubarsky, “CZF and Second Order Arithmetic”. *Ann. Pure Appl. Logic* 141 1-2 (2006), 29-34.
- [16] R. Lubarsky, M. Rathjen, “On the Constructive Dedekind Reals”. *Log. Anal.* 1, 2 (2008), 131-152.
- [17] P. Martin-Löf, *Intuitionistic Type Theory*. Notes by G. Sambin. Studies in Proof Theory. Bibliopolis, Napoli (1984).

- [18] E. Palmgren, “Predicativity problems in point-free topology”. In: V. Stoltenberg-Hansen et al. eds. Proceedings of the Annual European Summer Meeting of the Association for Symbolic Logic, held in Helsinki, Finland, August 14-20, 2003, *Lecture Notes in Logic* 24, ASL, AK Peters Ltd, 2006.
- [19] M.H. Stone, “Applications of the theory of Boolean rings to general topology”. *Trans. Amer. Math. Soc.* 41 (1937), 375-481.
- [20] M. Rathjen, “The strength of some Martin–Löf type theories”. *Arch. Math. Logic* 33 (1994), 347-385.
- [21] M. Rathjen, “Choice principles in constructive and classical set theories”. In: Z. Chatzidakis, P. Koepke, W. Pohlers (eds.): Logic Colloquium 2002, *Lecture Notes in Logic* 27 (A.K. Peters, 2006) 299-326.
- [22] M. Rathjen, “Realizability for constructive Zermelo-Fraenkel set theory”. In: J. Väänänen, V. Stoltenberg-Hansen (eds.): Logic Colloquium 2003. *Lecture Notes in Logic* 24 (A.K. Peters, 2006) 282-314.
- [23] T. Streicher, Realizability models for  $\text{CZF} + \neg \text{Pow}$ , unpublished note.
- [24] A. Troelstra, D. van Dalen, *Constructivism in mathematics, an introduction*. Volume I. Studies in logic and the foundation of mathematics, vol. 121, North-Holland.