Theory of cosmic ray and γ -ray production in the supernova remnant RX J0852.0-4622

E.G. Berezhko¹, G. Pühlhofer^{2,★}, and H.J. Völk³

- Yu.G. Shafer Institute of Cosmophysical Research and Aeronomy, 31 Lenin Ave., 677980 Yakutsk, Russia e-mail: berezhko@ikfia.ysn.ru
- ² Landessternwarte, Königstuhl, D-69117 Heidelberg, Germany e-mail: Gerd.Puehlhofer@lsw.uni-heidelberg.de
- Max-Planck-Institut für Kernphysik, Postfach 103980, D-69029 Heidelberg, Germany e-mail: Heinrich.Voelk@mpi-hd.mpg.de

Received month day, year; accepted month day, year

ABSTRACT

Aims. The properties of the Galactic supernova remnant (SNR) RX J0852.0-4622 are theoretically analysed. Methods. An explicitly time-dependent, nonlinear kinetic model of cosmic ray (CR) acceleration in SNRs is used to describe the properties of SNR RX J0852.0-4622, the accelerated CRs and the nonthermal emission. The source is assumed to be at a distance of ≈ 1 kpc in the wind bubble of a massive progenitor star. An estimate of the thermal X-ray flux in such a configuration is given. Results. We find that the overall synchrotron spectrum of RX J0852.0-4622 as well as the filamentary structures in hard X-rays lead to an amplified magnetic field $B > 100 \mu G$ in the SNR interior. This implies that the leptonic very high energy (VHE) γ -ray emission is suppressed, and that the VHE γ -rays are hadronically dominated. The energy spectrum of protons produced over the life-time of the remnant until now may well reach "knee" energies. The derived γ -ray morphology is consistent with the H.E.S.S. measurements. The amount of energy in energetic particles corresponds to about 35% of the hydrodynamic explosion energy. A remaining uncertainty concerns the thermal X-ray flux at 1 keV. A rough estimate, possibly not quite appropriate for the assumed wind bubble configuration, results in it being larger than the nonthermal flux at this energy.

Conclusions. It is concluded that this SNR expanding into the wind bubble of a massive star in a dense gas environment can be a hadronic γ -ray source that is consistent with all existing multi-wavelength constraints, except possibly the thermal X-ray emission.

Key words. (ISM:)cosmic rays – +acceleration of particles – shock waves – supernovae individual (SNR RX J0852.0-4622) – radiation mechanisms:non-thermal – gamma-rays:theory

1. Introduction

RX J0852.0-4622 (also known as G266.2-1.9) is a shell-type supernova remnant (SNR) with a diameter of 2°, located in the Galactic plane. The SNR was originally discovered in X-rays with the ROSAT satellite (Aschenbach, 1998; Aschenbach et al., 1999). In projection along the line of sight, RX J0852.0-4622 lies entirely within the still much larger Vela SNR and is only visible in hard X-rays, where the thermal radiation from the Vela SNR is no longer dominant. While nonthermal emission from the shell of RX J0852.0-4622 has been confirmed by several X-ray observatories (Slane et al., 2001; Bamba et al., 2005; Iyudin et al., 2005), a clear detection of the thermal X-ray emission from the shell or the interior was not yet possible because of confusion with the Vela SNR. This implies that the thermal emission is very weak¹.

The radio emission of RX J0852.0-4622 is also weak. In fact the SNR radio shell was only identified (Combi et al., 1999; Duncan & Green, 2000; Stupar et al., 2005) after its discovery in X-rays. Before that only a bright radio spot was known as "Vela Z" (Milne, 1968), which was usually identified with the

Send offprint requests to: H.J. Völk

Vela SNR. In addition, significant Galactic background variation over the size of the remnant cannot be excluded. The radio spectrum of RX J0852.0-4622 is therefore not well determined. Only for the northeastern rim a spectral index can be derived with moderate accuracy (Duncan & Green, 2000).

The shell of RX J0852.0-4622 was also detected in very high energy (VHE) γ -rays by the H.E.S.S. collaboration (Aharonian et al., 2005, 2007a), with a γ -ray flux at 1 TeV as large as that from the Crab Nebula. Emission from the northwestern rim had been detected already before by the CANGAROO experiment (Katagiri et al., 2005). The CANGAROO data have been revised since then (Enomoto et al., 2006).

RX J0852.0-4622 is the second SNR after RX J1713.7-3946 (e.g. Aharonian et al., 2007b), where morphologically a SNR shell was unambiguously identified to accelerate particles to TeV energies and beyond. As a third and fourth source of this character the objects RC W86 (Hoppe & Lemoine-Goumard, 2008; Aharonian et al., 2008) and SN 1006 (Naumann-Godo et al., 2009) have recently been detected. A few other TeV sources have been detected in the H.E.S.S. Galactic plane survey that are spatially coincident with radio shell-type SNRs. So far, however, the data do not permit to unambiguously identify the γ -ray emission with the SNR shells, e.g. using morphological arguments. In several cases, the TeV emission might also be associated with X-ray emitting pulsar wind nebula candidates.

^{*} now at Institut für Astronomie und Astrophysik, Universität Tübingen, Sand 1, 72076 Tübingen, Germany

¹ Very recently Uchiyama (2008) has indicated that the X-ray emission from RX J0852.0-4622 might show a thermal component.

Despite the scarcity of precise information from radio data and despite uncertainties about key astrophysical parameters, the prominence of RX J0852.0-4622 has therefore prompted us to model the acceleration of both electrons and protons together with their nonthermal emission in detail, applying explicitly time-dependent nonlinear kinetic theory. The theory couples particle acceleration on a kinetic level with the gas dynamical evolution of the system (Berezhko et al., 1996; Berezhko & Völk, 1997, 2000).

Compared to the other SNRs that were successfully described within the framework of this theory (e.g. Berezhko, 2005, 2008; Berezhko & Völk, 2006), the present uncertainties regarding RX J0852.0-4622 are quite large. Such important astronomical parameters as the distance and age are not well known. It is in fact not even clear, whether the source is in front or behind the Vela SNR. The latter object is generally considered to lie at a distance $d=250\pm30$ pc (Cha et al., 1999)². A position behind the Vela SNR could correspond to a solution with d=1 kpc (Slane et al., 2001), whereas another solution could correspond to the earlier distance estimate of d=200 pc (Aschenbach, 1998).

This prompted us originally to construct indeed two quite different source scenarios, one in front, and the other behind the Vela SNR. They were to correspond to earlier distance estimates: a "nearby" solution with d=200 pc (Aschenbach, 1998) in front of, and a "distant" solution with d=1 kpc (Slane et al., 2001) behind the Vela SNR. However, no "nearby" solution could be found that fulfilled all the observational constraints. Therefore we have abandoned the possibility of a small distance to RX J0852.0-4622 and will assume for the sequel that the source is at a distance d=1 kpc.

In agreement with Slane et al. (2001) but with independent additional arguments we shall assume that the observed non-thermal emission of RX J0852.0-4622 indicates that the SNR emerged from a core collapse explosion into the wind bubble of a massive progenitor star in a dense gas environment, possibly a molecular cloud. In this case the major part of the swept-up volume has originally been occupied by the highly diluted bubble gas that also has a minimal thermal emissivity. At the current epoch, however, we assume that the SNR shock already propagates into the dense shell of ambient *interstellar* medium (ISM) which has originally been compressed by the stellar wind. This also implies that the magnetic field upstream of the SNR shock is of interstellar origin.

We note here that this solution has a similar character as the solution suggested earlier for the object RX J1713.7-3946 (Berezhko & Völk, 2006). Indeed, in this sense the two SNRs RX J0852.0-4622 and RX J1713.7-3946 can be considered as twins.

The hydrodynamic state of the system, i.e. the given linear radius for known angular radius and distance together with the present age and shock velocity, etc., is basically determined by the choice of mechanical explosion energy, ejected mass, and external density.

The lack of knowledge of the radio spectral shape makes it impossible to derive from the radio synchrotron observations – and as far as the magnetic field is concerned, from the combined radio and X-ray synchrotron spectral observations – the most relevant pair of physical parameters for the acceleration theory, namely the proton injection rate and the effective magnetic field strength (Berezhko et al., 2002; Völk, 2004; Berezhko, 2005,

2008). Thus, even if we consider the circumstellar medium (CSM) structure to be given, it is not possible to directly *predict* the form of the overall synchrotron and the full VHE γ -ray emission from theory.

The observed overall nonthermal spectral shape – including the VHE γ -rays – and the small-scale filamentary structures in the nonthermal X-ray emission of RX J0852.0-4622 nevertheless provide evidence for effective nuclear cosmic ray (CR) acceleration, associated with considerable magnetic field amplification. This conclusion is possible because the overall nonthermal spectrum can be theoretically *fitted* with an appropriate proton injection rate, electron-to-proton ratio, and effective magnetic field strength (assumed to be uniform inside the shocked CSM cf. Berezhko & Völk (2004b)).

The main result of this paper is that the resulting VHE γ -ray flux is hadronically dominated. The well-known difference in the effectiveness of the basic radiation mechanisms then implies that the energy density of the nuclear component of the nonthermal charged-particle population in the SNR by far dominates that of the energetic electron component generated in the source. The energy in nonthermal particles at the present epoch amounts to ~ 35 percent of the assumed total mechanical energy $E_{\rm sn}=1.3\times 10^{51}$ erg, released in the explosion. Therefore, from the point of view of energetics, this solution for RX J0852.0-4622 more than fulfils the average requirement on a SNR source of the Galactic CRs. In this context we shall also qualitatively discuss the question of possible escape of the highest-energy particles, accelerated at an earlier phase of the SNR than the one we can observe at present.

In the next section the theoretical model is described. Section 3 presents our assumptions regarding the values of the physical parameters as they are suggested by the broadband data and by physical considerations. It contains also a discussion of the thermal emission, even though it has not been possible to estimate it appropriately for the assumed wind bubble and shell configuration. The results for the gas dynamics, the particle acceleration coupling with it, the γ -ray emission, and for the thermal X-ray emission are presented and discussed in section 4. In section 5 our conclusions are summarized.

2. Model

The theoretical model for the particle acceleration combined with the gas dynamics of the explosion has been described earlier, for instance in a recent analysis of SNR RX J1713.7-3946 (see Berezhko & Völk, 2006, and references therein).

The ejected mass $M_{\rm ej}$ has initially in its fastest moving outer parts a power law distribution $dM_{\rm ej}/du \propto u^{2-k}$ in flow velocity u, with 7 < k < 12 (Jones et al., 1981; Chevalier, 1982). We shall choose k = 8 (as in the case of SNR RX J1713.7-3946).

The interaction of the ejecta with the CSM creates a strong outward-propagating shock wave in the CSM at which particles are accelerated. Our nonlinear model is based on an explicitly time-dependent solution of the CR transport equations together with the gas dynamic equations in spherical symmetry. In particular the theory takes into account the adiabatic energy losses of thermal and nonthermal particles in the SNR interior, the diffusion of nonthermal particles from the outer shock into that interior, and the backreaction of CRs on the shock structure and dynamics. This backreaction decelerates the thermal gas already in front of the shock and leads to a smooth shock precursor that

² The distance to the Vela pulsar is estimated at $d=287^{+19}_{-17} {\rm pc}$ (Dodson et al., 2003), see also Caraveo et al. (2001).

reduces the Mach number of the subsequent collisionless plasma shock (the subshock), heating the inflowing gas³.

However, nuclear particles are only effectively injected into the acceleration process at those parts of the moving shock surface, where the locally mean magnetic field vector is quasiparallel to the shock normal. These regions are characterized by magnetic flux tubes within which the injection of moderately suprathermal particles proceeds.

In quasi-perpendicular shock regions, on the other hand, injection is instantaneously strongly suppressed or completely prohibited. In the extreme there are then separated regions at the shock surface, where particle injection is permanently either allowed or prohibited. Then also no nuclear particles are accelerated in the prohibited regions. An example for this extreme situation is SN 1006, where the X-ray emitting polar cap regions correspond to the quasi-parallel regions. In these allowed regions diffusive shock acceleration strongly proceeds on the Bohm diffusion level (see below) as a result of the effective injection of low-energy particles. However, since Bohm diffusion is isotropic, the energetic particles can also cross field lines and thus possibly reach quasi-perpendicular regions of the shock where they can also accelerate. The extent to which this happens depends on the spatial scales that separate quasi-perpendicular regions from quasi-parallel regions. For a homogeneous external magnetic field about 80% of the shock surface is quasiperpendicular in the above sense, and therefore only a fraction $f_{\rm re} \approx 0.2$ of the shock is efficiently accelerating (Völk et al., 2003). However we shall argue in section 3.1.4 that for a SNR, propagating into a stellar wind bubble with a radiatively cooling shell of high-density gas, the above spatial scales are probably so small that $f_{\rm re} \approx 1$. This has substantial implications especially for the resulting thermal X-ray emission, because the observed VHE γ -ray emission (which is by implication of hadronic origin) then requires a lower gas density.

For given magnetic field strength the electron injection rate in spherical symmetry is determined by the intensity of the observed overall synchrotron spectrum. For given injection rate of nuclear particles and magnetic field strength the ratio $K_{\rm ep}$ between the densities of nonthermal electrons and nuclear particle can then be calculated.

As a result of the streaming instability the accelerating CRs very effectively excite large-amplitude magnetic fluctuations upstream of the SN shock (Bell, 1978; Blandford & Ostriker, 1978; McKenzie & Völk, 1982). Since these fluctuations scatter CRs extremely strongly, we approximate the CR diffusion coefficient $\kappa(p)$ by its lower limit, corresponding to a scattering mean free path equal to the particle gyro radius. In this so-called Bohm limit $\kappa(p) = [mc^3/(3eB)](v/c)(p/mc)$, where e and m are the particle charge and mass, v and p denote the particle velocity and momentum, respectively, B is the effective magnetic field strength (see below), and c is the speed of light. Regarding the nuclear particles with the highest energies this Bohm limit may imply an underestimate for $\kappa(p)$ also for another reason, since for these particles the effective, amplified field has spatial scales that are smaller or equal to their gyro radius (Bell, 2004; Pelletier el al., 2006). To this extent our assumption of Bohm diffusion in the amplified field yields an upper limit to the maximum energy of the nuclear particle (e.g. Zirakashvili & Ptuskin, 2008). In addition we assume that the interior effective magnetic field has a uniform strength after its MHD compression in the thermal subshock. Practically speaking we assume this uniformity over a spatial scale that is large compared to the thickness of the observed X-ray filaments (Berezhko & Völk, 2004b). For a different point of view, see Pohl et al. (2005).

Another important nonlinear effect of the strong excitation of magnetic fluctuations by the accelerating particles themselves is the heating of the thermal plasma in the shock precursor that is generated by the accelerating particles. Combined analytical and numerical efforts, using plasma theory to give a nonlinear description of the magnetic field evolution Lucek & Bell (2000); Bell & Lucek (2001); Ptuskin & Zirakashvili (2003); Bell (2004); Pelletier el al. (2006), concluded that a considerable amplification of the upstream magnetic field should occur in the acceleration process to what we call the effective magnetic field. The physical reason is that the beam of efficiently accelerated nuclear CR component excites in particular also a non-resonant magnetic mode in addition to the well-known resonant Alfvén waves (Bell, 2004). The latter have nevertheless been argued to contribute dominantly to the overall turbulent magnetic energy density in the shock precursor (Pelletier el al., 2006). However, the three-dimensional MHD simulations of the non-resonant instability by Bell (2004) and Zirakashvili et al. (2008) show that the nonlinear growth of the magnetic fluctuations is accompanied by the formation of internal shocks and correspondingly strong dissipation which heats the thermal plasma⁴. Analogous dissipation should occur in three dimensions for the wave modes of the resonant streaming instability. We approximate this physical process by assuming that the heat input into the thermal gas equals the (linear) growth of the turbulent field energy in the excited Alfvén waves in the already amplified effective field (see Berezhko, 2008; Völk et al., 2008, for the correspondence of this approach to existing theory and experiment).

As already mentioned in the Introduction, without a reliable spectral index for the observed spatially-integrated radio synchrotron emission we shall choose the strength of the effective field as well as the nuclear injection rate η , such as to optimally fit the calculated synchrotron spectrum (from radio to X-ray energies) together with the calculated VHE γ -ray spectrum to the observations. Subsequently this spectrally fitted magnetic field is compared with the field derived from the observed filamentary structure. The degree of agreement between these two field strengths is then used as a measure of the success and self-consistency of the model.

The filament-based magnetic field strength $B_{\rm d}$ downstream of the shock is determined by the observed width L of the X-ray filament – interpreted as the synchrotron cooling length behind the shock – through the relation

$$B_{\rm d} = [3m_e^2 c^4/(4er_0^2 l_2^2)]^{1/3} (\sqrt{1+\delta^2} - \delta)^{-2/3},\tag{1}$$

where $\delta^2=0.12c/(r_0\nu)[V_{\rm s}/(\sigma c)]^2,\ l_2\approx L/7$ is the radial width of the X-ray emissivity $q_\nu(\epsilon_\nu,r),\ \epsilon_\nu$ is the X-ray energy, corresponding to the frequency ν , and r_0 denotes the classical electron radius (Berezhko & Völk, 2004a). This field strength is clearly a lower limit.

Since we have already used the observed amplitude of the VHE γ -ray spectrum to determine η we do not predict this am-

 $^{^3}$ It is implicitely assumed here that any energy loss due to radiative gas cooling has no effect on the shock structure. Given the rather high internal temperature $T_{\rm sub}$ of the remnant at this stage (see section 4.2) this appears to be a safe assumption.

⁴ Particle-in-cell simulations by Niemiec et al. (2008) led these authors even to the extreme conclusion that the non-resonant instability amplitudes become never large as a result of the bulk acceleration of the thermal gas by the streaming CRs. We believe that in a quasi-steady shock configuration the CR current is being steadily driven through the upstream gas by the downstream overpressure, as implied above.

plitude, as one could do using a detailed knowledge of the integrated synchrotron spectrum. The above consistency condition for the effective field still needs to be fulfilled for the solution to be acceptable.

Moreover, we shall not only investigate whether the chosen values for B and η are consistent with the *Chandra* measurement of the X-ray filamentary structure (Bamba et al., 2005), but also to which extent they are consistent with the semi-empirical relation (Berezhko & Völk, 2006)

$$B_0^2/(8\pi P_c) \approx 5 \times 10^{-3}$$
 (2)

which connects the upstream magnetic field pressure $B_0^2/(8\pi)$ in the shock precursor (i.e. upstream, but amplified by the CR instability) with the pressure P_c of the accelerated particles that drives the field amplification in the first place. Eq.(2) holds for a number of SNRs that could be analyzed with the aid of a well-known radio spectrum for these sources (Völk et al., 2005). The degree of fulfilment of this relation is a further quality criterium for the model to judge its success in theoretically describing the TeV γ -ray source. Eq.(2), or a relation of a similar type (Bell & Lucek, 2001), is likely to hold for an individual object also during its time evolution. Yet, in order to avoid the introduction of a further theoretical parameter, we shall consider B_0 as constant in time, equal to its present value, in our evaluation of the models for RX J0852.0-4622. We shall come back to this point in section 4.3.

In this specific form the three theory "parameters" B, η , and $K_{\rm ep}$ are determined *quantitatively* by comparison with the observations at the present age of the source. Their time-dependence during the evolution of the SNR is disregarded in this paper.

The numerical solution of the dynamical equations at each instant of time yields the CR spectrum and the spatial distributions of CRs and thermal gas. This allows the calculation of the spectra of the expected fluxes of nonthermal emission produced by the accelerated CRs, the morphology of the emission, and the future evolution inasmuch the same physical processes continue to work at later times.

In the following we shall consider the wind bubble scenario for RX J0852.0-4622 in the general framework of this model.

3. Physical parameters of RX J0852.0-4622

In this section, we will describe the physical parameters of the model. Section 3.1 gives an overview over the available broadband data. Section 3.2 describes the setup of the scenario. Because of the scarcity – and sometimes ambiguity – of the available data, we will use some general arguments from the non-thermal particle emission not only to constrain the acceleration parameters but also the environmental parameter gas density and its spatial distribution. The values of all relevant physical parameters are given in Table 1.

3.1. Broadband data of RX J0852.0-4622

3.1.1. Morphology of RX J0852.0-4622 in general

RX J0852.0-4622 is a shell-type SNR with a shell diameter of 2° , as seen in hard X-rays (Aschenbach, 1998; Aschenbach et al., 1999; Slane et al., 2001) and VHE γ -rays (Aharonian et al., 2005, 2007a). Also the radio continuum emission correlates spatially well with the high energy emission (Duncan & Green, 2000; Stupar et al., 2005; Aharonian et al.,

Table 1. Key model parameters, and corresponding spectral, dynamical, and morphological values expected from the calculations

\overline{d}	1 kpc
$R_{ m s}$	17.5 pc
It _S	17.5 pc
$B_{ m d}$ from X-ray filaments	$139\mu\mathrm{G}$
$E_{ m sn}$	$1.3 \times 10^{51} {\rm erg}$
$M_{ m ej}$	$3.5M_{\odot}$
$N_{ m g}(R_{ m s})$	$0.24{\rm cm}^{-3}$
$N_{\rm g}(r=0)$	$0.003{\rm cm}^{-3}$
k	8
B_0	$20 \mu \mathrm{G}$
η	3×10^{-4}
$K_{ m ep}$	3×10^{-4}
$f_{ m re}$	1
$t_{ m sn}$	3745 yr
$V_{ m s}(t_{ m sn})$	$1316 \mathrm{km s^{-1}}$
$\sigma(t_{ m sn})$	5.2
$\sigma_{ m s}(t_{ m sn})$	3.1
$M_{ m s}(t_{ m sn})$	$25~M_{\odot}$
$E_{\rm c}(t_{ m sn})$	$4.6 \times 10^{50} { m erg}$
$B_{\rm d}(t_{\rm sn})(=\sigma B_0)$	$104\mu\mathrm{G}$
$P_{ m c}/(ho_0 V_{ m s}^2)$	0.145
$B_0^2/(8\pi P_{\rm c})$	6.5×10^{-3}

Parameter description: The quantities d and R_s denote the assumed distance and the radius of the source, respectively, $B_{\rm d}$ is the internal magnetic field strength, as determined from the thickness of observed X-ray filaments cf. eq. 1, and $E_{\rm sn}$ is the total hydrodynamic explosion energy; $M_{\rm ej}, M_{\rm s}(t_{\rm sn}), N_{\rm g}(R_{\rm s})$ and $N_{\rm g}(r=0)$ are the ejected mass, the sweptup mass, the circumstellar gas number density at the SNR shock, and the number density at the centre, respectively; k is the power law index of the ejecta velocity distribution; B_0 is the assumed amplified magnetic field strength in the upstream region of the shock precursor, while η and $K_{\rm ep}$ denote the assumed proton injection rate and energetic electron-toproton ratio, respectively; $t_{\rm sn}$ is the calculated age of the SNR; $V_{\rm s}(t_{\rm sn})$, $\sigma(t_{\rm sn}), \, \sigma_{\rm s}(t_{\rm sn}), \, E_{\rm c}(t_{\rm sn}), \, {\rm and} \, B_{\rm d}(t_{\rm sn})$ are the resulting values of the subshock velocity, the total compression ratio, the subshock compression ratio, the total nonthermal energy, and the downstream magnetic field strength, respectively. Finally, $P_{\rm c}$ and $\rho_0=m_{\rm p}N_{\rm g}(R_{\rm s})$ denote the postshock pressure of accelerated particles and postshock mass density of the gas, respectively.

2007a). This confirms the shell-type nature of the source, although confusion with the emission from the Vela SNR prevented a detection based on the radio data alone.

In X-rays, a central diffuse source of $\sim 9' \times 14'$ extension is seen NW of the geometrical centre of the SNR (Slane et al., 2001; Becker & Aschenbach, 2002; Becker et al., 2007). While Slane et al. (2001) argue that the hard spectrum of the source seen with ASCA hints at a Pulsar Wind Nebula (PWN), Becker et al. (2007) argue, based on XMM-Newton data, that the source is soft and reject a PWN nature. Hence, there is currently no agreement whether RX J0852.0-4622 is a centre-filled, composite SNR or not. Spatially-integrated fluxes would only marginally be affected. There is however the possibility that such a central PWN would slightly influence the VHE γ -ray radial profile (see Aharonian et al., 2007a, and Fig. 7).

3.1.2. Central compact object and pulsar association

At the geometrical centre of the shell of RX J0852.0-4622 lies the X-ray point source CXOU J085201.4-461753 (Aschenbach et al., 1999; Pavlov et al., 2001; Kargaltsev et al., 2002; Mereghetti, 2001; Mereghetti et al., 2002), a central compact object (CCO) similar to that detected in the centre of Cas A. CXOU J085201.4-461753 might be a neutron star, but the nature of the object and the possibly associated compact $H\alpha$ nebula (Pellizzoni et al., 2002) is still under debate (e.g. Reynoso et al., 2006; Mignani et al., 2007; Becker et al., 2007). No X-ray pulsations have been detected (Kargaltsev et al., 2002; Becker et al., 2007). The association of CXOU J085201.4-461753 with RX J0852.0-4622 is nevertheless suggestive and, if true, would allow conclusions on the nature of the progenitor of RX J0852.0-4622. Since the scenario discussed in this paper implies a core collapse SN, it does not exclude an association of CXOU J085201.4-461753 with RX J0852.0-4622.

In the literature also the possibility is discussed that RX J0852.0-4622 is associated with PSR J0855-4644 (Redman & Meaburn, 2005). This appears unlikely however, given the implications of this association on distance and age of RX J0852.0-4622 (Redman & Meaburn, 2005).

3.1.3. Non-thermal X-rays

The soft X-ray emission from RX J0852.0-4622 is heavily confused by thermal emission from the Vela SNR. While the emission from the Vela SNR seems consistently constrained to two thermal components ($T_{1,2} = 0.05, 1.2 \,\mathrm{keV}$) with an absorption column density of about 10^{20}cm^{-2} (Lu & Aschenbach, 2000), the surface brightness and temperatures of these components are variable enough to prevent a clean subtraction of the Vela SNR in the X-ray spectra of RX J0852.0-4622 (Slane et al., 2001; Iyudin et al., 2005; Aharonian et al., 2007a). In the spectra above $\sim 1 \, \mathrm{keV}$, the non-thermal emission from RX J0852.0-4622 nevertheless clearly dominates. We adopt the common interpretation that this component is synchrotron emission from relativistic electrons and use two estimates of the total X-ray synchrotron flux from RX J0852.0-4622 in our modelling (see Figs. 3, 5): The first (lower) estimate was derived using the averaged power-law spectra derived from the three brightest shell components (Slane et al., 2001), which we scaled up to match the total flux measured with ROSAT in the soft X-ray band (Aschenbach, 1998). The second (upper) estimate is a reanalysis of the ASCA data set as presented in Aharonian et al. (2007a).

High-resolution imaging of this synchrotron X-ray component with XMM-Newton (Iyudin et al., 2005, 2007) and especially with Chandra (Bamba et al., 2005; Pannuti et al., 2004) permits the derivation of synchrotron cooling times and therefore of an estimate of the effective magnetic field (e.g. Berezhko et al., 2003a; Berezhko & Völk, 2004a; Bamba et al., 2005). In order to test our model we will compare the B-field derived in this manner to the field value that is required to fit the spatially integrated synchrotron data (see section 4).

3.1.4. Thermal X-rays

The detection of thermal X-ray emission could help to constrain the gas density and/or its spatial distribution in the SNR. However, the interpretation of the X-ray spectra is impeded by the strong background emission from the Vela SNR; the latter is dominated by low-temperature (< 1 keV) X-ray emission. After subtraction of such emission the remaining emission (dominant

at higher energies) has been attributed either to gas at higher temperatures from RX J0852.0-4622 (Aschenbach et al., 1999), or to synchrotron emission plus a very small thermal contribution (Slane et al., 2001). The latter conclusion has basically been taken over in the recent *H.E.S.S.* paper (Aharonian et al., 2007a) that also contains a re-analysis of the ASCA data, even though the X-ray line features exhibited in the spectrum below 2 keV might be associated with either RX J0852.0-4622 or the Vela SNR, or to both. Similarly the residuals of the spectra to the ASCA data visible around 1 keV are suggestive that another component, which might originate from RX J0852.0-4622, is needed. The possible detection of a thermal emission component from RX J0852.0-4622 by Uchiyama (2008) is to be mentioned here again.

These are complex possibilities. The calculation of the thermal emission for the concrete case of RX J0852.0-4622 is compounded by the wind-bubble plus swept-up-shell geometry. Standard methods for calculating the thermal emission from SNRs approximate the configuration either by a plane shock geometry or by a classical Sedov solution, and they neglect the existence of the accelerated particle component. Neither of these approximations is appropriate for the case of RX J0852.0-4622 in its present phase. The plane approximation disregards the adiabatic gas cooling in the interior and the overall dynamic evolution of the system. The classical Sedov solution implies a uniform circumstellar medium. In the extreme case the radiatively cooled wind bubble shell is even denser - and therefore even thinner – than assumed in subsection 3.2.2. and it might only be reached recently before the present epoch. Then collisional electron heating has had little time to operate until now.

Another important aspect is the modification of the SNR shock by the accelerating CRs. As mentioned before, for a SNR shock propagating into a uniform medium with a uniform magnetic field this implies that the larger part of the shock surface corresponds to a quasi-perpendicular shock with a strongly reduced injection of nuclear particles (Völk et al., 2003)⁵. Suprathermal injection of ions is only possible in the quasi-parallel shock regions. If the spatial scales of the quasiperpendicular regions are large enough, then the cross-field diffusion of the highest-energy particles, accelerated in the magnetic flux tubes delineating the neighboring quasi-parallel shock regions, does not reach deeply into these quasi-perpendicular regions. In the corresponding magnetic flux tubes there are no nuclear particles to be accelerated, there is no magnetic field amplification, and the shock remains unmodified there. This means that in the quasi-perpendicular regions the shock dissipation and therefore the gas heating occurs in a locally unmodified shock with the overall shock speed, leading to a correspondingly high gas temperature and high thermal emission. In the case where a radiatively cooling shell of a wind bubble is the obstacle for the SNR expansion, the situation may be different. MHD instabilities and the radiative cooling of such a shell probably break it into many small regions with strongly varying field directions. Then the spatial scales separating the flux tubes originating from quasi-perpendicular regions from those of quasi-parallel shock regions may become small enough that cross-field diffusion can also fill the quasi-perpendicular flux tubes with accelerating particles and then particle acceleration plus magnetic field amplification occur practically everywhere over the shock surface (Völk et al., 2008; Völk, 2008). In the extreme this implies shock modification over the entire shock region and thus a re-

⁵ For earlier discussions of this question in a more general context, see Ellison et al. (1995); Malkov & Völk (1995).

duced gas heating due to the subshock dissipation only. The enhanced overall acceleration efficiency then also demands a lower density of the thermal gas for a given hadronic γ -ray flux, and thus a lower thermal emission. The low swept-up mass in a low-density wind bubble in addition lowers the overall thermal emissivity compared to that of a classical Sedov remnant with the same upstream gas density at the shock at the present epoch.

In section 4.4 a rough estimate of the resulting thermal X-ray emission will be given, based on the emission from a classical Sedov solution in a uniform ambient medium. According to this estimate the thermal emission of soft X-rays at 1 keV is larger than the corresponding nonthermal X-ray emission. However, the error in this estimate is not known and is likely to be quite large. Therefore an uncertainty remains which we cannot resolve at this point.

3.1.5. X-ray morphology absorption and relation to CO data

Absorption of soft X-rays by neutral hydrogen can be used to put constraints on the source distance. Slane et al. (2001) used CO data of the Vela Molecular Ridge (VMR) to reject a distance of RX J0852.0-4622 of more than 1-2 kpc, based on the lack of strong X-ray absorption variation that should have been detected across RX J0852.0-4622. This is broadly in agreement with the inference by Moriguchi et al. (2001) that there is an anticorrelation of X-ray emission and molecular gas traced in CO, especially with regard to the VMR at a distance of $1-2~{\rm kpc}$.

In the 2-10~keV band, that is expected not to exhibit absorption, an X-ray power-law spectrum can be derived. Assuming this power-law to continue down to $0.7~keV,\, Slane$ et al. (2001) have then derived an absorption column density of $(4.0\pm1.8)\cdot10^{21}~cm^{-2}$ for RX J0852.0-4622. Since this column density is much larger than the one towards the Vela SNR, they concluded that RX J0852.0-4622 should be at a much larger distance than the Vela SNR.

3.1.6. Radio spectrum

We assume that the radio emission is due to synchrotron radiation. There is no good radio spectrum available fvelaj61.dvior the entire remnant. We use the differential flux values given by Duncan & Green (2000) at 2.42 GHz and 1.40 GHz, the errors of which are representing the uncertainty of the background level. The spectral index between the two bands has quite a large error ($\alpha=0.4\pm0.5$), but for the north-western rim a better spectrum ($\alpha=0.40\pm0.15$) could be derived. If this value is representative for the entire remnant, as Duncan & Green (2000) argue, then this index is somewhat harder than what we expect from a modified SNR shock environment, though still compatible within a $2\,\sigma$ error range.

Gamma-ray and X-ray line emission from radioactive ⁴⁴Ti decay

The $^{44}\mathrm{Ti}$ production in a SN explosion depends on progenitor star mass and explosion-type, with a yield spanning two orders of magnitude (see, e.g. Renaud et al., 2006a, and references therein). Because of the short lifetime of ~ 80 years (e.g. Wietfeldt et al., 1999), the mere detection of hard X-ray and γ -ray de-excitation lines at 69.7, 78.4, and 1157 keV from the $^{44}\mathrm{Ti}$ radioactive decay products can be used to significantly constrain the SNR age. So far, however, these lines have only been

detected unambiguously from Cassiopeia A (age presumably ~ 330 years), with COMPTEL onboard CGRO (Iyudin et al., 1994), PDS onboard *BeppoSAX* (Vink et al., 2001), and the ISGRI imager onboard INTEGRAL (Renaud et al., 2006c).

For RX J0852.0-4622, the situation is unfortunately unresolved. From COMPTEL data, the detection of a γ -ray line at 1163 \pm 16 keV, consistent with the $^{44}{\rm Ti}$ γ -ray decay line, was claimed and predominantly attributed to RX J0852.0-4622 (Iyudin et al., 1998; Aschenbach et al., 1999). Using this $^{44}{\rm Ti}$ line flux, and using a rather high shock velocity, Aschenbach et al. (1999) derived an age of 680 ± 100 yrs, and a distance of 200 pc. Such γ -ray data also suggest a core collapse SN event.

However, Schönfelder et al. (2000) pointed out that the significance of the COMPTEL ⁴⁴Ti result is only marginal. And the COMPTEL result could so far not be confirmed with the INTEGRAL instruments SPI and ISGRI. A SPI upper limit is so far unconstraining (von Kienlin et al., 2005). Under a point-source assumption, the ISGRI non-detection of the 78 keV line would be in conflict with the COMPTEL result (Renaud et al., 2006a). However, an extended-source analysis has not yet been performed (Renaud et al., 2006b).

Therefore a nearby and rather recent event is not decisively excluded from these specific observations, even though the arguments for it are rather weak.

3.2. Model parameters for a SN explosion in 1 kpc distance

As discussed in the previous section, the main observational motivation to locate RX J0852.0-4622 at a distance of $\sim 1\,\mathrm{kpc}$ is the much larger column density in neutral hydrogen derived from the X-ray spectrum of RX J0852.0-4622, compared to the values for Vela SNR.

3.2.1. Reasons for a wind bubble scenario

The lack (or low level) of thermal X-ray emission and the strong X-ray synchrotron flux already led Slane et al. (2001) to the conclusion that RX J0852.0-4622 could be evolving into a wind bubble. Similarly, Duncan & Green (2000) argue that the unusual radio properties of RX J0852.0-4622 (bipolar shell morphology, low surface brightness) could be explained if the SNR has so far evolved in a low-density region.

We can use the non-thermal X-ray plus γ -ray emission from RX J0852.0-4622 to put these arguments for a wind bubble scenario on a more firm footing, anticipating for the moment that the VHE γ -ray emission is indeed dominated by hadronic emission. As we shall see in section 4, this latter conclusion follows from the fact that with the assumption of a significantly amplified magnetic field it is possible to fit all the nonthermal spectra as well as the morphology in nonthermal X-rays and gammarays – and that this amplified field does not exceed that derived from the observed X-ray filaments. Such amplification is however only possible by the nonthermal pressure of the nuclear particles.

Arguments similar to those given below have been used in Berezhko & Völk (2006) for the comparable case of SNR RX J1713.7-3946 to which we refer the reader here.

In order to yield the observed nonthermal X-ray luminosity in the case of a uniform ISM the shock speed should be sufficiently large, $V_s \gtrsim 10^3$ km/s (Berezhko & Völk, 2004b). Given that the SNR would already be in the Sedov phase the observed

size $R_{\rm s}\sim 20\,{\rm pc}$, corresponding to a distance of 1 kpc, would lead to the age constraint $t_{\rm sn}\approx 0.4R_{\rm s}/V_{\rm s}\lesssim 3\times 10^3$ yr. For a typical SN type Ia explosion energy $E_{\rm sn}=10^{51}$ erg and ejected mass of $1.4M_{\odot}$ this would then imply a very low ISM number density $N_{\rm H}\lesssim 10^{-2}~{\rm cm}^{-3}$. On the other hand, the peak TeV γ -ray luminosity, achieved during SNR evolution from a type Ia event, roughly scales as (Berezhko & Völk, 1997):

$$\epsilon_{\gamma} F_{\gamma}(\epsilon_{\gamma}) \approx 150 \left(\frac{N_{\rm H}}{1~{\rm cm}^{-3}}\right) \left(\frac{1~{\rm kpc}}{d}\right)^2 \frac{{\rm eV}}{{\rm cm}^2 {\rm s}},$$
 (3)

for $f_{\rm re}=1$. Here $F_{\gamma}(\epsilon_{\gamma})$ is the integral flux of γ -rays with energies greater than ϵ_{γ} . This expression shows that for $N_{\rm H}<10^{-2}\,{\rm cm}^{-3}$ we would have to expect an energy flux $\epsilon_{\gamma}F_{\gamma}(\epsilon_{\gamma})<1.5\,{\rm eV/(cm^2s)}$. This is an order of magnitude smaller than the observed flux (see section 4).

Therefore the nonthermal observations make it clear that SNR RX J0852.0-4622 can not correspond to a type Ia event, if the source distance is as large as $d=1~{\rm kpc}$. As a consequence we shall consider a core collapse SN event.

3.2.2. Wind bubble parameters

The progenitor stars of core collapse SNe that significantly modify the density of their environment are massive main-sequence stars with initial masses $M_{\rm i}>15M_{\odot}$ which have intense winds, e.g. (Abbot, 1982). In the mean, during their evolution in the surrounding uniform ISM of gas number density $\rho_0=m_{\rm p}N_{\rm ISM}$, they create a low-density bubble, surrounded by a shell of swept-up and compressed ISM of radius (Weaver et al., 1977; Chevalier & Liang, 1989)

$$R_{\rm sh} = 0.76(0.5\dot{M}V_{\rm w}^2 t_{\rm w}^3/\rho_0)^{1/5},\tag{4}$$

where M is the mass-loss rate of the progenitor star, $V_{\rm w}$ is the wind speed, and $t_{\rm w}$ is the duration of the wind phase. The values of these parameters are given in tabular form by Chevalier (1982) in terms of $M_{\rm i}$.

In order to determine the SNR shock dynamics inside the shell we model the gas number density distribution in the bubble and in the shell in the form (e.g. Berezhko & Völk, 2006):

$$N_{\rm g} = N_{\rm b} + (r/R_{\rm sh})^{3(\sigma_{\rm sh} - 1)} N_{\rm sh},$$
 (5)

where $N_{\rm sh}=\sigma_{\rm sh}N_{\rm ISM}$ is the peak number density per hydrogen atom in the shell, $N_{\rm b}$ is the gas number density inside the bubble, typically very small compared with the shell density, and $\sigma_{\rm sh}=N_{\rm sh}/N_{\rm ISM}$ is the shell compression ratio. We note that as a result of radiative cooling the compression ratio $\sigma_{\rm sh}$ can exceed the classical upper limit of 4. The possibility of a very thin wind bubble with $\sigma_{\rm sh}\gg 4$ has been indicated in subsection 3.1.4. above.

The mass of the bubble

$$M_{\rm b} = (4\pi R_{\rm sh}^3/3)m_{\rm p}N_{\rm b}$$
 (6)

is rather small, $M_{\rm b} < M_{\odot}$, in the case of moderate progenitor masses $M_{\rm i} < 20 M_{\odot}$, whereas the shell mass

$$M_{\rm sh} = 4\pi N_{\rm sh} m_{\rm p} \int_0^{R_{\rm sh}} dr r^2 (r/R_{\rm sh})^{3(\sigma-1)} = (4\pi R_{\rm sh}^3/3) N_{\rm ISM} m_{\rm p}$$

is many hundred solar masses. Therefore, during SNR shock propagation through the bubble, only a small fraction of its energy is given to gas of stellar origin. The main part of the explosion energy is deposited in the shell.

Here we use the gas number density distribution $N_{\rm g}(r)=\rho(r)/m_{\rm p}$ in the form

$$N_{\rm g} = \{0.003 + 0.24[r/(17.5\,{\rm pc})]^{12}\}\,{\rm cm}^{-3},$$
 (8)

which fixes the gas density at the present shock radius and provides a consistent fit for all existing data for SNR RX J0852.0-4622. Together with Eq.(4), this relation also connects the external density $N_{\rm ISM}$ with the progenitor mass.

Such a distribution corresponds to a bubble with $\sigma_{\rm sh}=5$ and $28 < R_{\rm sh} < 32$ pc created by the wind of a main-sequence star of initial mass $15 M_{\odot} < M_{\rm i} < 20 M_{\odot}$ in the surrounding ISM of hydrogen number density $11 < N_{\rm ISM} < 49~{\rm cm}^{-3}$, respectively (Chevalier & Liang, 1989). It implies that this bubble is located inside a region of dense gas.

3.2.3. Further parameters that determine the SN evolution and CR acceleration

We shall use the SNR parameters $E_{\rm sn}=1.3\times 10^{51}$ erg, $M_{\rm ej}=3.5M_{\odot}$, and k=8 which, as will be shown below, give a good fit for the observed SNR properties.

We shall also use an upstream effective magnetic field value $B_0=20~\mu\mathrm{G}$, which is required to provide the observed synchrotron flux in the radio and X-ray bands (see below). Such a value of B_0 is significantly higher than a merely MHD-compressed dense-gas magnetic field in the inner part of the shell, at densities $N_{\rm g}\leq0.24~\mathrm{cm}^{-3}$.

4. Results and discussion

In this section we shall discuss the physical characteristics of the wind bubble scenario in detail, and compare them with the observations.

The calculated dynamical characteristics of the SNR are shown in Fig. 1. From Fig. 1a one can see that for the assumed distance of 1 kpc the calculation fits the observed SNR size $R_{\rm s}\approx 17.5~{\rm pc}$ at the age $t_{\rm sn}=3745~{\rm yr}.$

To fit the observed synchrotron and γ -ray spectra (see below) we assume a proton injection rate $\eta=10^{-3}$. This leads to a moderate nonlinear modification of the shock which at the current age of $t_{\rm sn}=3745$ yrs has a total compression ratio $\sigma\approx5.2$ and a subshock compression ratio $\sigma_{\rm s}\approx3.1$ (Fig. 1b). All parameters used and the resulting model properties are summarized in Table 1.

For its adopted density, the wind bubble contains only a small amount of gas $M_{\rm b}\approx 0.3 M_{\odot}$. Therefore the SN shock deposits only about 20% of the explosion energy during the initial 1000 years of propagation through the bubble, as seen in Fig. 1. However, up to the current epoch the SN shock has already swept up a considerable mass $M_{\rm sw}\approx 25 M_{\odot}$. Therefore the ejecta have transformed about 85% of their initial energy into gas and CR energy (Fig. 1c). The acceleration process is therefore characterised by a high efficiency under the assumption of spherical symmetry: at the current time about 35% of the explosion energy have been transferred to CRs, and the relative CR energy content $E_{\rm c}$ continues to increase to a maximum of about $R_{\rm c}$ in the later phase (Fig. 1c), when particles start to leave the source.

Therefore, in absolute terms the CRs inside SNR RX J0852.0-4622 already contain

$$E_{\rm c} \approx 0.35 E_{\rm sn} \approx 4.6 \times 10^{50} \,\text{erg}.$$
 (9)




Fig. 1. Model parameters as a function of time: (a) Shock radius $R_{\rm s}$ and overall shock speed $V_{\rm s}$; (b) total shock (σ) and subshock $(\sigma_{\rm s})$ compression ratios; (c) ejecta $(E_{\rm ej})$ and CR $(E_{\rm c})$ energies in spherical symmetry. The vertical *dotted line* marks the current epoch of SNR evolution. The external gas density $N_{\rm g} = \rho_{\rm g}/m_{\rm p} = 1.4N_{\rm H}$ is also shown in *panel (a)*.

The volume-integrated (or overall) CR spectrum

$$N(p,t) = 16\pi^{2}p^{2} \int_{0}^{\infty} dr r^{2} f(r,p,t)$$
 (10)

has, for the case of protons, almost a pure power-law form $N \propto p^{-\gamma}$ over a wide momentum range from $10^{-2}m_{\rm p}c$ up to the cutoff momentum $p_{\rm max} \approx 6 \times 10^5 m_p c$, corresponding to a cutoff energy of $\approx 5.6 \times 10^{14}$ eV. (see Fig. 2). This value $p_{\rm max}$ is limited mainly by geometrical factors, which are the finite size and speed of the shock, its deceleration and the adiabatic cooling effect in the downstream region (Berezhko, 1996). Due to the shock modification the power-law index slowly varies from




Fig. 2. The overall (volume-integrated) CR spectrum as a function of particle momentum. *Solid and dashed lines* correspond to protons and electrons, respectively.




Fig. 4. Calculated nonthermal Bremsstrahlung (*NB*, dash-dotted line), inverse Compton (*IC*, dashed line), and π^0 -decay (solid line) γ -ray spectral energy distributions as functions of photon energy ϵ_{γ} for the high-injection, high-field model. The observed *H.E.S.S.* (Aharonian et al., 2007a) and *CANGAROO* (Enomoto et al., 2006) γ -ray fluxes are shown as well.

 $\gamma=2.4$ at $p\lesssim m_{\rm p}c$ to $\gamma=1.9$ at $p\sim 100m_{\rm p}c$. The shape of the overall electron spectrum $N_{\rm e}(p)$ deviates from that of the proton spectrum N(p) at high momenta $p>p_{\rm l}\approx 350m_{\rm p}c$, as a result of the synchrotron losses in the downstream region with a magnetic field strength $B_{\rm d}\sim 100~\mu{\rm G}$ which is assumed uniform ($B_{\rm d}=B_2=\sigma B_0$). Therefore within the momentum range $p_{\rm l}< p< p_{\rm max}^{\rm e}$ the electron spectrum is considerably steeper $N_{\rm e}\propto p^{-3}$.

Specifically, the synchrotron losses become important for electron momenta greater than (Berezhko et al., 2002)

$$\frac{p_{\rm l}}{m_{\rm p}c} \approx 1.3 \left(\frac{10^8 \text{ yr}}{t}\right) \left(\frac{10 \mu \text{G}}{B_{\rm d}}\right)^2. \tag{11}$$

Substituting the SN age t=3745 yr into this expression, we have $p_{\rm l}\approx 350m_{\rm p}c$, in agreement with the numerical result (see Fig. 2).

The maximum electron momentum can be roughly estimated by equating the synchrotron loss time with the acceleration time. This gives (e.g. Berezhko et al., 2002)

$$\frac{p_{\mathrm{max}}^{\mathrm{e}}}{m_{\mathrm{p}}c} = 6.7 \times 10^{4} \left(\frac{V_{\mathrm{s}}}{10^{3} \, \mathrm{km/s}}\right)$$




Fig. 3. Calculated broadband spectral energy density of RX J0852.0-4622, as function of photon energy ϵ_{γ} . In the γ -ray region the *solid line* shows the $\pi^0-decay$ emission and the *dashed line* denotes the inverse Compton (IC) emission. Radio fluxes were obtained with the *Parkes* telescope (Duncan & Green, 2000). For the X-ray synchrotron flux a lower boundary is given by the sum of the *ASCA* fluxes from the brightest parts of the SNR, as given by Slane et al. (2001), scaled up to match the total remnant's flux measured with *ROSAT* in the soft X-ray range (Aschenbach, 1998). The upper boundary comes from a re-analysis of the total *ASCA* data from the remnant, as given in Aharonian et al. (2007a). TeV data are from *CANGAROO* (Enomoto et al., 2006) and *H.E.S.S.* (Aharonian et al., 2007a).




Fig. 5. Calculated synchrotron emission flux as a function of frequency ν . The *solid line* corresponds to the high-injection, high-field model, the *dashed line* corresponds to a hypothetical very low injection, and therefore unmodified low-field model. *Parkes* radio data (Duncan & Green, 2000) and *ASCA* X-ray data (Slane et al., 2001; Aharonian et al., 2007a) are shown (see also the caption of Fig. 3).

$$\times \sqrt{\frac{(\sigma - 1)}{\sigma(1 + \sigma^2)} \left(\frac{10 \,\mu\text{G}}{B_0}\right)}.\tag{12}$$

At the current epoch $V_{\rm s}\approx 1316$ km/s, which leads to a maximum electron momentum $p_{\rm max}^{\rm e}\approx 10^4 m_{\rm p}c$, in agreement with the numerical results (Fig. 2).

As a result of the shock propagating through the wind shell (cf. Fig. 1) the SN shock speed decreases rather quickly during the period $t>10^3$ yr. Therefore, during previous evolutionary phases the shock has produced electron spectra with cutoff momenta $p_{\rm max}^{\rm e}$ larger than at the current epoch. Due to this factor the spatially integrated electron spectrum has a relatively smooth cutoff (see Fig. 2). Together with the synchrotron cooling this gives a good fit of the observed X-ray spectrum (see below).

The present-day parameters $B_{\rm d}=104~\mu{\rm G}$ and $K_{\rm ep}\approx 3\times 10^{-4}$ lead to good agreement between the calculated and the measured spectral energy distribution of the synchrotron emission in the radio to X-ray ranges at the present time (Fig. 3). The steepening of the electron spectrum at high energies due to synchrotron losses and the smooth cutoff of the overall electron spectrum together naturally yield a fit to the X-ray data with their soft spectrum. Such a smooth spectral behaviour is achieved in an assumed upstream field of $20~\mu{\rm G}$ (which leads to the above downstream field $B_{\rm d}$).

Fig. 3 also shows the calculated γ -ray spectral energy distributions due to π^0 -decay, IC emission, and nonthermal Bremsstrahlung, together with the existing experimental data.

According to the calculation, the hadronic γ -ray production exceeds the electron contribution by more than two orders of magnitude at all energies. In detail these γ -ray spectra are shown in Fig. 4. For energies $\epsilon_{\gamma}=1-100$ GeV the γ -ray spectrum is close to $dF_{\gamma}/d\epsilon_{\gamma} \propto \epsilon_{\gamma}^{-2}$, hardening from $\epsilon_{\gamma}=0.1-1$ TeV, whereas starting from $\epsilon_{\gamma}\approx 1$ TeV it has a smooth extended cutoff despite the comparatively much sharper cutoff of the proton energy spectrum, cf. Fig. 2.

Note that the γ -ray cutoff energy $\epsilon_{\gamma}^{max}\approx 0.1p_{\rm max}c$ is sensitive to the magnetic field strength $B_{\rm d}$, since the proton cut-

off momentum has a dependence $p_{\rm max} \propto R_{\rm s} V_{\rm s} B_{\rm d}$ (Berezhko, 1996). It is clearly seen from Fig. 4 that the calculated spectrum fits the H.E.S.S. measurements in an acceptable way, at least up to ≈ 5 TeV. However the four highest-energy points tend to lie below the theoretical curve. This can be the result of escape of the highest-energy protons during the deceleration of the shock in the shell. See section 4.1 for further details.

The hadronic dominance in VHE γ -ray emission which we predict here, could be further investigated in the near future by the *Fermi* instrument in the GeV region. Even though at such comparatively low γ -ray energies the γ -ray background from the diffuse Galactic γ -rays is quite significant, especially for such a large, low-surface brightness source as RX J0852.0-4622 (Drury et al., 1994), *Fermi* should be able to detect the overall very high γ -ray flux from RX J0852.0-4622. It is therefore to be expected that the *Fermi* instrument will confirm our prediction that the spatially-integrated γ -ray spectral energy density at 1 GeV is only a factor ≈ 1.5 *lower* than at 1 TeV cf. Fig. 4, as a result of the nonlinear modification of the acceleration process. If the nonlinear modification in the wind bubble is less strong than assumed here, then this difference in the spectral energy density between 1 GeV and 1 TeV should be even smaller.

In Fig. 5 we separately present the differential synchrotron spectrum, produced at the current epoch. For comparison we also show a synchrotron spectrum, which would correspond to a hypothetical acceleration scenario in which the proton injection rate is taken so small ($\eta = 10^{-5}$) that the accelerated nuclear CRs do not produce any significant shock modification or magnetic field amplification; the value $B_0 = 5 \mu G$ is used in this case. There are two small but distinct differences between the synchrotron spectra that correspond to these two scenarios. The high-injection, high-field scenario leads to a steep differential radio frequency spectrum $S_{\nu} \propto \nu^{-\alpha}$ with power law index $\alpha \approx 0.7$, whereas for the unmodified, low-field scenario $\alpha = 0.5$. Unfortunately, the low quality of the existing radio data does not allow us to distinguish these two scenarios in the radio range, in order to conclude from the radio spectrum alone whether or not we deal in the case of RX J0852.0-4622 with efficient CR acceleration leading to a significant shock modification and magnetic field amplification. The essentially different behaviour of these two spectra at X-ray frequencies around $\nu = 10^{18}$ Hz demonstrates on the other hand that in the case of strong CR production and amplified magnetic field $B_{\rm d} \approx 100 \ \mu \rm G$ the spectrum $S_{\nu}(\nu)$ naturally exhibits a smooth cutoff consistent with the experiment. In the simple, unmodified low-field case the spectrum $S_{\nu}(\nu)$ has too sharp a cutoff to be consistent with the experiment.

It is noted that the X-ray flux represented in Figs. 3 and 5 comes from two different analyses of the X-ray flux (see also section 3.1): the lower boundary was derived by summing up the *ASCA* fluxes from the brightest parts of the SNR (Slane et al., 2001) and scaling the result up to match the total SNR's flux as measured with ROSAT (Aschenbach, 1998). The upper boundary – a factor of two higher – comes from a re-analysis of the total *ASCA* data from the remnant, as given in Aharonian et al. (2007a).

The properties of small-scale structures of RX J0852.0-4622 seen in X-rays furnish even stronger evidence that the magnetic field inside the SNR is indeed considerably amplified. As in the case of other young SNRs (e.g. SN 1006, Cassiopeia A, Tycho's SNR) *Chandra* shows very fine filamentary structures in nonthermal X-rays in the very outer part of the remnant. The thinnest filament detected by Bamba et al. (2005) has an angular thickness $\Delta\psi\approx38''$ in the radial profile of the X-ray emission in




Fig. 6. The projected radial profile of the X-ray synchrotron emission for $\epsilon_{\nu}=1\,\mathrm{keV}$. The abscissa is normalized to the shock radius and corresponds therefore to an angular scale. The *solid line* corresponds to the high-injection, high-field model, the *dash-dotted line* corresponds to a hypothetical low-injection, low-field model.

the $2-10~\rm keV$ range. In order to find out whether this type of structure is consistent with our theory we present in Fig. 6 the projected radial profile

$$J(\epsilon, \rho) \propto \int dx q(\epsilon, r = \sqrt{\rho^2 + x^2}, x),$$
 (13)

calculated for the X-ray energy $\epsilon_{\nu}=1$ keV. The abscissa in Fig. 6 (and correspondingly also in Fig. 7) is scaled to the radius of the remnant, i.e. to 17.5 pc, as derived from the angular radius of the remnant of $\sim 1^{\circ}$. Here $q(\epsilon,r)$ is the luminosity in the nonthermal emission with photon energy ϵ . The integration is performed along the line of sight. One can see that the theory predicts the peak of the emission just behind the shock front with a thickness $\Delta \rho/R_{\rm s} \approx 10^{-2}$ which corresponds to an angular width $\Delta \psi \approx 36''$. It corresponds reasonably well to the *Chandra* observation. At the same time Fig. 6 also shows that the unmodified low-field solution cannot explain the filament structure seen with *Chandra*.

It has been already demonstrated for other young SNRs (Berezhko et al., 2003a; Berezhko & Völk, 2004a) that the measured width of the projected radial profile of the nonthermal X-ray emission gives the possibility to determine the internal magnetic field strength according to Eq.(1).

Substituting into this equation $l_2=L/7=8.2\times 10^{16}$ cm, $\sigma=5.2, V_{\rm s}=1316$ km/s, and $\nu=3\times 10^{17}$ Hz (i.e. X-ray energy $\epsilon_{\nu}\approx 1$ keV), we obtain $B_{\rm d}\approx 139~\mu{\rm G}$, which agrees within 30% with the value $B_{\rm d}=104~\mu{\rm G}$ used in our spectrum calculation. Such a difference in $B_{\rm d}$ corresponds to the uncertainty in the field determination in the other objects analysed up to now.

The magnetic field amplification is driven by the gradient of the (nuclear) CR pressure upstream of the outer shock, and we can check whether RX J0852.0-4622 belongs to the class of objects that fulfil Eq. 2. In the present case of d=1 kpc we have $P_c\approx 0.15\rho_0V_{\rm s}^2$, where $\rho_0=N_{\rm g}(R_{\rm s})m_{\rm p}$ is the ambient gas density at the current shock front position. Substituting $N_{\rm g}(R_{\rm s})=0.24~{\rm cm}^{-3}$ and $B_0=20~\mu{\rm G}$ we have $B_0^2/(8\pi P_c)\approx 6.5\times 10^{-3}$, in rather good agreement with the average number in Eq.(2).




Fig. 7. The γ -ray emissivities for the γ -ray energy $\epsilon_{\gamma}=1~{\rm TeV}$ as a function of projected, normalized radial distance $\rho/R_{\rm s}$. The calculated radial profile is represented by the *solid line*. Data points are from the Northern part of RX J0852.0-4622 as measured with *H.E.S.S.* (Aharonian et al., 2007a), with an analysis point spread function of Gaussian width 0.06° . The *dashed line* represents the calculated profile convolved with the same point spread function.

The line-of-sight integrated γ -ray emission profile as a function of projected radius ρ is calculated for $\epsilon_{\gamma}=1$ TeV and is presented in Fig. 7. Due to the large radial gradient of the gas and CR distributions inside the SNR, the theoretically predicted three-dimensional radial emissivity profile of TeV-emission is concentrated within a thin shell of width $\Delta r \sim 0.01 R_s$. As a result of the projection effect, the two-dimensional width is by a factor of seven larger than the width of the three-dimensional profile, i.e. $\Delta \rho \approx 0.1 R_s$ (solid lines in Fig. 7). Since the *H.E.S.S.* instrument in addition has a finite angular resolution we present as dashed lines in Fig. 7 also the modified radial profile convolved with a Gaussian point spread function with a $\sigma_{\rho} = \Delta \rho = 0.06 R_s$, corresponding to an angular resolution of $\sigma_{\psi} = 0.06^{\circ}$.

Fig. 7 shows that the expected radial profile of the TeV-emission, after taking into account the instrumental angular resolution, is much broader than the intrinsic projected profile and is characterised by a minimum-to-maximum intensity ratio $J_{\gamma}^{\rm min}/J_{\gamma}^{\rm max}\approx 0.35$. The radial profile measured by *H.E.S.S.* compares reasonably well with the theoretical prediction if we ignore the two data points in the central region. This might be justified because a central component (e.g. a PWN) cannot be fully excluded, as was also argued in Aharonian et al. (2007a).

To illustrate the expected time evolution of the nonthermal emission we give in Fig. 8 the fluxes of radio emission at the frequency $\nu=1$ GHz and X-ray emission at the energy $\epsilon_{\nu}=1$ keV, as well as the TeV flux, all as a function of time. Since the CR energy content still increases, the radio emission increases during the next centuries. However, due to the substantial shock deceleration the nonthermal X-ray emission is expected to decrease with time at a rate of about 0.04%/yr. The TeV γ -ray emission, on the other hand, will increase at a rate of 0.14 %/yr due to the continuing increase of the ambient gas density. Nevertheless, a change of the order of one percent in 7 years is probably very difficult to measure. Therefore we can not put much practical




Fig. 8. The time dependence of the fluxes of the synchrotron emission at frequency $\nu=1\,\mathrm{GHz}$ (dash-dotted line), synchrotron X-ray emission with energy $\epsilon_{\nu}=1\,\mathrm{keV}$ (dashed line) and TeV-energy γ -ray (solid line). The fluxes are normalised to their values F_0 at the current epoch.

weight on these expectations regarding the secular variation of nonthermal fluxes.

4.1. Escape

The decrease of the shock speed $V_{\rm s}$ during SNR evolution diminishes the maximum energy to which particles can be accelerated during any given phase. This has the well-known consequence that particles already accelerated to a higher maximum energy at earlier times can now begin to diffusively escape from the interior of the remnant, even without any change of the scattering mean free path, as it was predicted analytically (Berezhko & Krymsky, 1988) and confirmed numerically (Berezhko et al., 1996).

Compared with this escape – which is relatively slow in our model because the diffusion of CR particles with momenta $p > p_{\rm m}(t)$ higher than the current cutoff momentum $p_{\rm m}(t)$ is still described by Bohm diffusion - the actual CR escape is presumably much faster. Since the CRs with $p>p_{
m m}(t)$ develop a much smaller spatial gradient than those with momenta $p < p_{\rm m}(t)$ (which are still efficiently accelerated by the shock), their production of chaotic magnetic fields whose scales are approximately equal to their gyroradii goes down quickly with time. As a consequence their diffusion coefficient increases considerably so that they leave the SNR much more rapidly than through Bohm diffusion, and then these CRs contribute much less to the γ -ray flux of the source than calculated above. The γ -ray spectrum corresponding to the particles that remain confined in the SNR should therefore drop off faster with energy than given by the solid line in Fig.4. However, the H.E.S.S. data at the highest γ -ray energies are not suggesting that this effect is very significant in the present case.

Secondly, the field amplification is probably stronger during earlier phases of the remnant evolution, when $\rho_0 V_{\rm s}^2$ is larger than at later phases (Bell & Lucek, 2001). Our calculation has assumed the effective field strength to be constant. Correcting for this, the overall nuclear particle spectrum released into the Interstellar Medium by SNRs presumably forms the Galactic

CR spectrum up to an energy of 10^{17} eV (Berezhko & Völk, 2007). That protons alone should be able to reach energies $\gtrsim 10^{15}$ eV was shown for the case of a representative type Ia SN by Berezhko & Völk (2004b). This is likely to be true also for RX J0852.0-4622 and, by implication, also for its twin RX J1713.7-3946, even though their environmental conditions are drastically different from those of a type Ia SN.

We add here that the above maximum energies of energetic nuclei are calculated under the assumption of Bohm diffusion. To this extent they are upper limits, given the strength of the corresponding amplified magnetic field.

Except in special cases one should therefore expect that escape sets in when the source becomes older. Ptuskin & Zirakashvili (2003, 2005) have in addition argued that at late times the damping of the scattering magnetic fluctuations should increase the scattering mean free path. The decrease of the effective field strength on the one hand, and of the scattering strength (through the slowing-down of the shock and wave damping) on the other, go in the same direction to lower the cutoff energy of the accelerating population. The relative importance of these effects probably depends on the details of the source, in particular its evolutionary phase.

4.2. A rough estimate for the thermal X-ray emission

Using the results for the dynamical evolution of the system one can also attempt to estimate the thermal X-ray emission. As argued in section 3.1.4 this is only possible in an approximate way, even if the ejecta emission is disregarded with the argument that the ejected mass in the present phase is small compared to the swept-up mass. In fact, looking at Fig. 1a it can be seen that the SNR is well past the sweep-up phase and has entered a quasi-Sedov phase in the stellar wind shell, i.e. a roughly self-similar evolutionary phase, modified by strong particle acceleration relative to a purely gas dynamic evolution.

The approximation used is the following. The bubble case is compared with a SNR in a uniform medium in the classical Sedov phase without any CR acceleration, making four assumptions (i) the total hydrodynamic explosion energy is the same in both cases (ii) the downstream unmodified temperature $T_{\rm s}$, determining the overall shock velocity and the thermal emission, is the same, (iii) the present gas density upstream of the shock is the same, and (iv) the two objects are at the same distance of 1 kpc. Then the results of Hamilton et al. (1983) for the thermal X-ray flux from a classical Sedov SNR are used, employing the emission measure of the bubble remnant instead of that of the classical Sedov remnant with the same five parameters above. This means that the X-ray emissivity of the remnant is reduced by the ratio $R_{\rm em} = E M_{\rm b}/E M_{\rm S}$ of the emission measure $E M_{\rm b}$ for the bubble solution to the emission measure for the classical Sedov solution EM_S which corresponds to a uniform ambient gas density $N_{\rm g}(R_{\rm s})$.

The thermally relevant gas temperature in the presence of particle acceleration is the temperature $T_{\rm sub}$ downstream of the gas subshock. It corresponds to the temperature downstream of a shock that is nonlinearly modified by the internal energy and the pressure of the accelerated particles.

The CR-modified shock, i.e. the precursor-subshock system, is approximated by a plane parallel structure, which implies that the precursor size is small compared to the SNR shock radius. This is generally the case. Then conservation of mass, momentum and energy fluxes permits in a straightforward way to calculate the ratio between the actual downstream gas temperature

 $T_{
m sub}$ and the downstream gas temperature $T_{
m s}$ that would obtain without particle acceleration:

$$\frac{T_{\text{sub}}}{T_{\text{s}}} = \frac{[(\sigma_{\text{s}} - 1)(\gamma + 1) + 2](\gamma + 1)^{2}}{4\gamma(\gamma - 1)\sigma^{2}}.$$
(14)

Here σ and $\sigma_{\rm s}$ are the total and the subshock compression ratio, respectively, as given in Fig.1. The specific heat ratio γ for the nonrelativistic thermal gas may be taken as $\gamma=5/3$. For the present phase of RX J0852.0-4622 , i.e. $\sigma\approx5.2$ and $\sigma_{\rm s}\approx3.1$, one then obtains $T_{\rm sub}\approx0.45T_{\rm s}$.

Hamilton et al. (1983) give the gas temperature T_s by

$$T_{\rm s} = 10^7 (V_{\rm s}/839 \,\mathrm{km \, s^{-1}})^2 \mathrm{K}$$
 (15)

in terms of the overall shock velocity $V_{\rm s}$, assuming a highly ionized system. For RX J0852.0-4622 the latter quantity is given in our Fig.1a. Using the present value $V_{\rm s}=1316$ km/sec, the foregoing equations result in $T_{\rm s}\approx2.5\times10^7$ K and $T_{\rm sub}\approx1.1\times10^7$ K, corresponding to a thermal energy of about 1 keV.

For the emission measure EM we have $EM=\int_0^{R_{\rm s}} dr r^2 N_{\rm g}^2(r)$, if we disregard irrelevant numerical factors. In order to calculate $R_{\rm em}$ for the CR-modified shock the overall shock is approximated by a shock in a thermal gas with an adiabatic index $\gamma_{\rm cr}$ such that $(\gamma_{\rm cr}+1)/(\gamma_{\rm cr}-1)=\sigma$, where σ denotes, as before, the total compression ratio of the CR-modified shock. The corresponding solution for the gas dynamic quantities is approximated as being self-similar cf. Sedov (1959). If the ambient density profile has a radial dependence in the form of a power law $N_{\rm g}(r)=N_0(r/R_0)^\beta$, then the self-similar density distribution in the SNR, downstream of the shock, can be further approximated by a power-law profile, with the same swept-up mass as the full solution. It has then the form $N_{\rm g}(r)=\sigma N_0(r/R_{\rm s})^{\beta'}$ with $\beta'=3(\sigma-1)+\beta\sigma$, where $N_0=N_{\rm g}(R_{\rm s})$. Such a density profile gives the emission measure $EM=N_0^2R_{\rm s}^3\sigma^2/(2\beta'+3)$.

The classical gas dynamic Sedov solution for a SN explosion into a uniform medium has $\beta=0$ and $\sigma=4$. Using the same values for N_0 for both density profiles leads to $R_{\rm s}\approx 12.51\,{\rm pc}$ for the Sedov case and to $R_{\rm em}\approx 0.63$ for the bubble case $\beta=12$ and $\sigma=5.2$. The swept-up mass in the Sedov case is $M_{\rm S}\approx 50.3\,M_{\odot}$.

Using the differential thermal X-ray model spectra $dF/d\epsilon \approx 10^{-4}$ photons/(keV cm²s) from Hamilton et al. (1983) (see their Fig. 2) for their $\lg T_s = 7.75$ and $\lg T_s = 7.25$, $\eta = N_{\rm H}^2 E_{\rm sn} = 10^{49}$ erg cm⁻⁶, scaling it according to $dF/d\epsilon \propto \eta$ with a factor of ≈ 3.8 for $\eta \approx 3.8 \times 10^{49}$ erg cm⁻⁶, multiplying it by the factor $\theta^2 [E_{\rm sn}/(10^{51}{\rm erg})]^{-1/2} \approx 0.65 \times 10^4$, as required, where $\theta \approx 86$ arcmin for the angular size of the classical Sedov remnant corresponding to the above parameters (Hamilton et al., 1983), and multiplying it finally also by the factor $R_{\rm em}$, results in a thermal spectral energy density $\epsilon^2 dF/d\epsilon \approx 1560~{\rm eV}~{\rm cm}^{-2}~{\rm sec}^{-1}$ for $\epsilon = 1~{\rm keV}$.

This must be compared with the observed nonthermal X-ray energy flux at 1 keV (see Fig.3) $\epsilon^2 dF/d\epsilon \approx 100$ eV cm $^{-2}$ sec $^{-1}$. Therefore, at 1 keV, the thermal flux, calculated in this form, comes out to be larger than the nonthermal flux by a factor of about 16.

We note however that the above X-ray energy flux has to be considered as a rough upper limit estimate because of the following reasons:

First of all, the SN shock into the bubble and shell interacts with a strongly rising gas density profile. Yet for the estimate the peak value of the gas density was used.

Moreover, in the case of a modified shock the actual gas temperature $T_{\rm sub}$ is at least by a factor of 2 lower than the temperature T_s used in our estimate. From a general point of view the actual thermal X-ray emission is expected to be lower due to lower temperature. A rough correction can be done in the following way. The shock radius corresponding to the Sedov solution depends on the relevent parameters according to the relation $R_s \propto (E_{SN}/(T_sN_{\rm H}))^{1/3}$. According to this relation the shock with the same radius but with two times lower postshock temperature corresponds to a two times lower explosion energy. Such a shock is characterised by the parameter value $\eta \approx 1.9 \times 10^{49}$ erg cm⁻⁶ that gives a 1 keV thermal X-ray flux which is lower by a factor of two compared with the above consideration.

Third, as it was already mentioned, the calculation in section 4 was performed for a rather moderate value of the injection rate, which leads to a moderate shock modification. Higher injection rates can not be excluded given the present data which can lead to a considerably higher shock modification with significantly lower gas energy and gas temperature. For a comparatively extreme position, see the recent paper by Drury et al. (2008).

Fourth, for a fixed $\gamma\text{-ray flux }F_{\gamma} \propto N_H E_{SN}/d^2$ the required gas number density scales as $N_H \propto d^2$ with source distance d. Consequently a decrease of d requires a decrease of the gas density. The thermal X-ray luminosity is quite sensitive to the gas density and to the total gas energy because roughly $dF/d\epsilon \propto N_H^2 E_{SN}$. A somewhat smaller distance than the d = 1 kpc, assumed here, can not be excluded.

Finally, the bubble solution developed in this paper has quite a different temperature profile than any "equivalent" classical Sedov solution used to estimate the thermal emission. This arises from the approximate uniformity of the total pressure in the SNR interior which roughly implies $T_{\rm sub} \propto N_{\rm H}^{-1}$. The density profiles are indeed quite different in both cases.

As a result, we believe that the uncertainties regarding the thermal X-ray emission are quite large. This situation also implies some systematic error in the overall model presented, although this can hardly change its key properties. In our opinion the uncertainty in the thermal emission has to be resolved by future work that extends the efforts of Hamilton et al. (1983) to more general circumstellar density profiles and ejected masses for core-collapse supernovae.

5. Conclusions

We have argued in the last section that a solution behind the Vela SNR, involving the core collapse of a massive star in its own wind bubble, describes the available data reasonably well. The physical characteristics of this solution are very similar to those of the well-known SNR RX J1713.7-3946. In this sense RX J0852.0-4622 and SNR RX J1713.7-3946 are twins.

The calculated γ -ray spectrum has a smooth cutoff at higher energies. The presently observable lower limit for the magnetic field amplification – in terms of thin shock filaments in hard Xrays – is consistent with the one deduced from a theoretical fit to the observed synchrotron spectra. If anything, the spectrally deduced field amplification is lower than the filament-deduced field amplification. Overall, the field amplification is also consistent with the semi-empirical relation Eq.(2) between the CR pressure $P_{\rm c}$ and the magnetic pressure, where $P_{\rm c}$ stems from the proton injection rate required to fit the observed γ -ray emission. We conclude from these consistency arguments that the observational lack of a detailed radio synchrotron spectrum does not preclude the determination of a consistent amplified field.

The magnetic field amplification results in a significant depression of the density of ultra-high energy electrons and reduces the IC and NB contributions of these electrons to less than one percent of the π^0 -decay γ -ray emission.

A remaining uncertainty is connected with the thermal emission properties. For the wind bubble configuration this may indeed not be too critical, since gas heating should occur primarily at the subshock alone. In addition, the amount of swept-up mass is rather small compared to a classical Sedov remnant in a uniform circumstellar medium of equal present preshock density.

Apart from this uncertainty the main result is the hadronic dominance in the γ -ray emission spectrum. As a consequence of the nonlinear modification of the shock the spatially integrated γ -ray spectral energy distribution at 1 GeV is predicted to be at best a factor 1.5 lower than at 1 TeV.

Acknowledgements. This work has been supported in part by the Russian Foundation for Basic Research (grants 06-02-96008, 07-02-0221). The authors thank V.S. Ptuskin and V.N. Zirakashvili and the anonymous referee for discussions on the thermal emission properties. EGB acknowledges the hospitality of the Max-Planck-Institut für Kernphysik, where part of this work was carried out.

References Abbott, D.C. 1982, ApJ, 263, 723 Aharonian, F.A. et al. (H.E.S.S. Collaboration) 2005, A&A, 437, L7 Aharonian, F.A. et al. (H.E.S.S. Collaboration) 2007a, ApJ, 661, 236 Aharonian, F.A. et al. (H.E.S.S. Collaboration) 2007b, A&A, 464, 235 Aharonian, F.A. et al. (H.E.S.S. Collaboration) 2008, to appear in A&A; arXiv:0810.2689 Aschenbach, B. 1998, Nature, 396, 141 Aschenbach, B., Iyudin, A. F. & Schönfelder, V. 1999, A&A, 350, 997. Bamba, A., Yamazaki, R. & Hiraga, J.S. 2005, ApJ, 632, 294 Becker, W. & Aschenbach, B., Proc. of the 270. WE-Heraeus Seminar on Neutron Stars, Pulsars, and Supernova Remnants, Ed.: W. Becker, H. Lesch & J. Trümper, MPE report 278, 64, available from ADS and astro-ph/0208492 Becker, W., Hui, C. Y., Aschenbach, B. & Iyudin, A. F. 2007, submitted to A&A, astro-ph/0607081 Bell, A.R. 1978, MNRAS, 182, 147 Bell, A. R. & Lucek, S.G. 2001, MNRAS, 321, 433 Bell, A. R. 2004, MNRAS, 327, 433 Berezhko, E.G. 1996, Astropart. Phys., 5, 367 Berezhko, E.G. 2005, Adv. Space Res., 35, 1031 Berezhko, E.G. 2008, Adv. Space Res., 41, 429 Berezhko, E.G. & Krymsky, G.F. 1988, Soviet Phys.-Uspekhi, 12, 155 Berezhko, E.G., Elshin, V.K. & Ksenofontov, L.T. 1996 JETPh, 82, 1 Berezhko, E.G. & Völk, H.J. 1997, Astropart. Phys., 7, 183

Berezhko, E.G. & Völk, H.J. 2000, A&A, 357, 183 Berezhko, E. G., Ksenofontov, L. T. & Völk, H. J. 2002, A&A, 395, 943

Berezhko, E. G., Ksenofontov, L. T. & Völk, H. J. 2003, A&A, 412, L11 Berezhko, E. G., Pühlhofer, G. & Völk, H. J. 2003, A&A, 400, 971

Berezhko, E.G. & Völk, H.J. 2004a, A&A, 419, L27 Berezhko, E.G. & Völk, H.J. 2004b, A&A, 427, 525

Berezhko, E.G. & Ksenofontov, L.T. 1999, JETPh 89, 391

Berezhko, E.G. & Völk, H.J. 2006, A&A, 451,981 Berezhko, E.G. & Völk, H.J. 2007, ApJ, 661, L175

Blandford, R.D. & Ostriker, J.P. 1978, ApJ, 221, L29

Borkowski, K.J., Lyerly, W.J. & Reynolds, S.P. 2001, ApJ, 548, 820

Caraveo, P.A., De Luca, A., Mignani, R.P., & Bignami, G.F. 2001, ApJ, 561, 930 Cha, A. N., Sembach, K. R. & Danks, A. C. 1999, ApJ, 1999, 515, L25

Chevalier, R.A. 1982, ApJ, 258, 790

Chevalier, R., & Liang, P. 1989, ApJ, 344, 332

Combi, J. A., Romero, G. E. & Benaglia, P. 1999, ApJ, 519, L177

Dodson, R., Legge, D., Reynolds, J.E., & McCulloch, P.M. 2003, ApJ, 596, 1137

Drury, L.O'C., Aharonian, F.A. & Völk, H.J. 1994, A&A, 287, 959

Drury, L.O'C., Aharonian, F.A., Malyshev, D. & Gabici, S. 2008, A&A, 496, 1

Duncan, A.R. & Green, D.A. 2000, A&A, 364, 732

Ellison, D.C., Baring, M.G., & Jones, F.C. 1995, ApJ, 453, 873

Enomoto, R., Tanimory, T., Naito, T., et al. 2002, Nature, 416, 823

Enomoto, R., Watanabe, S., Tanimori, T., et al. 2006, ApJ, 652, 1268 Hamilton, A.J.S., Sarazin, C.L. & Chevalier, R.A. 1983, ApJS, 41, 115

Hamilton, A.J.S. & Sarazin, C.L. 1984, ApJ, 284, 601

Hoppe, S. & Lemoine-Goumard, M. (HESS Collaboration) 2008, Proc. 30th ICRC (Merida, 2007), 2, 255; arXiv:0709.4103(astro-ph)

```
Iyudin, A. F., Diehl, R., Bloemen, H., et al., A&A, 1994, 284, L1
```

Iyudin, A. F., Schönfelder, V., Bennett, K., et al. 1998, Nature, 396, 142

Iyudin, A. F., Aschenbach, B., Becker, W.et al. 2005, A&A, 429, 225

Iyudin, A. F., Aschenbach, B., Burwitz, V., et al. 2007, astro-ph/0702404

Jones, E.M., Smith, B.W. & Straka, W.C. 1981, ApJ, 249, 185

Kargaltsev, O., Pavlov, G. G., Sanwal, D. & Garmire, G. P. 2002, ApJ, 580, 1060

Katagiri, H., Enomoto, R., Ksenofontov, L. T., et al. (CANGAROO collaboration) 2005, ApJ, 619, L163

von Kienlin, A., et al., in Proc 5th INTEGRAL Science workshop, 2005 (ESA SP-552; Noordwijk: ESA), astro-ph/0407129

Lu, F. J. & Aschenbach, B. 2000, A&A, 362, 1083

Lucek, S.G. & Bell, A.R. 2000, MNRAS, 314, 65

Malkov, M.A. & Völk, H.J. 1995, A&A, 300, 605

McKenzie, J.F. & Völk, H.J. 1982, A&A, 116, 191

Mereghetti, S., ApJ, 2001, 548, L213

Mereghetti, S. & Pellizzoni, A., De Luca, A., in *Neutron Stars in Supernova Remnants*, ASP Conf. Ser. 271, ed. P. O. Slane & B. M. Gaensler (San Francisco: ASP), 289, astro-ph/0112152

Mignani, R. P., De Luca, A., Zaggia, S., et al. 2007, submitted to A&A, astro-ph/0707.0937

Milne, D. K. 1968, Australian J. Phys., 21, 201

Moriguchi, Y., Yamaguchi, N., Onshi, T., et al. 2001, Publ. Astron. Soc. Japan, 53, 1025

Naumann-Godo, M. et al. (H.E.S.S. Collaboration) 2009, in *High Energy Gamma-Ray Astronomy* (Eds. F.A. Aharonian, W. Hofmann, F.M. Rieger), Melville, New York, 2009, AIP Conf. Proc. 1085, p. 304 ff.

Niemic, J., Pohl, M., Stroman, T., et al. 2008, ApJ, 684, 1174

Pastorello, A., Zampieri, L., Turatto, M. et al. 2004, MNRAS, 347, 74

Pavlov, G. G., Sanwal, D., Kiziltan, B. & Garmire, G. P. 2001, ApJ, 559, L131

Pannuti, T. G., Keohane, J. W., Allen, G. E. et al., Goldstein, G., in X-ray and Radio connections (eds. L. O. Sjouwerman and K. K. Dyer), published electronically by NRAO, http://www.aoc.nrao.edu/events/xraydio, 2004, available also at ADS.

Pelletier, G., Lemoine, M., Marcowith, A. 2006, A&A, 453, 181

Pellizzoni, A., Mereghetti, S. & De Luca, A. 2002, A&A, 393, L65

Pohl, M., Yan, H. & Lazarian, A. 2005, ApJ, 626, L101

Ptuskin, V.S. & Zirakashvili, V.N. 2003, A&A, 410, 189

Ptuskin, V.S. & Zirakashvili, V.N. 2005, A&A, 429, 755

Redman, M. P. & Meaburn, J. 2005, MNRAS, 356, 969

Renaud, M., Vink, J., Decourchelle, A. et al. 2006a, New Astronomy Rev., 50, 540

Renaud, M., Gros, A., Lebrun, F., et al., A&A, 2006b, 456, 389

Renaud, M., Vink, J., Decourchelle, A., et al., ApJ., 2006c, 647, 41

Reynoso, E. M., Dubner, G., Giacani, E., et al. 2006, A&A, 449, 243

Schönfelder, V., Bloemen, H., Collmar, W., et al., 2000, In: AIP Conference Series, vol. 510, 54

Sedov, L.I. 1959, Similarity and Dimensional Methods in Mechanics, 4th Russian Edition (translated by M. Holt and M. Friedman), Academic Press Inc., New York, NY, Chapter IV ff.

Slane, P., Hughes, J.P., Edgar, R.J., et al. 2001, ApJ, 548, 814

Stupar, M., Filipović, M. D., Jones, P. A. & Parker, Q. A. 2005, Adv. Space Res., 35, 1047

Uchiyama, Y. 2008, Invited talk at 37th COSPAR Scientific Assembly, Montreal, Canada (2008), p. 3248

Vink, J., Laming, J. M., Kaastra, J. S., et al., 2001, ApJ, 560, L79

Völk, H.J., 2004, Proc. 28th ICRC (Tsukuba), Invited Papers, 8, 29

Völk, H. J. 2008, Invited talk at 37th COSPAR Scientific Assembly Montreal, Canada (2008), p. 3356

Völk, H.J., Berezhko, E.G., Ksenofontov, L.T. & Rowell, G.P. 2002, A&A, 396, 971

Völk, H.J., Berezhko, E.G. & Ksenofontov, L.T. 2003, A&A, 409, 563

Völk, H.J., Berezhko, E.B. & Ksenofontov, L.T. 2005, A&A, 433, 229

Völk, H.J., Berezhko, E.B. & Ksenofontov, L.T. 2008, A&A, 483, 529

Weaver, R., McCray, R., Castor, J., et al. 1977, ApJ, 218, 377

Wietfeldt, F. E., Schima, F. J., Coursey, B. M., & Hoppes, D. D. 1999, Phys. Rev. C, 59, 528

Zirakashvili, V.N. & Ptuskin, V.S. 2008, ApJ, 678, 939

Zirakashvili, V.N., Ptuskin, V.S. & Völk, H.J. 2008, ApJ 678, 225