AN ASYMPTOTIC VERSION OF A THEOREM OF KNUTH

JONATHAN NOVAK

1. INTRODUCTION

In this note we consider the asymptotics of the number S(d, N) of permutations in the symmetric group $\mathfrak{S}(N)$ which have no decreasing subsequence of length d+1, in the limit where $d \geq 2$ is a fixed but arbitrary positive integer and $N \to \infty$. This is a fundamental problem in the subject of pattern avoidance in permutations, see [1] and [10, §7]. In the interest of brevity, familiarity with Young diagrams, Young tableaux, and the Robinson-Schensted-Knuth (RSK) correspondence is assumed. The reader is referred to Stanley's survey [10] for the necessary background and further references. We adhere to the notation and terminology of [10] save for the following exceptions: the $d \times q$ rectangular Young diagram is denoted R(d,q) rather than (q^d) , and the number of standard Young tableaux of shape λ is denoted dim λ rather than f^{λ} . Recall that the dimension of a Young diagram may be computed from Frobenius' fomula:

(1)
$$\dim \lambda = \frac{\Gamma(\lambda_1 + \dots + \lambda_d + 1)}{\prod_{i=1}^d \Gamma(\lambda_i - i + d + 1)} \prod_{1 \le i < j \le d} (\lambda_i - \lambda_j + j - i),$$

where d is any number such that $\lambda_{d+1} = 0$ and $\Gamma(z)$ is the gamma function.

The following well-known exact formula for S(2, N) is attributed to Hammersley in [10], with the first published proof due to Knuth [6, §5.1.4].

Theorem 1.1 (Knuth). Permutations with no decreasing subsequence of length 3 are counted by the Catalan numbers:

$$S(2, N) = \dim R(2, N) = \frac{(2N)!}{N!(N+1)!}$$

For d > 2, there is no known closed formula for S(d, N). The main result of this note is the following asymptotic version of Knuth's theorem.

Theorem 1.2 (Asymptotic Knuth theorem). For d fixed and $n \to \infty$,

$$S(d, dn) \sim \dim R(d, 2n)$$

We will see below that $S(d, dn) > \dim R(d, 2n)$ as soon as d > 2, so that Theorem 1.2 is false for n finite.

Via the RSK correspondence, an equivalent formulation of Theorem 1.2 is the following.

Theorem 1.3. The number of permutations in $\mathfrak{S}(dn)$ with no decreasing subsequence of length d + 1 is asymptotically equal, as $n \to \infty$, to the number of involutions in $\mathfrak{S}(2dn)$ with longest decreasing subsequence of length exactly d and longest increasing subsequence of length exactly 2n.

By Frobenius' formula, we have

(2)
$$\dim R(d,q) = \frac{\Gamma(dq+1)}{\prod_{i=1}^{d} \frac{\Gamma(q+i)}{\Gamma(i)}}.$$

Thus

(3)
$$\dim R(d,q) \sim (2\pi)^{\frac{1-d}{2}} \left(\prod_{i=1}^{d} \Gamma(i)\right) d^{dq+\frac{1}{2}} q^{\frac{1-d^2}{2}}$$

as $q \to \infty$ with d fixed, by Stirling's formula. Setting q = 2N/d, Theorem 1.2 together with (3) immediately implies the following.

Corollary 1.4. For d fixed and $N \to \infty$,

$$S(d,N) \sim (2\pi)^{\frac{1-d}{2}} \left(\prod_{i=1}^{d} \Gamma(i)\right) d^{2N+\frac{d^2}{2}} (2N)^{\frac{1-d^2}{2}}.$$

Corollary 1.4 was first obtained by A. Regev [8, Formula F.4.5.2] in 1981 by a rather different method, which will be discussed below.

2. Decomposition of rectangular tableaux

Given a Young diagram $\mu \subseteq R(d,q)$, let

(4)
$$\mu^* = (q - \mu_d, \dots, q - \mu_1)$$

denote the complement of μ relative to R(d,q). Clearly,

(5)
$$\dim R(d, 2n) = \sum_{\substack{\mu \vdash dn \\ \mu \subseteq R(d, 2n)}} (\dim \mu) (\dim \mu^*) \\ = \sum_{\substack{\mu \vdash dn \\ \mu \subseteq R(d, 2n) \\ \mu = \mu^*}} (\dim \mu)^2 + \sum_{\substack{\mu \vdash dn \\ \mu \subseteq R(d, 2n) \\ \mu \neq \mu^*}} (\dim \mu) (\dim \mu^*).$$

On the other hand, by RSK, we have

(6)
$$S(d, dn) = \sum_{\substack{\lambda \vdash dn \\ \ell(\lambda) \le d}} (\dim \lambda)^2 \\ = \sum_{\substack{\mu \vdash dn \\ \mu \subseteq R(d, 2n) \\ \mu = \mu^*}} (\dim \mu)^2 + \sum_{\substack{\mu \vdash dn \\ \mu \subseteq R(d, 2n) \\ \mu \neq \mu^*}} (\dim \mu)^2 + \sum_{\substack{\nu \vdash dn \\ \nu_1 > 2n \\ \ell(\nu) \le d}} (\dim \nu)^2.$$

Substituting for the first group of terms in (6) using (5) and completing the square yields the following.

Proposition 2.1. $S(d, N) = \dim R(d, 2n) + E(d, n)$, where the error term is given by

$$E(d,n) = \frac{1}{2} \sum_{\substack{\mu \vdash dn \\ \mu \subseteq R(d,2n)}} (\dim \mu - \dim \mu^*)^2 + \sum_{\substack{\nu \vdash dn \\ \nu_1 > 2n \\ \ell(\nu) \le d}} (\dim \nu)^2.$$

Clearly E(2, n) = 0, in agreement with Knuth's theorem, while E(d, n) > 0 for d > 2. Nevertheless, the error term is negligible in the limit $n \to \infty$.

3. Asymptotics of the dimension function

In order for E(d, n) to be negligible, the sum S(d, dn) must be dominated in the $n \to \infty$ limit by self-complementary diagrams contained in the rectangle R(d, 2n). The canonical self-complementary diagram relative to R(d, 2n) is the $d \times n$ rectangle R(d, n). We consider the asymptotics of diagrams which deviate from R(d, n) on the scale \sqrt{n} ; this choice of scale emerges constructively in the proof of the following key Lemma.

Lemma 3.1. For any distinct real numbers

$$y_1 > \cdots > y_d$$

satisfying

$$y_1 + \dots + y_d = 0,$$

 $we\ have$

$$\lim_{n \to \infty} C_{d,dn} \dim(n + y_1 \sqrt{n}, \dots, n + y_d \sqrt{n}) = e^{-W(y_1,\dots,y_d)}$$

where

$$C_{d,dn} = (2\pi)^{\frac{d}{2}} \frac{n^{dn + \frac{d(d+1)}{4}}}{\Gamma(dn+1)e^{dn}}$$

and

$$W(y_1, \dots, y_d) = \frac{1}{2} \sum_{i=1}^d y_i^2 - \sum_{1 \le i < j \le d} \log(y_i - y_j).$$

Proof. Let $0 < \varepsilon < 1$, and consider deviations from R(d, n) on the scale n^{ε} . By the Frobenius formula, we have

$$\dim(n+y_1n^{\varepsilon},\ldots,n+y_dn^{\varepsilon}) = \frac{\Gamma(dn+1)}{\prod_{i=1}^d \Gamma(n+y_in^{\varepsilon}+d-i+1)} \prod_{1 \le i < j \le d} ((y_i-y_j)n^{\varepsilon}+j-i)$$

for n sufficiently large.

Let us first analyze the asymptotics of the product

$$\frac{1}{\prod_{i=1}^{d} \Gamma(n+y_i n^{\varepsilon} + d - i + 1)}.$$

We begin by noting that $\Gamma(n+y_in^{\varepsilon}+d-i+1) \sim n^{d-i+1}\Gamma(n+y_in^{\varepsilon})$, so that

$$\frac{1}{\prod_{i=1}^{d} \Gamma(n+y_i n^{\varepsilon}+d-i+1)} \sim \frac{1}{n^{\frac{d(d+1)}{2}} \prod_{i=1}^{d} \Gamma(n+y_i n^{\varepsilon})}$$

Taking logarithms yields

$$\log\left(\frac{1}{\prod_{i=1}^{d}\Gamma(n+y_{i}n^{\varepsilon}+d-i+1)}\right) \sim -\frac{d(d+1)}{2}\log n - \sum_{i=1}^{d}\log\Gamma(n+y_{i}n^{\varepsilon}).$$

Recall Stirling's formula:

$$\log \Gamma(N) \sim \frac{1}{2} \log 2\pi + (N - \frac{1}{2}) \log N - N$$

for N large. Thus

$$\sum_{i=1}^d \log \Gamma(n+y_i n^\varepsilon) \sim \frac{d}{2} \log 2\pi - dn + \sum_{i=1}^d (n+y_i n^\varepsilon - \frac{1}{2}) \log(n+y_i n^\varepsilon).$$

Now since

$$\log(n+y_i n^{\varepsilon}) = \log n + \log(1+y_i n^{\varepsilon-1}),$$

we have

$$\sum_{i=1}^{d} (n+y_i n^{\varepsilon} - \frac{1}{2}) \log(n+y_i n^{\varepsilon}) = dn \log n - \frac{d}{2} \log n + \sum_{i=1}^{d} (n+y_i n^{\varepsilon} - \frac{1}{2}) \log(1+y_i n^{\varepsilon-1}).$$

Using the expansion

$$\log(1 + y_i n^{\varepsilon - 1}) = y_i n^{\varepsilon - 1} - \frac{1}{2} y_i^2 n^{2\varepsilon - 2} + O(n^{3\varepsilon - 3})$$

for n sufficiently large, we find that

$$\sum_{i=1}^{d} (n+y_i n^{\varepsilon} - \frac{1}{2}) \log(1+y_i n^{\varepsilon-1}) \sim \frac{n^{2\varepsilon-1}}{2} \sum_{i=1}^{d} y_i^2$$

Putting this all together, we find that

$$\frac{1}{\prod_{i=1}^{d} \Gamma(n+y_i n^{\varepsilon}+d-i+1)} \sim \frac{e^{dn}}{(2\pi)^{\frac{d}{2}} n^{dn+\frac{d^2}{2}}} e^{-\frac{n^{2\varepsilon-1}}{2} \sum_{i=1}^{d} y_i^2}$$

as $n \to \infty$.

The second group of factors is much easier to handle:

$$\prod_{1 \le i < j \le d} \left((y_i - y_j) n^{\varepsilon} + j - i \right) \sim n^{\varepsilon \frac{d(d-1)}{2}} \prod_{1 \le i < j \le d} (y_i - y_j).$$

Thus when $\varepsilon = \frac{1}{2}$ we have

$$\dim(n+y_1\sqrt{n},\ldots,n+y_d\sqrt{n}) \sim \frac{\Gamma(dn+1)e^{dn}}{(2\pi)^{\frac{d}{2}}n^{dn+\frac{d(d+1)}{4}}}e^{-W(y_1,\ldots,y_d)}$$

as $n \to \infty$, as claimed.

4. RIEMANN SUM

Consider the generalized sum

(7)
$$S(d, dn; \beta) = \sum_{\substack{\lambda \vdash dn \\ \ell(\lambda) \le d}} (\dim \lambda)^{\beta}, \quad \beta > 0,$$

and its presentation as a sum in parameters around the rectangle R(d, n):

(8)
$$S(d, dn; \beta) = \sum_{\substack{(d-1)n \\ \sqrt{n}} \ge y_1 \ge \dots \ge y_{d-1} \ge \frac{-n}{\sqrt{n}}} \dim(n + y_1 \sqrt{n}, \dots, n + y_d \sqrt{n})^{\beta},$$

where $y_d := -(y_1 + \dots + y_{d-1})$. The lattice $(\frac{1}{\sqrt{n}}\mathbb{Z})^{d-1}$ partitions \mathbb{R}^{d-1} into cells

(9)
$$\left(\frac{k_1}{\sqrt{n}}, \frac{k_1+1}{\sqrt{n}}\right) \times \cdots \times \left(\frac{k_{d-1}}{\sqrt{n}}, \frac{k_{d-1}+1}{\sqrt{n}}\right), \quad k_i \in \mathbb{Z},$$

of volume $(\frac{1}{\sqrt{n}})^{d-1}$. Scaling by the mesh volume makes this into a Riemann sum, and by Lemma 3.1 we have

(10)
$$\lim_{n \to \infty} \left(\frac{1}{\sqrt{n}}\right)^{d-1} C^{\beta}_{d,dn} S(d,dn;\beta) = \int_{\Omega_{d-1}} e^{-\beta W(y_1,\dots,y_d)} dy,$$

4

where $C_{d,dn}$ and W are as in Lemma 3.1 and $\Omega_{d-1} \subset \mathbb{R}^{d-1}$ is the region

(11)
$$\Omega_{d-1} = \{(y_1, \dots, y_{d-1}) \in \mathbb{R}^{d-1} : y_1 > \dots > y_d := -(y_1 + \dots + y_{d-1})\}.$$

The details of this convergence can be checked and made rigorous using the dominated convergence theorem; we refer the reader to [7, 8, 9] for the full argument.

Consider now the sum

(12)

$$\sum_{\substack{\mu \vdash dn \\ \mu \subseteq R(d,2n)}} (\dim \mu)^{\alpha} (\dim \mu^{*})^{\beta-\alpha}$$

$$= \sum_{\substack{\frac{n}{\sqrt{n}} \geq y_{1} \geq \cdots \geq y_{d-1} \geq \frac{-n}{\sqrt{n}} \\ y_{i} \in \frac{1}{\sqrt{n}} \mathbb{Z}}} \dim (n+y_{1}\sqrt{n}, \dots, n+y_{d}\sqrt{n})^{\alpha} \dim (n-y_{d}\sqrt{n}, \dots, n-y_{1}\sqrt{n})^{\beta-\alpha},$$

where $0 \le \alpha \le \beta$ and as before we denote $y_d := -(y_1 + \cdots + y_{d-1})$. This can again be viewed as a Riemann sum, and by exactly the same argument we have

(13)
$$\lim_{n \to \infty} \left(\frac{1}{\sqrt{n}}\right)^{d-1} C^{\beta}_{d,dn} \sum_{\substack{\mu \vdash dn \\ \mu \subseteq R(d,2n)}} (\dim \mu)^{\alpha} (\dim \mu^{*})^{\beta-\alpha}$$
$$= \int_{\Omega_{d-1}} e^{-\alpha W(y_{1},\dots,y_{d})} e^{-(\beta-\alpha)W(-y_{d},\dots,-y_{1})} dy.$$

. .

5. Symmetry

Note that the function W has the symmetry

(14)
$$W(y_1, \dots, y_d) = W(-y_d, \dots, -y_1).$$

It follows that

(15)
$$\int_{\Omega_{d-1}} e^{-\beta W(y_1,\dots,y_d)} dy = \int_{\Omega_{d-1}} e^{-\alpha W(y_1,\dots,y_d)} e^{-(\beta-\alpha)W(-y_d,\dots,-y_1)} dy,$$

and thus we have proved the following.

Theorem 5.1. For any $\beta > 0$ and $0 \le \alpha \le \beta$, we have

$$S(d, dn; \beta) \sim \sum_{\substack{\mu \vdash dn \\ \mu \subseteq R(d, 2n)}} (\dim \mu)^{\alpha} (\dim \mu^*)^{\beta - \alpha}$$

as $n \to \infty$.

Theorem 1.2 is the special case $\beta = 2, \alpha = 1$ of this more general asymptotic equivalence.

6. CONCLUSION

The multidimensional integral

(16)
$$\Psi(d;\beta) = \int_{\mathbb{R}^d} e^{-\frac{\beta}{2}\sum_{i=1}^d x_i^2} \prod_{1 \le i < j \le d} |x_i - x_j|^\beta dx, \quad \beta > 0,$$

J. I. NOVAK

is known as *Mehta's integral*. It is the partition function of a Coulomb gas of d identical point charges $x_1 > \cdots > x_d$ on the real line at inverse temperature β , with energy functional

(17)
$$W(x_1, \dots, x_d) = \frac{1}{2} \sum_{i=1}^d x_i^2 - \sum_{1 \le i < j \le d} \log(x_i - x_j).$$

Dyson and Mehta [3] studied this integral and conjectured the formula

(18)
$$\Psi(d;\beta) = (2\pi)^{\frac{d}{2}}\beta^{-\frac{d}{2}-\beta\frac{d(d-1)}{4}} \prod_{i=1}^{d} \frac{\Gamma(1+i\frac{\beta}{2})}{\Gamma(1+\frac{\beta}{2})},$$

which they verified for $\beta \in \{1, 2, 4\}$ using properties of Hermite polynomials (see e.g. [5, §3.5.1] for the $\beta = 2$ case of this argument). Later, Bombieri observed that, for general β , (18) can be deduced from the Selberg integral formula. See [4] for the interesting history of this problem.

Regev [8, Lemma 4.3] showed that

(19)
$$\int_{\Omega_{d-1}} e^{-\beta W(y_1,\dots,y_d)} dy = \frac{1}{\Gamma(d+1)} \sqrt{\frac{\beta}{2\pi d}} \Psi(d;\beta),$$

and used this fact together with equation (10) above to determine the asymptotics of $S(d, N; \beta)$ from the known form (18) of $\Psi(d; \beta)$. In this article, we have evaluated the asymptotics of S(d, N) = S(d, N; 2) directly, without appealing to the exact value of $\Psi(d; 2)$. Thus, we may obtain

(20)
$$\Psi(d;2) = (2\pi)^{\frac{d}{2}} 2^{-\frac{d^2}{2}} \prod_{i=1}^{d+1} \Gamma(i)$$

by substituting the asymptotic form of S(d, N) (Corollary 1.4) in (10) and using (19). It would be interesting to know if the asymptotics of $S(d, N; \beta)$ can be determined directly in a similar way for general $\beta > 0$. If so, this would yield a new and elementary verification of (18). In particular, for this purpose one may assume that β is an even integer, see [3].

Finally, in this note we have only considered the asymptotics of S(d, N) in the single scaling limit where $N \to \infty$ with d fixed. Baik, Deift, and Johansson [2] have shown that

$$(21) S(d,N) \sim F(t)N!$$

in the double scaling limit where $d, N \to \infty$ at the rate $d \sim 2\sqrt{N} + tN^{1/6}$, with $t \in \mathbb{R}$ fixed. Here F(t) is the Tracy-Widom distribution function, see [2]. Since

(22)
$$S(d, dn) = \dim R(d, 2n) + E(d, n),$$

if the error term E(d, n) can be effectively estimated in the double scaling limit then concrete estimates for F(t) will follow.

7. Acknowledgements

I would like to thank Sho Matsumoto and Andrei Okounkov for helpful correspondence, and Michael Albert for pointing out an error in an early version of this paper.

References

- [1] Adv. in Appl. Math. 36(2) (2006), special issue on pattern avoiding permutations.
- [2] J. Baik, P. Deift, and K. Johansson, On the distribution of the length of the longest increasing subsequence in random permutations, J. Amer. Math. Soc. 12(4), 1999, 1119-1178.
- [3] F. J. Dyson and M. L. Mehta, Statistical theory of the energy levels of complex systems, V., J. Math. Phys. 4(5), 1963, 713-719.
- [4] P. J. Forrester and S. O. Warnaar, The importance of the Selberg integral, Bull. Amer. Math. Soc. (N.S.), 45(4), 2008, 489-534.
- [5] S. K. Lando and A. K. Zvonkin, Graphs on Surfaces and Their Applications, Encyclopaedia of Mathematical Sciences, Volume 141, Springer-Verlag, 2004.
- [6] D. E. Knuth, The Art of Computer Programming, Volume 3: Sorting and Searching. Addison-Wesley, Reading, MA, 1973; second edition, 1998.
- [7] S. Matsumoto. Jack deformations of Plancherel measures and traceless Gaussian random matrices. *Electron. J. Combin.* 15 (2008), #R149.
- [8] A. Regev. Asymptotic values for degrees associated with strips of Young diagrams. Adv. Math. 41 (1981), 115-136.
- [9] P. Śniady. Permutations without long decreasing subsequences and random matrices. *Electron. J. Combin.* 14 (2007), #R11.
- [10] R. P. Stanley. Increasing and decreasing subsequences and their variants, Proceedings of the International Congress of Mathematicians, Madrid, Spain, 2006.

University of Waterloo, Faculty of Mathematics, Department of Combinatorics and Optimization, Waterloo, Ontario, Canada

E-mail address: j2novak@math.uwaterloo.ca *URL*: www.math.uwaterloo.ca/~j2novak