

# ALTERNATING GROUP AND MULTIVARIATE EXPONENTIAL FUNCTIONS

A. KLIMYK AND J. PATERA

ABSTRACT. We define and study multivariate exponential functions, symmetric with respect to the alternating group  $A_n$ , which is a subgroup of the permutation (symmetric) group  $S_n$ . These functions are connected with multivariate exponential functions, determined as the determinants of matrices whose entries are exponential functions of one variable. Our functions are eigenfunctions of the Laplace operator. By means of alternating multivariate exponential functions three types of Fourier transforms are constructed: expansions into corresponding Fourier series, integral Fourier transforms, and multivariate finite Fourier transforms. Alternating multivariate exponential functions are used as a kernel in all these Fourier transforms. Eigenfunctions of the integral Fourier transforms are obtained.

2000 Math. Subject Classif.: Primary 42B05; Secondary 20B30, 33E99, 42B10

## 1. INTRODUCTION

Mathematical and theoretical physics regularly deal with functions on the Euclidean space  $E_n$  that are symmetric or antisymmetric with respect to the permutation (symmetric) group  $S_n$ . For example, such functions describe collections of identical particles. Symmetric and antisymmetric solutions appear in the theory of integrable systems.

The symmetric group  $S_n$  contains the alternating group  $A_n$ . It is an invariant subgroup of  $S_n$  of index 2 (that is, the group  $S_n/A_n$  has 2 elements). The alternating group  $A_n$  consists of transformations  $w$  of the Euclidean space  $E_n$  with  $\det w = 1$  and, therefore, is a subgroup of the rotation group  $SO(n)$  (note that  $S_n$  does not belong to  $SO(n)$ ). The group  $A_n$  is simple. For studying multivariate exponential functions, the group  $A_n$  is more fundamental than the symmetric group, since multivariate exponential functions symmetric or antisymmetric with respect to  $S_n$  can be constructed by means of the multivariate exponential functions symmetric with respect to  $A_n$ .

The aim of this paper is to describe and study multivariate exponential functions symmetrized by the alternating group and the corresponding Fourier transforms. We call these functions *alternating multivariate exponential functions* and denote them by  $E_\lambda(x)$ ,  $\lambda = (\lambda_1, \lambda_2, \dots, \lambda_n)$ ,  $x = (x_1, x_2, \dots, x_n) \in E_n$ . Such a function is a sum of terms in the determinant of an  $n \times n$  matrix (whose entries are usual exponential functions of one variable) which enter in the expression for the determinant with sign  $+$ . We call this sum a semideterminant.

Alternating multivariate exponential functions are connected with symmetric and antisymmetric multivariate exponential functions studied in [1]. Symmetric

and antisymmetric multivariate exponential functions can be considered as a generalization of cosine and sine functions of one variable, respectively, whereas alternating multivariate exponential functions are a generalization of the usual exponential function of one variable. The connection of symmetric and antisymmetric multivariate exponential functions with alternating multivariate exponential functions is the same as that of the cosine and sine functions with the exponential function of one variable (see section 2 below).

As in the case of the exponential functions of one variable, we may consider three types of alternating multivariate exponential functions:

(a) functions  $E_m(x)$  with  $m = (m_1, m_2, \dots, m_n)$ ,  $m_i \in \mathbb{Z}$ , which determine Fourier series expansions in alternating multivariate exponential functions;

(b) functions  $E_\lambda(x)$  with  $\lambda = (\lambda_1, \lambda_2, \dots, \lambda_n)$ ,  $\lambda_i \in \mathbb{R}$ , which determine integral multivariate Fourier transforms;

(c) functions  $E_\lambda(x)$ , where  $x = (x_1, x_2, \dots, x_n)$  take a finite set of values; they determine multivariate finite Fourier transforms.

Functions (b) are symmetric with respect to elements of the alternating group  $A_n$ . Since exponential functions  $e^{2\pi imx}$ ,  $m \in \mathbb{Z}$ , of one variable are invariant with respect to shifts  $x \rightarrow x + k$ ,  $k \in \mathbb{Z}$ , then symmetries of functions (a) are described by a wider group, which is called the *affine alternating group*  $A_n^{\text{aff}}$ . This group is a product of the group  $A_n$  and the group  $T_n$ , consisting of shifts in the space  $E_n$  by vectors  $r = (r_1, r_2, \dots, r_n)$ ,  $r_j \in \mathbb{Z}$ . A fundamental domain  $F(A_n^{\text{aff}})$  of the group  $A_n^{\text{aff}}$  is a certain bounded subset of  $\mathbb{R}^n$ .

Functions on the fundamental domain  $F(A_n^{\text{aff}})$  can be expanded into series in the functions (a). These expansions are an analogue of the usual Fourier series for functions of one variable. Functions (b) determine Fourier integral transforms on the fundamental domain  $F(A_n)$  of the alternating group  $A_n$ . This domain consists of points  $x \in E_n$  such that  $x_1, x_2 > x_3 > \dots > x_n$ , where  $x_1, x_2 > x_3$  means that  $x_1 > x_3$  and  $x_2 > x_3$ .

Functions (c) are used to determine finite (on a finite set) Fourier transforms. These Fourier transforms are given on grids consisting of points in the fundamental domain  $F(A_n^{\text{aff}})$ .

Alternating multivariate exponential functions are closely related to symmetric and antisymmetric exponential functions of [1]. The symmetric and antisymmetric exponential functions are connected with symmetric and antisymmetric orbit functions defined in [2], [3] and studied in detail in [4] and [5]. Discrete orbit function transforms, corresponding to Coxeter–Dynkin diagrams of low order, were studied in detail, and it was shown that they are very useful for applications [6]–[14].

The discrete Fourier transforms, determined by (anti)symmetric multivariate exponential functions, studied in [1], and by alternating multivariate exponential functions have a number of practically useful properties. In particular, continuous extension of the discrete transforms smoothly interpolate digital data in any dimension. Examples show that relative to the amount of available data, these transforms provide much smoother interpolation than the conventional Fourier transforms.

Symmetric and antisymmetric multivariate exponential functions, studied in [1], satisfy certain boundary conditions (antisymmetric exponential functions vanish on the boundary of the corresponding fundamental domain and the derivative of the symmetric exponential functions with respect to the normal to the boundary of the

fundamental domain vanishes on the boundary). This means that smooth functions, which are expanded in these functions, have to satisfy these conditions, that is, not each smooth function can be expanded in (anti)symmetric exponential functions. Alternating multivariate exponential functions satisfy no boundary conditions and any smooth function can be expanded in these exponential functions.

Alternating multivariate exponential functions are also related to the so-called  $E$ -orbit functions exposed in [2], [3], and [9]). The exposition of the theory of orbit functions strongly depends on the theory of Weyl groups, properties of root systems, etc. In this paper we avoid this dependence. We use only the alternating group and its affinization.

The best way to define alternating multivariate exponential functions is to use a semideterminant of a finite matrix. The semideterminant of a matrix is closely related to the determinant and the antideterminant of the same matrix. It is well-known that the determinant  $\det(a_{ij})_{i,j=1}^n$  of an  $n \times n$  matrix  $(a_{ij})_{i,j=1}^n$  is defined as

$$\begin{aligned} \det(a_{ij})_{i,j=1}^n &= \sum_{w \in S_n} (\det w) a_{1,w(1)} a_{2,w(2)} \cdots a_{n,w(n)} \\ &= \sum_{w \in S_n} (\det w) a_{w(1),1} a_{w(2),2} \cdots a_{w(n),n}, \end{aligned} \tag{1}$$

where  $S_n$  is the symmetric group of  $n$  symbols  $1, 2, \dots, n$ , the set  $(w(1), w(2), \dots, w(n))$  denotes the set  $w(1, 2, \dots, n)$ , and  $\det w$  denotes the determinant of the transform  $w$ , that is,  $\det w = 1$  if  $w$  is an even permutation and  $\det w = -1$  otherwise. The antideterminant  $\det^+$  of a matrix  $(a_{ij})_{i,j=1}^n$  is the sum of all terms, entering in the expression for the corresponding determinant taken with sign  $+$ ,

$$\det^+(a_{ij})_{i,j=1}^n = \sum_{w \in S_n} a_{1,w(1)} a_{2,w(2)} \cdots a_{n,w(n)} = \sum_{w \in S_n} a_{w(1),1} a_{w(2),2} \cdots a_{w(n),n}.$$

For the semideterminant  $\text{sdet}$  of a matrix  $(a_{ij})_{i,j=1}^n$  we have

$$\text{sdet} (a_{ij})_{i,j=1}^n = \frac{1}{2} (\det(a_{ij})_{i,j=1}^n + \det^+(a_{ij})_{i,j=1}^n).$$

Clearly,

$$\text{sdet} (a_{ij})_{i,j=1}^n = \sum_{w \in A_n} a_{1,w(1)} a_{2,w(2)} \cdots a_{n,w(n)} = \sum_{w \in A_n} a_{w(1),1} a_{w(2),2} \cdots a_{w(n),n}. \tag{2}$$

In the text, we use formulas of the type

$$N \geq a_1, a_2 \geq a_3 \geq a_4 \geq \cdots \geq a_n.$$

Here  $N \geq a_1, a_2 \geq a_3$  means that  $N \geq a_1 \geq a_3$  and  $N \geq a_2 \geq a_3$ . Formulas of the type  $a_1, a_2 > a_3 > \cdots > a_n$  have a similar sense.

## 2. ALTERNATING MULTIVARIATE EXPONENTIAL FUNCTIONS

An alternating multivariate exponential function  $E_\lambda(x)$  of  $x = (x_1, x_2, \dots, x_n)$  is defined as the function

$$\begin{aligned} E_\lambda(x) &\equiv E_{(\lambda_1, \lambda_2, \dots, \lambda_n)}(x) = \text{sdet} \left( e^{2\pi i \lambda_i x_j} \right)_{i,j=1}^n \\ &= \text{sdet} \begin{pmatrix} e^{2\pi i \lambda_1 x_1} & e^{2\pi i \lambda_1 x_2} & \dots & e^{2\pi i \lambda_1 x_n} \\ e^{2\pi i \lambda_2 x_1} & e^{2\pi i \lambda_2 x_2} & \dots & e^{2\pi i \lambda_2 x_n} \\ \dots & \dots & \dots & \dots \\ e^{2\pi i \lambda_n x_1} & e^{2\pi i \lambda_n x_2} & \dots & e^{2\pi i \lambda_n x_n} \end{pmatrix} \\ &\equiv \sum_{w \in A_n} e^{2\pi i \lambda_1 x_{w(1)}} e^{2\pi i \lambda_2 x_{w(2)}} \dots e^{2\pi i \lambda_n x_{w(n)}} = \sum_{w \in A_n} e^{2\pi i \langle \lambda, wx \rangle}, \end{aligned} \quad (3)$$

where  $\lambda = (\lambda_1, \lambda_2, \dots, \lambda_n)$  is a set of real numbers, which determines the function  $E_\lambda(x)$ , and  $\langle \lambda, x \rangle$  denotes the scalar product in the  $n$ -dimensional Euclidean space  $E_n$ ,  $\langle \lambda, x \rangle = \sum_{i=1}^n \lambda_i x_i$ . When  $\lambda_1, \lambda_2, \dots, \lambda_n$  are integers, we denote this set of numbers as  $m \equiv (m_1, m_2, \dots, m_n)$ ,

$$E_{m_1, m_2, \dots, m_n}(x) = \text{sdet} \left( e^{2\pi i m_i x_j} \right)_{i,j=1}^n. \quad (4)$$

From expression (3), it follows that the alternating exponential functions  $E_\lambda(x)$  satisfy the relation

$$E_\lambda(x_1 + a, x_2 + a, \dots, x_n + a) = e^{2\pi i (\lambda_1 + \lambda_2 + \dots + \lambda_n) a} E_\lambda(x). \quad (5)$$

It is therefore sufficient to consider the function  $E_\lambda(x)$  on the hyperplane

$$x_1 + x_2 + \dots + x_n = b,$$

where  $b$  is a fixed number (we denote this hyperplane by  $\mathcal{H}_b$ ). A transition from one hyperplane  $\mathcal{H}_b$  to another  $\mathcal{H}_c$  is fulfilled by multiplication by a usual exponential function  $e^{2\pi i |\lambda| (c-b)}$ , where  $|\lambda| = \lambda_1 + \lambda_2 + \dots + \lambda_n$ ,

$$E_\lambda(x)|_{x \in \mathcal{H}_b} = e^{2\pi i (\lambda_1 + \lambda_2 + \dots + \lambda_n) (c-b)} E_\lambda(x)|_{x \in \mathcal{H}_c}.$$

It is useful to consider the functions  $E_\lambda(x)$  on the hyperplane  $\mathcal{H}_0$ . For  $x \in \mathcal{H}_0$  we have the relation

$$E_{\lambda_1 + \nu, \lambda_2 + \nu, \dots, \lambda_n + \nu}(x) = E_\lambda(x). \quad (6)$$

We conclude from (2) that the expression of semideterminant  $\text{sdet}$  does not change when applying a permutation from  $A_n$  to rows or to columns. This means that for any permutation  $w \in A_n$  we have

$$E_{w\lambda}(x) = E_\lambda(x), \quad E_\lambda(wx) = E_\lambda(x). \quad (7)$$

Therefore, it is sufficient to consider only alternating exponential functions  $E_\lambda(x)$  with  $\lambda = (\lambda_1, \lambda_2, \dots, \lambda_n)$  such that

$$\lambda_1, \lambda_2 \geq \lambda_3 \geq \dots \geq \lambda_n.$$

Such  $\lambda$  are called *semidominant*. The set of all semidominant  $\lambda$  is denoted by  $D_+^e$ . Below, when considering alternating exponential functions  $E_\lambda(x)$ , we assume that  $\lambda \in D_+^e$ .

Alternating exponential functions are related to symmetric and antisymmetric exponential functions  $E_\lambda^+(x)$  and  $E_\lambda^-(x)$ , which are studied in [1]. They are determined by the formulas

$$E_\lambda^+(x) = \det^+ \left( e^{2\pi i \lambda_i x_j} \right)_{i,j=1}^n, \quad E_\lambda^-(x) = \det \left( e^{2\pi i \lambda_i x_j} \right)_{i,j=1}^n,$$

where  $\lambda$  and  $x$  are such as in (3). This relation will be considered in section 4. Here we consider the case  $n = 2$ .

**The case  $n = 2$ .** We examine the alternating multivariate exponential functions  $E_\lambda(x)$  for  $n = 2$  and  $\lambda_2 = -\lambda_1 \equiv \lambda$ . Then

$$E_{(\lambda, -\lambda)}(x_1, x_2) = e^{2\pi i \lambda(x_1 - x_2)}, \quad E_{(-\lambda, \lambda)}(x_1, x_2) = e^{-2\pi i \lambda(x_1 - x_2)}.$$

For antisymmetric and symmetric multivariate exponential functions with these  $n$  and  $\lambda_2$  we have

$$\begin{aligned} E_{(\lambda, -\lambda)}^-(x_1, x_2) &= e^{2\pi i \lambda(x_1 - x_2)} - e^{-2\pi i \lambda(x_1 - x_2)} = 2i \sin 2\pi i \lambda(x_1 - x_2), \\ E_{(\lambda, -\lambda)}^+(x_1, x_2) &= e^{2\pi i \lambda(x_1 - x_2)} + e^{-2\pi i \lambda(x_1 - x_2)} = 2 \cos 2\pi i \lambda(x_1 - x_2). \end{aligned}$$

Thus,

$$E_{(\lambda, -\lambda)}^-(x_1, x_2) = E_{(\lambda, -\lambda)}(x_1, x_2) - E_{(-\lambda, \lambda)}(x_1, x_2), \quad (8)$$

$$E_{(\lambda, -\lambda)}^+(x_1, x_2) = E_{(\lambda, -\lambda)}(x_1, x_2) + E_{(-\lambda, \lambda)}(x_1, x_2). \quad (9)$$

Therefore, for  $n = 2$  and  $\lambda_2 = -\lambda_1$  the alternating multivariate exponential function is the usual exponential function of one variable, whereas antisymmetric and symmetric multivariate exponential functions are sine and cosine functions of one variable. As we shall see below, the relations (8) and (9) can be generalized for any  $n \in \mathbb{Z}_+$ .

### 3. AFFINE ALTERNATING GROUP AND FUNDAMENTAL DOMAINS

The functions  $E_\lambda(x)$  are symmetric with respect to the alternating group  $A_n$ , that is,  $E_\lambda(wx) = E_\lambda(x)$ ,  $w \in A_n$ . The functions  $E_m(x)$  with integral  $m = (m_1, m_2, \dots, m_n)$  admit additional symmetries related to the periodicity of the exponential functions  $e^{2\pi i r y}$ ,  $r \in \mathbb{Z}$ ,  $y \in \mathbb{R}$ . These symmetries are described by the discrete group of shifts in the space  $E_n$  by vectors

$$r_1 \mathbf{e}_1 + r_2 \mathbf{e}_2 + \dots + r_n \mathbf{e}_n, \quad r_i \in \mathbb{Z},$$

where  $\mathbf{e}_1, \mathbf{e}_2, \dots, \mathbf{e}_n$  are the unit vectors along the corresponding coordinate axes. We denote this group by  $T_n$ .

Permutations of  $A_n$  and shifts of  $T_n$  generate a group which is denoted as  $A_n^{\text{aff}}$  and referred to as the *affine alternating group*. The group  $A_n^{\text{aff}}$  is a semidirect product of its subgroups  $A_n$  and  $T_n$ ,

$$A_n^{\text{aff}} = A_n \circledast T_n,$$

where  $T_n$  is an invariant subgroup, that is,  $wtw^{-1} \in T_n$  for  $w \in A_n$  and  $t \in T_n$ .

An open connected simply connected set  $F \subset \mathbb{R}^n$  is called a *fundamental domain* for the group  $A_n^{\text{aff}}$  (for the group  $A_n$ ) if it does not contain equivalent points (that is, points  $x$  and  $x'$  such that  $x' = wx$ , where  $w$  belongs to  $A_n^{\text{aff}}$  or  $A_n$ , respectively) and if its closure contains at least one point from each  $A_n^{\text{aff}}$ -orbit (from each  $A_n$ -orbit). Recall that an  $A_n^{\text{aff}}$ -orbit of a point  $x \in \mathbb{R}^n$  is the set of points  $wx$ ,  $w \in A_n^{\text{aff}}$ .

It is evident that the set  $D_{++}^e$  of all points  $x = (x_1, x_2, \dots, x_n)$  such that

$$x_1, x_2 > x_2 > \dots > x_n,$$

is a fundamental domain for the group  $A_n$  (we denote it as  $F(A_n)$ ). The set of points  $x = (x_1, x_2, \dots, x_n) \in D_{++}^e$  such that

$$1 > x_1, x_2 > x_2 > \dots > x_n > 0$$

is a fundamental domain for the affine group  $A_n^{\text{aff}}$  (we denote it as  $F(A_n^{\text{aff}})$ ).

As previously seen, the functions  $E_\lambda(x)$  are symmetric with respect to the alternating group  $A_n$ . This means that it is sufficient to consider the functions  $E_\lambda(x)$  only on the closure of the fundamental domain  $F(A_n)$ . Values of  $E_\lambda(x)$  on other points are obtained by using the symmetry.

The symmetry of functions  $E_m(x)$ ,  $m = (m_1, m_2, \dots, m_n)$ ,  $m_i \in \mathbb{Z}$ , with respect to the affine alternating group  $A_n^{\text{aff}}$ ,

$$E_m(wx + r) = E_m(x), \quad w \in A_n, \quad r \in T_n, \quad (10)$$

means that we may consider  $E_m(x)$  only on the closure of the fundamental domain  $F(A_n^{\text{aff}})$ , that is, on the set of points  $x$  such that  $1 \geq x_1, x_2 \geq x_2 \geq \dots \geq x_n \geq 0$ . Values of  $E_m(x)$  on other points are obtained by using the relation (10).

#### 4. RELATION TO SYMMETRIC AND ANTISYMMETRIC EXPONENTIAL FUNCTIONS

The alternating multivariate exponential functions  $E_\lambda(x)$  are related to symmetric and antisymmetric multivariate exponential functions  $E_\lambda^+(x)$  and  $E_\lambda^-(x)$  defined in [1] and determined as

$$E_\lambda^-(x) = \det \left( e^{2\pi i \lambda_i x_j} \right)_{i,j=1}^n, \quad E_\lambda^+(x) = \det^+ \left( e^{2\pi i \lambda_i x_j} \right)_{i,j=1}^n,$$

where  $\det^+ \left( e^{2\pi i \lambda_i x_j} \right)_{i,j=1}^n$  is the antideterminant of the matrix  $\left( e^{2\pi i \lambda_i x_j} \right)_{i,j=1}^n$ .

It follows from the definitions of alternating and symmetric and antisymmetric multivariate exponential functions that for  $\lambda$  such that  $\lambda_1 > \lambda_2 > \lambda_3 > \dots > \lambda_n$  we have

$$E_\lambda^-(x) = E_\lambda(x) - E_{r_{12}\lambda}(x), \quad (11)$$

$$E_\lambda^+(x) = E_\lambda(x) + E_{r_{12}\lambda}(x), \quad (12)$$

where  $r_{12}$  means the permutation of  $\lambda_1$  and  $\lambda_2$ . It follows from here that

$$E_\lambda(x) = \frac{1}{2}(E_\lambda^+(x) + E_\lambda^-(x)), \quad (13)$$

$$E_{r_{12}\lambda}(x) = \frac{1}{2}(E_\lambda^+(x) - E_\lambda^-(x)). \quad (14)$$

It is directly derived from (11)–(14) that

$$\begin{aligned} (E_\lambda^+(x))^2 - (E_\lambda^-(x))^2 &= 4E_\lambda(x)E_{r_{12}\lambda}(x), \\ (E_\lambda^+(x))^2 + (E_\lambda^-(x))^2 &= 2(E_\lambda(x))^2 + 2(E_{r_{12}\lambda}(x))^2. \end{aligned}$$

If there are two coinciding numbers in the set  $\lambda_1, \lambda_2, \dots, \lambda_n$ , we get

$$E_\lambda(x) = E_\lambda^+(x). \quad (15)$$

#### 5. PROPERTIES

The symmetry of alternating multivariate exponential functions  $E_\lambda(x)$  with respect to the alternating group  $A_n$  is a main property of these functions. However, they possess many other interesting properties.

**Continuity.** The functions  $E_\lambda(x)$  are finite sums of multivariate exponential functions. Therefore, they are continuous functions of  $x_1, x_2, \dots, x_n$  and have continuous derivatives of all orders in  $\mathbb{R}^n$ .

**Complex conjugation.** Let there be two coinciding numbers in the set  $\lambda = (\lambda_1, \lambda_2, \dots, \lambda_n)$ . Since in this case  $E_\lambda(x) = E_\lambda^+(x)$ , then due to the behavior of  $E_\lambda^+(x)$  under complex conjugation (see section 4 in [1]) we obtain

$$E_{(\lambda_1, \lambda_2, \dots, \lambda_n)}(x) = \overline{E_{-(\lambda_n, \lambda_{n-1}, \dots, \lambda_1)}(x)}. \quad (16)$$

Let  $\lambda = (\lambda_1, \lambda_2, \dots, \lambda_n)$  be such that  $\lambda_1 > \lambda_2 > \dots > \lambda_n$ . Then due to relations (17) and (18) in [1] and formulas (13) and (14) we derive that

$$E_{(\lambda_1, \lambda_2, \dots, \lambda_n)}(x) = \overline{E_{-(\lambda_n, \lambda_{n-1}, \dots, \lambda_1)}(x)}, \quad (17)$$

$$E_{r_{12}(\lambda_1, \lambda_2, \dots, \lambda_n)}(x) = \overline{E_{-r_{12}(\lambda_n, \lambda_{n-1}, \dots, \lambda_1)}(x)} \quad (18)$$

for  $n = 4k, 4k + 1$  and

$$E_{(\lambda_1, \lambda_2, \dots, \lambda_n)}(x) = \overline{E_{-r_{12}(\lambda_n, \lambda_{n-1}, \dots, \lambda_1)}(x)}, \quad (19)$$

$$E_{r_{12}(\lambda_1, \lambda_2, \dots, \lambda_n)}(x) = \overline{E_{-(\lambda_n, \lambda_{n-1}, \dots, \lambda_1)}(x)} \quad (20)$$

for  $n = 4k - 1, 4k - 2$ , where  $k$  is a positive integer.

**Scaling symmetry.** For  $c \in \mathbb{R}$ , let  $c\lambda = (c\lambda_1, c\lambda_2, \dots, c\lambda_n)$ . Then

$$E_{c\lambda}(x) = \sum_{w \in A_n} e^{2\pi i \langle cw\lambda, x \rangle} = \sum_{w \in A_n} e^{2\pi i \langle w\lambda, cx \rangle} = E_\lambda(cx).$$

The equality  $E_{c\lambda}(x) = E_\lambda(cx)$  expresses the *scaling symmetry of exponential functions*  $E_\lambda(x)$ .

**Duality.** Due to invariance of the scalar product  $\langle \cdot, \cdot \rangle$  with respect to the alternating group  $A_n$ ,  $\langle w\mu, wy \rangle = \langle \mu, y \rangle$ , we have

$$E_\lambda(x) = \sum_{w \in A_n} e^{2\pi i \langle \lambda, w^{-1}x \rangle} = \sum_{w \in A_n} e^{2\pi i \langle x, w\lambda \rangle} = E_x(\lambda).$$

This relation expresses the *duality* of alternating exponential functions.

**Orthogonality on the fundamental domain  $F(A_n^{\text{aff}})$ .** Alternating exponential functions  $E_m(x)$  with  $m = (m_1, m_2, \dots, m_n) \in D_{++}^e$ ,  $m_j \in \mathbb{Z}$ , are orthogonal on  $F(A_n^{\text{aff}})$  with respect to the Euclidean measure,

$$|F(A_n^{\text{aff}})|^{-1} \int_{F(A_n^{\text{aff}})} E_m(x) \overline{E_{m'}(x)} dx = |A_n| \delta_{mm'},$$

where  $|A_n|$  means the number of elements in the set  $A_n$ ,  $\overline{F(A_n^{\text{aff}})}$  is the closure of  $F(A_n^{\text{aff}})$ , and  $|F(A_n^{\text{aff}})|$  is the area of the fundamental domain  $F(A_n^{\text{aff}})$ . This relation follows from the equality

$$\int_{\mathbb{T}} E_m(x) \overline{E_{m'}(x)} dx = |A_n| \delta_{mm'} \quad (21)$$

(where  $\mathbb{T}$  is the torus in  $E_n$  consisting of points  $x = (x_1, x_2, \dots, x_n)$ ,  $0 \leq x_i < 1$ ), which is a consequence of the orthogonality of the exponential functions  $e^{2\pi i \langle \mu, x \rangle}$  (entering into the definition of  $E_m(x)$ ) for different sets  $\mu$ .

Assuming that an area of  $\mathbb{T}$  is equal to 1,  $|\mathbb{T}| = 1$ , we have  $|F(A_n^{\text{aff}})| = |A_n|^{-1}$ , and formula (21) takes the form

$$\int_{F(A_n^{\text{aff}})} E_m(x) \overline{E_{m'}(x)} dx = \delta_{mm'}. \quad (22)$$

There can be coinciding summands in expression (3) for alternating exponential functions. For this reason, for functions  $E_m(x)$  with  $m = (m_1, m_2, \dots, m_n) \in D_+^e$ ,  $m_j \in \mathbb{Z}$ , the relation (22) is replaced by

$$\int_{F(A_n^{\text{aff}})} E_m(x) \overline{E_{m'}(x)} dx = |G_m| \delta_{mm'}. \quad (23)$$

where  $|G_m|$  is the number of elements in the subgroup  $G_m$  of  $A_n$  consisting of elements  $w \in A_n$  such that  $wm = m$ .

**Solutions of the Laplace equation.** The Laplace operator on the Euclidean space  $E_n$  in the Cartesian coordinates  $x = (x_1, x_2, \dots, x_n)$  takes the form

$$\Delta = \frac{\partial^2}{\partial x_1^2} + \frac{\partial^2}{\partial x_2^2} + \dots + \frac{\partial^2}{\partial x_n^2}.$$

Taking any summand in the expression for the alternating multivariate exponential function  $E_\lambda(x)$ , we get

$$\Delta e^{2\pi i((w(\lambda))_1 x_1 + \dots + (w(\lambda))_n x_n)} = -4\pi^2 \langle \lambda, \lambda \rangle e^{2\pi i((w(\lambda))_1 x_1 + \dots + (w(\lambda))_n x_n)},$$

where  $\lambda = (\lambda_1, \lambda_2, \dots, \lambda_n)$  determines  $E_\lambda(x)$ . Since this action of  $\Delta$  does not depend on a summand from the expression for the alternating exponential function, we have

$$\Delta E_\lambda(x) = -4\pi^2 \langle \lambda, \lambda \rangle E_\lambda(x), \quad (24)$$

The formula (24) can be generalized in the following way. Let  $\sigma_k(y_1, y_2, \dots, y_n)$  be the  $k$ -th elementary symmetric polynomial of degree  $k$ , that is,

$$\sigma_k(y_1, y_2, \dots, y_n) = \sum_{1 \leq k_1 < k_2 < \dots < k_n \leq n} y_{k_1} y_{k_2} \dots y_{k_n}.$$

Then for  $k = 1, 2, \dots, n$  we have

$$\sigma_k \left( \frac{\partial^2}{\partial x_1^2}, \frac{\partial^2}{\partial x_2^2}, \dots, \frac{\partial^2}{\partial x_n^2} \right) E_\lambda(x) = (-4\pi^2)^k \sigma_k(\lambda_1^2, \lambda_2^2, \dots, \lambda_n^2) E_\lambda(x). \quad (25)$$

Note that  $n$  differential equations (25) are algebraically independent.

Symmetric and antisymmetric multivariate exponential functions also satisfy these equations. They satisfy the certain boundary conditions (antisymmetric exponential functions vanish on the boundary of the corresponding fundamental domain and the derivative of the symmetric exponential functions with respect to the normal to the boundary of the fundamental domain vanishes on the boundary). Alternating exponential functions do not satisfy these conditions.

## 6. EXPANSIONS IN ALTERNATING EXPONENTIAL FUNCTIONS ON $F(A_n^{\text{aff}})$

Alternating exponential functions determine symmetric (with respect to  $A_n$ ) multivariate Fourier transforms that generalize the usual Fourier transform. There are three types of such transforms:

- (a) Fourier transforms related to the functions  $E_m(x)$  with  $m = (m_1, m_2, \dots, m_n)$ ,  $m_j \in \mathbb{Z}$  (Fourier series);
- (b) Fourier transforms related to  $E_\lambda(x)$  with  $\lambda \in D_+^e$ ;
- (c) Multivariate finite Fourier transforms.



In this section, we consider expansions in alternating exponential functions on the fundamental domain  $F(A_n^{\text{aff}})$ . These expansions are constructed in the same way as in the case of (anti)symmetric exponential functions in [1].

Let  $f(x)$  be a symmetric (with respect to the group  $A_n^{\text{aff}}$ ) continuous function on the  $n$ -dimensional Euclidean space  $E_n$  which has continuous derivatives. We may consider this function on the torus  $\mathbb{T}$  which can be identified with a closure of the union of the sets  $wF(A_n^{\text{aff}})$ ,  $w \in A_n$ . The function  $f(x)$ , as a function on  $\mathbb{T}$ , can be expanded in exponential functions  $e^{2\pi i m_1 x_1} e^{2\pi i m_2 x_2} \dots e^{2\pi i m_n x_n}$ ,  $m_i \in \mathbb{Z}$ . We have

$$f(x) = \sum_{m_i \in \mathbb{Z}} c_m e^{2\pi i m_1 x_1} e^{2\pi i m_2 x_2} \dots e^{2\pi i m_n x_n}, \quad (26)$$

where  $m = (m_1, m_2, \dots, m_n)$ . It follows from the symmetry  $f(wx) = f(x)$ ,  $w \in A_n$ , that

$$\begin{aligned} f(wx) &= \sum_{m_i \in \mathbb{Z}} c_m e^{2\pi i m_1 x_{w(1)}} \dots e^{2\pi i m_n x_{w(n)}} = \sum_{m_i \in \mathbb{Z}} c_m e^{2\pi i m_{w^{-1}(1)} x_1} \dots e^{2\pi i m_{w^{-1}(n)} x_n} \\ &= \sum_{m_i \in \mathbb{Z}} c_{wm} e^{2\pi i m_1 x_1} \dots e^{2\pi i m_n x_n} = f(x) = \sum_{m_i \in \mathbb{Z}} c_m e^{2\pi i m_1 x_1} \dots e^{2\pi i m_n x_n}. \end{aligned}$$

Therefore, the coefficients  $c_m$  satisfy the conditions  $c_{wm} = c_m$ ,  $w \in A_n$ . Collecting exponential functions in (26) at the same  $c_{wm}$ ,  $w \in A_n$ , we obtain the expansion

$$f(x) = \sum_{m \in P_+^e} c_m E_m(x), \quad (27)$$

where  $P_+^e = D_+^e \cap \mathbb{Z}^n$ . Thus, *any symmetric (with respect to  $A_n$ ) continuous function  $f$  on  $\mathbb{T}$  which has continuous derivatives (that is, any continuous function on  $D_+^e$  with continuous derivatives) can be expanded in alternating exponential functions  $E_m(x)$ ,  $m \in P_+^e$ .*

By the orthogonality relation (23), the coefficients  $c_m$  in the expansion (27) are determined by the formula

$$c_m = |G_m|^{-1} \int_{F(A_n^{\text{aff}})} f(x) \overline{E_m(x)} dx, \quad (28)$$

where, as before,  $|G_m|$  is the number of elements in the subgroup  $G_m$  of  $A_n$  consisting of  $w \in A_n$  such that  $wm = m$ . Moreover, the Plancherel formula holds:

$$\sum_{m \in P_+^e} |c_m|^2 = |G_m|^{-1} \int_{F(A_n^{\text{aff}})} |f(x)|^2 dx. \quad (29)$$

Formula (28) is the symmetrized (with respect to the group  $A_n$ ) Fourier transform of the function  $f(x)$ . Formula (27) gives an inverse transform. Formulas (27) and (28) give the *multivariate Fourier transforms* corresponding to the alternating exponential functions  $E_m(x)$ ,  $m \in P_+^e$ .

Let  $\mathcal{L}^2(\overline{F(A_n^{\text{aff}})})$  denote the Hilbert space of functions on the domain  $\overline{F(A_n^{\text{aff}})}$  with the scalar product

$$\langle f_1, f_2 \rangle = \int_{F(A_n^{\text{aff}})} f_1(x) \overline{f_2(x)} dx.$$

The formulas (27)-(29) show that *the set of alternating multivariate exponential functions  $E_m(x)$ ,  $m \in P_+^e$ , forms an orthogonal basis of  $\mathcal{L}^2(\overline{F(A_n^{\text{aff}})})$ .*

7. FOURIER TRANSFORMS ON THE FUNDAMENTAL DOMAIN  $F(A_n)$ 

The expansions (27) of functions on the fundamental domain  $F(A_n^{\text{aff}})$  are expansions in the exponential functions  $E_m(x)$  with integral  $m = (m_1, m_2, \dots, m_n)$ . The exponential functions  $E_\lambda(x)$  with  $\lambda$  lying in the fundamental domain  $F(A_n)$  (and not obligatory integral) are not invariant with respect to the corresponding affine alternating group  $A_n^{\text{aff}}$ . They are invariant only with respect to the alternating group  $A_n$ . A closure of the fundamental domain of  $A_n$  coincides with the set  $D_+^e$  consisting of points  $x$  such that  $x_1, x_2 \geq x_3 \geq \dots \geq x_n$ . The functions  $E_\lambda(x)$ ,  $\lambda \in D_+^e$ , determine Fourier transforms on  $D_+^e$ .

We begin with the usual Fourier transforms on  $\mathbb{R}^n$ :

$$\tilde{f}(\lambda) = \int_{\mathbb{R}^n} f(x) e^{2\pi i \langle \lambda, x \rangle} dx, \quad (30)$$

$$f(x) = \int_{\mathbb{R}^n} \tilde{f}(\lambda) e^{-2\pi i \langle \lambda, x \rangle} d\lambda. \quad (31)$$

Let the function  $f(x)$  be invariant with respect to the alternating group  $A_n$ ,  $f(wx) = f(x)$ ,  $w \in A_n$ . It is easy to verify that  $\tilde{f}(\lambda)$  is also invariant with respect to  $A_n$ . By replacing  $\lambda$  by  $w\lambda$ ,  $w \in A_n$ , in (30) and summing up both sides over  $w \in A_n$ , we obtain, instead of (30), the equality

$$\tilde{f}(\lambda) = \int_{D_+^e} f(x) E_\lambda(x) dx, \quad \lambda \in D_+^e, \quad (32)$$

where we have taken into account that  $f(x)$  is invariant with respect to  $A_n$ .

Similarly, starting from (31), we obtain the inverse formula:

$$f(x) = \int_{D_+^e} \tilde{f}(\lambda) \overline{E_\lambda(x)} d\lambda. \quad (33)$$

For the transforms (32) and (33), the Plancherel formula

$$\int_{D_+^e} |f(x)|^2 dx = \int_{D_+^e} |\tilde{f}(\lambda)|^2 d\lambda$$

holds. Formulas (32) and (33) determine the symmetric (with respect to  $A_n$ ) multivariate Fourier transforms on the domain  $F(A_n)$ , which are referred to as the *alternating Fourier transforms*.

Let  $\mathcal{L}^2(D_+^e)$  denote the Hilbert space of functions on the domain  $D_+^e$  with the scalar product

$$\langle f_1, f_2 \rangle = \int_{D_+^e} f_1(x) \overline{f_2(x)} dx$$

The formulas (32) and (33) show that *the set of alternating multivariate exponential functions  $E_\lambda(x)$ ,  $\lambda \in D_+^e$ , form a full system of functions of  $\mathcal{L}^2(D_+^e)$ .*

## 8. EIGENFUNCTIONS OF THE ALTERNATING FOURIER TRANSFORM

Let  $H_n(x)$ ,  $n = 0, 1, 2, \dots$ , be the well-known Hermite polynomials of one variable. They satisfy the relation

$$\int_{-\infty}^{\infty} e^{2\pi i p x} e^{-\pi p^2} H_m(\sqrt{2\pi} p) dp = i^{-m} e^{-\pi x^2} H_m(\sqrt{2\pi} x) \quad (34)$$

(see, for example, subsection 12.2.4 in [15]).

We generate polynomials of many variables

$$H_{\mathbf{m}}(\mathbf{x}) \equiv H_{m_1, m_2, \dots, m_n}(x_1, x_2, \dots, x_n) := H_{m_1}(x_1)H_{m_2}(x_2) \cdots H_{m_n}(x_n). \quad (35)$$

The functions

$$e^{-|\mathbf{x}|^2/2} H_{\mathbf{m}}(\mathbf{x}), \quad m_i = 0, 1, 2, \dots, \quad i = 1, 2, \dots, n, \quad (36)$$

where  $|\mathbf{x}|$  is a length of the vector  $\mathbf{x}$ , form an orthogonal basis of the Hilbert space  $\mathcal{L}^2(\mathbb{R}^n)$  with the scalar product  $\langle f_1, f_2 \rangle := \int_{\mathbb{R}^n} f_1(\mathbf{x})\overline{f_2(\mathbf{x})}d\mathbf{x}$ , where  $d\mathbf{x} = dx_1 dx_2 \cdots dx_n$ .

We symmetrize the functions

$$\mathcal{H}_{\mathbf{m}}(\mathbf{x}) := e^{-\pi|\mathbf{x}|^2} H_{\mathbf{m}}(\sqrt{2\pi}\mathbf{x})$$

(obtained from (36) by replacing  $\mathbf{x}$  by  $\sqrt{2\pi}\mathbf{x}$ ) by means of alternating multivariate exponential functions:

$$\int_{\mathbb{R}^n} E_{\lambda}(\mathbf{x})e^{-\pi|\mathbf{x}|^2} H_{\mathbf{m}}(\sqrt{2\pi}\mathbf{x})d\mathbf{x} = i^{-|\mathbf{m}|} e^{-\pi|\lambda|^2} \mathbf{H}_{\mathbf{m}}(\sqrt{2\pi}\lambda), \quad (37)$$

It is easy to see that the polynomials  $\mathbf{H}_{\mathbf{m}}$  are indeed symmetric with respect to the group  $A_n$ ,

$$\mathbf{H}_{\mathbf{m}}(w\lambda) = \mathbf{H}_{\mathbf{m}}(\lambda), \quad w \in A_n.$$

Therefore, we may consider  $\mathbf{H}_{\mathbf{m}}(\lambda)$  for values of  $\lambda = (\lambda_1, \lambda_2, \dots, \lambda_n)$  such that  $\lambda_1, \lambda_2 \geq \lambda_3 \geq \cdots \geq \lambda_n$ . The polynomials  $\mathbf{H}_{\mathbf{m}}$  are of the form

$$\mathbf{H}_{\mathbf{m}}(\lambda) = \text{sdet} (H_{m_i}(\lambda_j))_{i,j=1}^n, \quad (38)$$

that is, it is sufficient to consider the polynomials  $\mathbf{H}_{\mathbf{m}}(\lambda)$  for integer  $n$ -tuples  $\mathbf{m}$ , such that  $m_1, m_2 \geq m_3 \geq \cdots \geq m_n$ .

Let us apply the alternating Fourier transform (32) (we denote it as  $\mathfrak{F}$ ) to the functions (38). Taking into account formula (37) we derive

$$\begin{aligned} \mathfrak{F} \left( e^{-\pi|\mathbf{x}|^2} \mathbf{H}_{\mathbf{m}}(\sqrt{2\pi}\mathbf{x}) \right) &= \frac{1}{|A_n|} \int_{\mathbb{R}^n} E_{\lambda}(\mathbf{x})e^{-\pi|\mathbf{x}|^2} \mathbf{H}_{\mathbf{m}}(\sqrt{2\pi}\mathbf{x})d\mathbf{x} \\ &= i^{-|\mathbf{m}|} e^{-\pi|\lambda|^2} \mathbf{H}_{\mathbf{m}}(\sqrt{2\pi}\lambda), \end{aligned}$$

that is, the functions  $e^{-\pi|\mathbf{x}|^2} \mathbf{H}_{\mathbf{m}}(\sqrt{2\pi}\mathbf{x})$  are eigenfunctions of the alternating Fourier transform  $\mathfrak{F}$ . Since these functions for  $m_i = 0, 1, 2, \dots, i = 1, 2, \dots, n$ , such that  $m_1, m_2 \geq m_3 \geq \cdots \geq m_n$ , form an orthogonal basis of the Hilbert space  $\mathcal{L}^2(D_+^e)$ , they constitute a complete set of eigenfunctions of this transform. Thus, this transform has only four eigenvalues  $i, -i, 1, -1$  in  $\mathcal{L}^2(D_+^e)$ . This means that we have  $\mathfrak{F}^4 = 1$ .

## 9. FINITE ALTERNATING FOURIER TRANSFORMS

Along with the integral Fourier transform in one variable, there exists a discrete Fourier transform in one variable. It is given by the kernel

$$e_{mn} := N^{-1/2} \exp(2\pi imn/N), \quad m, n = 1, 2, \dots, N, \quad (39)$$

where  $N$  is a fix positive integer. The matrix  $(e_{mn})_{m,n=1}^N$  is unitary, that is,

$$\sum_k e_{mk} \overline{e_{nk}} = \delta_{mn}, \quad \sum_k e_{km} \overline{e_{kn}} = \delta_{mn}. \quad (40)$$

Let  $f(n)$  be a function of  $n \in \{1, 2, \dots, N\}$ . Then the function

$$\tilde{f}(m) = \sum_{n=1}^N f(n) e_{mn} \equiv N^{-1/2} \sum_{n=1}^N f(n) \exp(2\pi i mn/N) \quad (41)$$

is a finite Fourier transform of  $f(n)$ . The inverse transform is given by

$$f(n) = N^{-1/2} \sum_{m=1}^N \tilde{f}(m) \exp(-2\pi i mn/N). \quad (42)$$

The Plancherel formula  $\sum_{m=1}^N |\tilde{f}(m)|^2 = \sum_{n=1}^N |f(n)|^2$  holds for transforms (41) and (42). This means that the finite Fourier transform conserves the norm introduced in the space of functions on  $\{1, 2, \dots, N\}$ .

In order to derive the finite alternating multivariate Fourier transform, we use the discrete exponential function (39) in the form

$$e_m(s) := N^{-1/2} \exp(2\pi i ms), \quad s \in F_N \equiv \{\frac{1}{N}, \frac{2}{N}, \dots, \frac{N-1}{N}, 1\}, \quad m \in \mathbb{Z}^{\geq 0}, \quad (43)$$

and generate a multivariate discrete exponential function by taking a product of  $n$  copies of these functions,

$$\begin{aligned} e_{\mathbf{m}}(\mathbf{s}) &:= e_{m_1}(s_1) e_{m_2}(s_2) \cdots e_{m_n}(s_n) \\ &= N^{-n/2} \exp(2\pi i m_1 s_1) \exp(2\pi i m_2 s_2) \cdots \exp(2\pi i m_n s_n) \end{aligned} \quad (44)$$

where  $\mathbf{s} = (s_1, s_2, \dots, s_n) \in F_N^n$  and  $\mathbf{m} = (m_1, m_2, \dots, m_n) \in (\mathbb{Z}^{\geq 0})^n$ . We introduce a scalar product in the space of linear combinations of the functions (44) by the formula

$$\langle e_{\mathbf{m}}(\mathbf{s}), e_{\mathbf{m}'}(\mathbf{s}) \rangle \equiv \prod_{i=1}^n \langle e_{m_i}(s_i), e_{m'_i}(s_i) \rangle := \prod_{i=1}^n \sum_{s_i \in F_N} e_{m_i}(s_i) \overline{e_{m'_i}(s_i)} = \delta_{\mathbf{m}\mathbf{m}'} \quad (45)$$

where  $m_i, m'_i \in \{1, 2, \dots, N\}$ . Here we used the relations (40).

We now take multivariate functions (44) for integers  $m_i$  such that

$$N \geq m_1, m_2 \geq m_3 \geq \cdots \geq m_n \geq 1$$

and symmetrize them with respect to the alternating group  $A_n$ . We obtain a finite version of the alternating exponential functions (3),

$$\tilde{E}_{\mathbf{m}}(\mathbf{s}) := |A_n|^{-1/2} \text{sdet}^+(e_{m_i}(s_j))_{i,j=1}^n = |A_n|^{-1/2} N^{-n/2} E_{\mathbf{m}}(\mathbf{s}), \quad (46)$$

where the discrete functions  $e_m(s)$  are given by (43).

The  $n$ -tuples  $\mathbf{s}$  in (46) run over  $F_N^n \equiv F_N \times \cdots \times F_N$  ( $n$  times). We denote by  $\check{F}_N^n$  the subset of  $F_N^n$  consisting of  $\mathbf{s} = (s_1, s_2, \dots, s_n) \in F_N^n$  such that

$$s_1, s_2 \geq s_2 \geq \cdots \geq s_n.$$

The set  $\check{F}_N^n$  is a finite subset of the closure of the fundamental domain  $F(A_n^{\text{aff}})$ .

Acting by permutations  $w \in A_n$  upon  $\check{F}_N^n$  we obtain the whole set  $F_N^n$ , where each point, having some coinciding coordinates  $m_i$ , is repeated several times. Namely, a point  $\mathbf{s}$  is contained  $|G_{\mathbf{s}}|$  times in  $\{w\check{F}_N^n; w \in A_n\}$ , where  $G_{\mathbf{s}}$  is the subgroup of  $A_n$  consisting of elements  $w \in A_n$  such that  $w\mathbf{s} = \mathbf{s}$ .

By  $\check{D}_N^n$  we denote the set of integer  $n$ -tuples  $\mathbf{m} = (m_1, m_2, \dots, m_n)$  such that

$$N \geq m_1, m_2 \geq m_3 \geq \cdots \geq m_n \geq 1.$$

**Proposition.** For  $\mathbf{m}, \mathbf{m}' \in \check{D}_N^n$  the discrete functions (46) satisfy the orthogonality relation

$$\langle \tilde{E}_{\mathbf{m}}(\mathbf{s}), \tilde{E}_{\mathbf{m}'}(\mathbf{s}) \rangle = |A_n| \sum_{\mathbf{s} \in \check{F}_M^n} |G_{\mathbf{s}}|^{-1} \tilde{E}_{\mathbf{m}}(\mathbf{s}) \overline{\tilde{E}_{\mathbf{m}'}(\mathbf{s})} = |G_{\mathbf{m}}| \delta_{\mathbf{m}\mathbf{m}'}. \quad (47)$$

**Proof.** Due to the definition of the scalar product we get

$$\begin{aligned} \langle \tilde{E}_{\mathbf{m}}(\mathbf{s}), \tilde{E}_{\mathbf{m}'}(\mathbf{s}) \rangle &= \sum_{\mathbf{s} \in F_N^n} \tilde{E}_{\mathbf{m}}(\mathbf{s}) \overline{\tilde{E}_{\mathbf{m}'}(\mathbf{s})} \\ &= |A_n|^{-1} |G_{\mathbf{m}}| \sum_{w \in A_n} \prod_{i=1}^n \sum_{s_i \in F_N} e_{m_{w(i)}}(s_i) \overline{e_{m'_{w(i)}}(s_i)} = |G_{\mathbf{m}}| \delta_{\mathbf{m}\mathbf{m}'}, \end{aligned}$$

where we have taken into account that some summands in the expression for sdet in (46) can coincide.

Since functions  $\tilde{E}_{\mathbf{m}}(\mathbf{s})$  are symmetric with respect to  $A_n$ , then

$$\sum_{\mathbf{s} \in F_N^n} \tilde{E}_{\mathbf{m}}(\mathbf{s}) \overline{\tilde{E}_{\mathbf{m}'}(\mathbf{s})} = |A_n| \sum_{\mathbf{s} \in \check{F}_N^n} |G_{\mathbf{s}}|^{-1} \tilde{E}_{\mathbf{m}}(\mathbf{s}) \overline{\tilde{E}_{\mathbf{m}'}(\mathbf{s})},$$

where we have taken into account that under the action by  $A_n$  upon  $\check{F}_N^n$  a point  $\mathbf{s}$  appears  $|G_{\mathbf{s}}|$  times in  $F_N^n$ . This proves the proposition.

Let  $f$  be a function on  $\check{F}_N^n$  (or a symmetric function on  $F_N^n$ ). Then it can be expanded in functions (46) as

$$f(\mathbf{s}) = \sum_{\mathbf{m} \in \check{D}_N^n} a_{\mathbf{m}} \tilde{E}_{\mathbf{m}}(\mathbf{s}). \quad (48)$$

The coefficients  $a_{\mathbf{m}}$  are determined by the formula

$$a_{\mathbf{m}} = |A_n| |G_{\mathbf{m}}|^{-1} \sum_{\mathbf{s} \in \check{F}_N^n} |G_{\mathbf{s}}|^{-1} f(\mathbf{s}) \overline{\tilde{E}_{\mathbf{m}}(\mathbf{s})}. \quad (49)$$

The expansions (48) and (49) follow from the facts that numbers of elements in  $\check{D}_N^n$  and in  $\check{F}_N^n$  are the same and from the orthogonality relation (47) (see also [16]). We refer to expansions (48) and (49) as the *alternating multivariate finite Fourier transforms*.

**Acknowledgements.** The research of the first author was partially supported by Grant 14.01/016 of the State Foundation of Fundamental Research of Ukraine. We acknowledge also partial support for this work by the National Science and Engineering Research Council of Canada, MITACS, the MIND Research Institute.

## REFERENCES

- [1] Klimyk A. U. and Patera J., (Anti)symmetric multivariate exponential functions and corresponding Fourier transforms, *J. Phys. A: Math. Theor.*, **40** (2007), 10473-10489.
- [2] Patera J., C-functions of compact semisimple Lie groups as special functions, *Proc. Inst. Math. Nat. Acad. Sci. of Ukraine*, **30** (2004), 1152-1160.
- [3] Patera J., Compact simple Lie groups and their C-, S-, and E-transforms, *Symmetry, Integrability and Geometry: Methods and Applications*, **1** (2005), paper 025, 6 pages.
- [4] Klimyk A. U. and Patera J., Orbit functions, *Symmetry, Integrability and Geometry: Methods and Applications*, **2** (2006), paper 06, 60 pages.
- [5] Klimyk A. U. and Patera J., Antisymmetric orbit functions, *Symmetry, Integrability and Geometry: Methods and Applications*, **3** (2007), paper 023, 83 pages.

- [6] Atoyán A. and Patera J., Properties of continuous Fourier extension of the discrete cosine transform and its multidimensional generalization, *J. Math. Phys.* **45** (2004), 2468–2491.
- [7] Patera J. and Zaratsyan A., Discrete and continuous cosine transform generalized to Lie groups  $SU(2) \times SU(2)$  and  $O(5)$ , *J. Math. Phys.* **46** (2005), 053514, 17 pages.
- [8] Patera J. and Zaratsyan A., Discrete and continuous cosine transform generalized to Lie groups  $SU(2)$  and  $G_2$ , *J. Math. Phys.* **46** (2005), 113506, 25 pages.
- [9] Kashuba I. and Patera J., Discrete and continuous exponential transforms of simple Lie groups of rank two, *J. Phys. A: Math. Theor.* **40** (2007), 1751–1774.
- [10] Atoyán A. and Patera J., Continuous extension of the discrete cosine transform, and its applications to data processing, in Group Theory and Numerical Analysis, *CRM Proc. Lecture Notes*, Vol. 39, Amer. Math. Soc., Providence, RI, 2005, 1–15.
- [11] Atoyán A., Patera J., Sahakian V., and Akhperjanian A., Fourier transform method for imaging atmospheric Cherenkov telescopes, *Astroparticle Phys.* **23** (2005), 79–95.
- [12] Patera J., Zaratsyan A., and Zhu H.-M., New class of interpolation methods based on discretized Lie group transform, *SPIE Electronic Imaging*, **6064A-06** (2006), S1.
- [13] Germain M., Patera J., and Zaratsyan A., Multiresolution analysis of digital images using the continuous extension of discrete group transform, *SPIE Electronic Imaging*, **6064A-03** (2006), S2.
- [14] Germain M., Patera J. and Allard Y., Cosine transform generalized to Lie groups  $SU(2) \times SU(2)$ ,  $O(5)$ , and  $SU(2) \times SU(2) \times SU(2)$ : application to digital image processing, *Proc. SPIE* **6065** (2006), 387–395.
- [15] Vilenkin N. Ja. and Klimyk A. U., *Representations of Lie Groups and Special Functions*, Vol. 2, Kluwer, Dordrecht, 1993.
- [16] Moody R. V. and Patera J., Orthogonality within the families of  $C$ -,  $S$ -, and  $E$ -functions of any compact semisimple Lie group, *Symmetry, Integrability and Geometry: Methods and Applications*, **2** (2006), paper 76, 14 pages.

BOGOLYUBOV INSTITUTE FOR THEORETICAL PHYSICS, KIEV 03680, UKRAINE  
*E-mail address:* `aklimyk@bitp.kiev.ua`

CENTRE DE RECHERCHES MATHÉMATIQUES, UNIVERSITÉ DE MONTRÉAL, C.P.6128-CENTRE VILLE, MONTRÉAL, H3C 3J7, QUÉBEC, CANADA  
*E-mail address:* `patera@crm.umontreal.ca`