

RESEARCH ARTICLE

The discrete potential Boussinesq equation and its multisoliton solutions

Ken-ichi Maruno^{a*} and Kenji Kajiwara^b

^a*Department of Mathematics, The University of Texas-Pan American, Edinburg, TX 78539-2999;* ^b*Faculty of Mathematics, Kyushu University, 744 Motoooka, Fukuoka 819-0395, Japan*

(March 12, 2022)

An alternate form of discrete potential Boussinesq equation is proposed and its multisoliton solutions are constructed. An ultradiscrete potential Boussinesq equation is also obtained from the discrete potential Boussinesq equation using the ultradiscretization technique. The detail of the multisoliton solutions is discussed by using the reduction technique. The lattice potential Boussinesq equation derived by Nijhoff et al. is also investigated by using the singularity confinement test. The relation between the proposed alternate discrete potential Boussinesq equation and the lattice potential Boussinesq equation by Nijhoff et al. is clarified.

Keywords: discrete potential Boussinesq equation; multisoliton solutions; bilinear equations

AMS Subject Classification: 35Q51; 37K10

1. Introduction

In this article, we propose an alternate form of discrete potential Boussinesq (BSQ) equation

$$U_{n-1}^{m+2}U_{n-1}^{m-1}(-\delta_1U_n^{m-1} + \delta_2U_n^{m+2}) = U_n^{m+1}U_{n-1}^m(-\delta_1U_{n-1}^{m-1} + \delta_2U_{n-1}^{m+2}), \quad (1)$$

and study the relation to the lattice potential BSQ equation proposed by Nijhoff et al. [1, 2]

$$\begin{aligned} & \frac{p^3 - q^3}{p - q + u_{n+1}^{m+1} - u_{n+2}^m} - \frac{p^3 - q^3}{p - q + u_n^{m+2} - u_{n+1}^{m+1}} \\ & - u_n^{m+1}u_{n+1}^{m+2} + u_{n+1}^m u_{n+2}^{m+1} + u_{n+2}^{m+2}(p - q + u_{n+1}^{m+2} - u_{n+2}^{m+1}) \\ & + u_n^m(p - q + u_{n+1}^{m+1} - u_{n+1}^m) \\ & = (2p + q)(u_{n+1}^m + u_{n+1}^{m+2}) - (p + 2q)(u_n^{m+1} + u_{n+2}^{m+1}). \end{aligned} \quad (2)$$

Here $U = U_n^m$ and $u = u_n^m$ are the dynamical field variables at the site (n, m) of a rectangular lattice, and δ_1, δ_2, p, q are the lattice parameters. These are integrable

*Corresponding author. Email: kmaruno@utpa.edu

arXiv:0908.1800v4 [nlin.SI] 10 Dec 2009

discrete analogues of the potential BSQ equation

$$3w_{tt} + 4c_0 w_{xx} - 6w_x w_{xx} - w_{xxx} = 0, \quad (3)$$

which leads to the BSQ equation[3, 4]

$$3u_{tt} + 4c_0 u_{xx} - 3(u^2)_{xx} - u_{xxx} = 0, \quad (4)$$

by $u = w_x$. Note that Eq.(2) is the lowest order member of a hierarchy of lattice equations which is called the lattice Gelfand-Dikii(GD) hierarchy[1]. See also recent works about the lattice potential BSQ equation (2)[5, 6].

In this paper, we propose an alternate form of discrete potential BSQ equation and present multisoliton solutions. It is shown that the multisoliton solution for the discrete potential BSQ equation can be constructed from one for the Hirota-Miwa (discrete KP) equation using the reduction technique. Using the proposed discrete potential BSQ equation, we can construct the ultradiscrete potential BSQ equation. We also study the relation between the alternate discrete potential BSQ equation (1) and the lattice potential BSQ equation (2). Bilinear equations of the lattice potential BSQ equation can be derived systematically using the singularity confinement (SC) test. This reveals the relationship with other discrete potential BSQ equations.

2. Multisoliton solutions for the Boussinesq equation

First, we review fundamental results about reductions of the Kadomtsev-Petviashvili (KP) equation.

It is well known that the solutions of the KP equation[7]

$$(-4u_t + 6uu_x + u_{xxx})_x + 3u_{yy} = 0, \quad (5)$$

can be expressed via a tau function $\tau(x, y, t)$ as $u(x, y, t) = 2\frac{\partial^2}{\partial x^2} \log \tau(x, y, t)$ where $\tau(x, y, t)$ satisfies Hirota's bilinear equation

$$(D_x^4 - 4D_x D_t + 3D_y^2)\tau \cdot \tau = 0, \quad (6)$$

where D_x, D_t, D_y are the Hirota D -operators. Soliton solutions of the bilinear equation can be written in terms of the Wronskian determinant [8, 9, 10, 11]

$$\tau(x, y, t) = \text{Wr}(f_1, \dots, f_N) = \det(f_n^{(n'-1)})_{1 \leq n, n' \leq N}, \quad (7)$$

with $f_n^{(j)} = \partial^j f_n / \partial x^j$, and where f_1, \dots, f_N are a set of linearly independent solutions of the linear system

$$\frac{\partial f}{\partial y} = \frac{\partial^2 f}{\partial x^2}, \quad \frac{\partial f}{\partial t} = \frac{\partial^3 f}{\partial x^3}. \quad (8)$$

For example, ordinary N -soliton solutions are obtained by taking $f_n = e^{\theta_{2n-1}} + e^{\theta_{2n}}$ for $n = 1, \dots, N$, where

$$\theta_m(x, y, t) = k_m x + k_m^2 y + k_m^3 t + \theta_{m;0}, \quad (9)$$

for $m = 1, \dots, 2N$, where the $4N$ parameters $k_1 < \dots < k_{2N}$ and $\theta_{1;0}, \dots, \theta_{2N;0}$ are real constants. For $N = 1$, one obtains the single-soliton solution of KP equation:

$$u_{i,j}(x, y, t) = \frac{1}{2}(k_j - k_i)^2 \operatorname{sech}^2 \left[\frac{1}{2}(\theta_i - \theta_j) \right], \tag{10}$$

where $i = 1$ and $j = 2$. The most general form of the N -soliton solution is given by

$$\tau(x, y, t) = \det(A\Theta K) = \sum_{1 \leq m_1 < \dots < m_N \leq M} V_{m_1, \dots, m_N} A_{m_1, \dots, m_N} \exp \theta_{m_1, \dots, m_N}, \tag{11}$$

where $A = (a_{n,m})$ is the $N \times M$ coefficient matrix, $\Theta = \operatorname{diag}(e^{\theta_1}, \dots, e^{\theta_M})$, $\theta_{m_1, \dots, m_N} = \theta_{m_1} + \dots + \theta_{m_N}$, and the $M \times N$ matrix K is given by $K = (k_m^{n-1})$ [12, 13, 14, 15, 16, 17, 18, 19]. V_{m_1, \dots, m_N} is the Vandermonde determinant $V_{m_1, \dots, m_N} = \prod_{1 \leq j < j' \leq N} (k_{m_{j'}} - k_{m_j})$, and A_{m_1, \dots, m_N} is the $N \times N$ -minor whose n -th column is respectively given by the m_n -th column of the coefficient matrix for $n = 1, \dots, N$. The only time dependence in the tau function comes from the exponential phases θ_{m_1, \dots, m_N} . Also, for all $G \in \operatorname{GL}(N, \mathbb{R})$, the coefficient matrices A and $A' = GA$ produce the same solution of the KP equation. Thus without loss of generality one can consider A to be in row-reduced echelon form (RREF). One can also multiply each column of A by an arbitrary positive constant which can be absorbed in the definition of $\theta_{1;0}, \dots, \theta_{M;0}$.

Real nonsingular (positive) solutions of the KP equation are obtained if $k_1 < \dots < k_M$ and all minors of A are nonnegative. Under these assumptions and some fairly general irreducibility conditions on the coefficient matrix, Eq.(23) produces (N_-, N_+) -soliton solutions of the KP equation with $N_- = M - N$ and $N_+ = N$, as in the simpler case of fully resonant solutions. Asymptotic line solitons are given by Eq.(10) with the indices i and j labeling the phases θ_i and θ_j being swapped in the transition between two dominant phase combinations along the line $\theta_i = \theta_j$. Asymptotic solitons can thus be uniquely characterized by an index pair $[i, j]$ with $1 \leq i < j \leq M$. Recently, line soliton solutions of the KP II equation were classified using this formulation [12, 13, 14, 15, 17, 18]. Elastic 2-soliton solutions are classified into three classes: ordinary (O-type), asymmetric (P-type) and resonant (T-type). The coefficient matrices corresponding to these classes have the following RREFs:

$$A_O = \begin{pmatrix} 1 & a & 0 & 0 \\ 0 & 0 & 1 & b \end{pmatrix}, \quad A_P = \begin{pmatrix} 1 & 0 & 0 & -b \\ 0 & 1 & a & 0 \end{pmatrix}, \quad A_T = \begin{pmatrix} 1 & 0 & -c & -d \\ 0 & 1 & a & b \end{pmatrix}, \tag{12}$$

where $a, b, c, d > 0$ are free parameters with $ad - bc > 0$. These three types of solutions cover disjoint sectors of the 2-soliton parameter space of amplitudes and directions. Moreover, their interaction properties are also different. This difference is obvious in the case of T-type solutions, but also applies to O-type and P-type solutions, since P-type solutions only exist for unequal amplitude, and the interaction phase shift has the opposite sign for O-type and P-type solutions. We remark that inelastic 2-soliton solutions fall into four categories, but we do not focus on them in this article.

It is known that the KP equation reduces to the Korteweg-de Vries (KdV) equation

$$-4u_t + 6uu_x + u_{xxx} = 0 \tag{13}$$

by the symmetry constraint $\partial u/\partial y = 0$ [13, 20, 21, 22]. In the bilinear form, the bilinear KP equation (6) is reduced to the bilinear KdV equation $(D_x^4 - 4D_x D_t)\tau \cdot \tau = 0$ by the constraint of omitting terms including D_y , which is the so-called 2-reduction. As mentioned in [13], this constraint implies that all the solitons of the KdV equation are parallel to the y -axis. Thus we must have a condition $k_j = -k_i$ for each $[i, j]$ -soliton. From the ordering $k_1 < k_2 < \dots < k_{2N}$, we must assume

$$k_1 < k_2 < \dots < k_N < 0, \quad k_{N+j} = -k_{N-j+1} \text{ for } j = 1, \dots, N.$$

This allows only P-type soliton solution in which A -matrix is

$$A = \begin{pmatrix} 1 & 0 & \dots & 0 & 0 & \dots & 0 & a_{1,2N} \\ 0 & 1 & \dots & 0 & 0 & \dots & a_{2,2N-1} & 0 \\ \vdots & \vdots & \dots & \vdots & \vdots & \dots & \vdots & \vdots \\ 0 & 0 & \dots & 1 & a_{N,N} & \dots & 0 & 0 \end{pmatrix}.$$

Let us consider multisoliton solutions of the BSQ equation. It is also known that the KP equation is reduced to the BSQ equation without the term u_{xx} , namely

$$3\delta_0 u_{t't'} - 3(u^2)_{xx} - u_{xxxx} = 0, \quad (14)$$

by the symmetry constraint $\partial u/\partial t = 0$ [20, 21, 22]. Here we introduced a new independent variable t' such that $y = \sqrt{-\delta_0}t'$, $\delta_0 = \pm 1$. In the bilinear form, the bilinear KP equation (6) is reduced to $(D_x^4 - 3\delta_0 D_{t'}^2)\tau \cdot \tau = 0$ by the constraint of omitting terms including D_t , which is the so-called 3-reduction. This condition implies that all the solitons of Eq.(14) are parallel to the t -axis. Thus we must have a condition $k_j^2 + k_j k_i + k_i^2 = 0$, i.e. $k_j = \omega k_i$ or $k_j = \omega^2 k_i$ ($\omega = -1/2 + i\sqrt{3}/2$, $\omega^2 = -1/2 - i\sqrt{3}/2$, $\omega^3 = 1$), for $[i, j]$ -soliton. In the BSQ equation, k_1, \dots, k_{2N} take complex values in general. Since k_1, \dots, k_{2N} are complex values, we cannot consider the ordering of k_1, \dots, k_{2N} which was assumed when we consider KP soliton solutions. However, from the constraint of k_j , we have a restriction to the A -matrix such that each row has only 2 or 3 nonzero elements and each column has only one element. Note that there are only 2 nonzero elements in each row and only one element in each column in the case of the KdV equation because of the constraint $k_j = -k_i$ [13]. Note also that there is no such restriction in the case of the KP equation. For 2-soliton solutions, 2 types of 2-soliton interactions are possible, i.e. 2 elastic soliton interactions (O-type and P-type), other 2-soliton interactions are impossible because some columns in A -matrix have 2 elements. However, there is no distinction between O-type and P-type solitons because we do not have the ordering of k_1, \dots, k_{2N} . Thus 2-soliton interaction of the BSQ equation is actually only of one type. To get real solutions, we must remove imaginary numbers by the following ways.

The case of 2 elements:

(i) Suppose that we have 2 elements in the i -th row of the A -matrix. Let the corresponding wave numbers of these elements be k_{j_1}, k_{j_2} ; (ii) Let $k_{j_2} = \omega k_{j_1}$ (or $k_{j_2} = \omega^2 k_{j_1}$); (iii) Using the gauge invariance of τ -function, each element in the Wronskian determinant can be $f_i \sim 1 + a_{i,j_2} \exp((\omega - 1)k_{j_1}x + \sqrt{-\delta_0}(\omega^2 - 1)k_{j_1}^2 t' + \theta_{j_2,0})$ (or $f_i \sim 1 + a_{i,j_2} \exp((\omega^2 - 1)k_{j_1}x + \sqrt{-\delta_0}(\omega - 1)k_{j_1}^2 t' + \theta_{j_2,0})$); (iv) Reparametrize k_{j_1} and k_{j_2} by $\kappa_i = (\omega - 1)k_{j_1}$ and $\Omega_i = \sqrt{-\delta_0}(\omega^2 - 1)k_{j_1}^2$ (or $\kappa_i = (\omega^2 - 1)k_{j_1}$ and $\Omega_i = \sqrt{-\delta_0}(\omega - 1)k_{j_1}^2$). Then we have $f_i \sim 1 + a_{i,j_2} \exp(\kappa_i x + \Omega_i t' + \theta_{j_2,0})$ with linear dispersion relations $\kappa_i^4 - 3\delta_0 \Omega_i^2 = 0$; (vi) So a set of $f_i \sim$

$1 + a_{i,j_2} \exp(\kappa_i x + \Omega_i t' + \theta_{j_2,0})$ with linear dispersion relations $\kappa_i^4 - 3\delta_0 \Omega_i^2 = 0$ gives the real and regular multisoliton solutions.

The case of 3 elements:

(i) Suppose that we have 3 elements in the i -th row of the A -matrix. Let the corresponding wave numbers of these elements be $k_{j_1}, k_{j_2}, k_{j_3}$; (ii) Let $k_{j_2} = \omega k_{j_1}$ and $k_{j_3} = \omega^2 k_{j_1}$; (iii) Using the gauge invariance of τ -function, each element in the Wronskian determinant can be $f_i \sim 1 + a_{i,j_2} \exp((\omega - 1)k_{j_1}x + (\omega^2 - 1)k_{j_1}^2 y + \theta_{j_2,0}) + a_{i,j_3} \exp((\omega^2 - 1)k_{j_1}x + (\omega - 1)k_{j_1}^2 y + \theta_{j_3,0})$; (iv) Reparametrize parameters k_{j_1}, k_{j_2} and k_{j_3} by $\kappa_{i,2} = k_{j_2} - k_{j_1}, \kappa_{i,3} = k_{j_3} - k_{j_1}, \Omega_{i,2} = \sqrt{-\delta_0}(k_{j_2}^2 - k_{j_1}^2)$ and $\Omega_{i,3} = \sqrt{-\delta_0}(k_{j_3}^2 - k_{j_1}^2)$. Then we have $f_i \sim 1 + a_{i,j_2} \exp(\kappa_i x + \Omega_i t' + \theta_{j_2,0}) + a_{i,j_3} \exp(\kappa_i x + \Omega_i t' + \theta_{j_3,0})$ with linear dispersion relations $\kappa_{i,j}^4 - 3\delta_0 \Omega_{i,j}^2 = 0$ for $j = 1, 2$; For example, consider $N = 1$. This gives Y-shape soliton resonance interaction.

The KP equation is reduced to the BSQ equation

$$3\delta_0 u_{t't'} + 4c_0 u_{xx} - 3(u^2)_{xx} - u_{xxxx} = 0, \tag{15}$$

by the symmetry constraint $\frac{\partial}{\partial t} = c_0 \frac{\partial}{\partial x}$. Here we introduced a new independent variable t' such that $y = \sqrt{-\delta_0} t'$. In the bilinear form, the bilinear KP equation (6) is reduced to $(D_x^4 - 4c_0 D_x^2 - 3\delta_0 D_{t'}^2)\tau \cdot \tau = 0$ by the constraint of replacing D_t by $c_0 D_x$, which is the so-called 3-pseudo reduction[22]. To realize this constraint in the multisoliton solutions, we can add the constraint $k_j^3 - k_i^3 = c_0(k_j - k_i)$, i.e.

$k_j = \frac{1}{2} \left(-k_i \pm \sqrt{4c_0 - 3k_i^2} \right)$, to the soliton solutions of the KP equation. Note that k_j can be real if $4c_0 > 3k_i^2$. So we can assume the ordering of $k_1 < \dots < k_{2N}$. From the constraint we have a restriction to the A -matrix such that each row has only 2 or 3 nonzero elements and each column has only one element. For 2-soliton solutions, 2 types of 2-soliton interactions are possible, i.e. 2 elastic soliton interactions (O-type and P-type), other 2-soliton interactions are impossible because some columns in A -matrix have 2 elements.

Let us consider explicit real and regular multisoliton solutions. For the case of $\delta_0 = -1$, (i) Suppose that we have 2 elements in the i -th row of the A -matrix. Let the corresponding wave numbers of these elements be k_{j_1}, k_{j_2} ; (ii) Let $k_{j_2} = \frac{1}{2} \left(-k_{j_1} \pm \sqrt{4c_0 - 3k_{j_1}^2} \right)$; (iii) Each element in the Wronskian determinant can be $f_i \sim a_{i,j_1} \exp(k_{j_1}x + k_{j_1}^2 t' + \theta_{j_1,0}) + a_{i,j_2} \exp(k_{j_2}x + k_{j_2}^2 t' + \theta_{j_2,0})$. This gives multisoliton solutions for the BSQ equation.

If we have 3 elements in the i -th row of the A -matrix, each element in the Wronskian determinant can be $f_i \sim a_{i,j_1} \exp(k_{j_1}x + k_{j_1}^2 t' + \theta_{j_1,0}) + a_{i,j_2} \exp(k_{j_2}x + k_{j_2}^2 t' + \theta_{j_2,0}) + a_{i,j_3} \exp(k_{j_3}x + k_{j_3}^2 t' + \theta_{j_3,0})$. For $N = 1$, this gives Y-shape resonant soliton solution.

For the case of $\delta_0 = 1$, (i) Suppose that we have 2 elements in the i -th row of the A -matrix. Let the corresponding wave numbers of these elements be k_{j_1}, k_{j_2} ; (ii) Let $k_{j_2} = \frac{1}{2} \left(-k_{j_1} \pm \sqrt{4c_0 - 3k_{j_1}^2} \right)$; (iii) Using the gauge invariance of τ -function, each element in the Wronskian determinant can be $f_i \sim 1 + a_{i,j_2} \exp((k_{j_2} - k_{j_1})x + i(k_{j_2}^2 - k_{j_1}^2)t' + \theta_{j_2,0})$; (iv) Reparametrize k_{j_1} and k_{j_2} by $\kappa_i = k_{j_2} - k_{j_1}$ and $\Omega_i = i(k_{j_2}^2 - k_{j_1}^2)$. Then we have $f_i \sim 1 + a_{i,j_2} \exp(\kappa_i x + \Omega_i t' + \theta_{j_2,0})$ with linear dispersion relations $\kappa_i^4 - 4c_0 \kappa_i^2 - 3\delta_0 \Omega_i^2 = 0$.

Next, we consider the case of complex parameters. Suppose that a parameter k_i corresponds to the pivot of A -matrix. If $4c_0 < 3k_i^2$, then we have a complex parameter $k_j = \frac{1}{2} \left(-k_i \pm i\sqrt{3k_i^2 - 4c_0} \right)$ among k_1, \dots, k_{2N} . Since some of k_1, \dots, k_{2N} are complex values, we cannot consider the ordering of k_1, \dots, k_{2N} which was assumed

when we consider KP soliton solutions. However, from the constraint we have a restriction to the A -matrix such that each row has only 2 or 3 nonzero elements and each column has only one element. For 2-soliton solutions, 2 types of 2-soliton interactions are possible, i.e. 2 elastic soliton interactions (O-type and P-type), other 2-soliton interactions are impossible because some columns in A -matrix have 2 elements. However, there is no distinction between O-type and P-type solitons because we don't have the ordering of k_1, \dots, k_{2N} . Thus 2-soliton interaction of the BSQ equation is actually only one type again. To get real solutions, we must remove imaginary numbers by the following ways.

The case of 2 elements:

(i) Suppose that we have 2 elements in the i -th row of the A -matrix. Let the corresponding wave numbers of these elements be k_{j_1}, k_{j_2} ; (ii) Let $k_{j_2} = \frac{1}{2} \left(-k_{j_1} \pm i\sqrt{3k_{j_1}^2 - 4c_0} \right)$; (iii) Using the gauge invariance of τ -function, each element in the Wronskian determinant can be $f_i \sim 1 + a_{i,j_2} \exp((k_{j_2} - k_{j_1})x + (k_{j_2}^2 - k_{j_1}^2)y + \theta_{j_2,0})$; (iv) Reparametrize k_{j_1} and k_{j_2} by $\kappa_i = k_{j_2} - k_{j_1}$ and $\Omega_i = \sqrt{-\delta_0}(k_{j_2}^2 - k_{j_1}^2)$. Then we have $f_i \sim 1 + a_{i,j_2} \exp(\kappa_i x + \Omega_i t' + \theta_{j_2,0})$ with linear dispersion relations $\kappa_i^4 - 4c_0\kappa_i^2 - 3\delta_0\Omega_i^2 = 0$; (vi) So a set of $f_i \sim 1 + a_{i,j_2} \exp(\kappa_i x + \Omega_i t' + \theta_{j_2,0})$ with linear dispersion relations $\kappa_i^4 - 4c_0\kappa_i^2 - 3\Omega_i^2 = 0$ gives the real and regular multisoliton solutions.

The case of 3 elements:

(i) Suppose that we have 3 elements in the i -th row of the A -matrix. Let the corresponding wave numbers of these elements be $k_{j_1}, k_{j_2}, k_{j_3}$; (ii) Let $k_{j_2} = \frac{1}{2} \left(-k_{j_1} + i\sqrt{3k_{j_1}^2 - 4c_0} \right)$ and $k_{j_3} = \frac{1}{2} \left(-k_{j_1} - i\sqrt{3k_{j_1}^2 - 4c_0} \right)$; (iii) Using the gauge invariance of τ -function, each element in the Wronskian determinant can be $f_i \sim 1 + a_{i,j_2} \exp((k_{j_2} - k_{j_1})x + \sqrt{-\delta_0}(k_{j_2}^2 - k_{j_1}^2)t' + \theta_{j_2,0}) + a_{i,j_3} \exp((k_{j_3} - k_{j_1})x + \sqrt{-\delta_0}(k_{j_3}^2 - k_{j_1}^2)t' + \theta_{j_3,0})$; (iv) Reparametrize k_{j_1}, k_{j_2} and k_{j_3} by $\kappa_{i,2} = k_{j_2} - k_{j_1}$, $\kappa_{i,3} = k_{j_3} - k_{j_1}$, $\Omega_{i,2} = \sqrt{-\delta_0}(k_{j_2}^2 - k_{j_1}^2)$ and $\Omega_{i,3} = \sqrt{-\delta_0}(k_{j_3}^2 - k_{j_1}^2)$. Then we have $f_i \sim 1 + a_{i,j_2} \exp(\kappa_{i,2}x + \Omega_{i,2}t' + \theta_{j_2,0}) + a_{i,j_3} \exp(\kappa_{i,3}x + \Omega_{i,3}t' + \theta_{j_3,0})$ with linear dispersion relations $\kappa_{i,j}^4 - 4c_0\kappa_{i,j}^2 - 3\delta_0\Omega_{i,j}^2 = 0$ for $j = 1, 2$; For example, consider $N = 1$. This gives Y-shape soliton resonance interaction.

3. Discrete analogues of the potential Boussinesq equation

3.1. The discrete potential Boussinesq equation

Here, we present the main theorem in this article.

Theorem 3.1: *The difference-difference equation*

$$U_{n-1}^{m+2} U_{n-1}^{m-1} (-\delta_1 U_n^{m-1} + \delta_2 U_n^{m+2}) = U_n^{m+1} U_{n-1}^m (-\delta_1 U_{n-1}^{m-1} + \delta_2 U_{n-1}^{m+2}), \quad (16)$$

where $\delta_1 = a_2(a_1 - a_3)$ and $\delta_2 = a_3(a_1 - a_2)$ and a_1, a_2, a_3 are arbitrary real constants, is an integrable discrete analogue of the potential BSQ equation (3).

Moreover, the discrete potential BSQ equation has multisoliton solutions

$$U_n^m = \frac{\tau_{n+1}^m}{\tau_n^m}, \quad \tau_n^m = \det(\mathcal{A}\Theta P), \quad (17)$$

where $\mathcal{A} = (\alpha_{n,m})$ is the $N \times 2N$ coefficient matrix, $\Theta = \text{diag}(e^{\theta_1}, \dots, e^{\theta_{2N}})$, $e^{\theta_j} = p_j^n (1 - p_j a_1)^{-n} (1 - p_j a_2)^{-m}$, and the $2N \times N$ matrix P is given by $P = (p_m^{n-1})$. where the $2N$ parameters p_1, \dots, p_{2N} are real constants. The \mathcal{A} -matrix has a restriction

such that each row has only 2 or 3 nonzero elements and each column has only one element. In the case having 2 elements (i, j_1) and (i, j_2) in the i -th row of the \mathcal{A} -matrix, p_{j_2} must satisfy a reduction condition

$$p_{j_2} = \frac{1}{a_2} + \frac{(1 - a_3 p_{j_1})}{2a_3} \pm \frac{\sqrt{a_2(1 - a_3 p_{j_1})(a_2 - 4a_3 + 3a_2 a_3 p_{j_1})}}{2a_2 a_3}. \quad (18)$$

Proof: The Hirota-Miwa (discrete KP) equation is written as

$$\begin{aligned} & a_1(a_2 - a_3)\tau(n_1 + 1, n_2, n_3)\tau(n_1, n_2 + 1, n_3 + 1) \\ & + a_2(a_3 - a_1)\tau(n_1, n_2 + 1, n_3)\tau(n_1 + 1, n_2, n_3 + 1) \\ & + a_3(a_1 - a_2)\tau(n_1, n_2, n_3 + 1)\tau(n_1 + 1, n_2 + 1, n_3) = 0, \end{aligned} \quad (19)$$

where τ depends on three discrete independent variables n_1, n_2 and n_3 , and a_1, a_2 and a_3 are the difference intervals for n_1, n_2 and n_3 , respectively [23].

The Casorati determinant solution for the Hirota-Miwa equation (19) is as follows[24]:

$$\tau(n_1, n_2, n_3) = \det(\psi_i(n_1, n_2, n_3; s + j - 1))_{1 \leq i, j \leq N}, \quad (20)$$

where ψ_1, \dots, ψ_N are a set of linearly independent solutions of the linear system

$$\Delta_{n_j} \psi_i(n_1, n_2, n_3; s) = \psi_i(n_1, n_2, n_3; s + 1), \quad (j = 1, 2, 3).$$

Here Δ_{n_j} are the backward difference operators:

$$\Delta_{n_j} f(n_j) \equiv \frac{f(n_j) - f(n_j - 1)}{a_j}, \quad (j = 1, 2, 3). \quad (21)$$

For example, ordinary N -soliton solutions are obtained by taking

$$\begin{aligned} \psi_i(n_1, n_2, n_3; s) &= \alpha_{2i-1} p_{2i-1}^s (1 - p_{2i-1} a_1)^{-n_1} (1 - p_{2i-1} a_2)^{-n_2} (1 - p_{2i-1} a_3)^{-n_3} \\ &+ \alpha_{2i} p_{2i}^s (1 - p_{2i} a_1)^{-n_1} (1 - p_{2i} a_2)^{-n_2} (1 - p_{2i} a_3)^{-n_3}, \end{aligned} \quad (22)$$

for $n = 1, \dots, N$ where the $4N$ parameters $p_1 < \dots < p_{2N}$ and $\alpha_1, \dots, \alpha_{2N}$ are positive real constants. The most general form of the N -soliton solution is given by

$$\tau(n_1, n_2, n_3) = \det(\mathcal{A}\Theta P) = \sum_{1 \leq m_1 < \dots < m_N \leq 2N} V_{m_1, \dots, m_N} \mathcal{A}_{m_1, \dots, m_N} \exp \theta_{m_1, \dots, m_N}, \quad (23)$$

where $\mathcal{A} = (\alpha_{n,m})$ is the $N \times 2N$ coefficient matrix, $\Theta = \text{diag}(e^{\theta_1}, \dots, e^{\theta_{2N}})$, $e^{\theta_j} = p_j^s (1 - p_j a_1)^{-n_1} (1 - p_j a_2)^{-n_2} (1 - p_j a_3)^{-n_3}$, and the $2N \times N$ matrix P is given by $P = (p_m^{n-1})$. V_{m_1, \dots, m_N} is the Vandermonde determinant $V_{m_1, \dots, m_N} = \prod_{1 \leq j < j' \leq N} (p_{m_{j'}} - p_{m_j})$, and $\mathcal{A}_{m_1, \dots, m_N}$ is the $N \times N$ -minor whose n -th column is respectively given by the m_n -th column of the coefficient matrix for $n = 1, \dots, N$. For all $G \in \text{GL}(N, \mathbb{R})$, the coefficient matrices \mathcal{A} and $\mathcal{A}' = G\mathcal{A}$ produce the same solution of the Hirota-Miwa equation. Thus without loss of generality one can consider \mathcal{A} to be in RREF.

Let us consider the 3-reduction condition

$$\tau(n_1, n_2, n_3) \asymp \tau(n_1, n_2 - 2, n_3 - 1), \tag{24}$$

for arbitrary n_1, n_2 and n_3 . Applying this reduction condition, we can omit the dependence on n_3 and obtain the bilinear form

$$\begin{aligned} & a_1(a_2 - a_3)\tau(n_1 + 1, n_2)\tau(n_1, n_2 - 1) + a_2(a_3 - a_1)\tau(n_1, n_2 + 1)\tau(n_1 + 1, n_2 - 2) \\ & + a_3(a_1 - a_2)\tau(n_1, n_2 - 2)\tau(n_1 + 1, n_2 + 1) = 0. \end{aligned} \tag{25}$$

After the change of variables $n_1 \rightarrow n, n_2 \rightarrow m, \tau(n_1, n_2) \rightarrow \tau_n^m$, we obtain

$$a_1(a_2 - a_3)\tau_{n+1}^{m+1}\tau_n^m + a_2(a_3 - a_1)\tau_n^{m+2}\tau_{n+1}^{m-1} + a_3(a_1 - a_2)\tau_n^{m-1}\tau_{n+1}^{m+2} = 0, \tag{26}$$

which is the bilinear form of the discrete potential BSQ equation.

Now we impose a constraint on the parameters of the solution so that the reduction condition is satisfied. For simplicity, we consider the case in which ψ_1, \dots, ψ_N have 2 terms. Then we observe

$$\begin{aligned} & \psi_i(n_1, n_2 + 2, n_3 + 1; s) \\ & = p_{j_1}^s (1 - p_{j_1} a_1)^{-n_1} (1 - p_{j_1} a_2)^{-n_2 - 2} (1 - p_{j_1} a_3)^{-n_3 - 1} \\ & \quad + p_{j_2}^s (1 - p_{j_2} a_1)^{-n_1} (1 - p_{j_2} a_2)^{-n_2 - 2} (1 - p_{j_2} a_3)^{-n_3 - 1} \\ & = p_{j_1}^s (1 - p_{j_1} a_1)^{-n_1} (1 - p_{j_1} a_2)^{-n_2 - 2} (1 - p_{j_1} a_3)^{-n_3 - 1} \\ & \quad \times \left[1 + C_i \left(\frac{p_{j_2}}{p_{j_1}} \right)^s \left(\frac{1 - p_{j_2} a_1}{1 - p_{j_1} a_1} \right)^{-n_1} \left(\frac{1 - p_{j_2} a_2}{1 - p_{j_1} a_2} \right)^{-n_2} \left(\frac{1 - p_{j_2} a_3}{1 - p_{j_1} a_3} \right)^{-n_3} \right], \end{aligned} \tag{27}$$

where

$$C_i = \left(\frac{1 - p_{j_2} a_2}{1 - p_{j_1} a_2} \right)^{-2} \left(\frac{1 - p_{j_2} a_3}{1 - p_{j_1} a_3} \right)^{-1}.$$

If we apply the reduction condition

$$(1 - p_{j_1} a_2)^2 (1 - p_{j_1} a_3) = (1 - p_{j_2} a_2)^2 (1 - p_{j_2} a_3), \tag{28}$$

i.e.

$$p_{j_2} = \frac{1}{a_2} + \frac{(1 - a_3 p_{j_1})}{2a_3} \pm \frac{\sqrt{a_2(1 - a_3 p_{j_1})(a_2 - 4a_3 + 3a_2 a_3 p_{j_1})}}{2a_2 a_3}, \tag{29}$$

we obtain

$$\psi_i(n_1, n_2, n_3; s) = (1 - p_{j_2} a_2)^{-2} (1 - p_{j_2} a_3)^{-1} \psi_i(n_1, n_2 - 2, n_3 - 1; s), \tag{30}$$

Finally we have a reduction condition

$$\tau(n_1, n_2, n_3) = \prod_{k=1}^N (1 - p_k a_2)^{-2} (1 - p_k a_3)^{-1} \tau(n_1, n_2 - 2, n_3 - 1). \tag{31}$$

In a similar way discussed in section 2, we can consider the general multisoliton solutions of the discrete potential BSQ equation. \square

The reality condition is $p_{j_1} < 1/a_3$, $p_{j_1} > (-a_2 + 4a_3)/(3a_2a_3)$ for $a_2 < a_3$, and $p_{j_1} > 1/a_3$, $p_{j_1} < (-a_2 + 4a_3)/(3a_2a_3)$ for $a_2 > a_3$. With parameters satisfying these conditions, we can construct real and regular multisoliton solutions using the formula in Theorem 3.1.

Otherwise, we must use the technique of reparametrization which was used in section 2. The procedure is as follows. (i) Suppose that we have 2 elements in the i -th row of the A -matrix. Let the corresponding complex wave numbers of these elements be p_{j_1} and p_{j_2} ; (iii) Using the gauge invariance of τ -function, each element in the Wronskian determinant can be $f_i \sim 1 + \alpha_{i,j_2} K_i^{-n} \Omega_i^{-m}$ where $K_i = (1 - a_1 p_{j_2}) / (1 - a_1 p_{j_1})$, $\Omega_i = (1 - a_2 p_{j_2}) / (1 - a_2 p_{j_1})$; (iv) Choose new parameters K_i and Ω_i to be real numbers.

It is easy to take the ultradiscrete limit in Eq.(16).

Theorem 3.2: *The ultradiscrete potential BSQ equation is*

$$V_{n-1}^{m+2} + V_{n-1}^{m-1} + L_n^{m+2} = V_n^{m+1} + V_{n-1}^m + L_{n-1}^{m+2}, \quad (32)$$

$$V_n^{m+2} = \max(L_n^{m+2}, V_n^{m-1} + c_1) - c_2. \quad (33)$$

Proof: Use the standard procedure of ultradiscretization. Equation (16) is rewritten in the form of

$$U_{n-1}^{m+2} U_{n-1}^{m-1} I_n^{m+2} = U_n^{m+1} U_{n-1}^m I_{n-1}^{m+2}, \quad (34)$$

$$I_n^{m+2} = -\delta_1 U_n^{m-1} + \delta_2 U_n^{m+2}. \quad (35)$$

Introduce new variables $U_n^m = \exp(V_n^m/\epsilon)$, $I_n^m = \exp(L_n^m/\epsilon)$, $\delta_1 = \exp(c_1/\epsilon)$, $\delta_2 = \exp(c_2/\epsilon)$. Then take the limit $\epsilon \rightarrow 0^+$ using the formula $\lim_{\epsilon \rightarrow 0^+} \epsilon \ln(\exp(A/\epsilon) + \exp(B/\epsilon)) = \max(A, B)$ for $A, B \in \mathbb{R}_{\geq 0}$ [25]. \square

Remark 1:

Applying the gauge transformation $U_n^m \rightarrow (-1)^{n+m} U_n^m$ to Eq.(16), we have

$$U_{n-1}^{m+2} U_{n-1}^{m-1} I_n^{m+2} = U_n^{m+1} U_{n-1}^m I_{n-1}^{m+2}, \quad (36)$$

$$I_n^{m+2} = \delta_1 U_n^{m-1} + \delta_2 U_n^{m+2}. \quad (37)$$

This form gives a more symmetric form of the ultradiscrete BSQ equation

$$V_{n-1}^{m+2} + V_{n-1}^{m-1} + L_n^{m+2} = V_n^{m+1} + V_{n-1}^m + L_{n-1}^{m+2}, \quad (38)$$

$$L_n^{m+2} = \max(V_n^{m-1} + c_1, V_n^{m+2} + c_2). \quad (39)$$

Remark 2:

Date et al. proposed another discrete potential BSQ equation[26]

$$\begin{aligned} & v_{n-1}^m v_n^{m-1} (a_1 (a_2 - a_3) v_{n+1}^m + a_2 (a_3 - a_1) v_n^{m+1}) \\ &= v_{n-1}^{m-1} v_{n+1}^{m+1} (a_1 (a_2 - a_3) v_{n+1}^{m-1} + a_2 (a_3 - a_1) v_{n-1}^m). \end{aligned} \quad (40)$$

By the transformation $v_n^m = \tau_{n+1}^{m+1}/\tau_n^m$, we obtain a bilinear equation

$$a_1(a_2 - a_3)\tau_{n+1}^m\tau_{n-1}^m + a_2(a_3 - a_1)\tau_n^{m+1}\tau_n^{m-1} + a_3(a_1 - a_2)\tau_{n-1}^{m-1}\tau_{n+1}^{m+1} = 0. \tag{41}$$

This bilinear equation is obtained by adding the reduction condition

$$\tau(n_1, n_2, n_3) \simeq \tau(n_1 - 1, n_2 - 1, n_3 - 1), \tag{42}$$

which gives yet another 3-reduction.

For this discrete potential BSQ equation, we can also make an ultradiscrete analogue of the potential BSQ equation

$$X_{n-1}^m + X_n^{m-1} + Y_n^{m+1} = X_{n-1}^{m-1} + X_{n+1}^{m+1} + Y_{n-1}^m, \tag{43}$$

$$Y_n^m = \max(X_{n+1}^{m-1} + c_1, X_n^m + c_2), \tag{44}$$

taking the ultradiscrete limit after setting $v_n^m = \exp(X_n^m/\epsilon)$, $w_n^m = \exp(Y_n^m/\epsilon)$, $\alpha_1 = \exp(c_1/\epsilon)$, $\alpha_2 = \exp(c_2/\epsilon)$ where $w_n^m = \alpha_1 v_{n+1}^{m-1} + \alpha_2 v_n^m$, $\alpha_1 = a_1(a_2 - a_3)$, $\alpha_2 = a_2(a_3 - a_1)$.

3.2. The lattice potential Boussinesq equation

Singularity confinement (SC) test was proposed by Grammaticos et al. as a detector of integrability in discrete systems[27]. This property has been applied to several problems [28, 29, 30, 31, 32]. The SC test is also powerful tool for constructing solutions for discrete integrable systems[30, 31, 32]. In this section, we apply the SC test to the lattice potential BSQ equation of Nijhoff et al. and obtain bilinear equations using the result of SC test.

We start from the slightly simplified form of the lattice potential BSQ equation

$$\begin{aligned} & \frac{p^3 - q^3}{p - q + u_{n+1}^{m+1} - u_{n+2}^m} - \frac{p^3 - q^3}{p - q + u_n^{m+2} - u_{n+1}^{m+1}} \\ &= (p + 2q + u_{n+1}^m - u_{n+2}^{m+2})(p - q + u_{n+1}^{m+2} - u_{n+2}^{m+1}) \\ & \quad - (p + 2q + u_n^m - u_{n+1}^{m+2})(p - q + u_n^{m+1} - u_{n+1}^m). \end{aligned} \tag{45}$$

Introducing new variables

$$I_n^m = p - q + u_{n-1}^{m+1} - u_n^m, \quad V_n^m = p + 2q + u_{n-1}^{m-2} - u_n^m, \tag{46}$$

Eq.(45) is written as

$$\frac{p^3 - q^3}{I_{n+2}^m} - \frac{p^3 - q^3}{I_{n+1}^{m+1}} = V_{n+2}^{m+2} I_{n+2}^{m+1} - V_{n+1}^{m+2} I_{n+1}^m, \tag{47}$$

$$I_n^m + V_{n+1}^{m+2} = I_{n+1}^{m+2} + V_n^{m+3}. \tag{48}$$

After the transformation of the independent variables $m + n \rightarrow m$, Eqs.(45),(46)

and (47), (48) are rewritten in the following form:

$$\begin{aligned} & \frac{p^3 - q^3}{p - q + u_{n+1}^{m+2} - u_{n+2}^{m+2}} - \frac{p^3 - q^3}{p - q + u_n^{m+2} - u_{n+1}^{m+2}} \\ &= (p + 2q + u_{n+1}^{m+1} - u_{n+2}^{m+4})(p - q + u_{n+1}^{m+3} - u_{n+2}^{m+3}) \\ & \quad - (p + 2q + u_n^m - u_{n+1}^{m+3})(p - q + u_n^{m+1} - u_{n+1}^{m+1}), \end{aligned} \tag{49}$$

$$I_n^m = p - q + u_{n-1}^m - u_n^m, \quad V_n^m = p + 2q + u_{n-1}^{m-3} - u_n^m, \tag{50}$$

and

$$\frac{p^3 - q^3}{I_{n+2}^{m+2}} - \frac{p^3 - q^3}{I_{n+1}^{m+2}} = V_{n+2}^{m+4} I_{n+2}^{m+3} - V_{n+1}^{m+3} I_{n+1}^{m+1}, \tag{51}$$

$$I_n^m + V_{n+1}^{m+3} = I_{n+1}^{m+3} + V_n^{m+3}. \tag{52}$$

After some calculation, we get the form which is suitable to perform the SC test:

$$V_n^m = \frac{1}{I_n^{m-1}} \left(V_{n-1}^{m-1} I_{n-1}^{m-3} + \frac{p^3 - q^3}{I_n^{m-2}} - \frac{p^3 - q^3}{I_{n-1}^{m-2}} \right), \tag{53}$$

$$I_n^m = I_{n-1}^{m-3} + V_n^m - V_{n-1}^m. \tag{54}$$

Now we perform the SC test. Suppose that we have some initial data which can be evolved using the above system. Let us assume that during the successive applications of the above system the value of I at $(m - 1, n)$ becomes zero. The point where this occurs depends on the initial data. Then we have the following pattern of 0 and ∞ :

$$\{I_n^{m-1}, I_n^m, I_{n+1}^m, I_{n+1}^{m+1}\} \rightarrow \{0, \infty, \infty, 0\}, \quad \{V_n^m, V_{n+1}^{m+3}\} \rightarrow \{\infty, \infty\}.$$

One can see nonzero finite values for all dependent variables in further steps, so the singularity is perfectly confined. Suppose that this pattern was created by a function F_n^m which has a zero at (m, n) .

Using the above singularity pattern, we obtain the independent variable transformation

$$I_n^m = \alpha \frac{F_n^{m+1} F_{n-1}^{m-1}}{F_n^m F_{n-1}^m}. \tag{55}$$

Since u_n^m and I_n^m are related by Eq.(50), u_n^m should have F_n^m in the denominator. Thus we set

$$u_n^m = \beta \frac{G_n^m}{F_n^m}. \tag{56}$$

From Eqs.(46) and (56), we obtain

$$I_n^m = p - q + \beta \frac{G_{n-1}^m}{F_{n-1}^m} - \beta \frac{G_n^m}{F_n^m}, \quad V_n^m = p + 2q + \beta \frac{G_{n-1}^{m-3}}{F_{n-1}^{m-3}} - \beta \frac{G_n^m}{F_n^m}. \tag{57}$$

Using Eqs.(53), (54), (55) and (57), we obtain the following equations:

$$(p - q)F_{n-1}^{m-1}F_n^{m-1} + \beta G_{n-1}^{m-1}F_n^{m-1} - \beta F_{n-1}^{m-1}G_n^{m-1} = \alpha F_n^m F_{n-1}^{m-2}, \quad (58)$$

$$\begin{aligned} & \frac{p^3 - q^3}{\alpha} F_n^{m+1} F_{n+1}^m - \alpha(p + 2q)F_{n+1}^{m+2} F_n^{m-1} \\ & - \alpha\beta G_{n+1}^{m-1} F_{n+1}^{m+2} + \alpha\beta F_n^{m-1} G_{n+1}^{m+2} = \gamma F_n^m F_{n+1}^{m+1}, \end{aligned} \quad (59)$$

$$\begin{aligned} & \alpha \frac{F_n^m F_{n-1}^{m-2}}{F_n^{m-1} F_{n-1}^{m-1}} - \frac{p^3 - q^3}{\alpha^2} \frac{F_{n-1}^{m+1} F_n^m}{F_n^{m+2} F_{n-1}^{m-1}} + \frac{\gamma}{\alpha} \frac{F_{n-1}^m F_n^{m+1}}{F_n^{m+2} F_{n-1}^{m-1}} \\ & = \alpha \frac{F_{n+1}^{m+3} F_n^{m+1}}{F_{n+1}^{m+2} F_n^{m+2}} - \frac{p^3 - q^3}{\alpha^2} \frac{F_{n+1}^{m+1} F_n^m}{F_{n+1}^{m+2} F_n^{m-1}} + \frac{\gamma}{\alpha} \frac{F_n^m F_{n+1}^{m+1}}{F_{n+1}^{m+2} F_n^{m-1}}, \end{aligned} \quad (60)$$

where γ is a decoupling constant. Note that V is written in the following form:

$$V_n^m = (p + 2q) + \beta \frac{G_{n-1}^{m-2}}{F_{n-1}^{m-2}} - \beta \frac{G_n^{m+1}}{F_n^{m+1}} = \frac{p^3 - q^3}{\alpha^2} \frac{F_{n-1}^m F_n^{m-1}}{F_n^{m+1} F_{n-1}^{m-2}} - \frac{\gamma}{\alpha} \frac{F_{n-1}^{m-1} F_n^m}{F_n^{m+1} F_{n-1}^{m-2}}. \quad (61)$$

Assuming $\gamma = 0$, we obtain

$$(p - q)F_{n-1}^{m-1}F_n^{m-1} + \beta G_{n-1}^{m-1}F_n^{m-1} - \beta F_{n-1}^{m-1}G_n^{m-1} = \alpha F_n^m F_{n-1}^{m-2}, \quad (62)$$

$$\begin{aligned} & \frac{p^3 - q^3}{\alpha} F_n^{m+1} F_{n+1}^m - \alpha(p + 2q)F_{n+1}^{m+2} F_n^{m-1} - \alpha\beta G_n^{m-1} F_{n+1}^{m+2} \\ & + \alpha\beta F_n^{m-1} G_{n+1}^{m+2} = 0, \end{aligned} \quad (63)$$

$$\alpha F_n^{m-1} F_{n+1}^{m+3} - \frac{p^3 - q^3}{\alpha^2} F_n^{m+2} F_{n+1}^m = \delta F_n^m F_{n+1}^{m+2}, \quad (64)$$

where δ is a decoupling constant. After changing back to original independent variables ($m - n \rightarrow m$), we have

$$(p - q)F_{n-1}^{m-1}F_n^{m-1} + \beta G_{n-1}^{m-1}F_n^{m-1} - \beta F_{n-1}^{m-1}G_n^{m-1} = \alpha F_n^m F_{n-1}^{m-1}, \quad (65)$$

$$\begin{aligned} & \frac{p^3 - q^3}{\alpha} F_n^{m+1} F_{n+1}^{m-1} - \alpha(p + 2q)F_{n+1}^{m+1} F_n^{m-1} - \alpha\beta G_n^{m-1} F_{n+1}^{m+1} \\ & + \alpha\beta F_n^{m-1} G_{n+1}^{m+1} = 0, \end{aligned} \quad (66)$$

$$\alpha F_n^{m-1} F_{n+1}^{m+2} - \frac{p^3 - q^3}{\alpha^2} F_n^{m+2} F_{n+1}^{m-1} = \delta F_n^m F_{n+1}^{m+1}. \quad (67)$$

Note that Eq.(67) can be derived by eliminating G from Eqs.(65) and (66). This is a discrete analogue of a bilinear form of the potential BSQ equation (3). Thus we have the following theorem.

Theorem 3.3: *Solutions of Eq.(45), i.e. Eqs.(47) and (48), are expressed in the following form:*

$$\begin{aligned} u_n^m &= \frac{G_n^m}{F_n^m}, \quad I_n^m = \frac{F_n^{m+1} F_{n-1}^{m-1}}{F_n^m F_{n-1}^m} = p - q + \frac{G_{n-1}^{m+1}}{F_{n-1}^{m+1}} - \frac{G_n^m}{F_n^m}, \\ V_n^m &= p + 2q + \frac{G_{n-1}^{m-2}}{F_{n-1}^{m-2}} - \frac{G_n^m}{F_n^m}, \end{aligned}$$

where F_n^m and G_n^m satisfy Eqs.(65), (66) and (67). Moreover, F_n^m is given by τ_n^m of the discrete potential BSQ equation (16) in Theorem 3.1 when parameters have relations

$$\frac{\alpha}{\delta} = \frac{a_3(a_1 - a_2)}{a_1(a_3 - a_2)}, \quad \frac{p^3 - q^3}{\alpha^2\delta} = \frac{a_2(a_1 - a_3)}{a_1(a_3 - a_2)}.$$

Furthermore, we have the following theorem.

Theorem 3.4: *The above τ -functions F_n^m and G_n^m are given by*

$$F_n^m = \tau_n^m, \quad G_n^m = \rho_n^m + \left[\left(\frac{p}{\beta} - \frac{1}{a_1} \right) n + \left(\frac{q}{\beta} - \frac{1}{a_2} \right) m \right] \tau_n^m. \quad (68)$$

τ_n^m is defined by Eq.(69) (τ_n^m is also given in Theorem 3.1), ρ_n^m is defined by Eq.(70). Here parameters should be chosen as $a_3 = -a_2/2$,

$$\alpha = \left(\frac{1}{a_1} - \frac{1}{a_2} \right) \beta, \quad \delta = -\frac{3\beta}{a_2}, \quad \beta = \left[\frac{(p^3 - q^3)a_1^3 a_2^3}{(2a_1 + a_2)(a_1 - a_2)^2} \right]^{\frac{1}{3}}.$$

Proof: Let us introduce the following τ -functions using the Freeman-Nimmo notation [9, 24, 32, 33]:

$$\begin{aligned} \tau_n^m(k) &= \begin{vmatrix} \psi_1(n, m, k; s) & \psi_1(n, m, k; s + 1) & \cdots & \psi_1(n, m, k; s + N - 1) \\ \psi_2(n, m, k; s) & \psi_2(n, m, k; s + 1) & \cdots & \psi_2(n, m, k; s + N - 1) \\ \vdots & \vdots & \cdots & \vdots \\ \psi_N(n, m, k; s) & \psi_N(n, m, k; s + 1) & \cdots & \psi_N(n, m, k; s + N - 1) \end{vmatrix} \\ &= |\mathbf{0}, \mathbf{1}, \dots, \mathbf{N} - \mathbf{2}, \mathbf{N} - \mathbf{1}|, \end{aligned} \quad (69)$$

$$\begin{aligned} \rho_n^m(k) &= \begin{vmatrix} \psi_1(n, m, k; s) & \cdots & \psi_1(n, m, k; s + N - 2) & \psi_1(n, m, k; s + N) \\ \psi_2(n, m, k; s) & \cdots & \psi_2(n, m, k; s + N - 2) & \psi_2(n, m, k; s + N) \\ \vdots & \cdots & \vdots & \vdots \\ \psi_N(n, m, k; s) & \cdots & \psi_N(n, m, k; s + N - 2) & \psi_N(n, m, k; s + N) \end{vmatrix} \\ &= |\mathbf{0}, \mathbf{1}, \dots, \mathbf{N} - \mathbf{2}, \mathbf{N}|, \end{aligned} \quad (70)$$

where ψ_1, \dots, ψ_N are a set of linearly independent solutions of the linear system

$$\Delta_n \psi_i(n, m, k; s) = \Delta_m \psi_i(n, m, k; s) = \Delta_k \psi_i(n, m, k; s) = \psi_i(n, m, k; s + 1).$$

Here $\Delta_n, \Delta_m, \Delta_k$ are the backward difference operators

$$\Delta_n f(n) \equiv \frac{f(n) - f(n - 1)}{a_1}, \quad \Delta_m f(m) \equiv \frac{f(m) - f(m - 1)}{a_2}, \quad \Delta_k f(k) \equiv \frac{f(k) - f(k - 1)}{a_3}.$$

Then we notice the following difference formulas [24, 33]

$$\begin{aligned}
 a_1 \tau_{n+1}^m(k) &= |\mathbf{0}, \mathbf{1}, \dots, \mathbf{N} - \mathbf{3}, \mathbf{N} - \mathbf{2}, \mathbf{N} - \mathbf{2}_{n+1}|, \\
 a_2 \tau_n^{m+1}(k) &= |\mathbf{0}, \mathbf{1}, \dots, \mathbf{N} - \mathbf{3}, \mathbf{N} - \mathbf{2}, \mathbf{N} - \mathbf{2}_{m+1}|, \\
 a_3 \tau_n^m(k+1) &= |\mathbf{0}, \mathbf{1}, \dots, \mathbf{N} - \mathbf{3}, \mathbf{N} - \mathbf{2}, \mathbf{N} - \mathbf{2}_{k+1}|, \\
 (a_1 - a_2) \tau_{n+1}^{m+1}(k) &= |\mathbf{0}, \mathbf{1}, \dots, \mathbf{N} - \mathbf{3}, \mathbf{N} - \mathbf{2}_{m+1}, \mathbf{N} - \mathbf{2}_{n+1}|, \\
 (a_2 - a_3) \tau_{n+1}^{m+1}(k+1) &= |\mathbf{0}, \mathbf{1}, \dots, \mathbf{N} - \mathbf{3}, \mathbf{N} - \mathbf{2}_{k+1}, \mathbf{N} - \mathbf{2}_{m+1}|, \\
 a_1 \rho_{n+1}^m(k) - \tau_{n+1}^m(k) &= |\mathbf{0}, \mathbf{1}, \dots, \mathbf{N} - \mathbf{3}, \mathbf{N} - \mathbf{1}, \mathbf{N} - \mathbf{2}_{n+1}|, \\
 a_2 \rho_n^{m+1}(k) - \tau_n^{m+1}(k) &= |\mathbf{0}, \mathbf{1}, \dots, \mathbf{N} - \mathbf{3}, \mathbf{N} - \mathbf{1}, \mathbf{N} - \mathbf{2}_{m+1}|, \\
 a_3 \rho_n^m(k+1) - \tau_n^m(k+1) &= |\mathbf{0}, \mathbf{1}, \dots, \mathbf{N} - \mathbf{3}, \mathbf{N} - \mathbf{1}, \mathbf{N} - \mathbf{2}_{k+1}|.
 \end{aligned}$$

From Plücker relations and these difference formulas, we obtain the following three bilinear equations

$$\begin{aligned}
 &\rho_n^{m+1}(k) \tau_{n+1}^m(k) - \rho_{n+1}^m(k) \tau_n^{m+1}(k) \\
 &= \left(\frac{1}{a_1} - \frac{1}{a_2} \right) (\tau_{n+1}^{m+1}(k) \tau_n^m(k) - \tau_n^{m+1}(k) \tau_{n+1}^m(k)), \tag{71}
 \end{aligned}$$

$$\begin{aligned}
 &\rho_n^m(k+1) \tau_{n+1}^m(k) - \rho_{n+1}^m(k) \tau_n^m(k+1) \\
 &= \left(\frac{1}{a_1} - \frac{1}{a_3} \right) (\tau_{n+1}^m(k+1) \tau_n^m(k) - \tau_n^m(k) \tau_{n+1}^m(k+1)), \tag{72}
 \end{aligned}$$

$$\begin{aligned}
 &a_1(a_2 - a_3) \tau_{n+1}^m(k) \tau_n^{m+1}(k+1) + a_2(a_3 - a_1) \tau_n^{m+1}(k) \tau_{n+1}^m(k+1) \\
 &+ a_3(a_1 - a_2) \tau_n^m(k+1) \tau_{n+1}^{m+1}(k) = 0. \tag{73}
 \end{aligned}$$

Applying the 3-reduction condition $\tau_n^m(k+1) \simeq \tau_n^{m-2}(k)$ and setting $f_n^m = \tau_n^m(k)$, $g_n^m = \rho_n^m(k)$, we obtain

$$g_n^{m+1} f_{n+1}^m - g_{n+1}^m f_n^{m+1} = \left(\frac{1}{a_1} - \frac{1}{a_2} \right) (f_{n+1}^{m+1} f_n^m - f_n^{m+1} f_{n+1}^m), \tag{74}$$

$$g_n^{m-2} f_{n+1}^m - g_{n+1}^m f_n^{m-2} = \left(\frac{1}{a_1} - \frac{1}{a_3} \right) (f_{n+1}^{m-2} f_n^m - f_n^{m-2} f_{n+1}^m), \tag{75}$$

$$a_1(a_2 - a_3) f_{n+1}^m f_n^{m-1} + a_2(a_3 - a_1) f_n^{m+1} f_{n+1}^{m-2} + a_3(a_1 - a_2) f_n^{m-2} f_{n+1}^{m+1} = 0. \tag{76}$$

To compare these bilinear equations with bilinear equations (65)-(67), we set

$$g_n^m = h_n^m + \left(\frac{n}{a_1} + \frac{m}{a_2} \right) f_n^m, \quad a_3 = -\frac{a_2}{2}, \quad C_n^m = H_n^m + \frac{pn + qm}{\beta} F_n^m.$$

Then bilinear equations (74)-(76) and (65)-(67) are rewritten as

$$h_n^{m+1} f_{n+1}^m - h_{n+1}^m f_n^{m+1} = \left(\frac{1}{a_1} - \frac{1}{a_2} \right) f_{n+1}^{m+1} f_n^m, \tag{77}$$

$$h_n^{m-2} f_{n+1}^m - h_{n+1}^m f_n^{m-2} = \left(\frac{1}{a_1} + \frac{2}{a_2} \right) f_{n+1}^{m-2} f_n^m, \tag{78}$$

$$3a_1 f_{n+1}^m f_n^{m-1} - (2a_1 + a_2) f_n^{m+1} f_{n+1}^{m-2} - (a_1 - a_2) f_n^{m-2} f_{n+1}^{m+1} = 0, \tag{79}$$

and

$$\beta H_n^{m+1} F_{n+1}^m - \beta H_{n+1}^m F_n^{m+1} = \alpha F_{n+1}^{m+1} F_n^m, \tag{80}$$

$$\beta H_n^{m-2} F_{n+1}^m - \beta H_{n+1}^m F_n^{m-2} = \frac{p^3 - q^3}{\alpha^2} F_{n+1}^{m-2} F_n^m, \tag{81}$$

$$\delta F_{n+1}^m F_n^{m-1} + \frac{p^3 - q^3}{\alpha^2} F_n^{m+1} F_{n+1}^{m-2} - \alpha F_n^{m-2} F_{n+1}^{m+1} = 0. \tag{82}$$

Assuming $F_n^m = f_n^m, H_n^m = h_n^m$ and comparing coefficients of these equations, we have the relations of parameters

$$\alpha = \left(\frac{1}{a_1} - \frac{1}{a_2} \right) \beta, \quad \delta = -\frac{3\beta}{a_2}, \quad \beta = \left[\frac{(p^3 - q^3)a_1^3 a_2^3}{(2a_1 + a_2)(a_1 - a_2)^2} \right]^{\frac{1}{3}}.$$

□

Note that Hietarinta and Zhang gave the Casorati determinant form of the τ -functions F_n^m and G_n^m [6]. We also note that if we introduce the auxiliary independent variable x by $\partial_x \psi_i(n, m, k; s) = \psi_i(n, m, k; s + 1)$, then ρ_n^m can be expressed as $\rho_n^m = \partial_x \tau_n^m$.

Remark 3:

In the case of $p^3 - q^3 = 0$, Eq.(59) is

$$-\alpha(p + 2q)F_{n+1}^{m+1}F_n^{m-1} - \alpha\beta G_n^{m-1}F_{n+1}^{m+1} + \alpha\beta F_n^{m-1}G_{n+1}^{m+1} = \gamma F_n^m F_{n+1}^m. \tag{83}$$

Equation (60) is written as

$$\alpha \frac{F_n^m F_{n-1}^{m-1}}{F_n^{m-1} F_{n-1}^m} + \frac{\gamma}{\alpha} \frac{F_{n-1}^{m+1} F_n^{m+1}}{F_n^{m+2} F_{n-1}^m} = \alpha \frac{F_{n+1}^{m+2} F_n^{m+1}}{F_{n+1}^{m+1} F_n^{m+2}} + \frac{\gamma}{\alpha} \frac{F_n^m F_{n+1}^m}{F_{n+1}^{m+1} F_n^{m-1}}. \tag{84}$$

After some calculation, we have

$$\frac{\alpha F_{n-1}^{m-1} F_{n+1}^{m+1} - \frac{\gamma}{\alpha} F_{n+1}^m F_{n-1}^m}{F_n^{m-1} F_n^{m+1}} = \frac{\alpha F_{n+1}^{m+2} F_{n-1}^m - \frac{\gamma}{\alpha} F_{n-1}^{m+1} F_{n+1}^{m+1}}{F_n^m F_n^{m+2}}. \tag{85}$$

After decoupling, we obtain

$$\alpha F_{n-1}^{m-1} F_{n+1}^{m+1} - \frac{\gamma}{\alpha} F_{n+1}^m F_{n-1}^m = \delta F_n^{m-1} F_n^{m+1}, \tag{86}$$

which is nothing but Eq.(41). Thus we conclude that this special case gives a discrete potential BSQ equation which has the same τ -function to Eq.(40).

4. Conclusion

We have proposed an alternate form of discrete potential BSQ equation. We have constructed the bilinear equations and multisoliton solutions for the discrete potential BSQ equation. The bilinear equations and multisoliton solutions have been constructed by one of 3-reductions of the Hirota-Miwa equation. Using the discrete potential BSQ equation, we have presented the ultradiscrete potential BSQ equation. We have also studied the lattice potential Boussinesq equation of Nijhoff et al. using the singularity confinement test. Although the lattice potential Boussinesq equation of Nijhoff et al. is in very complicated form, we can find bilinear equations easily by using singularity confinement test. We have investigated the relationships among our alternate discrete potential BSQ equation, the discrete potential BSQ equation of Date et al. and the lattice BSQ equation by Nijhoff et al.

An interesting problem is to present explicit forms of soliton solutions of the ultradiscrete potential BSQ equation. Since 3-reduction condition is very complicated, it is not easy to see which solutions can survive in the ultradiscrete limit. We will address this problem in the near future.

Acknowledgement(s)

The authors would like to thank Professor Masayuki Oikawa for stimulating discussions. The authors would like to acknowledge helpful comments and suggestions of the referees. The authors are grateful for the hospitality of the Isaac Newton Institute for Mathematical Sciences (INI) in Cambridge where this article was completed during the programme Discrete Integrable Systems (DIS). This work was partially supported by the JSPS Grant-in-Aid for Scientific Research No.21656027 and No. 19340039.

References

- [1] F. W. Nijhoff, V. G. Papageorgiou, H. W. Capel and G. R. W. Quispel, *The lattice Gel'fand-Dikii hierarchy*, Inverse Problems, **8** (1992), pp. 597–621.
- [2] F. W. Nijhoff, *Discrete Painlevé Equations and Symmetry Reductions on the Lattice*, in *Discrete Integrable Geometry and Physics* eds. A. Bobenko and R. Seiler (Clarendon Press, Oxford, 1999) pp. 209–234.
- [3] J. Boussinesq, *Theorie des ondes et des remous qui se propagent le long d'un canal rectangulaire horizontal, en communiquant au liquide contenu dans ce canal des vitesses sensiblement parallèles de la surface au fond*, J. Math. Pures. Appli. Ser. 2, **17** (1872), pp. 55–108.
- [4] M. J. Ablowitz and H. Segur, *Solitons and the Inverse Scattering Transform* (SIAM, Philadelphia, 1981).
- [5] A. Tongas and F. Nijhoff, *The Boussinesq integrable system: Compatible lattice and continuum structures*, Glasgow Math. J. **47A** (2005), pp. 205–219.
- [6] J. Hietarinta and D. J. Zhang, *Multisoliton solutions to the lattice Boussinesq equation*, arXiv:0906.3955 (2009).
- [7] B. B. Kadomtsev and V. I. Petviashvili, *On stability of waves in weakly dispersive media*, Sov. Phys. Dokl. **15** (1970), pp. 539–541.
- [8] N. C. Freeman, *Soliton interactions in two dimensions*, Adv. Appl. Mech. **20** (1980), pp. 1–37.
- [9] N. C. Freeman and J. J. C. Nimmo, *Soliton solutions of the KdV and KP equations: the Wronskian technique*, Phys. Lett. A, **95** (1983), pp. 1–3.
- [10] R. Hirota, *The direct method in soliton theory*, (Cambridge University Press, Cambridge, 2004).
- [11] J. Satsuma, *N-Soliton Solution of the Two-Dimensional Korteweg-deVries Equation*, J. Phys. Soc. Jpn **40** (1976), pp. 286–290.

- [12] G. Biondini and Y. Kodama, *On a family of solutions of the Kadomtsev-Petviashvili equation which also satisfy the Toda lattice hierarchy*, J. Phys. A 36 (2003), pp. 10519–10536.
- [13] Y. Kodama, *Young diagrams and N -soliton solutions of the KP equation*, J. Phys. A 37 (2004), pp. 11169–11190.
- [14] G. Biondini and S. Chakravarty, *Soliton solutions of the Kadomtsev-Petviashvili II equation*, J. Math. Phys. 47 (2006), 033514.
- [15] G. Biondini and S. Chakravarty, *Elastic and inelastic line-soliton solutions of the Kadomtsev-Petviashvili II equation*, Math. Comp. Sim. 74 (2007), pp. 237–250.
- [16] G. Biondini, *Line Soliton Interactions of the Kadomtsev-Petviashvili Equation*, Phys. Rev. Lett. 99 (2007), 064103: 1–4.
- [17] S. Chakravarty and Y. Kodama, *Classification of the line-soliton solutions of KP II*, J. Phys. A 41 (2008), pp. 275209 (33pp).
- [18] S. Chakravarty and Y. Kodama, *Soliton solutions of the KP equation and application to shallow water waves*, Stud. App. Math. 123 (2009), pp. 83–151.
- [19] M. Boiti, F. Pempinelli, A. K. Pogrekov and B. Prinari, *Towards an inverse scattering theory for non-decaying potentials of the heat equation*, Inverse Problems 17 (2001), pp. 937–957.
- [20] E. Date, M. Jimbo, M. Kashiwara and T. Miwa, *Transformation groups for soliton equations. Euclidean Lie algebras and reduction of the KP hierarchy*, Publ. RIMS. Kyoto Univ. 18 (1982), pp. 1077–1110.
- [21] M. Jimbo and T. Miwa, *Solitons and infinite dimensional Lie algebras*, Publ. RIMS. Kyoto Univ. 19 (1983), pp. 943–1001.
- [22] R. Hirota, *Reduction of soliton equations in bilinear form*, Physica D 18 (1986), pp. 161–170.
- [23] T. Miwa, *On Hirota's difference equations*, Proc. Japan Acad. A 58 (1982), pp. 9–12.
- [24] Y. Ohta, R. Hirota, S. Tsujimoto and T. Imai, *Casorati and Discrete Gram Type Determinant Representations of Solutions to the Discrete KP Hierarchy*, J. Phys. Soc. Jpn 62 (1993), pp. 1872–1886.
- [25] T. Tokihiro, D. Takahashi, J. Matsukidaira and J. Satsuma, *From soliton equations to integrable cellular automata through a limiting procedure*, Phys. Rev. Lett. 76 (1996) pp. 3247–3250.
- [26] E. Date, M. Jimbo and T. Miwa, *Method for Generating Discrete Soliton Equation. III*, J. Phys. Soc. Jpn 52 (1993) pp. 388–393.
- [27] B. Grammaticos, A. Ramani and V. G. Papageorgiou, *Do integrable mappings have the Painlevé property?*, Phys. Rev. Lett. 67 (1991) pp. 1825–1828.
- [28] A. Ramani, B. Grammaticos and J. Hietarinta, *Discrete versions of the Painlevé equations*, Phys. Rev. Lett. 67 (1991), pp. 1829–1832.
- [29] V. G. Papageorgiou, B. Grammaticos and A. Ramani, *Integrable lattices and convergence acceleration algorithms*, Phys. Lett. A 179 (1993), pp. 111–115.
- [30] A. Ramani, B. Grammaticos and J. Satsuma, *Bilinear discrete Painlevé equations*, J. Phys. A: Math. Gen. 28 (1995), pp. 4655–4665.
- [31] K. Maruno, K. Kajiwara, S. Nakao and M. Oikawa, *Bilinearization of discrete soliton equations and singularity confinement*, Phys. Lett. A 229 (1997), pp. 173–182.
- [32] K. Maruno, K. Kajiwara and M. Oikawa, *Casorati determinant solution for the discrete-time relativistic Toda lattice equation*, Phys. Lett. A 241 (1998), pp. 335–343.
- [33] Y. Ohta, K. Kajiwara, J. Matsukidaira and J. Satsuma, *Casorati determinant solution for the relativistic Toda lattice equation*, J. Math. Phys. 34 (1993), pp. 5190–5204.