Generalized calculus in radiobiology: Physical implications

O. Sotolongo-Grau¹, D. Rodriguez-Perez¹, J. C. Antoranz^{1,2}, and Oscar Sotolongo-Costa²

¹Departamento de Física Matemática y de Fluidos, Universidad Nacional de Educación a Distancia (UNED), Spain

²Cátedra de Sistemas Complejos "Henri Poincaré", Universidad de La Habana, Cuba

Abstract

Non-extensive statistical physics has allowed to generalize mathematical functions such as exponential and logarithms. The same framework is used to generalize sum and product so that the operations allow a more fluid way to work with mathematical expressions emerging from non-additive formulation of statistical physics. In this work we employ the generalization of the exponential, logarithm and product to obtain a formula for the survival fraction corresponding to the application of several radiation doses on a living tissue. Also we provide experimental recommendations to determine the universal characteristics of living tissues in interaction with radiation. These results have a potential application in radiobiology and radiation oncology.

Entropy, Statistical mechanics, Radiation, Non-extensivity

1 Introduction

Biological systems are the paradigm of open non equilibrium systems characterized by long range correlations and by constantly occurring non markovian processes. This makes its description in terms of equilibrium statistical physics a very hard task. Thus a formulation of processes in this kind of systems must be made in a non orthodox statistical framework in which the concept of entropy is expressed in a more general way.

Tsallis's definition of entropy has been successfully used, in the last years, to extend the classical statistical mechanics to several problems involving very complex correlations and properties. Its expression for the entropy (in units of the Boltzmann's constant) [1] is

$$S_q = \frac{1}{q-1} \left(1 - \int_0^\infty p^q(x) dx \right),\tag{1}$$

where p(x)dx is the probability of the magnitude x to adopt values between x and x + dx and q is the "index of non-extensivity". This, together with the constraints that the probability density satisfies the normalization condition and the finiteness of the "q-mean value" $\int_0^\infty p^q(x)xdx = \langle x \rangle_q < \infty$, has been applied in different problems with excellent results even when statistical physics based on Boltzmann-Gibbs (BG) entropy had not succeed. On the other hand, Tsallis entropy reduces to the BG one when q = 1, so that in this sense the former entropy can be interpreted as a generalization of the last one.

Our objective in this work is to employ this new formulation of statistical physics and the consequent generalization of several mathematical functions and operations, provided in the framework of this new formulation, to obtain a simple universal formula for the survival fraction of tissue cells under radiation.

We apply the maximum entropy principle using the Tsallis entropy to obtain an expression for the survival fraction in terms of the radiation dose. Then using a generalization of the definition of logarithm and exponential function and a generalized expression for the product the resultant survival fraction of several radiation sessions is obtained.

2 Materials and methods

To apply the maximum entropy principle in the Tsallis version to the problem of finding the survival factor of a living tissue [2] that receives a radiation, we postulate the existence of some amount of absorbed radiation $\Delta < \infty$ (or its equivalent "limit effect", $\Omega = \alpha_0 \Delta$) after which no cell survives. The application of the maximum entropy principle performs like the usual one but with a few modifications.

The Tsallis entropy becomes:

$$S_q = \frac{1}{q-1} \left(1 - \int_0^\Omega p^q(E) dE \right),\tag{2}$$

where p(E) is the cell death probability, dependent on the "radiation effect" E, which is a function of the applied dose D. The way E depends on D is a controversial matter, being generally accepted a linear relation $E = \alpha D$ and also the "linear quadratic model" $E = \alpha D + \beta D^2$.

The normalization condition is in this case $\int_0^{\Omega} p(E)dE = 1$ and the *q*-mean value becomes $\int_0^{\Omega} p^q(E)EdE = \langle E \rangle_q < \infty$. With this definition, all properties of the tissue and its characteristics of the interaction with radiation become included in $\langle E \rangle_q$ and therefore in Ω . This is the only parameter (besides *q*) entering in our description. It is clear that the determination of $\langle E \rangle_q$ for the different tissues under different conditions of radiation would give the necessary information for the characterization of the survival factor.

To calculate the maximum of (2) under the above conditions the well known

method of Lagrange multipliers [3] is applied, obtaining

$$\Omega = \frac{2-q}{1-q} \left(\frac{\langle E \rangle_q}{2-q}\right)^{\frac{1}{2-q}} \tag{3}$$

 and

$$p(E) = \left(\frac{2-q}{\langle E \rangle_q}\right)^{\frac{1}{2-q}} \left(1 - \frac{1-q}{2-q} \left(\frac{2-q}{\langle E \rangle_q}\right)^{\frac{1}{2-q}} E\right)^{\frac{1}{1-q}}.$$
 (4)

Then the survival factor is

$$F_s(E) = \int_E^{\Omega} p(x) dx = \left(1 - \frac{E}{\Omega}\right)^{\frac{2-q}{1-q}}$$
(5)

with q < 1 for $E < \Omega$ and zero otherwise. It is not hard to see that when $q \to 1$ then $\Omega \to \infty$ and $\langle E \rangle_q \to \langle E \rangle$.

Equation (5) can be written

$$F_s(D) = \begin{cases} \left(1 - \frac{D}{\Delta}\right)^{\gamma} & \forall D < \Delta\\ 0 & \forall D \geqslant \Delta \end{cases}$$
(6)

where we introduced $E = \alpha_0 D$ and $\gamma = \frac{2-q}{1-q}$.

Equation (6) represents the survival fraction in terms of the measurable quantities D (radiation dose) and Δ . All the information about the kind of radiation, radiation rate, etc is contained in the phenomenological term Δ , whereas tissues are characterized by γ . This makes (6) a very general expression with universal characteristics since the phase transition described by (6) is homomorphic with the phase transition of ferromagnets near the Curie point. The exponent γ in this case, as in ferromagnetic phase transitions, determines the universality class. Then γ in our case deals only with the kind of tissue that interacts with radiation [4].

The value of Δ characterizes a critical point for cell survival probability such that for $D < \Delta$ the probabilities of cell survival and death coexist but when $D > \Delta$ a phase transition takes place and no cell survives. Note that we have postulated a linear relation between E and D. With this, the curious property that the tissue effect E is additive while the survival fraction is not multiplicative emerges.

The linear model for the tissue effect [5], however, gives that if the dose is additive the corresponding survival fraction is multiplicative if their mean values are the same. Though this property belongs only to the linear model and not to more general descriptions like the LQ model [5] and others, we think it is worth to find a link between the additivity property of the dose and the probabilistic properties of the cell survival fraction.

To motivate this let us first observe that the Tsallis definition of entropy, though gives a more general expression than the BG one, lacks one of the important properties of this entropy: extensivity. Indeed, let us consider a system composed by two independent subsystems A and B. Extensivity of the entropy means that the entropy of the whole system is the sum of the entropies of the subsystems: $S(A \cup B) = S(A) + S(B)$. But in the Tsallis case $S_q(A \cup B) = S_q(A) + S_q(B) + (q-1)S_q(A)S_q(B)$.

As Tsallis entropy can be defined through the introduction of the q-logarithm:

$$S_q = \int_0^\infty p^q(x) lq\left(p(x)\right) dx \tag{7}$$

where $lq(x) = \frac{x^{1-q}-1}{1-q}$, it is possible to generalize the operation of multiplication introducing the q-product \otimes_q such that $x \otimes_q y = exp_q [lq(x) + lq(y)]$ [6] where

$$exp_q(x) = [1 + (1 - q)x]^{\frac{1}{1 - q}}.$$
 (8)

Taking into account that for the survival fraction we obtain expression (6) its analogy with (8) is evident. Then, let us define the $exp_{\gamma}(x)$ function

$$exp_{\gamma}(x) = \left[1 + \frac{x}{\gamma}\right]^{\gamma} \tag{9}$$

and the $l\gamma(x)$ its inverse function:

$$l\gamma(exp_{\gamma}(x)) = x. \tag{10}$$

Let us introduce the γ -product of two numbers x and y as:

$$x \otimes_{\gamma} y = exp_{\gamma} \left[l\gamma(x) + l\gamma(y) \right] = \left[x^{\frac{1}{\gamma}} + y^{\frac{1}{\gamma}} - 1 \right]^{\gamma}.$$
(11)

Note that definitions (9) and (10) are not essentially different from $e_q(x)$ and lq(x). We are just introducing these definitions to simplify the calculations.

3 Results

Let us now define the "effect potential" as $\chi = -l\gamma(F_s)$. We demand this potential to satisfy the additive property. Then the survival fraction expressed as

$$F_s = exp_{\gamma}(-\chi) = exp_{\gamma}(-\frac{\gamma D}{\Delta}), \qquad (12)$$

where the effect potential depends on $\langle E \rangle_q$, becomes γ -multiplicative. This implies that the statistical independence of the survival fractions is only possible when $\gamma \to \infty \ (q \to 1)$.

Then the survival fraction for the sum of two potentials *i.e.*, the survival fraction of a dose B and a dose A is:

$$F_s(\chi_A) \otimes_{\gamma} F_s(\chi_B) = \left[1 - \frac{\chi_A + \chi_B}{\gamma}\right]^{\gamma}$$
(13)

and for n doses

$$F_s(n\chi) = \left[1 - \sum_{i=1}^n \frac{\chi_i}{\gamma}\right]^\gamma = \left[1 - \sum_{i=1}^n \frac{D_i}{\Delta_i}\right]^\gamma = exp_\gamma \left[-\sum_{i=1}^n \chi_i\right] = \left[\bigotimes_{i=1}^n\right]_\gamma F_s(\chi_i)$$
(14)

where $[\bigotimes_{i=1}^{n}]_{\gamma}$ denotes the iterated application of the γ -product.

4 Discussion

The obtained result permits to calculate directly the result of the application of different radiation doses if γ and Δ are known. Then the importance of experimental determination of γ and Δ becomes evident.

To determine Δ means to measure the minimal dose for which, in a given tissue, no cell survives. This can be made by a progressive application of a given kind of radiation. Once Δ is known in those radiation conditions, a log-log plot of F_s versus $1 - D/\Delta$ must be fitted with the straight line whose slope is γ . Once γ is found the tissue is already characterized. Any other kind of radiations or the same radiation applied in different conditions to the same tissue can be fitted to the same straight line.

This way, the introduction of the rules for generalization of the product as in [6] permits an easy interpretation of the properties of the survival fraction and a formula to calculate it for several radiation doses: the survival fraction of several radiation doses is the γ -product of the separate survival fractions corresponding to each dose.

5 Acknowledgements

The authors wish to thank the Ministerio de Industria, Proyecto CD-TEAM, CENIT. One of us (OSC) wants to acknowledge Prof. Ute Labonté and Nike Fakiner for warm hospitality during his stay in Frankfurt where this work was elaborated.

References

- C Tsallis. Possible generalization of boltzmann-gibbs statistics. Journal of Statistical Physics, 52:479–487, 1988.
- [2] GG Steel. Basic Clinical Radiobiology for Radiation Oncologists. Edward Arnold Publishers, London, 1993.
- [3] A Platino and AR Plastino. Tsallis entropy and jaynes' information theory formalism. Brazilian Journal of Physics, 29:50-60, 1999.

- [4] O Sotolongo-Grau, D Rodriguez-Perez, JA Santos-Miranda, JC Antoranz, and Oscar Sotolongo-Costa. Statistical mechanics formulation of radiobiology. http://arxiv.org/abs/0907.5551v2, 2009.
- [5] M Tubiana. Introduction to Radiobiology. Taylor & Francis, London, 1990.
- [6] C Tsallis. Introduction to nonextensive statistical mechanics. Springer, New York, 2009.