

# Protein micro patterned lattices to probe a fundamental lengthscale involved in cell adhesion.

Hervé Guillou,<sup>1</sup> Benoit Vianay,<sup>1</sup> Jacques Chaussy,<sup>1</sup> and Marc R. Block<sup>2</sup>

<sup>1</sup>*Institut Néel, CNRS UPR2940 et Université  
Joseph Fourier, BP166, 38042 Grenoble, France*

<sup>2</sup>*INSERM U823, Equipe DySAD, Institut Albert Bonniot,  
Site Santé, BP170, 38047 LaTronche, France\**

## Abstract

Cell adhesion, a fundamental process of cell biology is involved in the embryo development and in numerous pathologies especially those related to cancers. We constrained cells to adhere on extracellular matrix proteins patterned in a micro lattices. The actin cytoskeleton is particularly sensitive to this constraint and reproducibly self organizes in simple geometrical patterns. Such highly organized cells are functional and proliferate. We performed statistical analysis of spread cells morphologies and discuss the existence of a fundamental lengthscale associated with active processes required for spreading.

---

\*author's fax: 33+456 38 70 87; Electronic address: herve.guillou@grenoble.cnrs.fr

In vivo, cells proliferate on the 3D structure of extracellular proteins that forms a highly organized scaffold: the extracellular matrix (ECM) is an arrangement of different proteins[1]. Cell adhesion onto the ECM is a fundamental process involved in the embryo development as well as in numerous pathologies especially those related to cancers [2]. In many cell types, the integrin family of transmembrane proteins is responsible for cell interactions with ECM proteins. This specific interaction among other micro environmental physicochemical parameters such as support compliance [3] or shape [4, 5, 6] has been shown to be relevant for the regulation of cell proliferation and to the accomplishment of their functions.

Following [4, 5, 6] we use adequately tailored micro lattice of proteins to obtained very reproducible and simple actin cytoskeleton organization. The microfabrication of the protein array and the simplified actin cytoskeleton morphology that we systematically observe are described next. The result of the statistical analysis of cell shapes are presented in a second part. In particular, we discuss results indicating the existence of a characteristic length. This length is associated with the actin based, active protrusive processes needed to explore the environment and to establish new adhesive contacts required for spreading and migration.

According to amphiphilic properties of globular proteins they tend to accumulate and adsorb at interfaces [7]. Hydrophobic surfaces yield protein films of larger density than hydrophilic surfaces [8]. The principle of fabrication of the protein lattice is based on such non specific protein immobilization on hydrophobic surfaces through a patterned photo-resist mask followed by a lift-off. The photo-resist is insolated using optical UV lithography. Patterns with a critical size of  $1\mu\text{m}$  are thus achievable. In this study, although several proteins where successfully tested (fibronectin, fragment of fibronectin, antibodies, vitronectin, gelatin) best results in terms of spatial resolution were obtained with a fragment of fibronectin made of domains of type III 7 to 10[9]. Fig. 1 shows the adsorption kinetic of the fragment at  $20\mu\text{g/ml}$ . The adsorption isotherm on hydrophobic surface is shown in the inset of fig. 1. Both graphs indicate that the adsorption is limited as expected [10]. After adsorption, the photo-resist mask is lifted-off. The yet photo-resist protected surface must be blocked to prevent further adsorption of protein from the culture medium that would otherwise rapidly blur out the protein pattern. This major problem in protein patterning and has been solved [11, 12] by using a commercially available polymer [13] that spontaneously adsorb on hydrophobic surface in the same manner as proteins. We were able to maintain consistent cell




Figure 1: Kinetic of fibronectin adsorption on hydrophobic glass substrates. Lower inset: adsorption isotherm on hydrophobic surface. Upper inset: durability of protein patterns with cells cultured on fibronectin square islands.

patterns and to keep the repellent properties of the surface for more than 4 days as shown in the images of figure 1: the square adhesive islands kept their geometry despite confluent cell crowding. The lift-off of proteins on hydrophobic surfaces followed by the immobilization of blocking polymer is a very efficient way to produce high resolution adhesive patterns of controlled geometry. Nevertheless it will never reach the resolution achieved in [14] where patterning at the molecular level is realised. However, in contrast with microstamping it produces patterns of very high quality and allows easy alignment with existing structures. We tested the biocompatibility of the protein lattice by studying cell spreadings. Cells were harvested and plated onto the patterns and spread for 4 hours before fixation in paraformaldehyde. The actin cytoskeleton was revealed using rhodamine labelled phalloidin and focal contacts by primary antibody directed toward vinculin. Cover slips were mounted




Figure 2: Typical single cell shapes observed for various protein lattice. Red is F-actin cytoskeleton, green is vinculin and blue is fibronectin. Scale : white bar is  $10\mu m$ . Non patterned substrate (a), lattice pitch  $4\mu m$  (b),  $6\mu m$  (c),  $10\mu m$ (d),  $14\mu m$  (e).

in mowiol for fluorescence microscopy. More details on the methods are given in [15]. For non patterned substrates (fig. 2a) no particular actin cytoskeleton organization can be observed. For small distances between adhesive dots (fig. 2b), the cytoskeleton organisation is similar to the one observed on non patterned substrates. However, as the distance between adhesive spots is increased, the shape of cells is very quickly simplified into elementary geometries (fig. 2c-d). It is remarkable how the complex cytoskeleton organization of figs 2a-b is simplified: stress fibers are exclusively located on the edges of the cell body and linked to well identified focal contacts at their ends.

Each actin cytoskeleton organization was classified according to its area and compacity [16]. Fig. 3 shows the population of cells in each class for a lattice pitch of  $10\mu m$ . Each peak corresponds to a simple geometrical compact shape that can be drawn between adhesive




Figure 3: Distribution of shape obtained for a dot spacing of  $10\mu m$ . The colorcode represent the number of cells observed in a particular configuration. The maxima of occurrence are obtained for the 4 dots square and 6 dots rectangle shown in the inset.

islands. It shows that the most probable organization is the 4 dots square and that cell shapes can be classified.

On a non patterned substrate the ECM layer is dense and almost every membrane protrusion leads to engagement of integrins. Very quickly cells adopt a migrating phenotype in which the cytoskeleton is continuously rearranging. Cell shapes cannot be classified. By contrast, on a protein lattice, a finite distance between neighbouring adhesive spots is introduced and the membrane protrusion are less efficient in finding new ligands. Fig. 4 supports this idea and compares the mean area of spread cells as a function of the lattice pitch with the distribution of the length of filopodia. The mean area decreases very rapidly once the distance is increased. As shown on fig. 4 membrane protrusions are able to explore only the close environment of cells. The probability that a protrusion exceeds a




Figure 4: Shape selection through protrusion length. Left axis, open squares: average mean area of single spread cells on proteins arrays with different pitches. Right axis, open circles: probability to find a filopodia of a given length (averaged on 20 different cells and 400 filopodia).

distance of  $10\mu\text{m}$  is low. Moreover, because of the geometry of the lattice, all protrusions are not successful in finding new ligand, reducing further the probability to engage membrane receptors. The distance between adhesive spots introduce thus a length that can be compared with the average length of protrusion. By increasing the distance between adhesive spots the probability to change the set of adhesive constraint is gradually reduced. The actin cytoskeleton has thus time to relax and to self organize into very simple shapes. The protein lattice probe thus a fundamental length related to the protrusion length.

In this letter we studied single cell spreading onto micro fabricated patterns of controlled geometries. Protrusions are the limiting process resulting in a greatly reduced spread cell area as soon as the pitch is greater than their typical length. As a result stress fibers are exclusively located at the edges of the cell body and attached to two focal adhesion at their

ends.

- 
- [1] B. Alberts, D. Bray, J. Lewis, M. Raff, K. Roberts, and J.D. Watson. *The Molecular Biology of the Cell*. Garland Publishing, New-York, 1994.
- [2] J.P. Thiéry. Cell adhesion in cancer. *C.R. Phys.*, 4:289, 2003.
- [3] D.E. Discher, P. Janmey, and Y.L. Wang. Tissue cells feel and respond to the stiffness of their substrate. *Science*, 310:p1139, 2005.
- [4] S. Huang and D.E. Ingber. The structural and mechanical complexity of cell-growth control. *Nat. Cell Bio.*, 1:pE131–8, 1999.
- [5] C.S. Chen, M. Mrksich, S. Huang, G.M. Whiteside, and D.E. Ingber. Geometric control of cell life and death. *Science*, 276:1425, 1997.
- [6] M. Théry, V. Racine, M. Piel, A. Pépin, A. Dimitrov, Y. Chen, J.B. Sibarita, and M. Bornens. Anisotropy of cell adhesive microenvironment governs cell internal organization and orientation of polarity. *Proc. Natl. Acad. Sci. USA*, 103:p19771, 2006.
- [7] J.N. Israelachvili. *Intermolecular and surface forces*. Academic Press, 1991.
- [8] H. Sorribas, C. Padeste, and Tiefenauer L. Photolithographic generation of protein micropatterns for neuron culture applications. *Biomaterials*, 23:p893, 2002.
- [9] D.J. Leahy, I. Aukhil, and H.P. Erickson. 2.0 a crystal structure of a four-domain segment of human fibronectin encompassing the rgd loop and synergy region. *Cell*, 84:pp155–164, 1996.
- [10] L. Guemouri, J. Ogier, Z. Zekhini, and J.J. Ramsden. The architecture of fibronectin at surfaces. *J. Chem. Phys.*, 113:p8183–8186, 2000.
- [11] M. Zhang, T. Desai, and M. Ferrari. Proteins and cells on peg immobilized silicon surfaces. *Biomaterials*, 19:p953–960, 1998.
- [12] V.A. Liu, W.E. Jastromb, and S. N. Bhatia. Engineering protein and cell adhesivity using peo-terminated triblock polymers. *J. Biomed. Mater. Res.*, 60:p126, 2002.
- [13] Pluronic is a BASF copolymer. BASF, Germany.
- [14] E.A. Cavalcanti-Adam, T. Volberq, A. Micoulet, H. H. Kessler, B. Geiger, and J.P. Spatz. Cell spreading and focal adhesion dynamics are regulated by spacing of integrin ligands. *Biophys. J.*, 92:2964, 2007.

- [15] H. Guillou, A. Depraz-Depland, E. Planus, B. Vianay, J. Chaussy, A. Grichine, C. Albigès-Rizo, and M.R. Block. Lamellipodia nucleation by filopodia depends on integrin occupancy and downstream rac1 signaling. *Exp. Cell Res.*, 314:478, 2008.
- [16] compacity is defined as in the IMAQ library of the Labview programming software from National Instrument, Austin, Texas, USA.