arXiv:0909.1383v3 [q-fin.RM] 15 Dec 2009

Hidden Noise Structure and Random Matrix Models of Stock Correlations

Ivailo I. Dimov,^{1,*} Petter N. Kolm,^{1,†} Lee Maclin,^{1,‡} and Dan Y. C. Shiber^{1,§}

¹Courant Institute of Mathematical Sciences, New York University Corresponding author: Dan Y. C. Shiber

We find a novel correlation structure in the residual noise of stock market returns that is remarkably linked to the composition and stability of the top few significant factors driving the returns, and moreover indicates that the noise band is composed of multiple subbands that do not fully mix. Our findings allow us to construct effective generalized random matrix theory market models [3, 4] that are closely related to correlation and eigenvector clustering [6, 12]. We show how to use these models in a simulation that incorporates heavy tails. Finally, we demonstrate how a subtle purely stationary risk estimation bias can arise in the conventional cleaning prescription [3].

Introduction: Originally started in the context of nuclear physics [1], random matrix theory (RMT) has thereafter found numerous applications in a variety of fields such as number theory, disordered systems, neural networks, and signal processing [1, 2]. Recently the pioneering work of Laloux et al [3], as well as much subsequent research [4, 5], have shown that RMT can also be a valuable tool for analyzing stock market correlations, where noise can account for more than 2/3 of the eigenvalue spectrum, and a typical large portfolio has size comparable to the measurement time frame. Thus, much of the empirical eigenvalues are spurious and represent measurement noise and biases. The remarkable insight provided by Laloux et al was to show that a suitable fit to RMT can clean these spurious contributions, and moreover identify the statistically significant signal, or common market risk factors that drive the individual stock returns. The most prominent such non-idiosyncratic factor is the nearly equal-weight top eigenvector, whose eigenvalue is more than 20 times bigger than the average spectrum. Secondary factors, are long-short portfolios of certain liquidity [3] and industry structure [6, 7], but their contribution is typically an order of magnitude smaller. Most of the rest of the eigenvectors are unstable in time, appear random, and their spectral contribution can be fitted to the Marcenko-Pasteur (MP) distribution [8] derived in the context of Gaussian RMT (GRMT). The noisy eigenvalue correlations [4] also agree with theory [1]. These results have been verified over many stock selections, as well as return frequencies [3, 4, 5].

Despite the apparent success of the theory, subsequent research suggests several empirical aspects that the original RMT cleaning may not account for properly. (1) Tails and their correlations have non-trivial effects, and are known to both broaden the spectrum above the upper noise-band edge, as well sharpen it near the lower edge [4, 9, 10], thus making the fit to the MP distribution problematic. The above redistribution of spectral weight appears in conjunction with an enhancement of the inverse participation ratios around both ends of the noise spectrum, the so-called localization effect [4], unlike GRMT where the participations are flat [1]. (2) In addition to being partially localized, the band itself may be split due to the same separation of correlation scales [11] that is thought to give rise to clustering of stocks between industries [12]. So far this effect has not been observed, however, due to the large amount of mixing that depletes the stability of all the noisy eigenvectors. It is important to empirically distinguish between the single and multiple band cases. (3) Non-stationarity effects are insufficiently understood. They are suggested [3, 4] to be the source of a residual bias in the risk estimates obtained after RMT cleaning. However, in light of the abovementioned considerations, it is not clear that the original cleaning procedures are unbiased to begin with.

In this work, we consider both N = 484.2 minute S&P500 TAO midguote returns between June 20 - Sep 20, 2007, as well as N = 451 daily S&P500 returns between Jan 2001-Dec 2007 [13]. (1) We reveal a novel correlation structure of the residuals that is linked to the structure and stability of the top few empirical factors. Mainly, we find that the inverse participations of the localized edge-eigenmodes of the band are dominated by the outlier stocks in the composition of the top few factors, thus indicating that most of the noise there is due to these stocks. The upper edge fluctuations are mainly due to weakly correlated stocks with the smallest relative weight in the market portfolio while lower edge fluctuations are due to strongly correlated stocks identified as the outliers in the secondary factors. The groups in the lower edge belong to major industrial sectors [4, 12], while the upper edge contains a large diversified portfolio of medium to small liquidity stocks. Moreover, because we find these groups to be disjoint, we conclude that as long as the top few factors are stable and distinct, the noise band is composed of multiple subbands that do not fully mix. (2) We pinpoint the effective positivedefinite cleaned matrices that exhibit the multi-residual and factor structure above to be the hierarchical RMT models [11], which are closely related to coarse-grained "real space" models of market clustering [12], and fundamentally arise out of correlation scale separation. (3) We use these effective models to perform a one-factor stochastic volatility [9] simulation in order to take into account the effect of tails and their correlations. (4) We show how conventional cleaning can give rise to a subtle purely stationary risk-estimation bias.

<u>Empirical Results</u>: Given our set of N stock series $S_i(t)$, t = 1, ..., T, from their log returns $x_{t,i} = \log(S_{t,i}/S_{t-1,i})$ we calculate the empirical correlation $C_E = (\langle x_i x_j \rangle - \langle x_i \rangle \langle x_j \rangle)/\sigma_i \sigma_j$ where $\sigma_i = \sqrt{\langle x_i^2 \rangle - \langle x_i \rangle^2}$. According to



Figure 1: *Main*: A fit of the MP distribution to 2 min data for N = 484, stocks in the S&P 500 with T = 3N yields $Q_{eff} = 2.25$, and $\sigma_{eff} = 0.67$. The top three eigenvalues, $\lambda_1 = 152.9$, $\lambda_2 = 8.2$, $\lambda_3 = 7.6$, $\lambda_4 = 5.3$, $\lambda_5 = 5.2$ were omitted from the plot due to their scale. *Inset*: The top three eigenvectors, e_{ki} , k = 1, 2, 3, with their entries *i* sorted by decreasing liquidity (from left to right). Note the significant outliers in each e_k as emphasized by the horizontal lines.

RMT [1, 2, 3], if C_E were obtained from a purely random signal of bounded variance σ whose marginal tails are not too heavy [9], then in the limit $N \to \infty$ with $Q \equiv N/T$ fixed, the correlations will self-average and will have an asymptotically deterministic eigenvalue spectrum given by the MP distribution [8]:

$$p_{MP}(\lambda) = \frac{Q}{2\pi\sigma^2} \frac{\sqrt{(\lambda_+ - \lambda)(\lambda - \lambda_-)}}{\lambda}.$$
 (1)

In the above, the eigenvalues λ are restricted to lie within the hard-edge spectral band, $\lambda \in [\lambda_-, \lambda_+]$, with $\lambda_{\pm} = \lambda_{\pm}(Q, \sigma) = \sigma^2(1 \pm \sqrt{1/Q})^2$. One can interpret (1) as the the finite T/N noise-induced broadening and bias away from the underlying trivial spectrum $p_{clean}(\lambda) = \delta(\lambda - 1)$ that $p_{MP}(\lambda)$ reduces to in the limit $Q \to \infty$.

Of course, stock market correlations are not purely random, so a fit for σ and Q is necessary in (3) if one wants to identify the trully residual part of the spectrum [3, 4]. In Fig. 1 we show such a fit to the noisy region of the 2 min data that yields $\sigma_{eff} = 0.67$, $Q_{eff} = 2.25$. Note that much of the spectrum lies outside the MP band. In the inset of Fig 1 we plot the composition, of the top three eigenvectors $\{e_k\}$, k = 1, 2, 3 sorted by decreasing liquidity. Unlike the prediction of GRMT where e_k should be a mean-zero unit Gaussian, there are clear deviations from such behavior in all three eigenvectors, as emphasized in the inset. In fact, e_1 has nonzero mean, $\langle e_{1i} \rangle_i = 0.044 \simeq N^{-1/2}$, representing a longonly market portfolio, while e_2 and e_3 represent long-short portfolios. All three e_k can be interpreted as significant common factors. Furthermore, as is clear from the plot, the outliers in the factor composition have certain liquidity structure. In the case of e_2 and e_3 in of Fig 1, these outliers can be identified with major sectors such as financials, oil, and utilities, whose correlations are relatively stable in time [6, 7, 12].

Despite the appearance of factors, one expects the random residual spectral contribution to be well fitted to RMT. However, there are important issues with the fit to p_{MP} that one needs to address. Heavy tails in the multivariate distribution of $x_{t,i}$ have non-trivial effects. They are known to broaden the spectral weight above the upper edge, as well as sharpen it near the lower edge [4, 9, 10], both features readily noticeable in Fig. 1 as well as in daily data [3]. Moreover, such tails tend to induce outliers in the composition of the principal components near the band edge [9] causing deviations from the standard Gaussian distribution of the composition expected by GRMT and inducing localization. Indeed, just as in daily data [4] we see in the 2 min returns that the eigenvectors are localized at both ends of the noise spectrum by computing the *inverse participation ratio* [4], $I_k = \sum_{i=1}^{N} [e_{ki}]^4$ for each eigenvector e_k . Intuitively, the participation $P_k \equiv 1/I_k$ scales as the number of non-trivial entries in a normalized e_k : $P_k = N$ for equal weight vectors, while $P_k = 1$ for a single non-trivial weight. As evident in Fig 2 (a), the participation is strongly localized near the band edges indicating that the eigenvectors there are dominated by outliers.

A nice trick that avoids estimating the effects of tails is to clean the noisy eigenvalues $\Lambda_E = diag(\lambda_1, \ldots, \lambda_K; \{\lambda_{noise}\})$ of $C_E = S_E \Lambda_E S'_E$ with a flat band with scale proportional to σ_{eff} while working in the original eigenvector basis $S_E = (e_1, \ldots, e_N)$ [3], thus obtaining a "filtered" matrix [4]. In fact, unless the empirically measured tail fluctuations significantly break the rotational invariance implied by cleaning with a flat band, σ_{eff} can be thought of as the overall scale of the residual noise that one can tune even without fitting to the Gaussian formula. The feasibility of this "filtering" procedure can also be justified with the resulting significant improvement of the portfolio risk estimates that one obtains with cleaning [3, 6].

We will now show, however, that because of a novel structure of the eigenvectors, cleaning with a flat band σ_{eff} is inconsistent with their symmetry. Suppose we look at the following group of stocks, $\{G_1, G_k^{\pm}, G^{\perp}\}, k = 2, 3$ selected so that G_k^{\pm} contains the top/bottom outliers $|e_{ki}| > \tilde{e}_k$ of e_k above/below a certain threshold \tilde{e}_k (see Fig 1 inset), G_1 are the outliers of smallest absolute weight in e_1 , and G^{\perp} are all the other stocks. We find that for reasonable threshold values, $\tilde{e}_k \simeq 1.5\sigma_{e_k}, G_1$ is disjoint from $G_{23} \equiv \bigcup G_k^{\pm}$. Moreover, the relative contribution of each group G to the inverse participation,

$$R_k^{(G)} = \frac{\sum_{i \in G} [e_{ki}]^4}{I_k} \equiv \frac{I_k^{(G)}}{I_k},$$
 (2)

is inhomogeneously distributed across the noise band as shown in Fig 2 (b), so that G_1 contributes mostly to the upper



Figure 2: *Top*: The participation P_k of the eigenvectors e_k , $k \ge 2$ for the data in Fig 1 exhibits localization. Flat horizontal line represents $P_k^{GMRT} = N/3$. [1]. *Bottom*: Relative inverse participation $R_k^{(G)}$ for the groups G_1 (red), G_{23} (green) and G^{\perp} (blue) defined in the text. Flat lines represent $R_{k,RMT}^{(G)}$.

edge while G_{23} contributes mostly to the lower edge. Because this behavior is inconsistent with homogeneous cleaning, we interpret it as an indication that *the noise band is composed* of multiple subbands that do not mix. In particular, the three groups above form a partition of all the stocks, where the number of assets in each subgroup, or the group degeneracies, for the data in Fig 1 are $\{D_1, D_{23}, D^{\perp}\} = \{46, 61, 377\}$.

<u>Constructing Coarse-Grained Effective Models</u>: The partition above is reminiscent of partitions previously obtained by "real space" hierarchical clustering of stocks into industries [12], which can be thought of as arising from a coarsegrained separation of correlation scales, also known to give rise to multiple subbands in the spectrum [11]. Moreover, it has been observed [6] that clustering of significant *eigenvector components* results in the same industries as those obtained from the real space procedure, and arises out of a mean-field duality relation between the two approaches [17]. Therefore, we interpret the multiple subband structure above as arising from a particular type of underlying separation of correlation scales apparent at the time scales of measurement of C_E . We have observed the above multiple band structure in both 2 min and daily data.

Let us gain insight into the details of this correlation structure for the case of the data in Fig 1. By clustering analysis [12] we find that G_{23} separates further, $G_{23} = \{G_{23}^+, G_{23}^-\}$, into two nearly-equally large groups of distinctly higher/lower mean average correlation with degeneracies $\{D_{23}^+, D_{23}^-\} =$ $\{29, 32\}$ respectively. For this sample, we find that G_{23}^+ contains Electric Utilities, as well as Oil & Gas Drilling & Exploration stocks, while G_{23}^- contains Oil & Gas Exploration, as well as some Financial stocks. From the average correlation between all four groups, we thus construct the following "minimal" coarse-grained *effective model*:



Figure 3: The same quantities as in Fig 2, except all the data was *simulated* from the effective model (3) by taking into account tail effects in the multivariate distribution via the one-factor stochastic volatility model [9] with tail index $\nu = 3$.

$$C_{eff}^{N \times N} = \begin{pmatrix} C_{23+}^{D_{23}^+ \times D_{23}^+} & 0.36 & 0.35 & 0.22 \\ 0.36 & C_{23-}^{D_{23}^- \times D_{23}^-} & 0.33 & 0.20 \\ 0.35 & 0.33 & C_{\perp}^{D^+ \times D^\perp} & 0.20 \\ 0.22 & 0.20 & 0.20 & C_1^{D_1 \times D_1} \end{pmatrix}$$
(3)

which can be readily checked to be positive definite. The entries of each $D \times D$ diagonal block above are $C^{D \times D} = (1 - \rho_G)I_D + \rho_G^{D \times D}$, with $\{\rho_{23}^+, \rho_{23}^-, \rho_\perp, \rho_1\} = \{0.59, 0.48, 0.32, 0.13\}$ respectively being the average correlation of each of the four groups G and $\rho_G^{D \times D}$ is a block whose entries are ρ_G . One can check that (3) also gives rise to 4 distinct factors and 4 distinct subbands.

Note that unlike the strongly correlated ones in G_{23} , the stocks in G_1 are typically not easily detectable with conventional hierarchical clustering approaches [12], although they are distinctly visible if one looks at the top factor (see Fig 1 inset). Indeed, being weakly correlated between each other as well as with the rest of the market, these stocks will not appear in localized real-space clusters but instead will group with other stocks in later stages of the hierarchy. At the same time, both the degeneracy D_1 and overal risk contribution of G_1 are comparable to those of the localized sectors, as also directly suggested by Fig 2 (b). To properly account for the separation of correlation scales in markets, one must also include the contribution of the weakly correlated stocks.

<u>Simulating with tails</u>: A check of the validity of the effective model (3) is ultimately provided if one can reproduce the empirical spectrum and participations through simulation. To do so, one must properly take into account heavy tailed behavior of actual returns. It is known that such tails can be induced by heteroskedasticities of the underlying stock volatilities [14], although the details of the correlations of such volatility dynamics are not well understood. We thus use the simplest multivariate conditional Gaussian model with onefactor variance-gamma volatilities, which is known to produce a Student-t type of series [9] for the joint returns. We use an inverse-gamma tail index of $\nu \simeq 3$. The resulting spectrum [9, 10] agrees well with the empirical one. Moreover, comparing Figs 2 and 3, we see that the inverse participations are also in good agreement.

A Subtle Stationary Bias: The discussion so far suggests that even for stationary data, RMT cleaning could produce biased risk estimates. Let us demonstrate this for the simplest case of multivariate Gaussian returns simulated with the effective model (3). Without loss of generality we normalize the returns to mean zero unit variance. Using the notation in [4], the predicted risk of a portfolio $w = (w_1, \ldots, w_N)$ is $\Omega_p^2 = w \cdot C_{clean} \cdot w^T$. The portfolios we look at are equal-weight average representatives of different subbands K, $w_K = \sum_{k \in K} e_k$, where $\{e_k\}$ are the eigenvectors of the effective model (3). Moreover, instead of a "budget constraint" [4], we impose a "risk constraint" by normalizing w_K to unit norm. We then compute at every forecasting period the relative difference between realized and predicted risk, $\delta r = (\Omega_r^2 - \Omega_n^2)/\Omega_n^2$. For the subbands $\{K_1, K^{\perp}, K_{23}\}$ corresponding to the groups of stocks that enter in Fig 2 (b), we find respective biases $\delta r_i = \{26 \pm 4\%, 2 \pm 4\%, -17 \pm 2\%\}.$ Note that although δr_1 and δr_{23} are significant, they are of opposite sign. Indeed, we have checked that all three contributions nearly cancel when one looks at the relative realized versus predicted risk of the entire noise band, $\delta r_{all} = 2 \pm 3\%$. Finally, we also observe significant biases δr_1 and δr_{23} in the actual data. However, in this case, there are subtleties in disentangling the effects of multiple bands, tails, and nonstationarity. We postpone discussing these effects, as well as multi-residual generalizations of the RMT cleaning procedure to later work [17].

<u>Summary:</u> In conclusion, we have found strong evidence that instead of homogeneous, the stock market correlation residuals are composed of multiple subbands that do not fully mix. This structure is manifested through an asymmetry in the relative inverse participations of the eigenvectors within the noise band, which is inconsistent with purely symmetric cleaning that doesn't distinguish between different parts of

the noise spectrum. The multi-residual picture above natrually emerges from market models with multiple correlation scales, that we have identified and simulated. As a direct consequence, the scale separation within the noise band also produces inhomogeneities in the effective residual risk that in turn induce purely stationary biases of the original RMT cleaning.

We would like to thank Marco Avellaneda and Jim Gatheral for their insightful comments and discussion.

- * Electronic address: iid201@cims.nyu.edu
- [†] Electronic address: kolm@cims.nyu.edu
- [‡] Electronic address: lee.maclin@gmail.com
- [§] Electronic address: dcs255@cims.nyu.edu
- [1] M. L. Mehta, *Random Matrices* (Academic Press, Boston, 1991).
- [2] A. M. Sengupta and P. P. Mitra, Phys. Rev. E 60, 3389 (1999).
- [3] L. Laloux et al., Phys. Rev. Lett. 83, 1467 (1999).
- [4] V. Plerou *et al.*, Phys. Rev. Lett. 83, 1471 (1999); V. Plerou *et al.*, Phys. Rev. E 65, 066126 (2002).
- [5] A. Utsugi *et al.*, Phys. Rev. E **70**, 026110 (2004); V. Kulkarni and N. Deo, Europhys. J. **B**, **60**, 101 (2007); R. Pan and S. Sinha, Phys. Rev. E **76**, 046116 (2007); J. Shen and B. Zheng, Europhys. Lett. **86**, 28005 (2009).
- [6] P. Gopikrishnan et al., arXiv:cond-mat/0011145 (2000).
- [7] M. Avellaneda and J-H Lee, *Working Paper Series*. Available at SSRN: http://ssrn.com/abstract=1153505.
- [8] V. Marčenko and L. Pastur, Math. USSR-Sbornik 1, 457 (1967).
- [9] G. Biroli *et al.*, Europhys. Lett. **78** (No. 1), (2007); G. Biroli *et al.*, Acta Phys. Pol. B **38** (13), 4009 (2007).
- [10] Z. Burda *et al.*, Phys. Rev. E **71**, 026111 (2005); Burda *et al.*, Phys. Rev. E **74**, 041129 (2006).
- [11] F. Lillo and R. N. Mantegna, Phys. Rev. E 72, 016219 (2005).
- [12] R. N. Mantegna, Europhys. J. B. 11 (No. 1), 193 (1999).
- [13] We have performed all the necessary tests [14] to ensure that the marginal 2 min midquote returns have no residual autocorrelations.
- [14] J.-P. Bouchaud and M. Potters, *Theory of Financial Risk and Derivative Pricing* (Cambridge Univ. Press, Cambridge, 2003).
- [15] After fitting for Q_{eff} and σ_{eff} one typically normalizes so as to preserve the conservation of the trace of correlations [3, 4].
- [16] M. Potters et al., Acta Phys. Pol. B 36 (9), 2767 (2005).
- [17] I. Dimov, D. Shiber, P. Kolm, L. Maclin in preparation.