

# Average sensitivity and noise sensitivity of polynomial threshold functions

Ilias Diakonikolas      Prasad Raghavendra      Rocco A. Servedio  
Li-Yang Tan

February 6, 2020

## Abstract

We give the first non-trivial upper bounds on the average sensitivity and noise sensitivity of degree- $d$  polynomial threshold functions (PTFs). These bounds hold both for PTFs over the Boolean hypercube  $\{-1, 1\}^n$  and for multilinear PTFs over  $\mathbb{R}^n$  under the standard  $n$ -dimensional Gaussian distribution  $\mathcal{N}(0, I_n)$ . Our bound on the Boolean average sensitivity of PTFs represents progress towards the resolution of a conjecture of Gotsman and Linial [GL94], which states that the symmetric function slicing the middle  $d$  layers of the Boolean hypercube has the highest average sensitivity of all degree- $d$  PTFs. Via the  $L_1$  polynomial regression algorithm of Kalai et al. [KKMS08], our bounds on Gaussian and Boolean noise sensitivity yield polynomial-time agnostic learning algorithms for the broad class of constant-degree PTFs under these input distributions.

Our bounds on both average and noise sensitivity of multilinear PTFs in the Gaussian setting are obtained using fairly simple arguments; the main ingredients are tail bounds and anti-concentration bounds on low-degree polynomials in Gaussian random variables [Jan97, CW01]. To obtain our bound on the Boolean average sensitivity of PTFs, we generalize the “critical-index” machinery of [Ser07] (which in that work applies to halfspaces, i.e. degree-1 PTFs) to general PTFs. Together with the “invariance principle” of [MOO05], this lets us extend our techniques from the Gaussian setting to the Boolean setting. Our bound on Boolean noise sensitivity is achieved via a simple reduction from upper bounds on average sensitivity of Boolean PTFs to corresponding bounds on noise sensitivity.

# 1 Introduction

A degree- $d$  polynomial threshold function (PTF) over a domain  $X \subseteq \mathbb{R}^n$  is a Boolean-valued function  $f : X \rightarrow \{-1, +1\}$ ,

$$f(x) = \text{sign}(p(x_1, \dots, x_n))$$

where  $p : X \rightarrow \mathbb{R}$  is a degree- $d$  polynomial with real coefficients. The function  $f$  is said to be a *multilinear* PTF if the polynomial  $p$  is multilinear. When  $d = 1$  polynomial threshold functions are simply linear threshold functions (also known as halfspaces or LTFs), which play an important role in complexity theory, learning theory, and other fields such as voting theory. Low-degree PTFs (where  $d$  is greater than 1 but is not too large) are a natural generalization of LTFs which are also of significant interest in these fields.

Over more than twenty years much research effort in the study of Boolean functions has been devoted to different notions of the “sensitivity” of a Boolean function to small perturbations of its input, see e.g. [KKL88, BT96, BK97, Fri98, BKS99, Shi00, MO03, MOO05, OSSS05, OS07] and many other works. In this work we focus on two natural and well-studied measures of this sensitivity, the “average sensitivity” and the “noise sensitivity.” As our main results, we give the first non-trivial upper bounds on average sensitivity and noise sensitivity of low-degree PTFs. These bounds have several applications in learning theory and complexity theory as we describe later in this introduction.

We now define the notions of average and noise sensitivity in the setting of Boolean functions  $f : \{-1, 1\}^n \rightarrow \{-1, 1\}$ . (Our paper also deals with average sensitivity and noise sensitivity of functions  $f : \mathbb{R}^n \rightarrow \{-1, 1\}$  under the Gaussian distribution, but the precise definitions are more involved than in the Boolean case so we defer them until later.)

## 1.1 Average Sensitivity and Noise Sensitivity

The *sensitivity* of a Boolean function  $f : \{-1, 1\}^n \rightarrow \{-1, 1\}$  on an input  $x \in \{-1, 1\}^n$ , denoted  $s_f(x)$ , is the number of Hamming neighbors  $y \in \{-1, 1\}^n$  of  $x$  (i.e. strings which differ from  $x$  in precisely one coordinate) for which  $f(x) \neq f(y)$ . The *average sensitivity* of  $f$ , denoted  $\text{AS}(f)$ , is simply  $\mathbf{E}[s_f(x)]$  (where the expectation is with respect to the uniform distribution over  $\{-1, 1\}^n$ ). An alternate definition of average sensitivity can be given in terms of the influence of individual coordinates on  $f$ . For a Boolean function  $f : \{-1, 1\}^n \rightarrow \{-1, 1\}$  and a coordinate index  $i \in [n]$ , the *influence of coordinate  $i$  on  $f$*  is the probability that flipping the  $i$ -th bit of a uniform random input  $x \in \{-1, 1\}^n$  causes the value of  $f$  to change, i.e.  $\text{Inf}_i(f) = \mathbf{Pr}[f(x) \neq f(x^{\oplus i})]$  (where the probability is with respect to the uniform distribution over  $\{-1, 1\}^n$ ). The sum of all  $n$  coordinate influences,  $\sum_{i=1}^n \text{Inf}_i(f)$ , is called the *total influence* of  $f$ ; it is easily seen to equal  $\text{AS}(f)$ . Bounds on average sensitivity have been of use in the structural analysis of Boolean functions (see e.g. [KKL88, Fri98, Shi00]) and in developing computationally efficient learning algorithms (see e.g. [BT96, OS07]).

The average sensitivity is a measure of how  $f$  changes when a single coordinate is perturbed. In contrast, the noise sensitivity of  $f$  measures how  $f$  changes when a random collection

of coordinates are all perturbed simultaneously. More precisely, given a noise parameter  $0 \leq \epsilon \leq 1$  and a Boolean function  $f : \{-1, 1\}^n \rightarrow \{-1, 1\}$ , the *noise sensitivity of  $f$  at noise rate  $\epsilon$*  is defined to be

$$\text{NS}_\epsilon(f) = \Pr_{x,y}[f(x) \neq f(y)]$$

where  $x$  is uniform from  $\{-1, 1\}^n$  and  $y$  is obtained from  $x$  by flipping each bit independently with probability  $\epsilon$ . Noise sensitivity has been studied in a range of contexts including Boolean function analysis, percolation theory, and computational learning theory [BKS99, KOS04, MO03, SS, KOS08].

## 1.2 Main Results: Upper Bounds on Average Sensitivity and Noise Sensitivity

### 1.2.1 Boolean PTFs

In 1994 Gotsman and Linial [GL94] conjectured that the symmetric function slicing the middle  $d$  layers of the Boolean hypercube has the highest average sensitivity among all degree- $d$  PTFs. Since this function has average sensitivity  $\Theta(d\sqrt{n})$  for every  $1 \leq d \leq \sqrt{n}$ , this conjecture implies (and is nearly equivalent to) the conjecture that every degree- $d$  PTF  $f$  over  $\{-1, 1\}^n$  has  $\text{AS}(f) \leq d\sqrt{n}$ .

Our first main result is an upper bound on average sensitivity which makes progress toward this conjecture:

**Theorem 1.1** *For any degree- $d$  PTF  $f$  over  $\{-1, 1\}^n$ , we have*

$$\text{AS}(f) \leq 2^{O(d)} \cdot \log n \cdot n^{1-1/(4d+2)}.$$

Using a completely different set of techniques, we also prove a different bound which improves on [Theorem 1.1](#) for  $d \leq 4$ :

**Theorem 1.2** *For any degree- $d$  PTF  $f$  over  $\{-1, 1\}^n$ , we have*

$$\text{AS}(f) \leq 2n^{1-1/2^d}.$$

We give a simple reduction which translates any upper bound on average sensitivity for degree- $d$  PTFs over Boolean variables into a corresponding upper bound on noise sensitivity. Combining this reduction with [Theorems 1.1](#) and [1.2](#), we establish:

**Theorem 1.3** *For any degree- $d$  PTF  $f$  over  $\{-1, 1\}^n$  and any  $0 \leq \epsilon \leq 1$ , we have*

$$\begin{aligned} \text{NS}_\epsilon(f) &\leq 2^{O(d)} \cdot \epsilon^{1/(4d+2)} \log(1/\epsilon) \\ \text{NS}_\epsilon(f) &\leq O(\epsilon^{1/2^d}). \end{aligned}$$

### 1.2.2 Gaussian PTFs

Looking beyond the Boolean hypercube, there are well-studied notions of average sensitivity and noise sensitivity for Boolean-valued functions over  $\mathbb{R}^n$ , where we view  $\mathbb{R}^n$  as endowed with the standard multivariate Gaussian distribution  $\mathcal{N}(0, I_n)$  [Bog98, MOO05]. Let  $\text{GAS}(f)$  denote the Gaussian average sensitivity of a function  $f : \mathbb{R}^n \rightarrow \{-1, 1\}$ , and let  $\text{GNS}_\epsilon(f)$  denote the Gaussian noise sensitivity at noise rate  $\epsilon$ . (See Section 2 for precise definitions of these quantities; here we just note that these are natural analogues of their uniform-distribution Boolean hypercube counterparts defined above.) We prove an upper bound on Gaussian average sensitivity of low-degree multilinear PTFs:

**Theorem 1.4** *For any degree- $d$  multilinear PTF  $f$  over  $\mathbb{R}^n$ , we have*

$$\text{GAS}(f) \leq O(d \cdot \log n \cdot n^{1-1/2d})$$

We also prove an upper bound on the Gaussian noise sensitivity of degree- $d$  multilinear PTFs:

**Theorem 1.5** *For any degree- $d$  multilinear PTF  $f$  over  $\mathbb{R}^n$  and any  $0 \leq \epsilon \leq 1$ , we have*

$$\text{GNS}_\epsilon(f) \leq O(d \cdot \log(1/\epsilon) \cdot \epsilon^{1/2d}).$$

## 1.3 Application: agnostically learning constant-degree PTFs in polynomial time

Our bounds on noise sensitivity, together with machinery developed in [KOS04, KKMS08, KOS08], let us obtain the first efficient agnostic learning algorithms for low-degree polynomial threshold functions. In this section we state our new learning results; details are given in Section 8.

We begin by briefly reviewing the fixed-distribution agnostic learning framework that has been studied in several recent works, see e.g. [KKMS08, KOS08, BOW08, GKK08, KMV08, SSS09]. Let  $\mathcal{D}_X$  be a (fixed, known) distribution over an example space  $X$  such as the uniform distribution over  $\{-1, 1\}^n$  or the standard multivariate Gaussian distribution  $\mathcal{N}(0, I_n)$  over  $\mathbb{R}^n$ . Let  $\mathcal{C}$  denote a class of Boolean functions, such as the class of all degree- $d$  PTFs. An algorithm  $A$  is said to be an *agnostic learning algorithm for  $\mathcal{C}$  under distribution  $\mathcal{D}_X$*  if it has the following property: Let  $\mathcal{D}$  be any distribution over  $X \times \{-1, 1\}$  such that the marginal of  $\mathcal{D}$  over  $X$  is  $\mathcal{D}_X$ . Then if  $A$  is run on a sample of labeled examples drawn independently from  $\mathcal{D}$ , with high probability  $A$  outputs a hypothesis  $h : X \rightarrow \{-1, 1\}$  such that  $\Pr_{(x,y) \sim \mathcal{D}}[h(x) \neq y] \leq \text{opt} + \epsilon$ , where  $\text{opt} = \min_{f \in \mathcal{C}} \Pr_{(x,y) \sim \mathcal{D}}[f(x) \neq y]$ . In words,  $A$ 's hypothesis is nearly as accurate as the best hypothesis in  $\mathcal{C}$ .

Kalai et al. [KKMS08] gave an  $L_1$  polynomial regression algorithm and showed that it can be used for agnostic learning. More precisely, they showed that for a class  $\mathcal{C}$  of functions and

a distribution  $\mathcal{D}$ , if every function in  $\mathcal{C}$  has a low-degree polynomial approximator (in the  $L_2$  norm) under the marginal distribution  $\mathcal{D}_X$ , then the  $L_1$  polynomial regression algorithm is an efficient agnostic learning algorithm for  $\mathcal{C}$  under  $\mathcal{D}_X$ . They used this  $L_1$  polynomial regression algorithm together with the existence of low-degree polynomial approximators for halfspaces (under the uniform distribution on  $\{-1, 1\}^n$  and the standard Gaussian distribution  $\mathcal{N}(0, I_n)$  on  $\mathbb{R}^n$ ) to obtain  $n^{O(1/\epsilon^4)}$ -time agnostic learning algorithms for halfspaces under these distributions.

Using ingredients from [KOS04], one can easily convert upper bounds on Boolean noise sensitivity (such as Theorem 1.3) into results asserting the existence of low-degree  $L_2$ -norm polynomial approximators under the uniform distribution on  $\{-1, 1\}^n$ . We thus obtain the following agnostic learning result (a more detailed proof is given in Section 8):

**Theorem 1.6** *The class of degree- $d$  PTFs is agnostically learnable under the uniform distribution on  $\{-1, 1\}^n$  in time  $n^{(2^d/\epsilon)^{O(d)}}$ . For  $d \leq 4$ , this bound can be improved to  $n^{O(1/\epsilon^{2^{d+1}})}$ .*

Similarly, using ingredients from [KOS08], one can easily convert upper bounds on Gaussian noise sensitivity (such as Theorem 1.5) into results asserting the existence of low-degree  $L_2$ -norm polynomial approximators under  $\mathcal{N}(0, I_n)$ . We thus obtain

**Theorem 1.7** *The class of degree- $d$  multilinear PTFs is agnostically learnable under  $\mathcal{N}(0, I_n)$  in time  $n^{(d/\epsilon)^{O(d)}}$ .*

For  $\epsilon$  constant, these results are the first polynomial-time agnostic learning algorithms for constant-degree PTFs.

## 1.4 Other applications

The results and approaches of this paper have found other recent applications beyond the agnostic learning results presented above; we describe two of these below.

Gopalan and Servedio [GS09] have combined the average sensitivity bound given by Theorem 1.1 with techniques from [LMN93] to give the first sub-exponential time algorithms for learning  $AC^0$  circuits augmented with a small (non-constant) number of arbitrary threshold gates, i.e. gates that compute arbitrary LTFs which may have weights of any magnitude. (Previous work using different techniques [JKS02] could only handle  $AC^0$  circuits augmented with majority gates.)

In other recent work Diakonikolas et al. [DSTW09] have refined the approach used to prove Theorem 1.1 to establish a “regularity lemma” for low-degree polynomial threshold functions. Roughly speaking, this lemma says that any degree- $d$  PTF can be decomposed into a constant number of subfunctions, almost all of which are “regular” degree- $d$  PTFs. [DSTW09] apply this regularity lemma to extend the positive results on the existence of low-weight approximators for LTFs, proved in [Ser07], to low-degree PTFs.

**Related work.** Simultaneously and independently of this work, Harsha et al. [HKM09] have obtained very similar results on average sensitivity, noise sensitivity, and agnostic learning of low-degree PTFs using techniques very similar to ours.

## 1.5 Techniques

In this section we give a high-level overview of how Theorems 1.1, 1.4 and 1.5 are proved. (As mentioned earlier, Theorem 1.2 is proved using completely different techniques; see Section 6.) The arguments are simpler for the Gaussian setting so we begin with these.

### 1.5.1 The Gaussian case

We sketch the argument for the Gaussian noise sensitivity bound, Theorem 1.5; the Gaussian average sensitivity bound, Theorem 1.4, follows along similar lines.

Let  $f = \text{sign}(p)$  where  $p : \mathbb{R}^n \rightarrow \mathbb{R}$  is a degree- $d$  polynomial. The Gaussian noise sensitivity  $\text{GNS}_\epsilon(f)$  of  $f$  is equal to  $\Pr_{x,y}[f(x) \neq f(y)]$  where  $x$  is distributed according to  $\mathcal{N}(0, I_n)$  and  $y$  is an “ $\epsilon$ -perturbed” version of  $x$  (see Section 2 for the precise definition). Intuitively, the event  $f(x) \neq f(y)$  can only take place if either

- $x$  lies close to the boundary of  $p$ , i.e.  $|p(x)|$  is “small”, or
- $|p(x) - p(y)|$  is “large”.

We use an anti-concentration result for polynomials in Gaussian random variables, due to Carbery and Wright [CW01], to show that  $|p(x)|$  is “small” only with low probability. For the second bullet, it turns out that  $p(x) - p(y)$  can be expressed as a low-degree polynomial in independent Gaussian random variables, and thus we can apply tail bounds for this setting [Jan97] to show that  $|p(x) - p(y)|$  is “large” only with low probability. We can thus argue that  $\Pr_{x,y}[f(x) \neq f(y)]$  is low, and bound the Gaussian noise sensitivity of  $f$ .

### 1.5.2 The Boolean case

Our upper bound on Boolean average sensitivity uses ideas from the previous subsection but requires significant additional ingredients.

An important notion in the Boolean case is that of a “regular” PTF; this is a PTF  $f = \text{sign}(p)$  where every variable *in the polynomial*  $p$  has low influence. (See Section 2 for a definition of the influence of a variable on a real-valued function; note that the definition from Section 1.1 applies only for Boolean-valued functions.) If  $f$  is a regular PTF, then the “invariance principle” of [MOO05] tells us that  $p(x)$  (where  $x$  is uniform from  $\{-1, 1\}^n$ ) behaves much like  $p(\mathcal{G})$  (where  $\mathcal{G}$  is drawn from  $\mathcal{N}(0, I_n)$ ), and essentially the arguments from the Gaussian case can be used.

It remains to handle the case where  $f$  is not a regular PTF, i.e. some variable has high influence in  $p$ . To accomplish this, we generalize the notion of the “critical-index” of a halfspace (see [Ser07, DGJ<sup>+</sup>09]) to apply to PTFs. We show that a carefully chosen random restriction (one which fixes only the variables up to the critical index – very roughly speaking, only the highest-influence variables – and leaves the other ones free) has non-negligible probability of causing  $f$  to collapse down to a regular PTF. This lets us give a recursive bound on average sensitivity which ends up being not much worse than the bound that can be obtained for the regular case; see [Section 5.1](#) for a detailed explanation of the recursive argument.

## 1.6 Organization

Formal definitions of average sensitivity and noise sensitivity (especially in the Gaussian case), and tail bounds and anticoncentration results for low degree polynomials are presented in [Section 2](#). In [Section 3](#), we show an upper bound on the Gaussian average sensitivity of PTFs ([Theorem 1.4](#)). Upper bounds on Gaussian noise sensitivity ([Theorem 1.5](#)) are obtained in the section that follows ([Section 4](#)).

The main result of the paper – a bound on the boolean average sensitivity ([Theorem 1.1](#)) – is proved in [Section 5](#). In [Section 6](#), an alternate bound for boolean average sensitivity that is better for degrees  $d \leq 4$  ([Theorem 1.2](#)) is shown. This is followed by a reduction from boolean average sensitivity bounds to corresponding noise sensitivity bounds ([Theorem 7.1](#)) in [Section 7](#). Finally, we sketch the applications of these upper bounds to agnostic learning PTFs in [Section 8](#).

# 2 Definitions and Background

## 2.1 Basic Definitions

In this subsection we record the basic notation and definitions used throughout the paper. For  $n \in \mathbb{N}$ , we denote by  $[n]$  the set  $\{1, 2, \dots, n\}$ . We write  $\mathcal{N}$  to denote the standard univariate Gaussian distribution  $\mathcal{N}(0, 1)$ .

For a degree- $d$  polynomial  $p : X \rightarrow \mathbb{R}$  we denote by  $\|p\|_2$  its  $l_2$  norm,  $\|p\|_2 = \mathbf{E}_x[p(x)^2]^{1/2}$ , where the intended distribution over  $x \in \mathbb{R}^n$  (which will always be either uniform over  $\{-1, 1\}^n$ , or the  $\mathcal{N}^n$  distribution) will always be clear from context. We note that for multilinear  $p$  the two notions are always equal (see e.g. Proposition 3.5 of [MOO05]).

We now proceed to define the notion of influence for real-valued functions in a product probability space. Throughout this paper we consider either the uniform distribution on the hypercube  $\{\pm 1\}^n$  or the standard  $n$ -dimensional Gaussian distribution in  $\mathbb{R}^n$ . However, for the sake of generality, we adopt this more general setting.

Let  $(\Omega_1, \mu_1), \dots, (\Omega_n, \mu_n)$  be probability spaces and let  $(\Omega = \otimes_{i=1}^n \Omega_i, \mu = \otimes_{i=1}^n \mu_i)$  denote the corresponding product space. Let  $f : \Omega \rightarrow \mathbb{R}$  be any  $\mu$ -measurable function on  $\Omega$ . The influence of the  $i$ th coordinate on  $f$  [MOO05] is

$$\text{Inf}_i^\mu(f) \stackrel{\text{def}}{=} \mathbf{E}_\mu[\text{Var}_{\mu_i}[f]]$$

and the total influence of  $f$  is  $\text{Inf}^\mu(f) \stackrel{\text{def}}{=} \sum_{i=1}^n \text{Inf}_i^\mu(f)$ .

For a function  $f : \{-1, 1\}^n \rightarrow \mathbb{R}$  over the Boolean hypercube endowed with the uniform distribution, the influence of variable  $i$  on  $f$  can be expressed in terms of the Fourier coefficients of  $f$  as,

$$\text{Inf}_i(f) = \sum_{S \ni i} \widehat{f}(S)^2,$$

and as mentioned in the introduction it is easily seen that  $\text{AS}(f) = \text{Inf}(f)$  for boolean-valued functions  $f : \{-1, 1\}^n \rightarrow \{-1, 1\}$ .

In this paper we are concerned with variable influences for functions defined over  $\{-1, 1\}^n$  under the uniform distribution, and over  $\mathbb{R}^n$  under  $\mathcal{N}(0, I_n)$ ; we shall adopt the convention that  $\text{Inf}_i(f)$  denotes the former and  $\text{GI}_i(f)$  the latter. We also denote by  $\text{GAS}(f) = \sum_{i \in [n]} \text{GI}_i(f)$  the Gaussian average sensitivity.

Note that for a function  $f : \mathbb{R}^n \rightarrow \{-1, 1\}$ , the Gaussian influence  $\text{GI}_i(f)$  can be equivalently written as:  $\text{GI}_i(f) = 2 \Pr_{x, x^i}[f(x) \neq f(x^i)]$ , where  $x \sim \mathcal{N}^n$  and  $x^i$  is obtained by replacing the  $i^{\text{th}}$  coordinate of  $x$  by an independent random sample from  $\mathcal{N}$ .

We proceed to define the notion of noise sensitivity for boolean-valued functions in  $(\mathbb{R}^n, \mathcal{N}^n)$ . For the boolean setting, the notion has been defined already in the introduction. (We remark that “noise sensitivity” can be defined in a much more general setting and also for real-valued functions; but such generalizations are not needed here.)

**Definition 1 (Gaussian Noise Sensitivity)** *Given  $f : \mathbb{R}^n \rightarrow \{-1, 1\}$ , the “Gaussian noise sensitivity of  $f$  at noise rate  $\epsilon \in [0, 1]$ ” is*

$$\text{GNS}_\epsilon(f) \stackrel{\text{def}}{=} \Pr_{x, z}[f(x) \neq f(y)];$$

where  $x \sim \mathcal{N}^n$  and  $y \stackrel{\text{def}}{=} (1 - \epsilon)x + \sqrt{2\epsilon - \epsilon^2}z$  for an independent Gaussian noise vector  $z \sim \mathcal{N}^n$ .

## 2.2 Probabilistic Facts

In this subsection, we record the basic probabilistic tools we use in our proofs.

The first result is a standard concentration bound for low-degree polynomials over independent random signs or standard Gaussians. It can be proved (in both cases) using Markov’s inequality and hypercontractivity.

**Theorem 2.1 (“degree- $d$ ” Chernoff bound)** *Let  $p(x)$  be a degree- $d$  multilinear polynomial. Let  $x$  be drawn either from the uniform distribution in  $\{-1, 1\}^n$  or from  $\mathcal{N}^n$ . For any  $t > e^d$ , we have*

$$\Pr_x[|p(x)| \geq t\|p\|_2] \leq \exp(-\Omega(t^{2/d})).$$

The second fact is a powerful anti-concentration bound for low-degree polynomials over Gaussian random variables. (We note that this result does not hold in the boolean setting.)

**Theorem 2.2 ([CW01])** *Let  $p : \mathbb{R}^n \rightarrow \mathbb{R}$  be a degree- $d$  polynomial. Then for all  $\epsilon > 0$ , we have*

$$\Pr[|p(\mathcal{G})| \leq \epsilon\|p\|_2] \leq O(d\epsilon^{1/d})$$

where  $\mathcal{G} = (\mathcal{G}_1, \dots, \mathcal{G}_n)$  are independent standard Gaussians.

We also make essential use of a (weak) anti-concentration property of low-degree polynomials over the hypercube.

**Theorem 2.3 ([DFKO06, AH09])** *Let  $p : \{-1, 1\}^n \rightarrow \mathbb{R}$  be a degree- $d$  multilinear polynomial with  $\text{Var}[p] \equiv \sum_{0 < |S| \leq d} \widehat{p}(S)^2 = 1$  and  $\mathbf{E}[p] = \widehat{p}(\emptyset) = 0$ . Then we have*

$$\Pr[p(x) > 1/2^{O(d)}] > 1/2^{O(d)} \quad \text{and hence} \quad \Pr[|p(x)| \geq 1/2^{O(d)}] > 1/2^{O(d)}.$$

The following is a restatement of the invariance principle, specifically Theorem 3.19 under hypothesis **H4** in [MOO05].

**Theorem 2.4 ([MOO05])** *Let  $p(x) = \sum_{|S| \leq d} \widehat{p}(S)x_S$  be a degree- $d$  multilinear polynomial with  $\sum_{0 < |S| \leq d} \widehat{p}(S)^2 = 1$ . Suppose each variable  $i \in [n]$  has low influence  $\text{Inf}_i(p) \leq \tau$ , i.e.  $\sum_{S \ni i} \widehat{p}(S)^2 \leq \tau$ . Let  $X_1, \dots, X_n$  be independent uniformly random  $\pm 1$  bits. Then,*

$$\sup_{t \in \mathbb{R}} |\Pr[p(X_1, \dots, X_n) \leq t] - \Pr[p(\mathcal{G}_1, \dots, \mathcal{G}_n) \leq t]| \leq O(d\tau^{1/(4d+1)})$$

where  $\mathcal{G}_1, \dots, \mathcal{G}_n$  are independent standard Gaussians.

### 3 Gaussian Average Sensitivity

In this section, we will show an upper bound on the Gaussian average sensitivity of degree- $d$  multilinear PTFs (Theorem 1.4).

The following lemma relating the influence of a variable on  $f$  to its influence on the multilinear polynomial  $p$ , is central to the argument.

**Lemma 3.1** *Let  $p : \mathbb{R}^n \rightarrow \mathbb{R}$  be a degree- $d$  multilinear polynomial over Gaussian inputs with  $\|p\|_2 = 1$  and let  $f = \text{sign}(p)$ . Then for each  $i \in [n]$ ,*

$$\text{GI}_i(f) \leq O(d \cdot \text{GI}_i(p)^{1/(2d)} \cdot \log(1/\text{GI}_i(p))).$$

**Proof:**[of Lemma 3.1]

For notational convenience let us fix  $i = 1$ . Let  $\tau = \text{GI}_1(p)$ . We may assume that  $\text{GI}_1(p) = \tau < 1/4$  since otherwise the claimed bound holds trivially.

Let us express  $p(x)$  as a linear form in  $x_1$ , i.e.

$$p(x) = p(x_1, \dots, x_n) = a(x_2, \dots, x_n) + x_1 b(x_2, \dots, x_n).$$

By assumption we have  $\sum_S \widehat{b}(S)^2 = \tau$  and thus  $\sum_S \widehat{a}(S)^2 = 1 - \tau \geq 1/2$ .

The high-level idea of the proof is to show that with high probability, a draw of  $(g_2, \dots, g_n) \sim \mathcal{N}^{n-1}$  causes  $\text{sign}(p(g_1, g_2, \dots, g_n))$  (viewed as a function of  $g_1$  only) to be quite unbalanced. Let  $t = C^{d/2} \tau^{1/2} \log^{d/2}(1/\tau)$  and  $\gamma = d \cdot \tau^{1/(2d)} \log(1/\tau)$  where  $C$  is an absolute constant that will be defined later in [Claim 3.3](#). For these values of  $t$  and  $\gamma$ , the proof strategy is as follows:

- We use the “small ball probability” bound ([Theorem 2.2](#)) to argue that with high probability,  $a(g_2, \dots, g_n)$  is not too small: more precisely,  $\Pr_{(g_2, \dots, g_n) \sim \mathcal{N}^{n-1}}[|a(g_2, \dots, g_n)| \leq t\sqrt{2 \ln(1/\gamma)}] \leq O(\gamma)$  (see [Claim 3.2](#)).
- We use the concentration bound ([Theorem 2.1](#)) to argue that with high probability,  $b(g_2, \dots, g_n)$  is not too large: more precisely,  $\Pr_{(g_2, \dots, g_n) \sim \mathcal{N}^{n-1}}[|b(g_2, \dots, g_n)| \geq t] \leq O(\gamma)$  (see [Claim 3.3](#)).
- We use elementary properties of the  $N(0, 1)$  distribution to argue that if  $|a| \geq t\sqrt{2 \ln(1/\gamma)}$  and  $|b| \leq t$ , then the function  $\text{sign}(a + g_1 b)$  (a function of one  $N(0, 1)$  random variable  $g_1$ ) is  $O(\gamma)$ -close to the constant function  $\text{sign}(a)$  (see [Claim 3.4](#)).
- Thus we know that with probability at least  $1 - O(\gamma)$  over the choice of  $g_2, \dots, g_n$ , we have  $\text{Var}_{g_1}[\text{sign}(p(g_1, \dots, g_n))] \leq O(\gamma(1 - \gamma)) \leq O(\gamma)$ . For the remaining (at most)  $O(\gamma)$  fraction of outcomes for  $g_2, \dots, g_n$  we always have  $\text{Var}_{g_1}[\text{sign}(p(g_1, \dots, g_n))] \leq 1$ , so overall we get  $\text{GI}_1(\text{sign}(p)) \leq O(\gamma)$ .

Thus, to complete the proof of the lemma, it suffices to prove the three aforementioned claims.

**Claim 3.2** *Let  $a(g_2, \dots, g_n)$  be a degree- $d$  multilinear polynomial with  $\|a\|_2 \geq 1/2$ . Then  $\Pr_{g_2, \dots, g_n}[|a(g_2, \dots, g_n)| \leq t\sqrt{2 \ln(1/\gamma)}] \leq O(\gamma)$ .*

**Proof:** The claim follows directly from [Theorem 2.2](#) for the above choice of parameters. By applying [Theorem 2.2](#) to polynomial  $a(x)$  we get

$$\Pr_{g_2, \dots, g_n} \left[ |a(g_2, \dots, g_n)| \leq \frac{t\sqrt{2\ln(1/\gamma)}}{\|a\|} \cdot \|a\| \right] \leq O \left( d \left( \frac{t\sqrt{2\ln(1/\gamma)}}{\|a\|} \right)^{1/d} \right).$$

Recall that  $\|a\| \geq \frac{1}{2}$ ,  $t = C^{d/2}\tau^{1/2} \log^{d/2}(1/\tau)$  and  $\gamma = d \cdot \tau^{1/2d} \log(1/\tau)$ . Substituting these in the previous inequality yields the upper bound  $O(\gamma)$  on the right hand side. ■

**Claim 3.3** *Let  $b(g_2, \dots, g_n)$  be a degree- $(d-1)$  multilinear polynomial with  $\|b\|_2^2 = \tau$ . There exists an absolute constant  $C$  such that if  $t = C^{d/2}\tau^{1/2} \log^{d/2}(1/\tau)$  then  $\Pr_{g_2, \dots, g_n} [|b(g_2, \dots, g_n)| > t] \leq \gamma$ .*

**Proof:** Applying [Theorem 2.1](#) on the polynomial  $b$  we get

$$\Pr_{g_2, \dots, g_n} \left[ |b(g_2, \dots, g_n)| > \frac{t}{\|b\|} \cdot \|b\| \right] \leq 2 \exp \left( -\Omega \left( \left( \frac{t}{\|b\|} \right)^{2/d} \right) \right).$$

Substituting the value of  $\|b\| = \sqrt{\tau}$  and  $t = C^{d/2}\tau^{1/2} \log^{d/2}(1/\tau)$ , the right hand side is upper bounded by  $2 \exp(-\Omega(C \log(1/\tau)))$ . For a sufficiently large absolute constant  $C$ , this probability is less than  $\exp(-4 \log(1/\tau)) = \tau^4 \leq \gamma$ . ■

**Claim 3.4** *Fix any  $t > 0$  and  $\gamma < \frac{1}{3}$ . If  $|a| \geq t\sqrt{2\ln(1/\gamma)}$  and  $|b| \leq t$ , then  $\Pr_{g_1 \sim N(0,1)} [\text{sign}(a+g_1b) \neq \text{sign}(a)] < O(\gamma)$ .*

**Proof:** We may assume wlog that  $a, b > 0$ . We have  $\text{sign}(a+g_1b) \neq \text{sign}(a)$  only if  $g_1 < -a/b$ ; since  $-a/b \leq -\sqrt{2\ln(1/\gamma)}$ , the standard bound

$$\Pr_{g_1 \sim N(0,1)} [g_1 < c] \leq \frac{1}{\sqrt{2\pi}|c|} \exp(-c^2/2) \quad \text{for } c < 0$$

(see e.g. p.6 of Durrett's *Probability: Theory and Examples*) gives the claim. ■

The proof of [Lemma 3.1](#) is now complete. ■

We can now complete the proof of [Theorem 1.4](#).

**Proof:**[Proof of [Theorem 1.4](#)] Let us denote by  $\tau_i = \text{GI}_i(p)$  for  $i \in [n]$ . Note that since  $p$  is of degree  $d$ , we have

$$\sum_{i \in [n]} \tau_i = \sum_{i \in [n]} \sum_{S \ni i} \hat{p}(S)^2 = \sum_{|S| \leq d} |S| \cdot \hat{p}(S)^2 \leq d. \quad (1)$$

Let  $h_d(x) = dx^{1/2d} \log(1/x)$ . By [Lemma 3.1](#) the average sensitivity of  $f$  can be bounded as

$$\text{GAS}(f) = \sum_{i \in [n]} \text{GI}_i(f) \leq O\left(\sum_{i \in [n]} h_d(\tau_i)\right).$$

The function  $h_d(x)$  is monotone increasing and concave in  $[0, e^{-2d}]$ . In this light, we split the summation into terms greater than  $e^{-2d}$  and the rest. Let  $S = \{i | \tau_i \geq e^{-2d}\}$  and  $T = [n] \setminus S$ . From (1), we have  $|S| \leq de^{2d}$ . Observe that for  $n < (27d)^{2d}$ , [Theorem 1.4](#) holds trivially since  $\text{GAS}(f) \leq n \leq 27dn^{1-1/2d} \leq 27dn^{1-1/2d} \log n$ . Hence we may assume  $n \geq (27d)^{2d}$ , and consequently  $|T|$  is at least  $n/2$ . Using concavity and monotonicity of  $h_d$ , we can write

$$\sum_{i \in T} h_d(\tau_i) \leq |T| \cdot h_d\left(\left(\sum_{i \in T} \tau_i\right)/|T|\right) \leq nh_d\left(\frac{2d}{n}\right) \leq O(dn^{1-1/2d} \log n).$$

Therefore, the average sensitivity of  $f$  is bounded by

$$\begin{aligned} \text{GAS}(f) &= \sum_{i \in S} \text{GI}_i(f) + \sum_{i \in T} \text{GI}_i(f) \\ &\leq |S| + O\left(\sum_{i \in T} h_d(\tau_i)\right) \leq de^{2d} + O(dn^{1-1/2d} \log n). \end{aligned}$$

For all  $d \geq 1$  we have

$$de^{2d} < e^{3d} < (3d^{1/3})^{3d} < (27d)^d \leq n^{1/2}, \quad \text{since } n \geq (27d)^{2d}.$$

Consequently we have  $\text{GAS}(f) \leq n^{1/2} + O(dn^{1-1/2d} \log n) = O(dn^{1-1/2d} \log n)$ , and the proof is complete. ■

## 4 Gaussian Noise Sensitivity

In this section we prove an upper bound on the noise sensitivity of degree- $d$  multilinear PTFs.

**Proof:**[Proof of [Theorem 1.5](#)] Let  $f = \text{sign}(p)$ , where  $p$  is multilinear with  $\|p\|_2 = 1$ . Recall that  $\text{GNS}_\epsilon(f) = \Pr_{x,z}[f(x) \neq f(y)]$  where  $x \sim \mathcal{N}^n$ ,  $z \sim \mathcal{N}^n$ ;  $x$  and  $z$  are independent; and  $y = \alpha x + \beta z$ , with  $\alpha \stackrel{\text{def}}{=} 1 - \epsilon$  and  $\beta = \sqrt{2\epsilon - \epsilon^2}$ .

We can write

$$p(x) = \sum_{|S| \leq d} \hat{p}(S) x_S$$

and

$$p(y) = p(\alpha x + \beta z) = \sum_{|S| \leq d} \hat{p}(S) (\alpha x + \beta z)_S.$$

Let us define the function

$$q(x, z) = p(x) - p(y).$$

Note that  $q$  is a degree- $d$  multilinear polynomial over  $2n$  variables (with no constant term).

Fix a real number  $t^* > 0$ . It is easy to see that  $f(x) \neq f(y)$  only if at least one of the following two events hold:

$$(\text{Event } \mathcal{E}_1) \quad |p(x)| \leq t^* \quad \text{OR} \quad (\text{Event } \mathcal{E}_2) \quad |q(x, z)| \geq t^*$$

We will upper bound the probability of these two events for a carefully chosen  $t^*$ . We will bound the probability of the event  $\mathcal{E}_1$  using Carbery-Wright ([Theorem 2.2](#)), the probability of event  $\mathcal{E}_2$  using the tail bound for degree- $d$  polynomials ([Theorem 2.1](#)) and then apply a union bound.

The choice of  $t^*$  will be dictated by the [Theorem 2.1](#). More precisely, to apply [Theorem 2.1](#), a bound on  $\|q\|_2$  is needed. To this end, we show the following claim:

**Claim 4.1** *We have  $\|q\|_2 = O(d \cdot \sqrt{\epsilon})$ .*

**Proof:** Note that  $q$  is multilinear, hence its  $l_2$  norm is just the square root of the sum of its squared coefficients. Let us expand the term  $(\alpha x + \beta z)_S$  that appears in  $p(y)$  with coefficient  $\widehat{p}(S)$ . For notational convenience we assume that  $S = [k]$ , for some  $k \leq d$ . We have that

$$(\alpha x + \beta z)_S = \prod_{i=1}^k (\alpha x_i + \beta z_i) = \sum_{T \subseteq [k]} \alpha^{|T|} x_T \beta^{k-|T|} z_{\bar{T}} = \alpha^k x_{[k]} + \sum_{T \subsetneq [k]} \alpha^{|T|} x_T \beta^{k-|T|} z_{\bar{T}}$$

Now, we observe that in the polynomial  $q$  the monomial  $x_S = x_{[k]}$  will have coefficient  $\widehat{p}(S) \cdot (1 - \alpha^k) = O(d \cdot \epsilon) \cdot \widehat{p}(S)$ . The monomials  $x_T z_{\bar{T}}$  are all distinct, even across different  $S$ 's (this is important since it means there are no cancelations across different  $S$ 's). For a given  $S = [k]$ , we have that the sum of the squares of the coefficients of all these monomials is exactly

$$\widehat{p}(S)^2 \cdot \sum_{i=0}^{k-1} \binom{k}{i} \alpha^{2i} \beta^{2(k-i)} = \widehat{p}(S)^2 \cdot [(\alpha^2 + \beta^2)^k - \alpha^{2k}] = \widehat{p}(S)^2 \cdot [1 - (1 - \epsilon)^{2k}] = \widehat{p}(S)^2 \cdot O(d \cdot \epsilon).$$

By summing up across all  $S$ 's and taking square root we get the claim. ■

Fix  $t^* = \Theta(d\sqrt{\epsilon} \log^d(1/\epsilon))$ . By [Theorem 2.2](#), we have:

$$\Pr_{x \sim \mathcal{N}^n} [|p(x)| \leq t^*] = O(d \cdot (t^*)^{1/d}) = O(d \cdot \epsilon^{1/(2d)} \cdot \log(1/\epsilon)).$$

By [Theorem 2.1](#) and [Claim 4.1](#), we get

$$\Pr_{x, z \sim \mathcal{N}^{2n}} \left[ |q(x, z)| \geq \frac{t^*}{\|q\|_2} \cdot \|q\|_2 \right] \leq 2 \exp \left( -\Omega \left( \left( \frac{t^*}{\|q\|_2} \right)^{2/d} \right) \right) \leq \epsilon.$$

Hence, by a union bound the noise sensitivity is  $O(d \cdot \epsilon^{1/(2d)} \cdot \log(1/\epsilon))$ . This completes the proof of [Theorem 1.5](#). ■

## 5 Boolean Average Sensitivity

Let  $\text{AS}(n, d)$  denote the maximum possible average sensitivity of any degree- $d$  PTF over  $n$  boolean variables. In this section we prove the claimed bound in [Theorem 1.1](#):

$$\text{AS}(n, d) \leq 2^{O(d)} \cdot \log n \cdot n^{1-1/(4d+2)} \quad (2)$$

For  $d = 1$  (linear threshold functions) it is well known that  $\text{AS}(n, 1) = 2^{-n} \binom{n}{n/2} = \Theta(\sqrt{n})$ . Also, notice that the RHS of (2) has value  $\omega(n)$  for  $d = \omega(\sqrt{\log n})$ , yielding a trivial bound of  $\text{AS}(n, d) \leq n$ . Therefore throughout this section we shall assume  $d$  satisfies  $2 \leq d \leq O(\sqrt{\log n})$ .

### 5.1 Overview of proof

The high-level approach to proving [Theorem 1.1](#) is a combination of a case analysis and a recursive bound.

For certain types of PTFs (“ $\tau$ -regular” PTFs; see [Section 5.2](#) for a precise definition) we argue directly that the average sensitivity is small, using arguments similar to the Gaussian case together with the invariance principle. In particular, we show:

**Claim 5.1** *Suppose  $f = \text{sign}(p)$  is a  $\tau$ -regular degree- $d$  PTF where  $\tau \stackrel{\text{def}}{=} n^{-(4d+1)/(4d+2)}$ . Then,*

$$\text{AS}(f) \leq O(d \cdot n^{1-1/(4d+2)})$$

[Claim 5.1](#) follows directly from [Lemma 5.8](#), which we prove in [Section 5.4](#).

For PTFs that are not  $\tau$ -regular, we show that there is a not-too-large value of  $k$  (at most  $K \stackrel{\text{def}}{=} 2d \log n / \tau$ ), and a collection of  $k$  variables whose influence in  $p$  are largest, such that the following holds: if we consider all  $2^k$  subfunctions of  $f$  obtained by fixing the variables in all possible ways, a “large” (at least  $1/2^{O(d)}$ ) fraction of the restricted functions have low average sensitivity. More precisely, we show:

**Claim 5.2** *Let  $K \stackrel{\text{def}}{=} 2d \log n / \tau$  where  $\tau \stackrel{\text{def}}{=} n^{-(4d+1)/(4d+2)}$ . Suppose  $f = \text{sign}(p)$  is a degree- $d$  PTF that is not  $\tau$ -regular. Then for some  $1 \leq k \leq K$ , there is a set of  $k$  variables with the following property: for at least a  $1/2^{O(d)}$  fraction of all  $2^k$  assignments  $\rho$  to those  $k$  variables, we have*

$$\text{AS}(f_\rho) \leq O(d \cdot (\log n)^{1/4} \cdot n^{1-1/(4d+2)})$$

The proof of [Claim 5.2](#) is given in [Section 5.7](#). We do this by generalizing the “critical index” case analysis from [[Ser07](#)]. We define a notion of the  $\tau$ -critical index of a degree- $d$  polynomial;

a  $\tau$ -regular polynomial  $p$  is one for which the  $\tau$ -critical index is 0. If the  $\tau$ -critical index of  $p$  is some value  $k \leq 2d \log n / \tau$ , we restrict the  $k$  largest-influence variables (see Section 5.5). If the  $\tau$ -critical index is larger than  $2d \log n / \tau$ , we restrict the  $k = 2d \log n / \tau$  largest-influence variables in  $p$  (see Section 5.6).

### 5.1.1 Proof of main result (Theorem 1.1) assuming Claim 5.1 and Claim 5.2

Given these two claims it is not difficult to obtain the final result. In Claim 5.2, we note that the  $k$  restricted variables may each contribute at most 1 to the average sensitivity of  $f$  (recall that average sensitivity is equal to the sum of influences of each variable), and that the total influence of the remaining variables on  $f$  is equal to the expected average sensitivity of  $f_\rho$ , where the expectation is taken over all  $2^k$  restrictions  $\rho$ . Since each function  $f_\rho$  is itself a degree- $d$  PTF over at most  $n$  variables, we have the following recursive constraint on  $\text{AS}(n, d)$ :

$$\text{AS}(n, d) \leq \max\left\{ O(d \cdot n^{1-1/(4d+2)}), \max_{1 \leq k \leq K, 1/2^{O(d)} \leq \alpha \leq 1} \{k + \alpha \cdot O(d \cdot (\log n)^{1/4} \cdot n^{1-1/(4d+2)}) + (1 - \alpha)\text{AS}(n, d)\} \right\}.$$

It is easy to see that the maximum possible value of  $\text{AS}(n, d)$  subject to the above constraint is at most the maximum possible value of  $\text{AS}'(n, d)$  that satisfies the following weaker constraint:

$$\text{AS}'(n, d) \leq K + \left(1 - \frac{1}{2^{O(d)}}\right) \text{AS}'(n, d)$$

which is satisfied by  $\text{AS}'(n, d) \leq 2^{O(d)} \cdot \log n \cdot n^{1-1/(4d+2)}$ .

## 5.2 Regularity and the critical index of polynomials

In [Ser07] a notion of the “critical index” of a linear form was defined and subsequently used in [OS08, DS09, DGJ+09]. We now give a generalization of the critical index notion for polynomials.

**Definition 2** Let  $p : \{-1, 1\}^n \rightarrow \mathbb{R}$  and  $\tau > 0$ . Assume the variables are ordered such that  $\text{Inf}_i(f) \geq \text{Inf}_{i+1}(f)$  for all  $i \in [n - 1]$ . The  $\tau$ -critical index of  $f$  is the least  $i$  such that:

$$\frac{\text{Inf}_{i+1}(p)}{\sum_{j=i+1}^n \text{Inf}_j(p)} \leq \tau. \quad (3)$$

If (3) does not hold for any  $i$  we say that the  $\tau$ -critical index of  $p$  is  $+\infty$ . If  $p$  has  $\tau$ -critical index 0, we say that  $p$  is  $\tau$ -regular.

The following simple lemma will be useful for us. It says that the total influence  $\sum_{i=j+1}^n \text{Inf}_i(p)$  goes down exponentially as a function of  $j$  prior to the critical index:

**Lemma 5.3** *Let  $p : \{-1, 1\}^n \rightarrow \mathbb{R}$  and  $\tau > 0$ . Let  $k$  be the  $\tau$ -critical index of  $p$ . For  $0 \leq j \leq k$  we have*

$$\sum_{i=j+1}^n \text{Inf}_i(p) \leq (1 - \tau)^j \cdot \text{Inf}(p).$$

**Proof:** The lemma trivially holds for  $j = 0$ . In general, since  $j$  is at most  $k$ , we have that

$$\text{Inf}_j(p) \geq \tau \cdot \sum_{i=j}^n \text{Inf}_i(p),$$

or equivalently

$$\sum_{i=j+1}^n \text{Inf}_i(p) \leq (1 - \tau) \cdot \sum_{i=j}^n \text{Inf}_i(p)$$

which yields the claimed bound. ■

Let  $p : \{-1, 1\}^n \rightarrow \mathbb{R}$  be a degree- $d$  polynomial. We note here that the total influence of  $p$  is within a factor of  $d$  of the sum of squares of the non-constant coefficients of  $p$ :

$$\sum_{S \neq \emptyset} \widehat{p}(S)^2 \leq \sum_{i=1}^n \sum_{S \ni i} \widehat{p}(S)^2 = \sum_{i=1}^n \text{Inf}_i(p) = \sum_{S \subseteq [n]} |S| \cdot \widehat{p}(S)^2 \leq d \sum_{S \neq \emptyset} \widehat{p}(S)^2,$$

where the final inequality holds since  $\widehat{p}(S) \neq 0$  only for sets  $|S| \leq d$ .

### 5.3 Restrictions and the influences of variables in polynomials

Let  $p : \{-1, 1\}^n \rightarrow \mathbb{R}$  be a degree- $d$  polynomial. The goal of this section is to understand what happens to the influences of a variable  $x_\ell$ ,  $\ell > k$ , when we do a random restriction to variables  $x_1, \dots, x_k$ .

We start with the following elementary claim:

**Claim 5.4** *Let  $\rho$  be a randomly chosen assignment to the variables  $x_1, \dots, x_k$ . Fix any  $S \subseteq \{k+1, \dots, n\}$ . Then for any polynomial  $p : \{-1, 1\}^n \rightarrow \mathbb{R}$  we have*

$$\widehat{p}_\rho(S) = \sum_{T \subseteq [k]} \widehat{p}(S \cup T) \rho_T,$$

and so we have

$$\mathbf{E}_\rho[\widehat{p}_\rho(S)^2] = \sum_{T \subseteq [k]} \widehat{p}(S \cup T)^2. \tag{4}$$

In words, all the Fourier weight on sets of the form  $S \cup \{\text{some restricted variables}\}$  “collapses” down onto  $S$  in expectation. A corollary of this is that in expectation, the influence of an unrestricted variable  $x_\ell$  does not change when we do a restriction:

**Corollary 5.5** *Let  $\rho$  be a randomly chosen assignment to the variables  $x_1, \dots, x_k$ . Fix any  $\ell \in \{k+1, \dots, n\}$ . Then for any polynomial  $p : \{-1, 1\}^n \rightarrow \mathbb{R}$  we have*

$$\mathbf{E}_\rho[\text{Inf}_\ell(p_\rho)] = \text{Inf}_\ell(p).$$

**Proof:**

$$\begin{aligned} \mathbf{E}_\rho[\text{Inf}_\ell(p_\rho)] &= \mathbf{E}_\rho \left[ \sum_{\ell \in S \subseteq \{k+1, \dots, n\}} \widehat{p}_\rho(S)^2 \right] \\ &= \sum_{T \subseteq [k]} \sum_{\ell \in S \subseteq \{k+1, \dots, n\}} \widehat{p}(S \cup T)^2 \\ &= \sum_{U \ni \ell} \widehat{p}(U)^2 = \text{Inf}_\ell(p). \end{aligned}$$

■

### 5.3.1 Influences of low-degree polynomials behave nicely under restrictions

In this subsection we prove the following lemma: For a low-degree polynomial, a random restriction with very high probability does not cause any variable’s influence to increase by more than a polylog( $n$ ) factor.

**Lemma 5.6** *Let  $p(x_1, \dots, x_n)$  be a degree- $d$  polynomial. Let  $\rho$  be a randomly chosen assignment to the variables  $x_1, \dots, x_k$ . Fix any  $t > e^{2d}$  and any  $\ell \in [k+1, n]$ . With probability at least  $1 - \exp(-\Omega(t^{1/d}))$  over the choice of  $\rho$ , we have*

$$\text{Inf}_\ell(p_\rho) \leq t \cdot 3^d \text{Inf}_\ell(p).$$

*In particular, for  $t = \log^d n$ , we have that with probability at least  $1 - n^{-\omega(1)}$ , every variable  $\ell \in [k+1, n]$  has  $\text{Inf}_\ell(p_\rho) \leq (3 \log n)^d \cdot \text{Inf}_\ell(p)$ .*

**Proof:** Since  $\text{Inf}_\ell(p_\rho)$  is a degree- $2d$  polynomial in  $\rho$ , [Lemma 5.6](#) follows as an immediate consequence of [Theorem 2.1](#) if we can upper bound  $\|\text{Inf}_\ell(p_\rho)\|_2$ . We use the bound in [Lemma 5.7](#), stated and proven below. ■

**Lemma 5.7** *Let  $p(x_1, \dots, x_n)$  be a degree- $d$  polynomial. Let  $\rho$  be a randomly chosen assignment to the variables  $x_1, \dots, x_k$ , and let  $\ell \in [k+1, n]$ . Then  $\text{Inf}_\ell(p_\rho)$  is a degree- $2d$  polynomial in variables  $\rho_1, \dots, \rho_k$ , and*

$$\|\text{Inf}_\ell(p_\rho)\|_2 \leq 3^d \cdot \text{Inf}_\ell(p).$$

**Proof:** The triangle inequality tells us that we may bound the 2-norm of each squared-coefficient separately:

$$\|\text{Inf}_\ell(p_\rho)\|_2 \leq \sum_{\ell \in S \subseteq [k+1, n]} \|\widehat{p}_\rho(S)^2\|_2.$$

Since  $\widehat{p}_\rho(S)$  is a degree- $d$  polynomial, Bonami-Beckner (i.e.,  $(4, 2)$ -hypercontractivity) tells us that

$$\|\widehat{p}_\rho(S)^2\|_2 = \|\widehat{p}_\rho(S)\|_4^2 \leq 3^d \|\widehat{p}_\rho(S)\|_2^2,$$

hence

$$\|\text{Inf}_\ell(p_\rho)\|_2 \leq 3^d \sum_{\ell \in S \subseteq [k+1, n]} \|\widehat{p}_\rho(S)\|_2^2 = 3^d \cdot \text{Inf}_\ell(p)$$

where the last equality is by [Corollary 5.5](#). ■

## 5.4 The regular case

In this section we prove that regular degree- $d$  PTF's have low average sensitivity. In particular, we show:

**Lemma 5.8** *Fix  $\tau = n^{-\Theta(1)}$ . Let  $f$  be a  $\tau$ -regular degree- $d$  PTF. Then,*

$$\text{AS}(f) \leq O(d \cdot n \cdot \tau^{1/(4d+1)})$$

[Claim 5.1](#) follows directly from the above lemma, recalling we choose  $\tau \stackrel{\text{def}}{=} n^{-(4d+1)/(4d+2)}$ . However, the lemma will also be useful in the “small critical index” case for a slightly larger regularity parameter  $\tau$ .

**Proof:** Let  $f : \{-1, 1\}^n \rightarrow \mathbb{R}$  be a degree- $d$  PTF, i.e.  $f = \text{sign}(p)$  where  $p$  is  $\tau$ -regular. We may assume that  $p$  is normalized such that  $\sum_{0 < |S| \leq d} \widehat{p}(S)^2 = 1$ .

First we note that flipping the  $i$ -th bit of an input  $x \in \{-1, 1\}^n$  changes the value of  $p$  by the magnitude of its partial derivative with respect to  $i$ :

$$2D_i p(x) = 2 \sum_{S \ni i} \widehat{p}(S) x_{S-\{i\}}$$

It follows that:

$$\text{Inf}_i(f) \leq \mathbf{Pr}_{x \in \{-1, 1\}^n} [|p(x)| \leq |2D_i p(x)|]$$

Therefore, bounding from above the influence of variable  $i$  in  $f$  can be done by showing the following:

1.  $p(x)$  has small magnitude,  $|p(x)| \leq t$  for some threshold  $t$ , with small probability.
2.  $2D_i p(x)$  has large magnitude,  $|2D_i p(x)| \geq t$ , with small probability.

We bound the probability of the first event using the anti-concentration property of regular low-degree polynomials, as implied by the invariance principle along with [Theorem 2.2](#). For the second event we use the tail bound for degree- $d$  polynomials ([Theorem 2.1](#)).

We will take our threshold  $t$  to be  $t \stackrel{\text{def}}{=} \tau^{1/4}$ , where  $\tau$  is the regularity parameter of  $p$ .

#### 5.4.1 Bounding the probability of the first event

By the  $\tau$ -regularity of  $p$ , for all  $i \in [n]$  we have  $\text{Inf}_i(p) \leq \tau \cdot \text{Inf}(p) \leq d \cdot \tau$  where the last inequality follows by the assumed normalization. With this bound, the invariance principle ([Theorem 2.4](#)) tells us that  $\Pr_{x \in \{-1,1\}^n}[|p(x)| \leq \tau^{1/4}]$  differs from  $\Pr_{\mathcal{G}_1, \dots, \mathcal{G}_n}[|p(\mathcal{G})| \leq \tau^{1/4}]$  by at most  $O(d \cdot (d\tau)^{1/(4d+1)}) = O(d \cdot \tau^{1/(4d+1)})$ . Applying the anti-concentration bound of Carbery and Wright for polynomials in Gaussian random variables ([Theorem 2.2](#)), we get:

$$\begin{aligned} \Pr_x[|p(x)| \leq \tau^{1/4}] &\leq \Pr_{\mathcal{G}_1, \dots, \mathcal{G}_n}[|p(\mathcal{G})| \leq \tau^{1/4}] + O(d\tau^{1/(4d+1)}) \\ &\leq O(d \cdot \tau^{1/4d}) + O(d \cdot \tau^{1/(4d+1)}) \\ &= O(d \cdot \tau^{1/(4d+1)}). \end{aligned}$$

#### 5.4.2 Bounding the probability of the second event

Next we consider  $\Pr_x[|2D_i p(x)| \geq \tau^{1/4}]$ . Note that  $2D_i p$  is a degree- $(d-1)$  polynomial whose  $l_2$  norm is small:

$$\|2D_i p\| = 2 \sqrt{\sum_{S \ni i} \widehat{p}(S)^2} = 2\sqrt{\text{Inf}_i(p)} \leq 2\sqrt{d \cdot \tau}.$$

By ([Theorem 2.1](#)), we get that

$$\begin{aligned} \Pr_x[|2D_i p(x)| \geq \tau^{1/4}] &\leq \Pr_x[|2D_i p(x)| \geq \tau^{-1/4} / (2\sqrt{d}) \cdot \|2D_i p\|] \\ &\leq \exp(-\tau^{-1/(2d)} / (2\sqrt{d})^{2/d}) = \exp(-\Theta(1) \cdot \tau^{-1/(2d)}) \ll O(d \cdot \tau^{1/(4d+1)}). \end{aligned}$$

(In the second inequality, we were able to apply the concentration bound since, by our assumptions on  $d$  and  $\tau$ , we indeed have that  $\tau^{-1/4} / (2\sqrt{d}) > e^d$ .)

Hence, we have shown that:

$$\begin{aligned} \text{Inf}_i(f) &\leq \Pr_{x \in \{-1,1\}^n}[|p(x)| \leq |2D_i p(x)|] \\ &\leq \Pr_x[|p(x)| \leq \tau^{1/4}] + \Pr_x[|2D_i p(x)| \geq \tau^{1/4}] \\ &= O(d \cdot \tau^{1/(4d+1)}). \end{aligned}$$

Since this holds for all indices  $i \in [n]$ , we have the following bound on the average sensitivity of  $f = \text{sign}(p)$ :

$$\text{AS}(f) \leq O(d \cdot n \cdot \tau^{1/(4d+1)}).$$

■

## 5.5 The small critical index case

Let  $f = \text{sign}(p)$  be such that the  $\tau$ -critical index of  $p$  is some value  $k$  between 1 and  $K = 2d \log n / \tau$ . By definition, the sequence of influences  $\text{Inf}_{k+1}(p), \dots, \text{Inf}_n(p)$  is  $\tau$ -regular. We essentially reduce this case to the regular case for a regularity parameter  $\tau'$  somewhat larger than  $\tau$ .

Consider a random restriction  $\rho$  of all the variables up to the critical index. We will show the following:

**Lemma 5.9** *For a  $1/2^{O(d)}$  fraction of restrictions  $\rho$ , the sequence of influences  $\text{Inf}_{k+1}(p_\rho), \dots, \text{Inf}_n(p_\rho)$  is  $\tau'$ -regular, where  $\tau' \stackrel{\text{def}}{=} (3 \log n)^d \cdot \tau$ .*

By our choice of  $\tau = n^{-(4d+1)/(4d+2)}$ , we have that  $\tau' = n^{-\Theta(1)}$ , and so we may apply [Lemma 5.8](#) to these restrictions to conclude that the associated PTFs have average sensitivity at most  $O(d \cdot n \cdot (\tau')^{1/(4d+1)})$ .

**Proof:**

Since the sequence of influences  $\text{Inf}_{k+1}(p), \dots, \text{Inf}_n(p)$  is  $\tau$ -regular, we have

$$\frac{\text{Inf}_i(p)}{\sum_{j=k+1}^n \text{Inf}_j(p)} \leq \tau$$

for all  $i \in [k+1, n]$ .

We want to prove that for a  $1/2^{O(d)}$  fraction of all  $2^k$  restrictions  $\rho$  to  $x_1, \dots, x_k$  we have

$$\frac{\text{Inf}_i(p_\rho)}{\sum_{j=k+1}^n \text{Inf}_j(p_\rho)} \leq \tau'$$

for all  $i \in [k+1, n]$ .

To do this we proceed as follows: [Lemma 5.6](#) implies that, with very high probability over the random restrictions, we have  $\text{Inf}_i(p_\rho) \leq (3 \log n)^d \cdot \text{Inf}_i(p)$ , for all  $i \in [k+1, n]$ . We need to show that for a  $1/2^{O(d)}$  fraction of all restrictions the denominator of the fraction above is at least  $\sum_{j=k+1}^n \text{Inf}_j(p)$  (its expected value). The lemma then follows by a union bound.

We consider the degree- $2d$  polynomial  $A(\rho_1, \dots, \rho_k) \stackrel{\text{def}}{=} \sum_{j=k+1}^n \text{Inf}_j(p_\rho)$  in variables  $\rho_1, \dots, \rho_k$ . The expected value of  $A$  is  $\mathbf{E}_\rho[A] = \sum_{j=k+1}^n \text{Inf}_j(p) = \widehat{A}(\emptyset)$ . We apply the [Theorem 2.3](#) for  $B = A - \widehat{A}(\emptyset)$ . We thus get  $\Pr_\rho[B > 0] > 1/2^{O(d)}$ . We thus get  $\Pr_\rho[A > \mathbf{E}_\rho[A]] > 1/2^{O(d)}$  and we are done.  $\blacksquare$

## 5.6 The large critical index case

Finally we consider PTFs  $f = \text{sign}(p)$  with  $\tau$ -critical index greater than  $K = 2d \log n / \tau$ . Let  $\rho$  be a restriction of the first  $K$  variables  $\mathcal{H} = \{1, \dots, K\}$ ; we call these the “head” variables. We will show the following:

**Lemma 5.10** *For a  $1/2^{O(d)}$  fraction of restrictions  $\rho$ , the function  $\text{sign}(p_\rho(x))$  is a constant function.*

**Proof:** By Lemma 5.3, the surviving variables  $x_{K+1}, \dots, x_n$  have very small total influence in  $p$ :

$$\sum_{i=K+1}^n \text{Inf}_i(p) = \sum_{i=K+1}^n \sum_{S \ni i} \widehat{p}(S)^2 \leq (1 - \tau)^K \cdot \text{Inf}(p) \leq d/n^{2d}. \quad (5)$$

Therefore, if we let  $p'$  be the truncation of  $p$  comprising only the monomials with all variables in  $\mathcal{H}$ ,

$$p'(x_1, \dots, x_k) = \sum_{S \subset \mathcal{H}} \widehat{p}(S) x_S$$

we know that almost all of the original Fourier weight of  $p$  is on the coefficients of  $p'$ :

$$1 \geq \sum_{\substack{S \subset \mathcal{H} \\ |S| > 0}} \widehat{p}(S)^2 \geq 1 - \sum_{i=K+1}^n \text{Inf}_i(p) \geq 1 - d/n^{2d}$$

We now apply Theorem 2.3 to  $p'$ <sup>1</sup> and get:

$$\Pr_{x \in \{-1, 1\}^K} [|p'(x)| \geq 1/2^{O(d)}] \geq 1/2^{O(d)}.$$

In words, for a  $1/2^{O(d)}$  fraction of all restrictions  $\rho$  to  $x_1, \dots, x_K$ , the value  $p'(\rho)$  has magnitude at least  $1/2^{O(d)}$ .

For any such restriction, if the function  $f_\rho(x) = \text{sign}(p_\rho(x_{K+1}, \dots, x_n))$  is not a constant function it must necessarily be the case that:

$$\sum_{0 < |S| \subseteq \{x_{K+1}, \dots, x_n\}} |\widehat{p}_\rho(S)| \geq 1/2^{O(d)}$$

As noted in (5), each tail variable  $\ell > K$  has very small influence in  $p$ :

$$\text{Inf}_\ell(p) \leq \sum_{i=K+1}^n \text{Inf}_i(p) = d/n^{2d}$$

---

<sup>1</sup>after a very slight rescaling so the non-constant Fourier coefficients of  $p'$  have sum of squares equal to 1; this does not affect the bound we get because of the big-O.

Applying [Lemma 5.6](#), we get that for the overwhelming majority of the  $1/2^{O(d)}$  fraction of restrictions mentioned above, the influence of  $\ell$  in  $p_\rho$  is not much larger than the influence of  $\ell$  in  $p$ :

$$\text{Inf}_\ell(p_\rho) \leq (3 \log n)^d \cdot \text{Inf}_\ell(p) \leq d \cdot (3 \log n)^d / n^{2d} \quad (6)$$

Using Cauchy-Schwarz, we have

$$\begin{aligned} \sum_{S \ni \ell, S \subseteq \{x_{K+1}, \dots, x_n\}} |\widehat{p}_\rho(S)| &\leq n^{d/2} \cdot \sqrt{\sum_{S \ni \ell, S \subseteq \{x_{K+1}, \dots, x_n\}} \widehat{p}_\rho(S)^2} \\ &= n^{d/2} \sqrt{\text{Inf}_\ell(p_\rho)} \\ &\leq n^{-\Omega(1)} \end{aligned}$$

where we have used [\(6\)](#) (and our upper bound on  $d$ ). From this we easily get that

$$\sum_{0 < |S| \subseteq \{x_{K+1}, \dots, x_n\}} |\widehat{p}_\rho(S)| \leq n^{-\Omega(1)} \ll 1/2^{O(d)}$$

We have established that for a  $1/2^{O(d)}$  fraction of all restrictions to  $x_1, \dots, x_K$ , the function  $f_\rho = \text{sign}(p_\rho)$  is a constant function, and the lemma is proved.  $\blacksquare$

## 5.7 Proof of [Claim 5.2](#)

If  $f$  is a degree- $d$  PTF that is not  $\tau$ -regular, then its  $\tau$ -critical index is either in the range  $\{1, \dots, K\}$  or it is greater than  $K$ .

In the first case (small critical index case), as shown in [Section 5.5](#), we have that for a  $1/2^{O(d)}$  fraction of restrictions  $\rho$  to variables  $x_1, \dots, x_k$ , the total influence of  $f_\rho = \text{sign}(p_\rho)$  is at most

$$O(d \cdot n \cdot (\tau')^{1/(4d+1)}) = O(d \cdot (\log n)^{1/4} \cdot n^{1-1/(4d+2)}),$$

so the conclusion of [Claim 5.2](#) holds in this case.

In the second case (large critical index case), as shown in [Section 5.6](#), for a  $1/2^{O(d)}$  fraction of restrictions  $\rho$  to  $x_1, \dots, x_K$  the function  $f_\rho$  is constant and hence has zero influence, so the conclusion of [Claim 5.2](#) certainly holds in this case as well.  $\blacksquare$

## 6 A Fourier-Analytic Bound on Boolean Average Sensitivity

In this section, we present a simple proof of the following upper bound on the average sensitivity of a degree- $d$  PTF ([Theorem 1.2](#)):

$$\text{AS}(n, d) \leq 2n^{1-1/2^d}.$$

We recall here the definition of the formal derivative of a function  $f : \{-1, 1\}^n \rightarrow \mathbb{R}$ .

$$D_i p(x) = \sum_{S \ni i} \hat{p}_S x_{S-\{i\}}.$$

It is easy to see that,

$$D_i p(x) = \frac{1}{2} x_i [p(x) - p(x^{\oplus i})] = \frac{1}{2} \left( \frac{p(x) - p(x^{\oplus i})}{x_i} \right) \quad (7)$$

where “ $x^{\oplus i}$ ” means “ $x$  with the  $i$ -th bit flipped.”

For a Boolean function  $f$ , we have  $D_i f(x) = \pm 1$  iff flipping the  $i$ th bit flips  $f$ ; otherwise  $D_i f(x) = 0$ . So we have

$$\text{Inf}_i(f) = \mathbf{E}[|D_i f(x)|].$$

**Lemma 6.1** *Fix  $i \neq j \in [n]$ . Let  $f, g : \{-1, 1\}^n \rightarrow \mathbb{R}$  be functions such that  $f$  is independent of the  $i^{\text{th}}$  bit  $x_i$  and  $g$  is independent of the  $j^{\text{th}}$  bit  $x_j$ . Then*

$$\mathbf{E}_x[x_i x_j f(x) g(x)] \leq \frac{\text{Inf}_i(g) + \text{Inf}_j(f)}{2}.$$

**Proof:** First, note that the influence of  $i^{\text{th}}$  coordinate on a function  $f$  can be written as:

$$\text{Inf}_i(f) = \mathbf{E}_{x_{-i}}[\text{Var}_{x_i}[f(x)]] = \mathbf{E}_x \left[ \left( \frac{|f(x^{\oplus i}) - f(x)|}{2} \right)^2 \right] = \mathbf{E}_{x_{-i}} [|\mathbf{E}_{x_i}[x_i f(x)]|^2] \quad (8)$$

As  $f$  is independent of  $x_i$  and  $g$  is independent of  $x_j$ , we can write,

$$\begin{aligned} \mathbf{E}_x[x_i x_j f(x) g(x)] &= \mathbf{E}_{x_{-\{i,j\}}} \mathbf{E}_{x_i, x_j}[x_i x_j f(x) g(x)] \\ &= \mathbf{E}_{x_{-\{i,j\}}} [\mathbf{E}_{x_i}[x_i g(x)] \mathbf{E}_{x_j}[x_j f(x)]] \\ &\leq \mathbf{E}_{x_{-\{i,j\}}} \left[ \frac{1}{2} |\mathbf{E}_{x_i}[x_i g(x)]|^2 + \frac{1}{2} |\mathbf{E}_{x_j}[x_j f(x)]|^2 \right] \quad (\text{using } ab \leq \frac{1}{2}(a^2 + b^2)) \\ &\leq \frac{\text{Inf}_j(f) + \text{Inf}_i(g)}{2} \quad (\text{using Equation 8}) \end{aligned}$$

■

[Theorem 1.2](#) is shown using an inductive argument over the degree  $d$ . Central to this inductive argument is the following lemma relating the influences of a degree- $d$  PTF  $\text{sign}(p(x))$  to the degree- $(d-1)$  PTFs obtained by taking formal derivatives of  $p$ .

**Lemma 6.2** *For a PTF  $f = \text{sign}(p(x))$  on  $n$  variables and  $i \in [n]$ ,  $\text{Inf}_i(f) = \mathbf{E}[f(x)x_i\text{sign}(D_i p(x))]$ .*

The following simple claim will be useful in the proof of the above lemma.

**Claim 6.3** *For two real numbers  $a, b$ , if  $\text{sign}(a) \neq \text{sign}(b)$  then*

$$\text{sign}(\text{sign}(a) - \text{sign}(b)) = \text{sign}(a - b)$$

**Proof:** If  $\text{sign}(a) = 1$  and  $\text{sign}(b) = -1$  ( $a \geq 0, b < 0$ ) then  $a - b \geq 0$ . Hence in this case,  $\text{sign}(a - b) = 1 = \text{sign}(1 - (-1)) = \text{sign}(\text{sign}(a) - \text{sign}(b))$ . On the other hand, if  $\text{sign}(a) = -1$  and  $\text{sign}(b) = 1$ , then  $\text{sign}(a - b) = -1 = \text{sign}((-1) - 1) = \text{sign}(\text{sign}(a) - \text{sign}(b))$ . ■

**Proof:**[of [Lemma 6.2](#)] The influence of the  $i^{\text{th}}$  coordinate is given by,

$$\begin{aligned} \text{Inf}_i(f) &= \mathbf{E} \left[ \frac{1}{2} |f(x) - f(x^{\oplus i})| \right] \\ &= \mathbf{E} \left[ \frac{1}{2} (f(x) - f(x^{\oplus i})) \text{sign}(f(x) - f(x^{\oplus i})) \right] \end{aligned} \quad (9)$$

Consider an  $x$  for which  $f(x) \neq f(x^{\oplus i})$ . In this case, we can use [Claim 6.3](#) to conclude:

$$\begin{aligned} \text{sign}(f(x) - f(x^{\oplus i})) &= \text{sign}(p(x) - p(x^{\oplus i})) , \\ &= \text{sign}(2x_i D_i p(x)) = x_i \text{sign}(D_i p(x)) . \quad (\text{using } (7)) \end{aligned}$$

Hence for an  $x$  with  $f(x) \neq f(x^{\oplus i})$ ,

$$(f(x) - f(x^{\oplus i})) \text{sign}(f(x) - f(x^{\oplus i})) = (f(x) - f(x^{\oplus i})) x_i \text{sign}(D_i p(x)) .$$

On the other hand, if  $f(x) = f(x^{\oplus i})$  then the above equation continues holds since both the sides evaluate to 0. Substituting this equality into [Equation 9](#) yields,

$$\text{Inf}_i(f) = \frac{1}{2} \mathbf{E} [f(x)x_i\text{sign}(D_i p(x))] - \frac{1}{2} \mathbf{E} [f(x^{\oplus i})x_i\text{sign}(D_i p(x))] .$$

Notice that the  $i^{\text{th}}$  coordinate  $(x^{\oplus i})_i$  of  $x^{\oplus i}$  is given by  $-x_i$ . Since  $D_i p$  is independent of the  $i^{\text{th}}$  coordinate  $x_i$ , we have  $D_i p(x) = D_i p(x^{\oplus i})$ . Rewriting the above equation, we get

$$\begin{aligned} \text{Inf}_i(f) &= \frac{1}{2} \mathbf{E} [f(x)x_i\text{sign}(D_i p(x))] + \frac{1}{2} \mathbf{E} [f(x^{\oplus i})(x^{\oplus i})_i\text{sign}(D_i p(x^{\oplus i}))] , \\ &= \mathbf{E} [f(x)x_i\text{sign}(D_i p(x))] \quad ((x^{\oplus i}) \text{ is also uniformly distributed}) \end{aligned}$$

■

**Theorem 6.4** Let  $AS(n, d)$  denote the max possible average sensitivity of any degree- $d$  PTF on  $n$  variables. Then we have

$$AS(n, d) \leq \sqrt{n + n \cdot AS(n, d - 1)}.$$

**Proof:**

$$\begin{aligned} \text{Inf}(f) &= \sum_i \text{Inf}_i(f) \\ &= \sum_i \mathbf{E}[f(x)x_i \text{sign}(D_i p(x))] \quad (\text{by Lemma 6.2}) \\ &= \mathbf{E}[f(x) \sum_i x_i \text{sign}(D_i p(x))] \\ &\leq \sqrt{\mathbf{E}[f(x)^2]} \cdot \sqrt{\mathbf{E}[(\sum_i x_i \text{sign}(D_i p(x)))^2]} \end{aligned} \tag{10}$$

$$= 1 \cdot \sqrt{\mathbf{E}[\sum_{i,j} x_i x_j \text{sign}(D_i p(x)) \text{sign}(D_j p(x))]} \tag{11}$$

$$\leq \sqrt{\mathbf{E}[\sum_i x_i^2 \text{sign}(D_i p(x))^2] + \sum_{i \neq j} \text{Inf}_i(\text{sign}(D_j p(x)))} \tag{12}$$

$$= \sqrt{n + \sum_{i \neq j} \text{Inf}_i(\text{sign}(D_j p(x)))}. \tag{13}$$

Here (10) is the Cauchy-Schwarz inequality, (11) is expanding the square. Step (12) uses Lemma 6.1 which we may apply since  $D_j p(x)$  does not depend on  $x_i$ .

Observe that for any fixed  $j'$ , we have  $D_{j'} p(x)$  is a degree- $(d-1)$  polynomial and  $\text{sign}(D_{j'} p(x))$  is a degree- $(d-1)$  PTF. Hence, by definition we have,

$$\sum_{i \neq j'} \text{Inf}(\text{sign}(D_{j'} p(x))) \leq AS(n, d - 1),$$

for all  $j' \in [n]$ . Therefore the quantity  $\sum_{i \neq j} \text{Inf}(\text{sign}(D_j p(x))) \leq n \cdot AS(n, d - 1)$ , finishing the proof. ■

The bound on average sensitivity (Theorem 1.2) follows immediately from the above recursive relation.

**Proof:**[of Theorem 1.2] Clearly, we have  $AS(n, 0) = 0$ . For  $d = 1$ , Theorem 6.4 yields  $AS(n, 1) \leq \sqrt{n}$ . Now suppose  $AS(n, d) = 2n^{1-1/2^d}$  for  $d \geq 1$ , then by Theorem 6.4,

$$AS(n, d + 1) \leq \sqrt{n + n \cdot AS(n, d)} \leq \sqrt{4n^{2-1/2^d}} = 2n^{1-1/2^{d+1}},$$

finishing the proof. ■

## 7 Boolean average sensitivity vs noise sensitivity

Our results on Boolean noise sensitivity are obtained via the following simple reduction which translates any upper bound on average sensitivity for degree- $d$  PTFs over Boolean variables into a corresponding upper bound on noise sensitivity. This theorem is inspired by the proof of noise sensitivity of halfspaces by Peres [Per04].

**Theorem 7.1** *Let  $\text{NS}(\epsilon, d)$  denote the maximum noise sensitivity of a degree  $d$ -PTF at a noise rate of  $\epsilon$ . For all  $0 \leq \epsilon \leq 1$  if  $m = \lfloor \frac{1}{\epsilon} \rfloor$  then,*

$$\text{NS}(\epsilon, d) \leq \frac{1}{m} \text{AS}(m, d).$$

[Theorem 1.3](#) follows immediately from this reduction along with our bounds on Boolean average sensitivity ([Theorems 1.1](#) and [1.2](#)), so it remains for us to prove [Theorem 7.1](#).

### 7.1 Proof of [Theorem 7.1](#)

Let  $f(x) = \text{sign}(p(x))$  be a degree  $d$ -PTF. Let us denote  $\delta = \frac{1}{m}$ . As  $\delta \geq \epsilon$ , by the monotonicity of noise sensitivity we have  $\text{NS}_\epsilon(f) \leq \text{NS}_\delta(f)$ . In the following, we will show that  $\text{NS}_\delta(f) \leq \frac{1}{m} \text{AS}(m, d)$  which implies the intended result. Recall that  $\text{NS}_\delta(f)$  is defined as

$$\text{NS}_\delta(f) = \Pr_{x \sim_\delta y} [f(x) \neq f(y)],$$

where  $x \sim_\delta y$  denotes that  $y$  is generated by flipping each bit of  $x$  independently with probability  $\delta$ . An alternate way to generate  $y$  from  $x$  is as follows:

- Sample  $r \in \{1, \dots, m\}$  uniformly at random.
- Partition the bits of  $x$  into  $m = \frac{1}{\delta}$  sets  $S_1, S_2, \dots, S_m$  by independently assigning each bit to a uniformly random set. Formally, a partition  $\alpha$  is specified by a function  $\alpha : \{1, \dots, n\} \rightarrow \{1, \dots, m\}$  mapping bit locations to their partition numbers, i.e.,  $i \in S_{\alpha(i)}$ . A uniformly random partition is picked by sampling  $\alpha(i)$  for each  $i \in \{1, \dots, n\}$  uniformly at random from  $\{1, \dots, m\}$ .
- Flip the bits of  $x$  contained in the set  $S_r$  to obtain  $y$ .

Each bit of  $x$  belongs to the set  $S_r$  independently with probability  $\frac{1}{m} = \delta$ . Therefore, the vector  $y$  generated by the above procedure can equivalently be generated by flipping each bit of  $x$  with probability  $\delta$ .

Inspired by the above procedure, we now define an alternate equivalent procedure to generate the pair  $x \sim_\delta y$ .

- Sample  $a \in \{-1, 1\}^n$  uniformly at random.

- Sample a uniformly random partition  $\alpha : \{1, \dots, n\} \rightarrow \{1, \dots, m\}$  of the bits of  $a$ .
- Sample  $z \in \{-1, 1\}^m$  uniformly at random.
- Sample  $r \in \{1, \dots, m\}$  uniformly at random. Let  $\tilde{z} = z^{\oplus r}$  and

$$x_i = a_i z_{\alpha(i)} \qquad y_i = a_i \tilde{z}_{\alpha(i)}$$

Notice that  $x$  is uniformly distributed in  $\{-1, 1\}^n$ , since both  $a$  and  $z$  are uniformly distributed in  $\{-1, 1\}^n$  and  $\{-1, 1\}^m$  respectively. Furthermore,  $\tilde{z}_i = z_i$  for all  $i \neq r$  and  $\tilde{z}_r = -z_r$ . Therefore,  $y$  is obtained by flipping the bits of  $x$  in the coordinates belonging to the  $r^{\text{th}}$  partition. As the partition  $\alpha$  is generated uniformly at random, this amounts to flipping each bit of  $x$  with probability exactly  $\frac{1}{m} = \delta$ .

The noise sensitivity of  $f$  can be rewritten as,

$$\text{NS}_\delta(f) = \Pr_{a, \alpha, z, r} [f(x) \neq f(y)]$$

For a fixed choice of  $a$  and  $\alpha$ ,  $f(x)$  is a function of  $z$ . In this light, let us define the function  $f_{a, \alpha} : \{-1, 1\}^m \rightarrow \{-1, 1\}$  for each  $a, \alpha$  as  $f_{a, \alpha}(z) = f(x)$ . Returning to the expression for noise sensitivity we get:

$$\begin{aligned} \text{NS}_\delta(f) &= \Pr_{a, \alpha, z, r} [f_{a, \alpha}(z) \neq f_{a, \alpha}(\tilde{z})] , \\ &= \mathbf{E}_{a, \alpha, z, r} [\mathbf{1}[f_{a, \alpha}(z) \neq f_{a, \alpha}(z^{\oplus r})]] , \\ &= \mathbf{E}_{a, \alpha, z} \left[ \frac{1}{m} \sum_{r=1}^m \mathbf{1}[f_{a, \alpha}(z) \neq f_{a, \alpha}(z^{\oplus r})] \right] , \\ &= \mathbf{E}_{a, \alpha} \left[ \frac{1}{m} \sum_{r=1}^m \mathbf{E}_z [\mathbf{1}[f_{a, \alpha}(z) \neq f_{a, \alpha}(z^{\oplus r})]] \right] . \end{aligned}$$

In the above calculation, the notation  $\mathbf{1}[E]$  refers to the indicator function of the event  $E$ . Recall that, by definition of influences,

$$\text{Inf}_r(f_{a, \alpha}) = \mathbf{E}_z [\mathbf{1}[f_{a, \alpha}(z) \neq f_{a, \alpha}(z^{\oplus r})]] ,$$

for all  $r$ . Thus, we can rewrite the noise sensitivity of  $f$  as

$$\text{NS}_\delta(f) = \mathbf{E}_{a, \alpha} \left[ \frac{1}{m} \sum_{r=1}^m \text{Inf}_r(f_{a, \alpha}) \right] = \frac{1}{m} \mathbf{E}_{a, \alpha} [\text{Inf}(f_{a, \alpha})] . \quad (14)$$

We claim that  $f_{a, \alpha}$  is a degree  $d$ -PTF in  $m$  variables. To see this observe that

$$f_{a, \alpha}(z) = \text{sign}(p(x_1, \dots, x_n)) = \text{sign}(p(a_1 z_{\alpha(1)}, \dots, a_n z_{\alpha(n)})) ,$$

which for a fixed choice of  $a, \alpha$  is a degree  $d$ -PTF in  $z$ . Consequently, by definition of  $\text{AS}(m, d)$  we have  $\text{Inf}(f_{a, \alpha}) \leq \text{AS}(m, d)$  for all  $a$  and  $\alpha$ . Using this in (14), the result follows.

## 8 Application to Agnostic Learning

In this section, we outline the applications of the noise sensitivity bounds presented in this work to agnostic learning of PTFs. Specifically, we will present the proofs of [Theorem 1.6](#) and [Theorem 1.7](#). To begin with, we recall the main theorem of [\[KKMS08\]](#) about the  $L_1$  polynomial regression algorithm:

**Theorem 8.1** *Let  $\mathcal{D}$  be a distribution over  $X \times \{-1, 1\}$  (where  $X \subseteq \mathbb{R}^n$ ) which has marginal  $\mathcal{D}_X$  over  $X$ . Let  $\mathcal{C}$  be a class of Boolean-valued functions over  $X$  such that for every  $f \in \mathcal{C}$ , there is a degree- $d$  polynomial  $p(x_1, \dots, x_n)$  such that  $\mathbf{E}_{x \sim \mathcal{D}_X}[(p(x) - f(x))^2] \leq \epsilon^2$ . Then given independent draws from  $\mathcal{D}$ , the  $L_1$  polynomial regression algorithm runs in time  $\text{poly}(n^d, 1/\epsilon, \log(1/\delta))$  and with probability  $1 - \delta$  outputs a hypothesis  $h : X \times \{-1, 1\}$  such that  $\mathbf{Pr}_{(x,y) \sim \mathcal{D}}[h(x) \neq y] \leq \text{opt} + \epsilon$ , where  $\text{opt} = \min_{f \in \mathcal{C}} \mathbf{Pr}_{(x,y) \sim \mathcal{D}}[f(x) \neq y]$ .*

We first consider the case where  $\mathcal{D}_X$  is the uniform distribution over the  $n$ -dimensional Boolean hypercube  $\{-1, 1\}^n$ . Klivans et al. [\[KOS04\]](#) observed that Boolean noise sensitivity bounds are easily shown to imply the existence of low-degree polynomial approximators in the  $L_2$  norm under the uniform distribution on  $\{-1, 1\}^n$ :

**Fact 8.2** *For any Boolean function  $f : \{-1, 1\}^n \rightarrow \{-1, 1\}$  and any value  $0 \leq \gamma < 1/2$ , there is a polynomial  $p(x)$  of degree at most  $d = 1/\gamma$  such that  $\mathbf{E}[(p(x) - f(x))^2] \leq \frac{2}{1-e^{-2}} \text{NS}_\gamma(f)$ .*

[Theorem 1.6](#) follows directly from [Theorem 8.1](#), [Fact 8.2](#) and [Theorem 1.3](#).

Next we turn to the case where  $\mathcal{D}_X$  is the  $\mathcal{N}(0, I_n)$  distribution over  $\mathbb{R}^n$ . In [\[KOS08\]](#) observed that using entirely similar arguments to the Boolean case, Gaussian noise sensitivity bounds are imply the existence of low-degree polynomial approximators in the  $L_2$  norm:

**Fact 8.3** *For any Boolean function  $f : \{-1, 1\}^n \rightarrow \{-1, 1\}$  and any value  $0 \leq \gamma < 1/2$ , there is a polynomial  $p(x)$  of degree at most  $d = 1/\gamma$  such that  $\mathbf{E}_{\mathcal{G} \sim \mathcal{N}(0, I_n)}[(p(\mathcal{G}) - f(\mathcal{G}))^2] \leq \frac{2}{1-e^{-1}} \text{GNS}_\gamma(f)$ .*

[Theorem 1.7](#) follows directly from [Theorem 8.1](#), [Fact 8.3](#) and [Theorem 1.5](#).

## References

- [AH09] Per Austrin and Johan Håstad. Randomly supported independence and resistance. In *Proc. 41st Annual ACM Symposium on Theory of Computing (STOC)*, pages 483–492. ACM, 2009. [8](#)
- [BK97] J. Bourgain and G. Kalai. Influences of variables and threshold intervals under group symmetries. *GAF*, 7:438–461, 1997. [1](#)
- [BKS99] I. Benjamini, G. Kalai, and O. Schramm. Noise sensitivity of Boolean functions and applications to percolation. *Inst. Hautes Études Sci. Publ. Math.*, 90:5–43, 1999. [1](#), [2](#)
- [Bog98] V. Bogachev. *Gaussian measures*. Mathematical surveys and monographs, vol. 62, 1998. [3](#)
- [BOW08] E. Blais, R. O’Donnell, and K. Wimmer. Polynomial regression under arbitrary product distributions. In *Proc. 21st Annual Conference on Learning Theory (COLT)*, pages 193–204, 2008. [3](#)
- [BT96] N. Bshouty and C. Tamon. On the Fourier spectrum of monotone functions. *Journal of the ACM*, 43(4):747–770, 1996. [1](#)
- [CW01] A. Carbery and J. Wright. Distributional and  $L^q$  norm inequalities for polynomials over convex bodies in  $R^n$ . *Mathematical Research Letters*, 8(3):233–248, 2001. [0](#), [5](#), [8](#)
- [DFKO06] Irit Dinur, Ehud Friedgut, Guy Kindler, and Ryan O’Donnell. On the Fourier tails of bounded functions over the discrete cube. In *Proc. 38th ACM Symp. on Theory of Computing*, pages 437–446, 2006. [8](#)
- [DGJ<sup>+</sup>09] I. Diakonikolas, P. Gopalan, R. Jaiswal, R. Servedio, and E. Viola. Bounded independence fools halfspaces. To appear in *Proc. 50th IEEE Symposium on Foundations of Computer Science (FOCS)*, 2009. [6](#), [14](#)
- [DS09] I. Diakonikolas and R. Servedio. Improved approximation of linear threshold functions. In *Proc. 24th Annual IEEE Conference on Computational Complexity (CCC)*, pages 161–172, 2009. [14](#)
- [DSTW09] I. Diakonikolas, R. Servedio, L.-Y. Tan, and A. Wan. A regularity lemma, and low-weight approximators, for low-degree polynomial threshold functions. manuscript, 2009. [4](#)
- [Fri98] E. Friedgut. Boolean functions with low average sensitivity depend on few coordinates. *Combinatorica*, 18(1):474–483, 1998. [1](#)
- [GKK08] P. Gopalan, A. Kalai, and A. Klivans. Agnostically learning decision trees. In *Proc. 40th Annual ACM Symposium on Theory of Computing (STOC)*, pages 527–536, 2008. [3](#)

- [GL94] C. Gotsman and N. Linial. Spectral properties of threshold functions. *Combinatorica*, 14(1):35–50, 1994. 0, 2
- [GS09] P. Gopalan and R. Servedio. Learning threshold-of- $AC^0$  circuits. Manuscript, 2009. 4
- [HKM09] P. Harsha, A. Klivans, and R. Meka. Bounding the sensitivity of polynomial threshold functions. Manuscript, 2009. 5
- [Jan97] S. Janson. *Gaussian Hilbert Spaces*. Cambridge University Press, Cambridge, UK, 1997. 0, 5
- [JKS02] J. Jackson, A. Klivans, and R. Servedio. Learnability beyond  $AC^0$ . In *Proc. 34th Annual ACM Symposium on Theory of Computing (STOC)*, pages 776–784, 2002. 4
- [KKL88] J. Kahn, G. Kalai, and N. Linial. The influence of variables on boolean functions. In *Proc. 29th Annual Symposium on Foundations of Computer Science (FOCS)*, pages 68–80, 1988. 1
- [KKMS08] A. Kalai, A. Klivans, Y. Mansour, and R. Servedio. Agnostically learning halfspaces. *SIAM Journal on Computing*, 37(6):1777–1805, 2008. 0, 3, 27
- [KMV08] A. Kalai, Y. Mansour, and E. Verbin. On agnostic boosting and parity learning. In *Proc. 40th Annual ACM Symposium on Theory of Computing (STOC)*, pages 629–638, 2008. 3
- [KOS04] A. Klivans, R. O’Donnell, and R. Servedio. Learning intersections and thresholds of halfspaces. *Journal of Computer & System Sciences*, 68(4):808–840, 2004. 2, 3, 4, 27
- [KOS08] A. Klivans, R. O’Donnell, and R. Servedio. Learning geometric concepts via Gaussian surface area. In *Proc. 49th IEEE Symposium on Foundations of Computer Science (FOCS)*, pages 541–550, 2008. 2, 3, 4, 27
- [LMN93] N. Linial, Y. Mansour, and N. Nisan. Constant depth circuits, Fourier transform and learnability. *Journal of the ACM*, 40(3):607–620, 1993. 4
- [MO03] E. Mossel and R. O’Donnell. On the noise sensitivity of monotone functions. *Random Structures and Algorithms*, 23(3):333–350, 2003. 1, 2
- [MOO05] E. Mossel, R. O’Donnell, and K. Oleszkiewicz. Noise stability of functions with low influences: invariance and optimality. In *Proc. 46th Symposium on Foundations of Computer Science (FOCS)*, pages 21–30, 2005. 0, 1, 3, 5, 6, 7, 8
- [OS07] R. O’Donnell and R. Servedio. Learning monotone decision trees in polynomial time. *SIAM J. Comput.*, 37(3):827–844, 2007. 1

- [OS08] R. O’Donnell and R. Servedio. The Chow Parameters Problem. In *Proc. 40th Annual ACM Symposium on Theory of Computing (STOC)*, pages 517–526, 2008. [14](#)
- [OSSS05] R. O’Donnell, M. Saks, O. Schramm, and R. Servedio. Every decision tree has an influential variable. In *Proc. 46th Symposium on Foundations of Computer Science (FOCS)*, pages 31–39, 2005. [1](#)
- [Per04] Y. Peres. Noise stability of weighted majority, 2004. [25](#)
- [Ser07] R. Servedio. Every linear threshold function has a low-weight approximator. *Computational Complexity*, 16(2):180–209, 2007. [0](#), [4](#), [6](#), [13](#), [14](#)
- [Shi00] Y. Shi. Lower bounds of quantum black-box complexity and degree of approximating polynomials by influence of boolean variables. *Inform. Process. Lett.*, 75(1-2):79–83, 2000. [1](#)
- [SS] O. Schramm and J. Steif. Quantitative noise sensitivity and exceptional times for percolation. *Ann. Math.*, to appear. [2](#)
- [SSS09] S. Shalev Shwartz, O. Shamir, and K. Sridharan. Agnostically learning halfspaces with margin errors. TTI Technical Report, 2009. [3](#)