

Bianchi Type V String Cosmological Models in General Relativity

ANIL KUMAR YADAV¹, VINEET KUMAR YADAV², LALLAN YADAV³

¹Department of Physics, Anand Engineering College, Keetham, Agra -282 007, India

¹E-mail : abanilyadav@yahoo.co.in, akyadav@imsc.res.in

^{2,3}Department of Physics, D. D. U. Gorakhpur University, Gorakhpur -273 009, India

²E-mail : vineetky@yahoo.co.in

³E-mail : nisaly06@rediffmail.com

Abstract

Bianchi type V string cosmological models in general relativity are investigated. To get the exact solution of Einstein's field equations, we have taken some scale transformations followed by Camci *et al* (2001). It is shown that Einstein's field equations are solvable for any arbitrary cosmic scale function. Solution for particular form of cosmic scale functions are also obtained. Some physical and geometrical aspects of the models are discussed.

PACS: 98.80.cq, 98.80.-k

Key words: Cosmology, Bianchi type V universe, string theory.

1 Introduction

The study of Bianchi type V cosmological models play an important role in the study of universe and it create more interest as these models contains isotropic special cases and permit arbitrary small anisotropy levels at some instant of time. The string theory plays a significant role in the study of physical situation at the very early stages of the formation of the universe. It is generally assumed that after the big bang , the universe may have undergone a series of phase transitions as its temperature lowered down below some critical temperature as predicted by grand unified theories [1-6]. At the very early stages of evolution of the universe, it is believed that during phase transition the symmetry of the universe is broken spontaneously. It can give rise to topologically stable defects such as domain walls, strings and monopoles. In all these three cosmological structures, only cosmic strings have excited the most interesting consequence

¹corresponding author

[7] because they are believed to give rise to density perturbations which lead to formation of galaxies[8]. These cosmic strings can be closed like loops or open like a hair which move through time and trace out a tube or a sheet, according to whether it is closed or open. The string is free to vibrate and its different vibrational modes present different types of particles carrying the force of gravitation. This is why much interesting to study the gravitational effect that arises from strings by using Einstein's field equations.

The general relativistic treatment of strings has been initially given by Stachel [9] and Letelier [10,11]. Letelier [10] obtained the general solution of Einstein's field equations for a cloud of strings with spherical, plane and a particular case of cylindrical symmetry. Letelier [11] also obtained massive string cosmological models in Bianchi type-I and Kantowski-Sachs space-times. Benjerjee et al. [12] have investigated an axially symmetric Bianchi type I string dust cosmological model in presence and absence of magnetic field. Exact solutions of string cosmology for Bianchi type II, VI_0 , VIII and IX space-times have been studied by Krori et al. [13] and Wang [14,15]. Bali and Upadhaya [16] have presented LRS Bianchi type I string dust magnetized cosmological models. Singh and Singh [17] investigated string cosmological models with magnetic field in the context of space-time with G_3 symmetry. Singh [18,19] has studied string cosmology with electromagnetic fields in Bianchi type II, VIII and IX space-times.

Bianchi V universes are the natural generalization of FRW models with negative curvature. These open models are favored by the available evidences for low density universes (Gott et al [20]). Bianchi type V cosmological model where matter moves orthogonally to the hyper-surface of homogeneity, has been studied by Heckmann and Schucking [21]. Exact tilted solutions for the Bianchi type V space-time are obtained by Hawkings [22], Grishchuk et al. [23]. Ftaclas and Cohen [24] have investigated LRS Bianchi type V universes containing stiff matter with electromagnetic field. Lorentz [25] has investigated LRS Bianchi type V tilted models with stiff fluid and electromagnetic field. Pradhan et al have investigated The generation of Bianchi type V cosmological models with varying Λ term. Yadav et al.[27,28] have investigated bulk viscous string cosmological models in different space-times. Bali and Anjali [29], Bali [30] have obtained Bianchi type-I, and type V string cosmological models in general relativity. The string cosmological models with a magnetic field are discussed by Tikekar and Patel [31], Patel and Maharaj [32]. Ram and Singh [33] obtained some new exact solution of string cosmology with and without a source free magnetic field for Bianchi type I space-time in the different basic form considered by Carminati and McIntosh [34]. Yavuz et al. [35] have examined charged strange quark matter attached to the string cloud in the spherical symmetric space-time admitting one-parameter group of conformal motion. Kaluza-Klein cosmological solutions are obtained by Yilmaz [36] for quark matter attached to the string cloud in the context of general relativity. Recently, Baysal et al. [37], Kilinc and Yavuz [38], Pradhan [39] and Pradhan et al.[40,41], Yadav et al [42] have investigated some string cosmological models in cylindrically symmetric inhomogeneous universe.

In this paper, we have investigated string cosmological models based on generation technique and obtained a more realistic behavior of the universe. This paper is organised as follows. The field equations are presented in Section 2. In Section 3, we deal with generation technique for solution of field equation and finally the results are discussed in Section 4.

2 Field Equations

We consider the Bianchi type V metric of the form

$$ds^2 = dt^2 - A^2(t)dx^2 - e^{2\alpha x}[B^2(t)dy^2 + C^2(t)dz^2] \quad (1)$$

where α is a constant.

The energy momentum tensor for cloud of string is given by

$$T_i^j = \rho v_i v^j - \lambda x_i x^j \quad (2)$$

where v_i and x_i satisfy the condition

$$v^i v_j = -x^i x_j = 1, v^i x_i = 0. \quad (3)$$

Here ρ is the proper energy density of the cloud of string with particle attached to them. λ is the string tension density, v^i , the four velocity of the particles and x^i , the unit space vector representing the direction of strings. If the particle density of the configuration is denoted by ρ_p then we have

$$\rho = \rho_p + \lambda \quad (4)$$

For the energy momentum tensor (2) and Bianchi type V metric (1), Einstein's field equations

$$R_i^j - \frac{1}{2}Rg_{ij} = -8\pi T_{ij} \quad (5)$$

yield the following five independent equations

$$\frac{\ddot{A}}{A} + \frac{\ddot{B}}{B} + \frac{\dot{A}\dot{B}}{AB} - \frac{\alpha^2}{A^2} = 0 \quad (6)$$

$$\frac{\ddot{A}}{A} + \frac{\ddot{C}}{C} + \frac{\dot{A}\dot{C}}{AC} - \frac{\alpha^2}{A^2} = 0 \quad (7)$$

$$\frac{\ddot{B}}{B} + \frac{\ddot{C}}{C} + \frac{\dot{B}\dot{C}}{BC} - \frac{\alpha^2}{A^2} = -8\pi\lambda \quad (8)$$

$$\frac{\dot{A}\dot{B}}{AB} + \frac{\dot{A}\dot{C}}{AC} + \frac{\dot{B}\dot{C}}{BC} - \frac{3\alpha^2}{A^2} = -8\pi\rho \quad (9)$$

$$\frac{2\dot{A}}{A} - \frac{\dot{B}}{B} - \frac{\dot{C}}{C} = 0 \quad (10)$$

Here and what follows the dots overhead the symbol A, B, C denotes differentiation with respect to t.

The physical quantities expansion scalar θ and shear scalar σ^2 have the following expressions

$$\theta = \frac{\dot{A}}{A} + \frac{\dot{B}}{B} + \frac{\dot{C}}{C} \quad (11)$$

$$\sigma^2 = \frac{1}{2}\sigma_{ij}\sigma^{ij} = \frac{1}{3} \left[\theta^2 - \frac{\dot{A}\dot{B}}{AB} - \frac{\dot{A}\dot{C}}{AC} - \frac{\dot{B}\dot{C}}{BC} \right] \quad (12)$$

Integrating equation (10) and absorbing the integrating constant into B or C, we obtain

$$A^2 = BC \quad (13)$$

without loss of any generality.

From equations (6), (7) and (13), we obtain

$$2\frac{\ddot{B}}{B} + \left(\frac{\dot{B}}{B}\right)^2 = 2\frac{\ddot{C}}{C} + \left(\frac{\dot{C}}{C}\right)^2 \quad (14)$$

which on integration yields

$$\frac{\dot{B}}{B} - \frac{\dot{C}}{C} = \frac{k}{(BC)^{\frac{3}{2}}} \quad (15)$$

where k is the constant of integration. Hence for the metric function B or C from the above first order differential equation (15), some scale transformations permit us to obtain new metric function B or C.

Firstly, under the scale transformation $dt = B^{\frac{1}{2}}dT$, equation (15) takes the form

$$CB_T - BC_T = kC^{-\frac{1}{2}} \quad (16)$$

where the subscript represents derivative with respect to T. Considering equation (16) as a linear differential equation for B, where C is an arbitrary function, we obtain

$$(i) \quad B = k_1C + kC \int \frac{dT}{C^{\frac{3}{2}}}, \quad (17)$$

where k_1 is the the constant of integration. Similarly, using the transformation $dt = B^{\frac{3}{2}}d\bar{T}$, $dt = C^{\frac{1}{2}}d\tilde{t}$ and $dt = C^{\frac{3}{2}}d\tau$ in equation (15) after some algebra we obtain respectively.

$$(ii) \quad B(\bar{T}) = k_2C e^{\left(k \int \frac{d\bar{T}}{C^{\frac{3}{2}}}\right)}, \quad (18)$$

$$(iii) \quad C(\tilde{t}) = k_3B - kB \int \frac{d\tilde{t}}{B^{\frac{3}{2}}}, \quad (19)$$

$$(iv) \quad C(\tau) = k_4 B e^{\left(k \int \frac{d\tau}{B^{\frac{3}{2}}}\right)}, \quad (20)$$

where k_2 , k_3 and k_4 are constant of integration. Thus choosing any given function B or C in cases (i), (ii), (iii) and (iv), one can obtain B or C.

3 Generation technique for solution

We consider the following four cases

3.1 Case (i): $C = T^n$ (n is a real number satisfying $n \neq \frac{2}{5}$)

Equation (16) leads to

$$B = k_1 T^n + \frac{2k}{2-5n} T^{1-\frac{3n}{2}} \quad (21)$$

From equations (13) and (21), we obtain

$$A^2 = k_1 T^{2n} + \frac{2k}{2-5n} T^{1-\frac{n}{2}} \quad (22)$$

Hence metric (1) reduces to the following form

$$ds^2 = (k_1 T^n + 2l T^{l_1}) [dT^2 - T^n dx^2] - e^{2\alpha x} \left[(k_1 T^n + 2l T^{l_1})^2 dy^2 + T^{2n} dz^2 \right], \quad (23)$$

where $l = \frac{k}{2-5n}$ and $l_1 = 1 - \frac{3n}{2}$

In this case the physical parameters, i.e. the string tension density (λ), the energy density (ρ), the particle density (ρ_p) and the kinematical parameters, i.e. the scalar of expansion (θ), shear scalar (σ) and the proper volume (V^3) for model (23) are given by

$$8\pi\lambda = \left[-2k_1^2 n(n-1) T^{2n-2} - k_1 l n (10-13n) T^{-(l_1+2n)} - \frac{1}{2} l^2 (4+4n-11n^2) T^{-3n} \right] \\ (k_1 T^n + 2l T^{l_1})^{-3} + \alpha^2 T^{-n} (k_1 T^n + 2l T^{l_1})^{-1}, \quad (24)$$

$$8\pi\rho = \left[3k_1^2 n^2 T^{2n-2} + 3k_1 l n (2-n) T^{-(l_1+2n)} + \frac{1}{2} l^2 (4+4n-11n^2) T^{-3n} \right] \\ (k_1 T^n + 2l T^{l_1})^{-3} - 3\alpha^2 T^{-n} (k_1 T^n + 2l T^{l_1})^{-1}, \quad (25)$$

$$8\pi\rho_p = \left[n(5n-2) k_1^2 T^{2n-2} + 16k_1 l n (1-n) T^{-(l_1+2n)} + l^2 (4+4n-11n^2) T^{-3n} \right] \\ (k_1 T^n + 2l T^{l_1})^{-3} - 4\alpha^2 T^{-n} (k_1 T^n + 2l T^{l_1})^{-1} \quad (26)$$

$$\theta = 3 \left[k_1 n T^{n-1} + \frac{1}{2} l (2-n) T^{-\frac{3n}{2}} \right] (k_1 T^n + 2l T^{l_1})^{-\frac{3}{2}} \quad (27)$$

$$\sigma = \frac{1}{2} k T^{-\frac{3n}{2}} (k_1 T^n + 2l T^{l_1})^{-\frac{3}{2}} \quad (28)$$

$$V^3 = (k_1 T^{2n} + 2l T^{n+l_1})^{\frac{3}{2}} e^{2\alpha x} \quad (29)$$

Equations (27) and (28) leads to

$$\frac{\sigma}{\theta} = \frac{1}{6} k \left[k_1 n T^{n-l_1} + \frac{1}{2} l (2-n) \right]^{-1} \quad (30)$$

The energy condition $\rho \geq 0$ and $\rho_p \geq 0$ satisfy for model (23). The condition $\rho \geq 0$ and $\rho_p \geq 0$ leads to

$$\left[3k_1^2 n^2 T^{3n-2} + 3k_1 l n (2-n) T^{-(l_1+n)} + \frac{1}{2} l^2 (4+4n-11n^2) T^{-2n} \right] (k_1 T^n + 2l T^{l_1})^{-2} \geq 3\alpha^2 \quad (31)$$

$$\left[n(2-5n)k_1^2 T^{3n-2} + 16k_1 l n (n-1) T^{-(l_1+n)} + l^2 (11n^2 - 4n - 4) T^{-2n} \right] (k_1 T^n + 2l T^{l_1})^{-2} \geq 4\alpha^2 \quad (32)$$

respectively.

We observe that the string tension density $\lambda \geq 0$, leads to

$$\left[-2k_1^2 n (n-1) T^{3n-2} - k_1 l n (10-13n) T^{-(l_1+n)} - \frac{1}{2} l^2 (4+4n-11n^2) T^{-2n} \right] (k_1 T^n + 2l T^{l_1})^{-2} \geq -\alpha^2 \quad (33)$$

Generally the model (23) are expanding, shearing and approaches to isotropy at late time. For $k = 0$, the solution represents the shear free model of universe. We observe that as $T \rightarrow \infty$, $V^3 \rightarrow \infty$ and $\rho \rightarrow 0$ hence volume increases when T increases and the proper energy density of the cloud of string with particle attached to them decreases i. e. the proper energy density (ρ) is decreasing function of time.

3.2 case(ii): $C = \bar{T}^n$ (n is a real number satisfying $n \neq \frac{2}{3}$)

Equation (18) leads to

$$B = k_2 \bar{T}^n e^{M\bar{T}^{l_1}} \quad (34)$$

From equations (13), and (34), we obtain

$$A^2 = k_2 \bar{T}^{2n} e^{M\bar{T}^{l_1}} \quad (35)$$

where $M = \frac{k}{l_1}$. Hence the metric (1) reduces to the form

$$ds^2 = \bar{T}^{\frac{4(1-l_1)}{3}} \left[\bar{T}^{\frac{2(1-l_1)}{3}} e^{3M\bar{T}^{l_1}} d\bar{T}^2 - e^{M\bar{T}^{l_1}} dx^2 - e^{2\alpha x} \left(e^{2M\bar{T}^{l_1}} dy^2 + dz^2 \right) \right], \quad (36)$$

Here k_2 is constant and we have taken k_2 as unity with out any loss of generality. In this case the physical parameters, i.e. the string tension density (λ), the energy density (ρ), the particle density (ρ_p) and the kinematical parameters, i.e. the scalar of expansion (θ), shear scalar (σ) and the proper volume (V^3) for model (36) are given by

$$8\pi\lambda = 2n\bar{T}^{2(l_1-2)} + 3nk\bar{T}^{3l_1-4} + \frac{1}{2}k^2\bar{T}^{4(l_1-1)} + \alpha^2\bar{T}^{\frac{4(l_1-1)}{3}}e^{-3M\bar{T}^{l_1}}, \quad (37)$$

$$8\pi\rho = 3n^2\bar{T}^{2(l_1-2)} + 3nk\bar{T}^{3l_1-4} + \frac{1}{2}k^2\bar{T}^{4(l_1-1)} - 3\alpha^2\bar{T}^{\frac{4(l_1-1)}{3}}e^{-3M\bar{T}^{l_1}}, \quad (38)$$

$$8\pi\rho_p = n(3n-2)\bar{T}^{2(l_1-2)} - 4\alpha^2\bar{T}^{\frac{4(l_1-1)}{3}}e^{-3M\bar{T}^{l_1}} \quad (39)$$

$$\theta = 3 \left[n\bar{T}^{l_1-2} + \frac{1}{2}k\bar{T}^{2(l_1-1)} \right], \quad (40)$$

$$\sigma = \frac{1}{2}k\bar{T}^{2(l_1-1)}e^{-3M\bar{T}^{l_1}} \quad (41)$$

$$V^3 = \left(k_3 \bar{T}^{2n} e^{M\bar{T}^{l_1}} \right)^{\frac{3}{2}} e^{2\alpha x} \quad (42)$$

From equation(40) and (41) leads to

$$\frac{\sigma}{\theta} = \frac{k}{6(n\bar{T}^{-l_1} + \frac{1}{2}k)} \quad (43)$$

The energy condition $\rho \geq 0$ and $\rho_p \geq 0$ are satisfy for model (36). The condition $\rho \geq 0$ and $\rho_p \geq 0$ leads to

$$e^{3M\bar{T}^{l_1}} \left[3n^2\bar{T}^{\frac{2l_1-8}{3}} + 3nk\bar{T}^{\frac{5l_1-8}{3}} + \frac{1}{2}k^2\bar{T}^{\frac{8l_1-8}{3}} \right] \geq 3\alpha^2 \quad (44)$$

$$n(3n-2)e^{3M\bar{T}^{l_1}}\bar{T}^{\frac{(2l_1-8)}{3}} \geq 4\alpha^2 \quad (45)$$

respectively.

we observe that the string tension density $\lambda \geq 0$, leads to

$$e^{3M\bar{T}^{l_1}} \left[2n\bar{T}^{\frac{2l_1-8}{3}} + 3nk\bar{T}^{\frac{5l_1-8}{3}} + \frac{1}{2}k^2\bar{T}^{\frac{8l_1-8}{3}} \right] \geq -\alpha^2 \quad (46)$$

For $l_1 > 2$, model (36) is expanding and for $l_1 < 2$, model starts with big bang singularity. Generally the model is expanding, shearing and approaches to isotropy at late time. For $k = 0$, the solution represents the shear free model of universe. We observe that as $\bar{T} \rightarrow \infty$, $V^3 \rightarrow \infty$ and $\rho \rightarrow 0$ hence volume increases when \bar{T} increases and the proper energy density of the cloud of string with particle attached to them decreases i. e. the proper energy density (ρ) is decreasing function of time.

3.3 Case(iii): $B = \tilde{t}^n$ (n is a real number)

Equation (19) leads to

$$C = k_3\tilde{t}^n - 2l\tilde{t}^{l_1} \quad (47)$$

From Equations (13) and (47), we obtain

$$A^2 = k_3\tilde{t}^{2n} - 2l\tilde{t}^{l_1+n} \quad (48)$$

Thus the metric (1) reduces to the form

$$ds^2 = (k_3\tilde{t}^n - 2l\tilde{t}^{l_1}) [dt^2 - \tilde{t} dx^2] - e^{2\alpha x} [\tilde{t}^{2n} dy^2 + (k_3\tilde{t}^n - 2l\tilde{t}^{l_1})^2 dz^2], \quad (49)$$

In this case the physical parameters, i.e. the string tension density (λ), the energy density (ρ), the particle density (ρ_p) and the kinematical parameters, i.e. the scalar of expansion (θ), shear scalar (σ) and the proper volume (V^3) for model (49) are given by

$$8\pi\lambda = \left[-\frac{1}{2}l^2(11n^2 - 4n - 4)\tilde{t}^{-3n} + lk_3n(13n - 10)\tilde{t}^{l_1+n} - 2k_3^2n(n-1)\tilde{t}^{2n-2} \right] \\ (k_3\tilde{t}^n - 2l\tilde{t}^{l_1})^{-3} + \alpha^2\tilde{t}^{-n} (k_3\tilde{t}^n - 2l\tilde{t}^{l_1})^{-1}, \quad (50)$$

$$8\pi\rho = \left[-\frac{1}{2}l^2(11n^2 - 4n - 4)\tilde{t}^{-3n} - 3lk_3n(2-n)\tilde{t}^{l_1+n} + 3k_3^2n^2\tilde{t}^{2n-2} \right] \\ (k_3\tilde{t}^n - 2l\tilde{t}^{l_1})^{-3} - 3\alpha^2\tilde{t}^{-n} (k_3\tilde{t}^n - 2l\tilde{t}^{l_1})^{-1} \quad (51)$$

$$8\pi\rho_p = (k_3\tilde{t}^n - 2l\tilde{t}^{l_1})^{-3} [n(5n-2)k_3^2\tilde{t}^{2n-2} - lk_3n(12n-8)\tilde{t}^{l_1+n}] \\ - 4\alpha^2\tilde{t}^n (k_3\tilde{t}^n - 2l\tilde{t}^{l_1})^{-1} \quad (52)$$

$$\theta = 3 \left[\frac{1}{2} l(n-2) \tilde{t}^{-\frac{3n}{2}} + k_3 n \tilde{t}^{n-1} \right] (k_3 \tilde{t}^n - 2l \tilde{t}^{l_1})^{-\frac{3}{2}}, \quad (53)$$

$$\sigma = \frac{1}{2} k \tilde{t}^{-\frac{3n}{2}} (k_3 \tilde{t}^n - 2l \tilde{t}^{l_1})^{-\frac{3}{2}} \quad (54)$$

$$V^3 = (k_3 \tilde{t}^{2n} - 2l \tilde{t}^{l_1+n})^{\frac{3}{2}} e^{2\alpha x} \quad (55)$$

Equation(53) and (54) leads to

$$\frac{\sigma}{\theta} = \frac{k}{6} \left[k_3 n \tilde{t}^{-(l_1+n)} + \frac{l(n-2)}{2} \right]^{-1} \quad (56)$$

The energy condition $\rho \geq 0$ and $\rho_p \geq 0$ are satisfy for model (49). The condition $\rho \geq 0$ and $\rho_p \geq 0$ leads to

$$\left[-\frac{1}{2} l^2 (11n^2 - 4n - 4) \tilde{t}^{-2n} - 3l k_3 n (2-n) \tilde{t}^{l_1+2n} + 3k_3^2 n^2 \tilde{t}^{(3n-2)} \right] (k_3 \tilde{t}^n - 2l \tilde{t}^{l_1})^{-2} \geq 3\alpha^2 \quad (57)$$

$$(k_3 \tilde{t}^n - 2l \tilde{t}^{l_1})^{-2} [n(5n-2)k_3^2 \tilde{t}^{n-2} - l k_3 n (12n-8) \tilde{t}^{l_1}] \geq 4\alpha^2 \quad (58)$$

respectively.

We observe that the string tension density $\lambda \geq 0$, leads to

$$\left[-\frac{1}{2} l^2 (11n^2 - 4n - 4) \tilde{t}^{-2n} + l k_3 (13n - 10) \tilde{t}^{l_1+2n} - 2k_3^2 n (n-1) \tilde{t}^{3n-2} \right] (k_3 \tilde{t}^n - 2l \tilde{t}^{l_1})^{-2} \geq -\alpha^2 \quad (59)$$

Thus we see that the model (49) is generally expanding, shearing and approaches to isotropy at late time. For $k = 0$, the solution represents the shear free model of universe. We observe that as $\tilde{t} \rightarrow \infty$, $V^3 \rightarrow \infty$ and $\rho \rightarrow 0$ hence volume increases when \tilde{t} increases and the proper energy density of the cloud of string with particle attached to them decreases i. e. the proper energy density (ρ) is decreasing function of time.

3.4 Case(iv): $B = \tau^n$ (n is any real number)

Equation (20) leads to

$$C = k_4 \tau^n e^{\left(\frac{k}{l_1} \tau^{l_1}\right)} \quad (60)$$

From equation (13) and (60), we obtain

$$A^2 = k_4 \tau^{2n} e^{\left(\frac{k}{l_1} \tau^{l_1}\right)} \quad (61)$$

Hence the metric (1) reduces to

$$ds^2 = \tau^{2n} e^{\left(\frac{k}{l_1} \tau^{l_1}\right)} \left[\tau^n e^{\left(\frac{2k}{l_1} \tau^{l_1}\right)} d\tau^2 - dx^2 \right] - e^{2\alpha x} \left[dy^2 + e^{\left(\frac{2k}{l_1} \tau^{l_1}\right)} dz^2 \right], \quad (62)$$

Here k_4 is constant and we have taken k_4 as unity without any loss of generality. In this case we see that A, B, and C are exponential function as that of in case (ii). Thus the physical and geometrical properties of the model (62) are similar to model (36).

4 Conclusion

If we choose $\alpha = 0$, metric (1) becomes Bianchi type I metric, studied by several authors in different context. In this paper, we have applied the technique followed by Camci et al [43] for solving Einstein's field equations and found new solution for string cosmological model. It is shown that the Einstein's field equation are solvable for an arbitrary cosmic scale function. Starting from a particular cosmic function, new classes of spatially homogeneous and anisotropic cosmological models have been investigated for which the string fluid are rotation free but they do have expansion and shear. It is also observed that in all cases, the physical and geometrical behavior of models are similar. Generally the models are expanding, shearing and non rotating. All the models are isotropized at late time.

In case (i), for $n \geq 1$, Model (23) start expanding with big bang singularity and for $n \leq 0$, Bianchi type-V universe preserve expanding nature as $T \rightarrow 0, \theta \rightarrow 0$ and $T \rightarrow \infty, \theta \rightarrow \infty$. It is also observed that for $k = 0$, shear scalar (σ) vanishes and model becomes isotropic. In case (ii), we observed that for $l_1 < 2$, model (36) started with big bang singularity and expand through out the evolution of universe and for $l_1 > 2$, model (36) also preserve the expanding nature as $\bar{T} \rightarrow 0, \theta \rightarrow 0$ and $\bar{T} \rightarrow \infty, \theta \rightarrow \infty$. From equation (41), it is clear that when $k = 0$, the shear scalar (σ) vanishes and model becomes isotropic. In case (iii), it is observed that model (49) started with big bang singularity and expanding through the evolution of universe. From equation (54), it is clear that for $k = 0$, the shear scalar (σ) vanishes and model isotropizes. In case (iv), it is observed that the physical and geometrical properties of the model (62) are similar to model (36), i.e the case (ii). We note that in all cases, $\rho(t)$ is a decreasing function of time and it is always positive. Further it is observed that for sufficiently large time, ρ_p and λ tend to zero. Therefore the strings disappear from the universe at late time (i. e. present epoch). The same is predicted by current observations.

Acknowledgements

The authors are extremely grateful to the referee for his fruitful comments. Author (AKY) is thankful to IMSc, Chennai and HRI, Allahabad for providing

research facilities .

References

- [1] A E Everett, Phys. Rev. **24**, 858 (1981)
- [2] T W B Kibble, J. Phys. A **9**, 1387 (1976)
- [3] T W B Kibble, Phys. Rep. **67**, 183 (1980)
- [4] A Vilenkin, Phys. Rev. D **24**, 2082 (1981)
- [5] Ya B Zel'dovich, I Yu Kobzarev, and L B Okun, Zh. Eksp. Teor. Fiz. **67**, 3 (1975)
- [6] Ya B Zel'dovich, I Yu Kobzarev, and L B Okun, Sov. Phys.-JETP **40**, 1 (1975)
- [7] A Vilenkin, Phy. Rep. **121**, 263 (1985)
- [8] Ya B Zel'dovich, Mon. Mot. R. Astron. Soc. **192**, 663 (1980)
- [9] J Stachel, Phys. Rev. D **21**, 2171 (1980)
- [10] P S Letelier, Phys. Rev. D, **20**, 1249 (1979)
- [11] P S Letelier, Phys. Rev. D, **28**, 2414 (1983)
- [12] A Banerjee, A K Sanyal, and S Chakraborty, Pramana-J. Phys. **34**, 1 (1990)
- [13] K D Krori, T Chaudhury, C R Mahanta, and A Mazumder, Gen. Rel. Grav., **22** 123 (1990)
- [14] X X Wang, Chin. Phys. Lett. **20**, 615 (2003)
- [15] X X Wang, Chin. Phys. Lett., **20**, 1205 (2003)
- [16] R Bali and R D Upadhaya, Astrophys. Space Sci. **283**, 97 (2002)
- [17] G P Singh, and T Singh, Gen. Rel. Grav. **31**, 371 (1999)
- [18] G P Singh, Nuovo Cim. B **110**, 1463 (1995)
- [19] G P Singh, Pramana-J. Phys. **45**, 189 (1995)
- [20] J R Gott et al, Astrophys. J. **194**, 543 (1974)
- [21] O heckmann and E Schucking, In Gravitation: An Introduction to Current Research, (1962)
- [22] S W Hawking, Mon. Not. R. Astron. Soc. **142**, 129 (1969)

- [23] L P Grishchuck, A G Doroshkevich and I D Novikov, Sov. Phys. JETP. **28**, 1214 (1969)
- [24] C Ftacilas and J M Cohen, Phys. Rev. D, **18**, 4373 (1978)
- [25] D Lorentz, Gen. Rel. Grav. **13**, 795 (1981)
- [26] A Pradhan, A K Yadav and L Yadav, Czech. J. Phys. **55**, (2005) 503.
- [27] M K Yadav, A Rai, and A Pradhan, Int. J. Theor. Phys. **46**, 2677 (2007)
- [28] M K Yadav, A Pradhan and S K Singh, Astrophys. Space Sci., **311**, 423 (2007)
- [29] R Bali and Anjali, Astrophys. Space Sci. **302**, 201 (2006)
- [30] Raj Bali, Electronic journal of Theor. Phys. **5**, 105 (2008)
- [31] R. Tikekar and L.K.Patel, Gen. Rel. Grav. **24**, 397 (1992)
- [32] L K Patel and S D Maharaj, Pramana-J. Phys. **47**, 1 (1996)
- [33] S Ram and T K Singh. Gen. Rel. Grav., **27**, 1207 (1995)
- [34] J Carminati, and C B G McIntosh, J. Phys. A: Math. Gen. **13**, 953 (1980)
- [35] I Yavuz I Yilmaz and H Baysal, Int. J. Mod. Phys. D **14**, 1365 (2005)
- [36] I. Yilmaz Gen. Rel. Grav. **38**,(2006) 1397
- [37] H Baysal, I Yavuz,I Tarhan, U Camci and I Yilmaz, Turk. J. Phys. **25**, 283 (2001)
- [38] C B Kilinc and I Yavuz, Astrophys. Space Sci. **238**, 239 (1996)
- [39] A Pradhan, Fizika B (Zagreb) **16**, 205 (2007)
- [40] A Pradhan, A K Yadav, R P Singh and V K Singh, Astrophys. Space Sci, **312**, 145 (2007)
- [41] A Pradhan, M K Mishra and A K Yadav, Rom. J. Phys., **54**, 747 (2009); arXiv:0705.1765 [gr-qc].
- [42] A K Yadav, V K Yadav and L Yadav, Int. J. Theor. Phys. **48**, 568 (2009)
- [43] U Camci, I Yavuz, H Baysal, I Tarhan and I Yilmaz, Astrophys. Space Sci. **275**, 391 (2001)