

Competitive cluster growth on networks: complex dynamics and survival strategies

N. Nirmal Thyagu* and Anita Mehta†

*Department of Theoretical Studies, S. N. Bose
Centre for Basic Sciences, Kolkata-700078, India*

(Dated: February 2, 2019)

Abstract

We extend the study of a model of competitive cluster growth in an active medium to a basis of networks; this is done by adding nonlocal connections with probability p to sites on a regular lattice, thus enabling one to interpolate between regularity and full randomness. The model on networks demonstrates high sensitivity to small changes in initial configurations, which we characterize using damage spreading. The main focus of this paper is, however, the devising of survival strategies through selective networking, to alter the the fate of an arbitrarily chosen cluster: whether this be to revive a dying cluster to life, or to make a weak survivor into a stronger one. Although such goals are typically achieved by networking with relatively small clusters, our results suggest that it ought to be possible also to network with peers and larger clusters. The main indication of this comes from the probability distributions of mass differences between survivors and their immediate neighbours, which show an astonishing universality; they suggest strategies for winning against the odds.

*Electronic address: nirmal@physics.iitm.ac.in

†Electronic address: anita@bose.res.in

I. INTRODUCTION

The study of complex systems with interacting agents is of great interest in very many fields ranging from physics through quantitative biology to economics [1]. Modelling the dynamics of interacting species should capture the competition between the species for growth and survival, which underlies all evolutionary dynamics. One of the most famous models of this class is the Lotka-Volterra model used for studying predator-prey interactions [2], which is loosely predicated on the principle of the survival of the fittest [3].

Typically, little attention has been paid to the case where agents interact through an active medium; this is a case where the Lotka-Volterra model cannot be simply applied. For instance, in the case of interacting traders, a central exchange or reserve bank could represent such an active medium; this could selectively deplete or enrich individuals depending on taxation modes in the system. Our present study is based on such an example [4]; the detailed model of competing clusters on which it is based [5], forms the basis of our study here. Clusters in this model are embedded in an active medium; they can grow or die depending on their masses, and different geometries lead to different survival statistics. In a mean-field scenario, for instance, there is one survivor at most; on a finite-dimensional lattice with nearest-neighbour interactions, on the other hand, there can be many survivors who are isolated from each other by empty sites (whose original occupants have died).

From recent studies of complex networks in the real world [6, 7], it is becoming increasingly clear that interactions between agents are neither of the mean field type, where all of the species interact with all others, nor necessarily confined to local neighbourhoods. Recent models of real-world networks, such as small world and scale-free networks, have been devised for the proper modelling of, e.g. ecological networks and the world wide web [7, 8]; it is becoming increasingly clear that in fact networks are a far more appropriate basis for the modelling of most systems with their combination of globality and locality, as well as their freer choice of agent-agent contacts. One of the main aims of this paper is thus to study the competitive cluster model of [5] on networks.

The plan of the paper is as follows. First, we introduce the model (Sec. II) and do a deeper investigation of the problem in two dimensions in Sec. III. Here, we demonstrate the inherent complexity of the problem in finite-dimensional systems, where, contrary to the mean-field case, the largest cluster is not necessarily a survivor. Next, we investigate the

main features of the model on a network. Starting with a regular network (which here is a finite-dimensional lattice) with nearest-neighbour interactions, we add non-local links to the existing lattice, with an associated probability p , in the range $0 \leq p \leq 1$ [8]. We measure the survivor ratios as a function of the rewiring probability p , as p is increased to reflect the topologies of small world networks and fully random networks ($p = 1$). Our system turns out to exhibit great sensitivity to small changes in the initial distribution of clusters; we characterize this using the concept of damage spreading (Sec. V).

In Sec. VI, we probe deeper into a more interesting problem by asking whether the destiny of a selected cluster can be changed by networking. We evolve a strategy to do this by selectively networking a cluster of our choice with non-local clusters based on their initial masses. One of our most interesting observations in this paper is that a cluster which would die in its original neighbourhood, *is indeed able to change its fate*, becoming a survivor via selective networking. Next, in Sec. VII we calculate the pair-wise probability for a cluster to survive against larger and smaller neighbours; the probability distributions obtained show an interesting universality, independent of the mass considered. We discuss the implications of this and other results in our concluding section, Sec. VIII.

II. MODEL

The present model was first used in the context of cosmology to describe the mass accretion of black holes in the presence of a radiation field [9]. Its applications, however, are considerably more general; used in the context of economics [4], it manifested an interesting rich-get-richer behaviour. Here, we recapitulate some of its principal properties [5].

Consider an array of immobile clusters with time-dependent masses $m_i(t)$ located at equi-spaced regular lattice sites. The time evolution of masses of the clusters is given by the coupled deterministic first order equations,

$$\frac{dm_i}{dt} = \left(\frac{\alpha}{t} - \frac{1}{t^{1/2}} \sum_j g_{ij} \frac{dm_j}{dt} \right) m_i - \frac{1}{m_i}. \quad (1)$$

Here, the parameter α is called the mass accretion parameter and g_{ij} defines the coupling between the clusters m_i and m_j . The first term in the R. H. S of Eqn. 1 represents the gain term, which is composed of two terms - the free rate of growth of an individual cluster (proportional to the parameter $\alpha > 1/2$), and the rate of growth induced by all other

clusters coupled through the surrounding medium. The coupling g_{ij} between two clusters is proportional to the inverse square distance d_{ij}^2 between them at an initial time t_0 [5]. The negative second term, $-1/m_i$, represents the dissipation to the surrounding medium.

A logarithmic time is introduced in the study for convenience. We define a reduced time $s = \ln(t/t_0)$, where t_0 is the initial time. Similarly, for convenience, we define reduced masses $X_i = m_i/t^{1/2}$. Using the new variables, Eqn . 1 can be rewritten as,

$$\frac{dX_i}{ds} \equiv X'_i = \left(\frac{2\alpha - 1}{2} - \sum_j g_{ij} \left(\frac{X_j}{2} + X'_j \right) \right) \quad (2)$$

The primes denote the differentiation performed with respect to the reduced time variable s throughout.

Continuing our recapitulation of the results of the model [5], we consider a scenario where there is a single isolated cluster, whose initial reduced square mass is X_0 . Under the dynamics defined by Eqn. 2, the cluster will be a survivor only if $X_0 > X_\star (= \sqrt{\frac{2}{2\alpha-1}})$, else it will eventually die out (see Fig. 1) [5]. Next, consider a system of two clusters with equal initial masses; it has been shown [5] that there exists a critical coupling g_c such that for $g < g_c$ they both survive if their initial masses are greater than X_\star . In the case of two clusters with unequal masses, the smallest of them dies first, in reduced time s_1 , and the largest cluster will either survive (if $X(s_1) > X_\star$) or die (if $X(s_1) < X_\star$).

Consider the limit of an infinitely large number of clusters all connected to each other; this represents a limiting mean field regime, with fully collective behaviour involving long-range interactions. For $g > g_c$, all the clusters but the largest will eventually die out. In the weak coupling regime ($g < g_c$), which is the focus of this paper, the dynamics consists of two successive stages [5]. In Stage I, the clusters behave as if they were isolated; they grow (or die) quickly if their masses are greater (or less) than the threshold X_\star . In Stage II, slow collective dynamics leads to a scenario where, again, only the single largest cluster survives. This weakly interacting mean field regime shows the presence of two well-separated time scales, a characteristic feature of glassy systems [5].

Similar glassy dynamics also arise when the model is solved on a periodic lattice with nearest-neighbour interactions. The dynamical equations are [5]:

$$X'_n = \left(\frac{2\alpha - 1}{2} + g \sum_{\mathbf{m}} \left(\frac{1}{X_{\mathbf{m}}} - \alpha X_{\mathbf{m}} \right) \right) X_n - \frac{1}{X_n}. \quad (3)$$




FIG. 1: (color online) The plots show the evolution of individual non-interacting clusters with a range of initial masses obeying Eqn. 2. Clusters with initial mass X_0 greater than X_* live forever, or else they die in time. Here $\alpha = 1.0$, therefore $X_* = \sqrt{\frac{2}{2\alpha-1}} = \sqrt{2}$.

Here \mathbf{m} runs over the z nearest neighbours of the site \mathbf{n} , where for a one-dimensional ring topology $z = d$, for a two-dimensional lattice $z = 2d$, and so on for any space of dimension d . We summarise earlier results on the dynamics: In Stage I, clusters evolve independently; only those whose initial mass $X_i(s=0) > X_*$ survive this stage. In Stage II, the dynamics is slow and glassy; the survivors of this stage are fully isolated from each other by ‘dead’ sites, and their number asymptotes to a constant S_∞ (Fig. 2). These isolated clusters survive forever.

Following the mean field scenario, where the largest cluster is the only one to survive, we ask if this would also be the case for the finite-dimensional case: that is, are the survivors the largest in the locality? Somewhat surprisingly, this is not always the case. While one can certainly rule out the survival of a cluster whose initial mass is less than threshold (X_*), many-body interactions give rise to extremely complex dynamics in Stage II for clusters with $X > X_*$. The asymptotic configurations of survivors in the finite-dimensional case are thus often nontrivial and counter-intuitive (see Sec. III).




FIG. 2: The survival ratio $S(s)$ plotted as a function of reduced time s , for clusters distributed in a regular one-dimensional lattice of size 100,000. In Stage I, clusters grow independently, while in Stage II the growth is collective. Here, $S_1 = 0.8$ is the survival ratio at the end of Stage I and S_∞ is the asymptotic survival ratio.

III. NONTRIVIAL DYNAMICS IN TWO-DIMENSIONAL REGULAR LATTICES

In this section we do a deeper investigation of the cluster growth model [5], with individual clusters placed on the vertices of a regular two-dimensional lattice with nearest-neighbour interactions. While the results of Sec. V will show the extreme sensitivity of the model to the distribution of cluster masses, the results of this section will help us get a flavour of the complexity displayed by the model.

We choose a 50×50 regular lattice for our investigations in this paper, unless otherwise specified; we have checked that there are no observable finite size effects for this choice. Initial masses are characterized by an exponential distribution $P(X_0) = \mu \exp(-\mu X_0)$, where the mean is taken to be $\mu = -\log(S_{(1)})/X_*$, for different values of S_1 . The coupling parameter is set at $g = 0.001$ and the parameter α set to 1.0 [5].




FIG. 3: The left panels show the cluster configurations at time $s = 0$. The right panels show the asymptotic configurations. (a) The central cluster dominates its immediate neighbours and emerges as the survivor. (b) Three larger neighbours eventually dominate the central cluster to become survivors by killing it. Open and filled circles represent non-survivors and survivors, respectively.

First we consider two of the more obvious cases for demonstration (Fig. 3). In Fig. 3(a), the central cluster has the largest mass among its neighbours, and in Fig. 3(b), the central cluster has the smallest mass of those of its three neighbours which are larger than threshold (the fourth one will obviously not survive beyond Stage I).

These two cases show results similar to the mean field scenario [5]; the central cluster with the largest initial mass becomes a survivor (Fig. 3(a)) and the central cluster with the smallest initial mass dies off (Fig. 3(b)).

However, in general, such simplistic reasoning may not work. In a general scenario such as in Fig. 4, consider the cluster located at co-ordinates $(0, 0)$, which has three larger and one smaller neighbours. Reasoning naively, we might expect that the larger neighbours $X_{(1,0)}, X_{(-1,0)}, X_{(0,1)}$ will eventually be able to kill the smaller central cluster $X_{(0,0)}$. Contrary to this, we find that in the asymptotic limit, the central cluster $(0, 0)$ is a survivor.

Similar results are obtained for the clusters at the coordinates $(1, 1)$ and $(1, -1)$; although surrounded by larger neighbours, they nevertheless survive against the odds. Another apparently counter-intuitive case is illustrated in Fig. 4 for the cluster with coordinates $(-1, 0)$. To start with, all the neighbours are smaller than the central cluster, but eventually the two neighbours at $(0, 0)$ and $(-1, 1)$ survive at the expense of the central cluster, which dies.

The explanation for such apparent anomalies is the many-body nature of this problem; the evolution of neighbouring sites is influenced by their neighbours, and so on. Examining the dynamical equation given in Eq. 3,

$$X'_n = \left(\frac{2\alpha - 1}{2} + g \sum_{\mathbf{m}} \left(\frac{1}{X_{\mathbf{m}}} - \alpha X_{\mathbf{m}} \right) \right) X_n - \frac{1}{X_n}.$$

we see that the overall contribution of the $\mathbf{m} = 4$ immediate neighbours of a given cluster at any instant of time is expressed by the summation, $g \sum_{\mathbf{m}} \left(\frac{1}{X_{\mathbf{m}}} - \alpha X_{\mathbf{m}} \right)$, where X_m denotes the reduced masses of the m neighbours. This term should be positive for X'_n to be positive, which means that the overall rate of growth of its neighbours $\sum X'_m$ should be negative.

Therefore, for a cluster to become a survivor, it should grow constantly; its rate of growth X'_n should thus always be positive. Should the overall rate of growth of the neighbours (given by the summation term in Eqn. 3) exceed the rate of growth of the cluster under consideration, they can, during the collective dynamics of Stage II, kill it.

Next, we present plots showing the evolution of a given cluster and its immediate neighbours for the cases discussed in Fig. 4. Consider the evolution of the cluster at $(0, 0)$, shown in Fig. 5 (a). The immediate neighbours of this cluster are seen to die when the central cluster grows (see Fig. 5(b)). In Fig. 5(c) we plot the rate of growth of the central cluster (in ‘red’) and the overall growth of the immediate neighbours $\sum X'_{(i,j)}$ (in ‘green’). Initially, both $X'_{(0,0)}$ and $\sum X'_{(i,j)}$ are positive (Stage I), with the neighbours growing faster than the central site, in accord with intuitive expectations based on their initial masses. When the collective dynamics of Stage II sets in, $\sum X'_{(i,j)}$ falls and becomes negative; this is a demonstration of many-body interactions in action, where neighbours are influenced by their neighbours, and so on. Finally, soon after the neighbours begin to decay ($\sum X'_{(i,j)}$ gets negative), the rate of growth of the central site, $X'_{(0,0)}$, shoots up, making it a survivor that continues to grow forever.

Graphs similar to those in Fig. 5 have been obtained for the other survivors of Fig. 4, viz. $X_{(1,1)}$ and $X_{(1,-1)}$; we have not plotted them here, in the interests of brevity. To




FIG. 4: A small region of the two-dimensional lattice showing neighbouring clusters with comparable initial masses. The numbers show the initial masses; open and filled circles represent non-survivors and survivors, respectively, in the asymptotic limit. This is an illustration of the complex dynamics referred to in the text.

complete our analysis, we show however the graphs in a contrasting case, where a given cluster eventually dies, despite being surrounded by smaller neighbours. Choosing $X_{(1,-1)}$ in Fig. 4, we show the growth of the central site and its associated neighbours, respectively, in Fig. 6 (a) and (b). Here, contrary to expectations, the central cluster dies in a finite time, while two of its neighbours manage to survive. These dynamics are seen clearly in Fig. 6 (c); in the fast dynamics of Stage I, the combined rate of growth of the neighbours is faster than the central cluster. After the onset of slow dynamics in Stage II, the rate of growth of the central cluster goes negative, which contributes to the growth of its neighbours.

Clearly from the above, the simple rule of the mean field scenario, that the largest cluster is always a survivor, does not hold for lattice-based models with nearest-neighbour interactions. In fact, it is rare to be able to find cases like Figs. 3 (a)-(b), where we can ignore the effect of further neighbours. The generic case is much more as depicted in Fig. 4, where information on the immediate neighbourhood of a site does *not* provide a reliable indication




FIG. 5: (Color online) (a) The central cluster $X_{(0,0)}$ as a function of reduced time s . (b) The neighbouring clusters around $(0,0)$ as a function of s . (c) The rate of evolution of the central cluster and the combined rates of evolution of the immediate neighbours are compared - the central cluster wins the competition.

of its fate, highlighting the innate complexity of our model on a lattice.

IV. CLUSTER DYNAMICS IN COMPLEX NETWORKS

In the previous section we discussed the complex dynamics exhibited by the cluster growth model on a lattice. However, both lattices and mean field scenarios are topologically unrealistic - recent work indicates [6, 7] that most real-world systems are more appropriately solved on networks, which are spatially disordered arrays with non-trivial connections between their sites. A particularly interesting example is the class of small world networks, which are constructed by starting with a regular lattice, and adding links randomly to its sites with probability p [8]. Alternative constructions are also possible, where existing links




FIG. 6: (Color online) (a) The growth of the central cluster $X_{(-1,0)}$ as a function of reduced time s . (b) The growth of the neighbouring clusters around $(-1,0)$ as a function of s . (c) The rate of evolution of the central cluster and the combined rates of evolution of the immediate neighbours are compared - the central site loses.

are ‘rewired’; i.e. existing links are severed and then reconnected to randomly chosen lattice sites with probability p . Small world networks have the property that long-range interactions can co-exist with short-range interactions; they contain realistic elements like ‘hubs’, where certain sites are preferentially endowed with many connections, as is the case for example, in the aviation industry [10].

We shall in the following apply ‘static rewiring’ [11] to construct our networks; here, the rewiring is done and frozen for all time, rather than evolving continuously.




FIG. 7: (Color online) The plots of $S(s)$ as a function of reduced time s , for increasing values of rewiring probabilities p . The 1-cycle scheme is shown in (a) for a 1-dimensional ring and (c) for a 2-dimensional square lattice. The 5-cycle scheme is shown in (b) for a 1-dimensional ring and (d) for a 2-dimensional square lattice. The insets in all the figures show the asymptotic survival ratios S_∞ as a function of the probability p . Here, the system size for the 1-d ring is 2000 and for the 2-d square lattice it is 50×50 - all our data is averaged over 10 random network configurations.

A. One-dimensional ring and two-dimensional square lattices

First, we consider a regular one-dimensional ring lattice of size $N = 2000$. To start with, the clusters located on the lattice sites evolve according to Eqn. 3, where the interactions are only with nearest neighbours. Next, we modify the lattice by adding new links between sites chosen randomly with an associated probability p lying $p \in [0, 1]$. For $p = 0$, the network is undisturbed and therefore ordered, while for $p = 1$, the network becomes completely random.

In our first scheme, we add links probabilistically starting with site $i = 1$ and end with $i = N$, only once: we call this the 1-cycle scheme. The survival ratios of clusters as a function of reduced time s for different values of rewiring probability p are presented in Fig. 7(a). Consider the $p = 0$ case, which corresponds to a regular lattice; here the survival ratio $S(s)$ shows two stages, Stage I and Stage II, in its evolution. For all values of $0 < p \leq 1$, the existence of these well-separated Stages I and II is also observed. There is a noticeable fall in the survivor ratio as p is increased, though; this is clearly visible in the asymptotic values $S_\infty(p)$ plotted with respect to p in the inset of Fig. 7(a). As the probability p increases, the number of links increases leading to more interaction and competition between the clusters and hence a decrease in the number of survivors. As expected, this behaviour interpolates between the two characteristic behaviours relevant to the regular lattice and mean field scenarios.

Next we implement the 5-cycle scheme, where the rewiring is done five times. Figure 7(b) shows the survivor ratio $S(s)$ as a function of s for this scheme. We can clearly see the decrease in the number of survivors at all times, for increasing values of p . The asymptotic survival ratios S_∞ for the 1-cycle scheme and 5-cycle scheme are shown in the insets of Fig. 7(a) and (b), respectively, where the monotonic decrease of S_∞ with increasing p is very clear for both cases. In addition, for all values of p , the survivor ratio in the 5-cycle scheme is consistently smaller than in the 1-cycle scheme.

The same procedure for a two-dimensional square lattice of size 50×50 is followed, and survival ratios obtained as a function of s using the 1-cycle scheme and 5-cycle scheme (see Figs. 7(c)-(d)). The asymptotic survivor ratios for these two cases are shown in the insets of Figs. 7(c)-(d); they follow a decreasing trend with increasing p , similar to the one-dimensional case. Notice that the trends in the plots for the 5-cycle scheme are far smoother

than those for the corresponding 1-cycle plots, as expected.

V. DAMAGE SPREADING RESULTS

In this section, we measure the sensitivity of the system to slight perturbations in terms of damage spreading. The concept of damage spreading was introduced in the context of biological systems [12] and was widely used in the study of the dynamics of statistical mechanical systems [13, 14]. However, our definition of damage spreading is different from the usual definition based on the Hamming distance, used by workers on Ising models; we will elaborate on this below.

A. Sensitivity to small perturbations

Again, we consider a 1-dimensional ring lattice with only nearest-neighbour interactions. An exponential distribution of initial masses of clusters is put on the lattice, as before. Starting with two identical initial mass configurations Ω_A and Ω_B , we swap two randomly chosen clusters in the configuration Ω_B . We recall from the discussion above that the evolution of clusters in a finite-dimensional lattice consists of two stages, fast and slow, and that an initial cluster can be a survivor at the end of Stage I only if its mass $X_i > X_*$. Therefore, while perturbing the initial mass configuration Ω_B , we consider a variety of regimes that takes into account all possible combinations of clusters that are swapped. Table I summarizes all such combinations of X_i and X_j , scaled with respect to the threshold square mass X_* . Next, we evolve the two configurations Ω_A and Ω_B in time and find the absolute difference of the *total mass* of the two configurations, $|m_A - m_B|$, with time. Figure 8 shows the difference $|m_A - m_B|$ for all five regimes shown in the table. Note that irrespective of the regime, the quantity $|m_A - m_B|$ diverges with time. We conclude therefore that the system is very sensitive to the initial masses of the clusters, even for the regular lattice, where *no* random links are added.

We now summarize the above results. First, even for the regular square lattice, we have demonstrated the complexity of the dynamics: it is impossible to predict the fate of a randomly chosen cluster unless its initial mass is less than the critical mass X_* . In Sec. IV we have started our study of the model on a network structure by rewiring the square lattice;

TABLE I: The five regimes defined here are with respect to the threshold mass X_* . The clusters located at sites i and j are swapped in the configuration Ω_B .

Regime	X_i	X_j
Regime 1	$\ll X_*$	$\ll X_*$
Regime 2	$\leq X_*$	$\ll X_*$
Regime 3	$\leq X_*$	$> X_*$
Regime 4	$\geq X_*$	$> X_*$
Regime 5	$> X_*$	$> X_*(X_i \sim X_j)$

we have shown that as the probability of random connections p increases, the asymptotic number of survivors decreases monotonically. This decrease, attributable to the death of clusters with marginal masses when their contacts increase in number, matches with the results of the mean field scenario [5]. From a damage spreading point of view, we also see that the system displays great sensitivity to the distribution of mass within it; the average mass difference between a perturbed and parent cluster diverges in time, irrespective of the




FIG. 8: The absolute mass difference $|m_A - m_B|$ of two mass configurations Ω_A and Ω_B , as a function of the reduced time s . The case shown here is for a regular 1-dimensional regular ring lattice of size 1000.

masses of clusters that are swapped,

Given all of this, we might guess that it would be even more difficult to predict the future of a randomly chosen cluster (again, for $X_i > X_*$) when the model is solved on a network. In the next section, we attempt this in order to address the following question - can selective networking be used to change the fate of given cluster?

VI. NETWORKING STRATEGIES

Since the central issue in this model is the survival of clusters against the competition, the most interesting use of networks would be in the finding of a smart networking strategy which is able to change the fate of a cluster. This can be accomplished by making a marginal survivor into a strong one, and more dramatically, by seeing if a *dying cluster can be revived to life*.

We systematically investigate the effect of adding a finite number of nonlocal connections to a chosen central cluster. In Sec. III, it was shown that the growth or decay of a cluster is solely dictated by its relative rate of change versus the cumulative rate of change of its neighbours. The key to better survival should therefore lie in choosing appropriately decaying non-local clusters to network with; we will show in the following, that the fate of a cluster can indeed be modified if this is done.

Given the complexity of the problem, we cannot categorize the clusters based on their rates of growth; the most viable option is a categorization of the clusters based on their initial masses. We divide the nonlocal connections into two classes: viz., class A comprises eventual non-survivors ($X < X_*$), while class B comprises would-be survivors ($X > X_*$). In the next subsection, we look at the outcome of networking with members of class A.

A. Nonlocal connections with eventual non-survivors ($X_i < X_*$)

Recall that non-survivors ($< X_*$) die very early during Stage I. In connecting such clusters to a given cluster with $X > X_*$, we can be sure that they will never be able to compete with it, much less kill it. Could such ‘harmless’ clusters, nevertheless be useful in enhancing the survival chances of clusters with $X > X_*$?

Let us consider a central cluster that will be an eventual survivor on its own in a given




FIG. 9: (Color online) (a) The central cluster here is a survivor in its original configuration. (b) The asymptotic mass of the central cluster diverges with increasing connectivity with eventual non-survivors.

neighbourhood; one such case is shown in Fig. 9. We now let the central cluster network with eventual non-survivors from all over the lattice, and record its growth as a function of the number of the clusters it networks with; the results are shown in Fig. 9 (b). It is evident that there is a marked increase in the mass of the central cluster as more small clusters are connected to it; since none of these will survive past Stage I, their cumulative growth rate will be negative, leading to an increase in the rate of growth of the chosen cluster. This makes it a better (more massive) survivor asymptotically. As the number of nonsurviving contacts is increased, the mass of the chosen cluster increases (Fig. 9).

We now use this observation to try to change the fate of a dying cluster, and turn it into a survivor. Although the nonlinear, many-body nature of the problem makes it rather difficult to construct such cases, we have been able to network an erstwhile dying cluster with suitable nonlocal contacts, and returning it to life. Figure 10 shows our results: the central cluster would eventually have died in its original environment, but on adding 100 small clusters (whose $< X_*$) it manages to live. Further additions, e.g. 200 or 364 clusters evidently make it a better survivor (Fig. 10 (b)).




FIG. 10: (Color online) (a) The central cluster dies in its original configuration without extra connections. On addition of more and more clusters it becomes a survivor. (b) The phase transition of the central cluster from a nonsurvivor to a survivor is clearly shown.

B. Networking with would-be survivors $X_i > X_{\star}$

Here we are dealing with a far more complex problem than the case discussed in earlier subsection. When we choose clusters with masses larger than X_{\star} , their lifespan will exceed Stage I; thus during Stage I they will have a positive growth rate. Depending on their individual environments, they may survive through Stage II - thereby having a positive growth - or die as a result of a negative growth rate. Networking with such clusters is therefore a rather delicate matter.




FIG. 11: (a) In this configuration, the central cluster is networked with clusters that are (a) less massive, and (b) more massive than itself, respectively.

First, we consider nonlocal connections with would-be survivors ($X_i > X_*$) which are less massive than our chosen cluster ($X_i < X_{cm}$). Such would-be survivors will live beyond Stage I and will grow with a positive rate (see Fig. 1); from a mean field perspective, we would therefore expect to see a decrease in the chances of survival of the chosen cluster as it networks with more and more would-be survivors.

Figure 12 shows a sample result: increasing the number of nonlocal connections with would-be survivors leads to a fall in the absolute value of X_{cm} as well its rate of growth X'_{cm} . Beyond a certain number of networked contacts, the central cluster begins to decay, and eventually dies. This crossover from life to death happens when the cumulative rate of growth of the neighbours $\Sigma X'_{i,j}$ is larger than that of the central cluster X'_{cm} . Unfortunately the intricate many-body nature of this problem precludes a prediction of when such crossovers might occur in general. We will have more to say on this issue in the next section.

Finally, in the case where a given cluster networks with larger would-be survivors ($X_i > X_*$ and $X_i > X_{cm}$), one would expect a speedier death. Using the example of Fig. 13, we notice that in its original configuration the central cluster (with four neighbours, ($n = 4$)) is a survivor. As we increase the number of networked connections, its growth gets stunted; there is a substantial fall for 3 extra links ($n = 7$ in Fig. 13). Adding one more link ($n = 8$) does the final damage; the central cluster dies. The rates shown in Fig. 13 (b) and (c)




FIG. 12: (Color online) The central cluster of a configuration given in Fig. 11(a) is networked with clusters whose $X_n > X_*$. In (a) the growth of X_{cm} with an increasing number of connections is plotted - $n = 4$ corresponds to a regular lattice. There is a crossover seen when the number of connections increases from $n = 9$ to $n = 10$; for larger n values, the central site dies. This observation is supported by the rates of growth of X_{cm} and its neighbours X_n (b) when $n = 9$ and (c) when $n = 10$.

for $n = 7$ connections and $n = 8$ connections vividly capture the competition for survival, leading to life in one case and death in the other.

As expected, we observe that fewer connections (here, $n = 8$) are needed, compared to the earlier case with smaller would-be survivors ($n = 10$), to kill the chosen cluster. In closing, we should of course emphasise that these results are sample results chosen to illustrate a qualitative point: the choice of different nonlocal sites to network with would certainly modify our n values.




FIG. 13: (Color online) The central cluster of a configuration given in Fig.11(b) is connected with nonlocal clusters with $X_n > X_{cm}$. In (a) the growth of X_{cm} with an increasing number of connections is plotted - $n = 4$ corresponds to a regular lattice. There is a crossover seen when the number of connections increases from $n = 7$ to $n = 8$; for larger n values, the central site dies. This observation is supported by the rates of growth of X_{cm} and its neighbours X_n (b) when $n = 7$ and (c) when $n = 8$.

VII. UNIVERSALITY IN THE DISTRIBUTION OF SURVIVORS - A CLUE TO SURVIVAL AGAINST THE ODDS?

The results of the previous section indicate that if a cluster wants to change its fate, it should always network with non-survivors; however, it seems intuitively reasonable to suppose that it might also be able to do this if it networks with would-be survivors, provided that they are much smaller than itself. Finding such cases is possible ex post facto; their prediction is next to impossible given the complexity of the problem.

An interesting case to consider might be one where, against the odds, the most massive cluster in a given neighbourhood dies marginally, because of the unexpected growth of its

erstwhile smaller neighbours. If, now, it is able to network with potential survivors much smaller than itself (who, as a result of their own dynamics, would eventually perish in their own neighbourhoods), this would be a promising way of selectively networking itself to life. Similar issues are associated with survivors who survive against the odds; they could be killed off by suitable networking.

A systematic way of exploring this question is to look at the statistics associated with some of these rare events. We look first at the four immediate neighbours of a given cluster, and consider their pairwise interactions with it. Clearly, had such a pair been isolated, the larger cluster would have won [5]. However, many-body interactions in the lattice mean that this is not always true. We therefore ask the question: *what is the proportion of cases where the larger cluster wins?*

Each survivor has four neighbours; we first calculate the probability distribution of the initial mass differences in a pairwise fashion between a survivor and each of its neighbours. The initial mass difference are given by $\delta X_i = X_{cm} - X_i$ ($i = 1, 2, 3, 4$) corresponding to the four neighbours - right, left, bottom and top - of a survivor. The distribution of δX_i for all the survivors is shown in Fig. 14. Here, a positive (negative) δX_i means that the survivor is larger (smaller) than its neighbour; i.e. the fraction of cases where a smaller (larger) survivor wins over a larger (smaller) neighbour lies to the left (right) of zero. All four distributions corresponding to four neighbouring pairs overlap, suggesting the presence of an astonishing universality in the problem.

We also obtain the cumulative mass difference between survivors and all of their four neighbours viz. $4X_{cm} - \sum_{i=1}^4 X_i = \sum_{i=1}^4 \delta_i$ (see Fig. 15). The distributions of $\sum_{i=1}^4 \delta_i$ are plotted in Fig. 15 for different values of μ . For a positive cumulative mass difference we know that the survivor has a larger initial mass than the sum of its neighbours, matching our intuition based on the mean-field regime. The negative side of the distribution is more interesting, comprising survivors whose initial masses are smaller than the sum of the initial masses of their immediate neighbours.

We recapitulate these results before drawing tentative conclusions. Recall that the initial masses are characterized by an exponential distribution with mean $1/\mu$, where $\mu = -\log(S_1)/X_*$ [5]. Notice that both the survivor-neighbour pair distribution, (Fig. 14), and the survivor- all neighbours cumulative distribution (Fig. 15) get broader with increasing μ . In each case, the fraction of area under the negative side of the survivor pair-




FIG. 14: The plots show the distribution of pairwise mass differences between survivors and their four neighbours. They are obtained for exponential distributions of initial masses with different mean values $1/\mu$, where $\mu = -\text{Log}(S_1)/X_*$. The plots are for (a) $1/\mu = 3.92$ ($S_1 = 0.6$) (b) 5.607 ($S_1 = 0.7$) (c) 8.963 ($S_1 = 0.8$) and (d) 18.982 ($S_1 = 0.9$). The system size is 400×400 .

distribution gives an estimate of survivors against the odds – an example of some of the rare events alluded to at the beginning of this section.

Figure 16 shows this fraction, both in terms of individual survivor-pair distribution and cumulative distribution, as a function of the μ of the initial mass distribution, for different system sizes. The interpretation of the depicted results is as follows: increasing μ increases the number of potential survivors S_1 beyond Stage 1. This implies that there are more survivors, both with, and *against* the odds, hence an increase in the fraction plotted on the abscissa of Figure 16 for both pairwise and cumulative distributions. Focusing on the largest system size for concreteness, where both probability distributions have saturated, we

note the striking and intriguing universality evinced; as the system size increases, there is a universal curve for both the pairwise and the cumulative distribution of masses, indicating that the probability of finding a survivor against either pairwise or cumulative odds, is a *regular and universal function* of μ . These distributions are independent of individual masses, relying as they do on only mass differences. Finally, it is easier to be a winner against the odds if only two sites are considered, but more difficult if one is comparing a site with the total mass of its four neighbours: the *distribution of mass* amongst the four will determine whether or not the given site survives.

A major conclusion to be drawn from Figure 16 is the following: there are clusters which *die* against the odds (clusters which are more massive than the survivor in either a pairwise or cumulative sense). Recalling again that the distributions plotted are independent of mass, it should therefore be possible to pick dying clusters whose mass was greater than threshold and to network them with their peers so that they can rightfully survive, displacing erstwhile neighbours who had survived against the odds.




FIG. 15: (Color online) The plots show the distribution of mass differences between survivors and all of their four neighbours. They are obtained for exponential distributions of initial clusters with different mean values $1/\mu$, where $\mu = -\text{Log}(S_1)/X_*$. The plot in ‘red’ represents the case for $1/\mu = 3.92$ ($S_1 = 0.6$), ‘green’ represents $1/\mu = 5.607$ ($S_1 = 0.7$), ‘blue’ represents $1/\mu = 8.963$ ($S_1 = 0.8$), and ‘pink’ represents $1/\mu = 18.982$ ($S_1 = 0.9$). The system size is 400×400 .




FIG. 16: The plots show the probability of finding a successful central cluster with a larger neighbouring cluster, to start with. The probability increases with increase in S_1 (refer to Figs. 14 and 15). While the solid line show the combined cumulative probabilities (from Fig. 15), the rest of the lines represent the individual cumulative probabilities (from Fig. 14).

VIII. DISCUSSION AND CONCLUSION

In this paper we have investigated the cluster growth model of [5] on a square lattice. Our main emphasis has been on stochastic networking strategies: long-range connections are introduced with probability $0 < p < 1$ to existing lattice sites.

We find first of all that the qualitative features of the networked system remain the same as that of the regular case, in that the presence of two well-separated dynamical stages is retained; the glassy dynamics and metastable states of the earlier work [5] persist. However, the number of survivors decreases as expected with increasing p , in accordance with a mean

field perspective. Also, the system is very sensitive to the smallest changes in the initial mass configuration; the divergence is, interestingly, exponential irrespective of the magnitudes of the masses swapped. To quantify this, we have used the concept of damage spreading.

The central result of this paper is the use of smart networking strategies to modify the fate of an arbitrary cluster. We find that it is safest to network with non-survivors; their decay and eventual death lead to the transformation of the destiny of a given site, from death to survival, or from survival to stronger survival. Networking with peers or with those of larger initial mass in general leads to weakening, and an almost inevitable death.

However, the above is not immutable: the probability distributions in the last section of the paper indicate an interesting universality of survival ‘against the odds’. While in this paper, we remark on the astonishing universality shown in these and sketch out scenarios where networking strategies with peers can enhance survival, we leave their quantitative details for future work.

-
- [1] B. Blasius, A. Huppert, and L. Stone, *Nature* **399**, 354-359 (1999).
 - [2] K. Takeuchi, *Global dynamical properties of Lotka-Volterra Systems*, World Scientific, Singapore (1996).
 - [3] C. Darwin, *The origin of species by means of natural selection, or the preservation of favoured races in the struggle for life*, John Murray. 6th ed., London (1876).
 - [4] A. Mehta, A. S. Majumdar and J. M. Luck, pp. 199-204 in ‘Econophysics of Wealth Distributions’ edited by A. Chatterjee et al, Springer-Verlag Italia (2005).
 - [5] J. M. Luck, and A. Mehta, *Eur. Phys. J. B* **44**, 79 (2005).
 - [6] R. Albert and A.-L. Barabasi, *Rev. Mod. Phys.* **74**, 47 (2002).
 - [7] M. Newman A.-L. Barabasi, and D. J. Watts, *The structure and dynamics of networks*, Princeton Univ. Press, New Jersey (2006), and the references therein.
 - [8] D. J. Watts, and S. H. Strogatz, *Nature* **393**, 440 (1998).
 - [9] A. S. Majumdar, *Phys. Rev. Lett.* **90**, 031303 (2003); A. S. Majumdar, A. Mehta and J. M. Luck, *Phys. Lett. B* **607**, 219 (2005).
 - [10] S. Boccaletta, V. Latora, Y. Moreno, M. Chavez, and D.-U. Hwang, *Phys. Rep.* **424**, 175 (2006).

- [11] for instance, see Z. Jabeen and S. Sinha, Phys. Rev. E **78**, 066120 (2008).
- [12] S. A. Kauffman, J. Theor. Biol. **22**, 437 (1969).
- [13] B. Derrida and Y. Pomeau, Europhys. Lett. **1**, 45 (1986).
- [14] H. E. Stanley, D. Stauffer, J. Kertesz, and H. J. Herrmann, Phys. Rev. Lett. **59**, 2326 (1987).