

q -Deformed diamagnetism Landau problem embedded in D -dimensions

F.A. Brito and A.A. Marinho

Departamento de Física, Universidade Federal de Campina Grande,

58109-970 Campina Grande, Paraiba, Brazil

(Dated: February 7, 2020)

Abstract

We address the issue of generalizing the thermodynamic quantities via q -deformation, i.e., via the q -algebra that describes q -bosons and q -fermions. In this study with the application of q -deformation to the Landau diamagnetism problem in two dimensions, embedded in a D -dimensional space, we will attempt to get a better understanding of the q -deformation. We obtain new results for q -deformed internal energy, number of particles, magnetization and magnetic susceptibility, which recover the values already known in the literature in the limit $q \rightarrow 1$.

I. INTRODUCTION

The study of quantum groups and quantum algebras has attracted a great interest in recent years and stimulated intense research in several areas of physics [1], taking into account a wide spectrum of applications, since cosmic strings and black holes to the fractional quantum Hall effect and high-Tc superconductors [2] and theories of rational field, non-commutative geometry, the quantum theory of super-algebras and so on [3]. There is no satisfactory universally recognized definition of a quantum group.

The concept of quantum groups was motivated by problems from a large number of physical situations. The so-called q -deformed algebra [4] has been the object of interest in literature both in physics and mathematics over the past years. A great effort has been devoted to their understanding and development [2]. One of its main ingredients is a measure of deformation q , introduced in the commutation relations that define the Lie algebra of the system with the condition that the original Lie algebra, not deformed, is produced at the limit $q \rightarrow 1$.

From the seminal work of Biedenharn [5] and Macfarlane [6], it is clear that the q -calculus initially introduced at the beginning of the last century by Jackson [7] based on the study of the hypergeometric function [8], plays a central role in the quantum group representations with a deep physical significance and not merely a mathematical exercise. Indeed, it was shown that the q -deformed oscillators using the Jackson derivative (JD) or the so-called q -derivative operators [9] define a generalized q -deformed dynamic of the q -commutative phase space. Thus, it appeared recently a great interest in investigating the q -deformed thermodynamical systems [10].

One possible mechanism capable of generating a deformed version of classical statistical mechanics is to replace, the Gibbs-Boltzmann distribution, the standard exponential with its deformed version, accordingly postulate a deformed entropy involving a generalized thermodynamic theory. Thus, some generalizations of statistical mechanics were proposed [11, 12, 13, 14]. It has also been demonstrated in [15] that a natural realization of thermodynamics of q -deformed bosons and fermions can be found in the formalism of q -calculus. In fact, it was shown that the q -integration is related to the free energy of systems of spin [16] — see also [17].

In a more specific case, we focus our attention to studying of the Landau diamagnetism problem. The Landau diamagnetism problem continues to raise issues that have strong relevance today. These issues are related to the inherent quantum nature of the problem. It can be used as a phenomenon to illustrate the essential role of quantum mechanics in the surface and perimeter

corrections, the dissipation of statistical mechanics of non-equilibrium, and so on.

The paper is organized as follows. In Sec. II we present the q -deformed algebra and Jackson derivative. In Sec. III we develop the q -deformed Landau diamagnetism. In Sec. IV we make our final comments.

II. q -DEFORMED QUANTUM ALGEBRA

The q -deformed algebraic symmetry of the quantum oscillator is defined by the q -deformed Heisenberg algebra in terms of creation and annihilation operators c^\dagger, c and the quantum number N as [3, 18, 19]

$$[c, c]_k = [c^\dagger, c^\dagger]_k = 0 \quad , \quad cc^\dagger - kqc^\dagger c = q^{-N}, \quad (1)$$

$$[N, c^\dagger] = c^\dagger \quad , \quad [N, c] = -c, \quad (2)$$

where the deformation parameter q is real, being the constant $k = 1$ for q -bosons (with commutators) and $k = -1$ for q -fermions (with anticommutators). The basic q -deformed quantum number is defined as

$$[x] = \frac{q^x - q^{-x}}{q - q^{-1}}. \quad (3)$$

In addition, the operators obey the relations

$$[x, y]_k = xy - kyx, \quad cc^\dagger = [1 + kN]. \quad (4)$$

Note that for $q \neq 1$ the q -deformed quantum number $[x]$ does not meet additivity

$$[x + y] = [x] + [y] + (q - 1)[x][y], \quad (5)$$

whereas in the limit of $q \rightarrow 1$ the basic q -deformed quantum number $[x]$ is reduced to an ordinary number x .

The q -Fock space spanned by the orthonormalized eigenstates $|n\rangle$ is constructed according to

$$|n\rangle = \frac{(a^\dagger)^n}{\sqrt{[n]!}}|0\rangle \quad , \quad a|0\rangle = 0, \quad (6)$$

where the factorial of the basic q -deformed quantum number $[n]$ is defined as

$$[n]! = [n][n-1]\dots[1]. \quad (7)$$

The actions of c , c^\dagger and N on the states $|n\rangle$ in the q -Fock space are known to be

$$c^\dagger|n\rangle = [n+1]^{1/2}|n+1\rangle, \quad (8)$$

$$c|n\rangle = [n]^{1/2}|n-1\rangle, \quad (9)$$

$$N|n\rangle = n|n\rangle. \quad (10)$$

One may transform the q -Fock space into the configuration space (Bargmann holomorphic representation) [9, 20] as in the following

$$c^\dagger = x, \quad c = D_x^{(q)}, \quad (11)$$

where $D_x^{(q)}$ is the Jackson derivative (JD) [7]

$$D_x^{(q)}f(x) = \frac{f(qx) - f(q^{-1}x)}{x(q - q^{-1})}. \quad (12)$$

Note that it becomes an ordinary derivative as $q = 1$. Therefore, JD naturally occurs in quantum deformed structures. It turns out to be a crucial role in q -generalization of thermodynamical relations [15].

III. q -DEFORMED LANDAU DIAMAGNETISM EMBEDDED IN D -DIMENSIONS

To explain the phenomenon of diamagnetism, we have to take into account the interaction between the external magnetic field and the orbital motion of electrons. Disregarding the spin, the Hamiltonian of a particle of mass m and charge e in the presence of a magnetic field \mathbf{H} , is given by the expression

$$H = \frac{1}{2m} \left(\mathbf{p} - \frac{e}{c} \mathbf{A} \right)^2, \quad (13)$$

where \mathbf{A} is the vector potential associated with the magnetic field \mathbf{H} and c is the speed of light. Let us start to formalize the statistical mechanical problem by using the grand partition function in the form

$$\begin{aligned} \ln \Xi = & \frac{2eHL^2}{hc} \sum_{n=0}^{\infty} \frac{L}{2\pi} \int_{-\infty}^{\infty} dk_z \ln \left\{ 1 + z \exp \left[-\frac{\beta \hbar^2 k_z^2}{2m} \right. \right. \\ & \left. \left. - \frac{\beta \hbar e H}{mc} \left(n + \frac{1}{2} \right) \right] \right\}, \end{aligned} \quad (14)$$

where $k_z^2 = k_1^2 + k_2^2 + \dots + k_{(D-2)}^2$. Here, we have generalized the Landau diamagnetism problem, usually embedded in three dimensions, to embedded it in D -dimensions, i.e., we have now $D - 2$ extra-dimensions. Below, we further generalize this result by also applying the Jackson derivative.

However, our study is focused on the analysis of diamagnetism in the limit of high temperatures ($z \ll 1$) where we apply the q -deformed algebra. Thus, the partition function is written as follows

$$\ln \Xi = \frac{2eHL^D}{\lambda^{(D-2)}(2\pi)^{(D)}hc} z k \sum_{n=0}^{\infty} y_n, \quad (15)$$

where $z = \exp(\beta\mu)$ is the fugacity, $y_n = \exp\left[\left(\frac{-\beta\hbar eH}{mc}\right)\left(n + \frac{1}{2}\right)\right]$, $\lambda = \frac{h}{(2\pi mT)^{1/2}}$ is the thermal wavelength. For the sake of simplicity, in our calculations we simply use $y_n = \exp(-\beta\epsilon_n)$, where $\epsilon_n = \frac{\hbar eH}{mc}\left(n + \frac{1}{2}\right)$. With these substitutions we can rewrite the equation (15) as

$$\ln \Xi = z k C \sum_{n=0}^{\infty} y_n, \quad C = \frac{2L^{(D)}eH}{\lambda^{(D-2)}(2\pi)^{(D)}hc}. \quad (16)$$

Now we are ready to applying the q -deformed algebra via the introduction of JD, which is a key factor of q -deformed thermodynamics. This is achieved through the modification of the ordinary thermodynamical derivatives as follows

$$\frac{\partial}{\partial z} \rightarrow D_z^{(q)}, \quad (17)$$

where $D_z^{(q)}$ is the JD defined as

$$D_z^{(q)} = \frac{q-1}{\ln q} \partial_z^{(q)}. \quad (18)$$

A. q -Deformed thermodynamical quantities

As one knows the internal energy is defined in the form

$$U = -\frac{\partial}{\partial \beta} \ln \Xi. \quad (19)$$

Thus, to implement the JD is necessary to change the basis as follows

$$-\frac{\partial}{\partial \beta} = -\left(\frac{\partial y_n}{\partial \beta} \frac{\partial}{\partial y_n}\right), \quad (20)$$

such that

$$U = -\frac{\partial y_n}{\partial \beta} D_{y_n}^{(q)} \ln \Xi, \quad (21)$$

$$U = -z k C \sum_{n=0}^{\infty} \frac{\partial y_n}{\partial \beta} D_{y_n}^{(q)} y_n. \quad (22)$$

So using the JD definition in Eq. (18) we find

$$U = -z k C \frac{q-1}{\ln q} \frac{\partial}{\partial \beta} \sum_{n=0}^{\infty} y_n. \quad (23)$$

Now performing the summation we find the following

$$\sum_{n=0}^{\infty} y_n = \frac{1}{2 \sinh(\beta \mu_B H)}, \quad (24)$$

where $\mu_B = \frac{\hbar e}{2mc}$. Therefore the internal energy is written as

$$U = zkC \frac{q-1}{\ln q} \left(\frac{\mu_B \cosh(\beta \mu H)}{2 \sinh^2(\beta \mu H)} \right). \quad (25)$$

To calculate the number of particles we have

$$N = z \frac{\partial}{\partial z} \ln \Xi. \quad (26)$$

Now applying in Eq. (26) the JD we just get

$$N = zkC \frac{q-1}{\ln q} \frac{1}{2 \sinh(\beta \mu_B H)}. \quad (27)$$

B. The specific heat

To determine the specific heat, we first make use of the Jacobian

$$C_V = \frac{1}{N} \left(\frac{\partial U}{\partial T} \right)_{V,N} = \frac{-\kappa_B \beta^2}{N} \left(\frac{\partial U}{\partial \beta} \right)_{V,N}, \quad (28)$$

$$\begin{aligned} \left(\frac{\partial U}{\partial \beta} \right)_N &= \frac{\partial(U, N)}{\partial(\beta, N)} = \frac{\partial(U, N)}{\partial(\beta, z)} \frac{\partial(\beta, z)}{\partial(\beta, N)} \\ &= \left(\frac{\partial U}{\partial \beta} \right)_z - \left(\frac{\partial U}{\partial z} \right)_\beta \left(\frac{\partial N}{\partial z} \right)_\beta. \end{aligned} \quad (29)$$

Using the derivatives above, we find

$$\left(\frac{\partial U}{\partial \beta} \right)_N = \frac{-zkC}{2} \frac{q-1}{\ln q} \frac{\mu_B^2 H}{\sinh^3(\beta \mu_B H)}. \quad (30)$$

Applying the equation (30) in the equation (28) and substituting N obtained in the equation (27), we arrive at our specific heat

$$C_V = \kappa_B \left(\frac{\beta \mu_B H}{\sinh(\beta \mu_B H)} \right)^2. \quad (31)$$

Where κ_B is the Boltzmann constant. This coincides with the usual result.

C. q -Deformed magnetization and susceptibility

For determining the magnetization we carried out the thermodynamical derivative

$$M = -\frac{\partial \phi}{\partial H}, \quad (32)$$

where the grand potential ϕ is determined as

$$\phi = -\frac{1}{\beta} \ln \Xi, \quad \phi = \frac{-zkC}{2\beta \sinh(\beta\mu_B H)}. \quad (33)$$

However, to implement the JD we have to make the change as follows

$$-\frac{\partial \phi}{\partial H} = -\left(\frac{\partial y_n}{\partial H} \frac{\partial \phi}{\partial y_n}\right). \quad (34)$$

Thus we find

$$M = -\frac{\partial y_n}{\partial H} D_{y_n}^{(q)} \phi, \quad (35)$$

and

$$M = \frac{zkC^*}{\beta} \frac{q-1}{\ln q} \frac{\partial}{\partial H} \left(2H \sum_{n=0}^{\infty} y_n\right), \quad (36)$$

where $C^* = C/2H$, such that

$$M = \frac{zkC^*}{\beta} \frac{q-1}{\ln q} \frac{1}{\sinh(\beta\mu_B H)} \left(1 - \frac{\beta\mu_B H \cosh(\beta\mu_B H)}{\sinh(\beta\mu_B H)}\right). \quad (37)$$

We can also eliminate the chemical potential through the number of particles N and rewrite the magnetization as

$$M = \frac{N}{\beta H} \left(1 - \frac{\beta\mu_B H \cosh(\beta\mu_B H)}{\sinh(\beta\mu_B H)}\right), \quad (38)$$

or in terms of the Langevin function given by

$$\mathcal{L}(\beta\mu_B H) = \coth(\beta\mu_B H) - \frac{1}{\beta\mu_B H}, \quad (39)$$

we get to

$$M = -N\mu_B \mathcal{L}(\beta\mu_B H). \quad (40)$$

Let us now make the analysis in the weak field limit $\beta\mu_B H \ll 1$ below:

i) The use of this limit into the equation (40) give us

$$M = -\frac{N\mu_B^2 H}{3\kappa_B T}. \quad (41)$$

Furthermore the susceptibility is

$$\chi = \frac{\partial M}{\partial H}, \quad \chi = -\frac{N\mu_B^2}{3\kappa_B T}. \quad (42)$$

ii) The use of this limit for $q = 1$, gives us, the usual results:

$$M = -\frac{zkC\mu_B^3 H^2}{6(\kappa_B T)^2} \left(\left(\frac{\kappa_B T}{\mu_B H} \right)^2 - \frac{1}{6} \right), \quad (43)$$

$$\chi = -\frac{zkC\mu_B^3 H}{6(\kappa_B T)^2} \left(\left(\frac{\kappa_B T}{\mu_B H} \right)^2 - \frac{1}{6} \right). \quad (44)$$




Figure 1: q -deformed internal energy for $q = 0.1$ (blue), $q = 0.5$ (green), $q = 0.99999999$ (red)




Figure 2: q -deformed magnetization as a function of magnetic field H for $q = 0.1$ (blue), $q = 0.5$ (green), $q = 0.99999999$ (red)

IV. CONCLUSIONS

We apply the q -deformed quantum algebra in the Landau diamagnetism problem embedded in D -dimensions, in the limit of high temperatures. We obtain q -deformed values $q \neq 1$ for internal




Figure 3: q -deformed susceptibility as a function of magnetic field H for $q = 0.1$ (blue), $q = 0.5$ (green), $q = 0.99999999$ (red)

energy, number of particles, magnetization, susceptibility and specific heat. We see that the q -deformed thermodynamical quantities change, except the specific ones, i.e., those quantities per particles, such as specific heat. That means that all effect of the q -deformation is completely stored into the number of particles N . It seems to be related to *impurities* in a sample of a diamagnetic material. As expected, in the limit $q = 1$ we see that the results are identical to those in the literature [21, 22]. The thermodynamical quantities also depend on the number of D -dimensions since it is present in the constant C . This fact may find some applications in extra-dimensional physics such as modern cosmology, particle physics and string theory.

Acknowledgments

We would like to thank CNPq, CAPES, and PNP/PROCAD - CAPES, for partial financial support.

-
- [1] L.C. Biedenharn and M.A. Lohe, *Quantum Group Symmetry and Q-Tensor Algebras*, (World Scientific, Singapore, 1995).
- [2] F. Wilczek, *Fractional Statistics and Anyon Superconductivity*, (World Scientific, Singapore, 1990) and references therein.
- [3] M. Chaichian, R. Gonzales Felipe, C. Montonen, *J. Phys. A: Math. Gen.* **26**, 4017 (1993).
- [4] D. Galetti, *Braz. J. Phys.* **33**, 1 (2003).
- [5] L. Biedenharn, *J. Phys. A: Math. Gen.* **22**, L873 (1989).
- [6] A. Macfarlane, *J. Phys. A: Math. Gen.* **22**, 4581 (1989).
- [7] F. Jackson, *Mess. Math.* **38**, 57 (1909).
- [8] H. Exton, *q-Hypergeometric Functions and Applications*, (John Wiley and Sons, New York, 1983).
- [9] E.G. Floratos, *J. Phys. Math.* **24**, 4739 (1991).
- [10] A. Lavagno, A.M. Scarfone and N.P. Swamy, *J. Phys. A: Math. Theor.* **40**, (2007) 8635-8654.
- [11] C. Tsallis, *J. Stat. Phys.* **52**, 479 (1988); E.M. Curado and C. Tsallis, *J. Phys. A: Math. Gen.* **24**, L69 (1991); C. Tsallis, R.S. Mendes and A.R. Plastino, *Physica A* **261**, 534 (1998).
- [12] S. Abe, *Phys. Lett. A* **224**, 326 (1997).
- [13] G. Kaniadakis, *Phys. Rev. E* **66**, 056125 (2002).
- [14] G. Kaniadakis, M. Lissia and A.M. Scarfone, *Physica A* **340**, 41 (2004); G. Kaniadakis, M. Lissia and A.M. Scarfone, *Phys. Rev. E* **71**, 046128 (2005).
- [15] A. Lavagno and N.P. Swamy, *Phys. Rev. E* **61**, 1218 (2000); A. Lavagno and N.P. Swamy, *Phys. Rev. E* **65**, 036101 (2002).
- [16] A. Erzan, *Phys. Lett. A* **225**, 235 (1997).
- [17] G. Bimonte, C. Esposito, G. Marmo and C. Stornaiolo, *Phys. Lett. A* **318**, 313 (2003).
- [18] Y.J. Ng, *J. Phys. A* **23**, 1023 (1990).
- [19] Sakurai, J.J., *Modern Quantum Mechanics*, (Addison Wesley Reading, Mass., 1994).
- [20] R.J. Finkelstein, *Int. J. Mod. Phys. A* **13**, 1795 (1998).
- [21] Patthria, *Statistical Mechanics*, Pergamon press, Oxford (1972)
- [22] K.Huang, *Statistical Mechanics*, John Wiley & Sons, (1987).