## A SUP-HODGE BOUND FOR EXPONENTIAL SUMS

CHUNLEI LIU

ABSTRACT. The C-function of T-adic exponential sums is studeid. An explicit arithmetic bound is established for the Newton polygon of the C-function. This polygon lies above the Hodge polygon. It gives a sup-Hodge bound of the C-function of p-power order exponential sums.

### 1. INTRODUCTION

Let p be a prime number,  $\mathbb{F}_p = \mathbb{Z}/(p)$ ,  $\overline{\mathbb{F}}_p$  a fixed algebraic closure of  $\mathbb{F}_p$ , and  $\mathbb{F}_{p^k}$  the subfield of  $\overline{\mathbb{F}}_p$  with  $p^k$  elements.

Let q > 1 be a power of p, W the ring scheme of Witt vectors,  $\mathbb{Z}_q = W(\mathbb{F}_q)$ ,  $\mathbb{Q}_q$  the fraction field of  $\mathbb{Z}_q$ ,  $\overline{\mathbb{Q}}_p = \lim_{k \to \infty} \mathbb{Q}_{p^k}$ , and  $\mathbb{C}_p$  the *p*-adic completion of

 $\overline{\mathbb{Q}}_p.$ 

Let  $\Delta \supseteq \{0\}$  be an integral convex polytope in  $\mathbb{R}^n$ , and I the set of vertices of  $\Delta$  different from the origin. Let

$$f(x) = \sum_{u \in \Delta} (a_u x^u, 0, 0, \dots) \in W(\mathbb{F}_q[x_1^{\pm 1}, \dots, x_n^{\pm 1}]) \text{ with } \prod_{u \in I} a_u \neq 0,$$

where  $x^u = x_1^{u_1} \cdots x_n^{u_n}$  if  $u = (u_1, \cdots, u_n) \in \mathbb{Z}^n$ .

**Definition 1.1.** For  $k \in \mathbb{N}$ , the sum

$$S_f(k,T) = \sum_{x \in (\mathbb{F}_{q^k}^{\times})^n} (1+T)^{\operatorname{Tr}_{\mathbb{Z}_{q^k}/\mathbb{Z}_p}(f(x))} \in \mathbb{Z}_p[[T]]$$

is call a T-adic exponential sum. And the function

$$L_f(s,T) = \exp(\sum_{k=1}^{\infty} S_f(k,T) \frac{s^k}{k}) \in 1 + s\mathbb{Z}_p[[T]][[s]]],$$

as a power series in the single variable s with coefficients in the T-adic complete field  $\mathbb{Q}_p((T))$ , is called an L-function of T-adic exponential sums.

Let *m* be a positive integer,  $\zeta_{p^m}$  a primitive  $p^m$ -th root of unity, and  $\pi_m = \zeta_{p^m} - 1$ . Then  $S_f(k, \pi_m)$  is the exponential sum studied by Liu-Wei [LWe]. If m = 1, the exponential sum  $S_f(k, \pi_m)$  was studied by Adolphson-Sperber [AS,AS2]. And, if n = 1, the exponential sum  $S_f(k, \pi_m)$  was studied by Kumar-Helleseth-Calderbank [KHC] and Li [Li].

This research is supported by NSFC Grant No. 10671015.

**Definition 1.2.** The function

$$C_f(s,T) = C_f(s,T;\mathbb{F}_q) = \exp(\sum_{k=1}^{\infty} -(q^k-1)^{-n}S_f(k,T)\frac{s^k}{k}),$$

as a power series in the single variable s with coefficients in the T-adic complete field  $\mathbb{Q}_p((T))$ , is called a C-function of T-adic exponential sums.

We have

$$L_f(s,T) = \prod_{i=0}^{n} C_f(q^i s,T)^{(-1)^{n-i+1} \binom{n}{i}}$$

and

$$C_f(s,T) = \prod_{j=0}^{\infty} L_f(q^j s,T)^{(-1)^{n-1}\binom{n+j-1}{j}}.$$

So the C-function  $C_f(s,T)$  and the L-function  $L_f(s,T)$  determine each other. From the last identity, one sees that

$$C_f(s,T) \in 1 + s\mathbb{Z}_p[[T]][[s]].$$

Let  $C(\Delta)$  be the cone generated by  $\Delta$ , and  $M(\Delta) = M(\Delta) \cap \mathbb{Z}^n$ . There is a degree function deg on  $C(\Delta)$  which is  $\mathbb{R}_{\geq 0}$ -linear and takes the values 1 on each co-dimension 1 face not containing 0. For  $a \notin C(\Delta)$ , we define  $\deg(a) = +\infty$ .

**Definition 1.3.** A convex function on  $[0, +\infty]$  which is linear between consecutive integers with initial value 0 is called the infinite Hodge polygon of  $\triangle$  if its slopes between consecutive integers are the numbers deg(a),  $a \in M(\triangle)$ . We denote this polygon by  $H^{\infty}_{\triangle}$ .

Liu-Wan [LWa] also proved the following.

Theorem 1.4 (Hodge bound). We have

 $T - adic \ NP \ of \ C_f(s,T) \ge ord_p(q)(p-1)H^{\infty}_{\Delta},$ 

where NP is the short for Newton polygon.

Denote by  $\lceil x \rceil$  the least integer equal or greater than x, and by  $\{x\}$  the fractional part of x.

**Definition 1.5.** Let  $C \subseteq M(\triangle)$  be a finite subset. We define

 $r_C = \max_{\beta} (\#\{a \in C \mid \{\deg(pa)\}' \ge \beta\} - \#\{a \in C \mid \{\deg(a)\}' \ge \beta\}).$ 

**Definition 1.6.** Let  $a \subseteq M(\triangle)$ . We define

 $\varpi(a) = \lceil p \deg(a) \rceil - \lceil \deg(a) \rceil + r_{\{a \in M(\triangle) | \deg(a) < \deg(a)\} \cup \{a\}} - r_{\{a \in M(\triangle) | \deg(u) < \deg(a)\}}.$ 

**Definition 1.7.** The arithmetic polygon  $p_{\Delta}$  of  $\Delta$  is a convex function on  $[0, +\infty]$  which is linear between consecutive integers with initial value 0, and whose slopes between consecutive integers are the numbers  $\varpi_{\Delta}(a)$ ,  $a \in M(\Delta)$ .

One can prove the following.

Theorem 1.8. We have then

$$p_{\Delta} \ge (p-1)H_{\Delta}^{\infty}.$$

Moreover, they coincide at the point  $n!Vol(\triangle)$ .

Let D be the least positive integer such that  $\deg(M(\triangle)) \subseteq \frac{1}{D}\mathbb{Z}$ . The main result of this paper is the following.

### **Theorem 1.9.** If p > 3D, then

$$T - adic \ NP \ of \ C_f(s, T) \ge \operatorname{ord}_p(q) p_{\triangle}.$$

From the above theorem we shall deduce the following.

**Theorem 1.10.** If p > 3D, then, for  $t \in \mathbb{C}_p$  with  $0 \neq |t|_p < 1$ , we have  $t - adic \ NP \ of C_f(s, t) \ge \operatorname{ord}_p(q)p_{\triangle}$ .

# 2. The T-adic Dwork Theory

In this section we review the *T*-adic analogue of Dwork theory on exponential sums.

Let

$$E(t) = \exp(\sum_{i=0}^{\infty} \frac{t^{p^i}}{p^i}) = \sum_{i=0}^{+\infty} \lambda_i t^i \in 1 + t\mathbb{Z}_p[[t]]$$

be the *p*-adic Artin-Hasse exponential series. Define a new *T*-adic uniformizer  $\pi$  of  $\mathbb{Q}_p((T))$  by the formula  $E(\pi) = 1 + T$ . Let  $\pi^{1/D}$  be a fixed *D*-th root of  $\pi$ . Let

$$L = \{ \sum_{u \in M(\Delta)} c_u \pi^{\deg(u)} x^u : c_u \in \mathbb{Z}_q[[\pi^{1/D}]] \}.$$

Let  $a \mapsto \hat{a}$  be the Teichmüller lifting. One can show that the series

$$E_f(x) := \prod_{a_u \neq 0} E(\pi \hat{a}_u x^u) \in L.$$

Note that the Galois group of  $\mathbb{Q}_q$  over  $\mathbb{Q}_p$  can act on L but keeping  $\pi^{1/D}$  as well as the variable x fixed. Let  $\sigma$  be the Frobenius element in the Galois group such that  $\sigma(\zeta) = \zeta^p$  if  $\zeta$  is a (q-1)-th root of unity. Let  $\Psi_p$  be the operator on L defined by the formula

$$\Psi_p(\sum_{i\in M(\triangle)}c_ix^i)=\sum_{i\in M(\triangle)}c_{pi}x^i.$$

Then  $\Psi := \sigma^{-1} \circ \Psi_p \circ E_f$  acts on the *T*-adic Banach module

$$B = \{ \sum_{u \in M(\triangle)} c_i \pi^{\deg(u)} x^u \in L, \text{ ord}_T(c_u) \to +\infty \text{ if } \deg(u) \to +\infty \}.$$

We call it Dwork's *T*-adic semi-linear operator because it is semi-linear over  $\mathbb{Z}_{q}[[\pi^{\frac{1}{D}}]]$ .

Let  $b = \log_p q$ . Then the *b*-iterate  $\Psi^b$  is linear over  $\mathbb{Z}_q[[\pi^{1/D}]]$ , since

$$\Psi^b = \Psi^b_p \circ \prod_{i=0}^{b-1} E_f^{\sigma^i}(x^{p^i}).$$

One can show that  $\Psi$  is completely continuous in the sense of Serre [Se]. So  $\det(1 - \Psi^b s \mid B/\mathbb{Z}_q[[\pi^{\frac{1}{D}}]])$  and  $\det(1 - \Psi s \mid B/\mathbb{Z}_p[[\pi^{\frac{1}{D}}]])$  are well-defined. We now state the *T*-adic Dwork trace formula [LWa].

Theorem 2.1. We have

$$C_f(s,T) = \det(1 - \Psi^b s \mid B/\mathbb{Z}_q[[\pi^{\frac{1}{D}}]])$$

**Lemma 2.2.** The Newton polygon of det $(1 - \Psi^b s^b | B/\mathbb{Z}_q[[\pi^{\frac{1}{D}}]])$  coincides with that of det $(1 - \Psi s | B/\mathbb{Z}_p[[\pi^{\frac{1}{D}}]])$ .

*Proof.* Note that

$$\det(1-\Psi^b s \mid B/\mathbb{Z}_p[[\pi^{\frac{1}{D}}]]) = \operatorname{Norm}(\det(1-\Psi^b s \mid B/\mathbb{Z}_q[[\pi^{\frac{1}{D}}]])),$$

where Norm is the norm map from  $\mathbb{Z}_q[[\pi^{\frac{1}{D}}]]$  to  $\mathbb{Z}_p[[\pi^{\frac{1}{D}}]]$ . The lemma now follows from the equality

$$\prod_{\zeta^{b}=1} \det(1 - \Psi \zeta s \mid B / \mathbb{Z}_{p}[[\pi^{\frac{1}{D}}]]) = \det(1 - \Psi^{b} s^{b} \mid B / \mathbb{Z}_{p}[[\pi^{\frac{1}{D}}]]).$$

Write

$$\det(1 - \Psi s \mid B / \mathbb{Z}_p[[\pi^{\frac{1}{D}}]]) = \sum_{i=0}^{+\infty} (-1)^i c_i s^i.$$

**Theorem 2.3.** The *T*-adic Newton polygon of det $(1 - \Psi^b s \mid B/\mathbb{Z}_q[[\pi^{\frac{1}{D}}]])$  is the lower convex closure of the points

$$(m, ord_T(c_{bm})), m = 0, 1, \cdots$$

*Proof.* By Lemma 2.2, the *T*-adic Newton polygon of det $(1-\Psi^b s^b \mid B/\mathbb{Z}_q[[\pi^{\frac{1}{D}}]])$  is the lower convex closure of the points

$$(i, \operatorname{ord}_T(c_i)), \ i = 0, 1, \cdots$$

It is clear that  $(i, \operatorname{ord}_T(c_i))$  is not a vertex of that polygon if  $b \nmid i$ . So that Newton polygon is the lower convex closure of the points

$$(bm, \text{ord}_T(c_{bm})), \ m = 0, 1, \cdots$$

It follows that the *T*-adic Newton polygon of det $(1 - \Psi^b s \mid B/\mathbb{Z}_q[[\pi^{\frac{1}{D}}]])$  is the lower convex closure of the points

$$(m, \text{ord}_T(c_{bm})), \ m = 0, 1, \cdots$$

4

### 3. The arithmetic bound

In this section we prove the following.

Theorem 3.1. We have

$$\operatorname{ord}_T(c_{bm}) \ge p_{\triangle}(m).$$

*Proof.* First, we choose a basis of  $B \otimes_{\mathbb{Z}_p} \mathbb{Q}_p(\pi^{1/D})$  over  $\mathbb{Q}_p(\pi^{1/D})$  as follows. Fix a normal basis  $\bar{\xi}_i$ ,  $i \in \mathbb{Z}/(b)$  of  $\mathbb{F}_q$  over  $\mathbb{F}_p$ . Let  $\xi_i$  be their Terchmüller lift of  $\bar{\xi}_i$ . The system  $\xi_i$ ,  $i \in \mathbb{Z}/(b)$  is a normal basis of  $\mathbb{Q}_q$  over  $\mathbb{Q}_p$ . Then  $\{\xi_i x^u\}_{u \in M(\Delta), 1 \leq i \leq b}$  is a basis of  $B \otimes_{\mathbb{Z}_p} \mathbb{Q}_p(\pi^{1/D})$  over  $\mathbb{Q}_p(\pi^{1/D})$ .

Secondly, we write out the matrix of  $\Psi$  on  $B \otimes_{\mathbb{Z}_p} \mathbb{Q}_p(\pi^{1/D})$  with respect to the basis  $\{\xi_i x^u\}_{u \in M(\Delta), 1 \leq i \leq b}$ . Write

$$E_f(x) = \sum_{u \in M(\triangle)} \gamma_u x^u,$$

and

$$\sigma^{-1}(\xi_j \gamma_{pu-w}) = \sum_{i=1}^b \gamma_{(u,i),(w,j)} \xi_i.$$

Then

$$\Psi(\xi_j x^w) = \sum_{u \in M(\Delta)} \sigma^{-1}(\xi_j \gamma_u) \Psi_p(x^{u+w})$$
$$= \sum_{u \in M(\Delta)} \sigma^{-1}(\xi_j \gamma_{pu-w}) x^u$$
$$= \sum_{u \in M(\Delta)} \sum_{i=1}^b \gamma_{(u,i),(w,j)} \xi_i x^u.$$

So  $(\gamma_{(u,i),(w,j)})_{u,w\in M(\triangle),1\leq i,j\leq b}$  is the matrix of  $\Psi$  on  $B\otimes_{\mathbb{Z}_p} \mathbb{Q}_p(\pi^{1/D})$  with respect to the basis  $\{\xi_i x^u\}_{u\in M(\triangle),1\leq i\leq b}$ .

Thirdly, we claim that

$$\operatorname{ord}_T(\gamma_{(u,i),(w,j)}) \ge \lceil \operatorname{deg}(pu-w) \rceil.$$

In fact, this follows from the equality

$$\sigma^{-1}(\xi_j \gamma_{pu-w}) = \sum_{i=1}^b \gamma_{(u,i),(w,j)} \xi_i,$$

and the inequality  $\operatorname{ord}_T(\gamma_u) \ge \lceil \operatorname{deg}(u) \rceil$ .

Finally, we show that

$$\operatorname{ord}_T(c_{bm}) \ge p_{\triangle}(m).$$

Note that

$$c_{bm} = \sum_{A} \det((\gamma_{(u,i),(w,j)})_{(u,i),(w,j)\in A}),$$

where A runs over all subsets of  $M(\triangle) \times \mathbb{Z}/(b)$  with cardinality bm. So it suffices to show that

$$\operatorname{ord}_T(\det(\gamma_{(i,u),(j,\omega)})_{(i,u),(j,\omega)\in A}) \ge bp_{\triangle}(m).$$

Note that

$$\det(\gamma_{(i,u),(j,\omega)})_{(i,u),(j,\omega)\in A}) = \sum_{\tau\in S_A} \sum_{a\in A} \operatorname{ord}_{\pi}(\gamma_{a,\tau(a)}),$$

where  $S_A$  is the permutation group of A. So it suffices to show that

$$\sum_{a \in A} \operatorname{ord}_{\pi}(\gamma_{a,\tau(a)}) \ge bp_{\triangle}(m), \ \tau \in S_A.$$

Since

$$\operatorname{ord}_T(\gamma_{(u,i),(w,j)}) \ge \lceil \operatorname{deg}(pu-w) \rceil,$$

the theorem follows from the following.

**Theorem 3.2.** If p > 3D, A is a subset of  $M(\triangle) \times \mathbb{Z}/(b)$  with cardinality bm, and  $\tau \in S_A$ , then

$$\sum_{a \in A} \lceil \deg(p\nu(a) - \nu(\tau(a))) \rceil \ge bp_{\triangle}(m),$$

where  $\nu(u, i) = u$ .

*Proof.* We have

$$\begin{split} &\sum_{a \in A} \lceil \deg(p\nu(a) - \nu(\tau(a))) \rceil \geq \sum_{a \in A} \lceil \deg(p\nu(a)) - \nu(\tau(a)) \rceil \\ &\geq \sum_{a \in A} \lceil \deg(p\nu(a)) \rceil - \lceil \deg(\nu(\tau(a))) \rceil + 1_{\{\deg(p\nu(a))\}' > \{\deg(\nu(\tau(a)))\}'} \\ &\geq \sum_{a \in A} \lceil \deg(p\nu(a)) \rceil - \lceil \deg(\nu(a)) \rceil + 1_{\{\deg(p\nu(a))\}' > \{\deg(\nu(\tau(a)))\}'} . \end{split}$$

Choose a set B of cardinality |A| such that  $B \cap A$  is as big as possible under the condition that, for some  $\alpha$ ,

$$\{a \in M(\triangle) \times \mathbb{Z}/(b) \mid \deg(\nu(a)) < \alpha\} \subseteq B \subsetneq \{a \in M(\triangle) \times \mathbb{Z}/(b) \mid \deg(\nu(a)) \le \alpha\}.$$

Choose a permutation  $\tau_0$  on  $B \cap A$  which agrees with  $\tau$  on  $(B \cap A) \cap \tau^{-1}(B \cap A)$ . Extend it trivially to B. We have

$$\sum_{a \in A} \lceil \deg(p\nu(a)) \rceil - \lceil \deg(\nu(a)) \rceil \ge \sum_{a \in B} (\lceil \deg(p\nu(a)) \rceil - \lceil \deg(\nu(a)) \rceil) + 2\#(A \setminus B).$$

We also have

$$\sum_{a \in A} \mathbb{1}_{\{\deg(p\nu(a))\}' > \{\deg(\nu(\tau(a)))\}'} \ge \sum_{a \in B} \mathbb{1}_{\{\deg(p\nu(a))\}' > \{\deg(\nu(\tau_0(a)))\}'} - 2\#(A \setminus B).$$

It follows that

$$\sum_{a \in A} \lceil \deg(p\nu(a)) \rceil - \lceil \deg(\nu(a)) \rceil + 1_{\{\deg(p\nu(a))\}' > \{\deg(\nu(\tau(a)))\}}$$

$$\geq \sum_{a \in B} \lceil \deg(p\nu(a)) \rceil - \lceil \deg(\nu(a)) \rceil + 1_{\{\deg(p\nu(a))\}' > \{\deg(\nu(\tau_0(a)))\}} \\ \geq \sum_{a \in B} (\lceil \deg(p\nu(a)) \rceil - \lceil \deg(\nu(a)) \rceil) + r_B,$$

where

$$r_B = \max_{\beta} (\#\{a \in B \mid \{\deg(p\nu(a))\}' \ge \beta\}) - \#\{a \in B \mid \{\deg(\nu(a))\}' \ge \beta\}).$$

Choose a set C of cardinality m such that for some  $\alpha$ ,

$$\{a \in M(\triangle) \mid \deg(\nu(a)) < \alpha\} \subseteq C \subsetneq \{a \in M(\triangle) \mid \deg(\nu(a)) \le \alpha\}.$$

Recall that

$$r_C = \max_{\beta} (\#\{a \in C \mid \{\deg(pa)\}' \ge \beta\} - \#\{a \in C \mid \{\deg(a)\}' \ge \beta\}).$$

It is easy to see that  $r_B = br_C$ , and

$$\sum_{a \in B} (\lceil \deg(p\nu(a)) \rceil - \lceil \deg(\nu(a)) \rceil) + r_B$$
$$= b \sum_{a \in C} (\lceil \deg(pa) \rceil - \lceil \deg(a) \rceil) + br_C = bp_{\triangle}(m).$$

The theorem now follows.

#### References

- [AS] A. Adolphson and S. Sperber, Newton polyhedra and the degree of the L-function associated to an exponential sum, Invent. Math. 88(1987), 555-569.
- [AS2] A. Adolphson and S. Sperber, Exponential sums and Newton polyhedra: cohomology and estimates, Ann. Math., 130 (1989), 367-406.
- [BF] R. Blache and E. Férard, Newton straitification for polynomials: the open stratum, J. Number Theory, 123(2007), 456-472.
- [KHC] P. V. Kumar, T. Helleseth and A. R. Calderbank, An upper bound for some exponential sums over Galois rings with applications, IEEE Trans. Inform. Theory 41 (1995), no.2, 456-468.
- [Li] W.-C. W. Li, Character sums over p-adic fields, J. Number Theory 74 (1999), no.2, 181-229.
- [LWa] C. Liu and D. Wan, T-adic exponential sums, Algebra & Number theory, Vol. 3, No. 5 (2009), 489-509.
- [LWe] C. Liu and D. Wei, The L-functions of Witt coverings, Math. Z., 255 (2007), 95-115.
- [Se] J-P. Serre, Endomorphismes complétement continus des espaces de Banach p-adiques, Publ. Math., IHES., 12(1962), 69-85.
- [Zh1] J. H. Zhu, p-adic variation of L functions of one variable exponential sums, I. Amer. J. Math., 125 (2003), 669-690.
- [Zh2] J. H. Zhu, Asymptotic variation of L functions of one-variable exponential sums, J. Reine Angew. Math., 572 (2004), 219–233. 1529–1550.

DEPARTMENT OF MATHEMATICS, SHANGHAI JIAO TONG UNIVERSITY, SHANGHAI 200240, P.R. CHINA.

E-mail address: clliu@sjtu.edu.cn