# On Taylor series expansion of $(1 + z)^A$ for |z| > 1

### Akhila Raman

University of California at Berkeley, CA-94720. Email: akhila.raman@berkeley.edu. Ph: 510-540-5544

#### **Abstract**

It is well known that the Taylor series expansion of  $(1+z)^A$  does not converge for |z| > 1 where A is a real number which is not equal to zero or a positive integer. A limited series expansion of this expression is obtained in this paper for |z| > 1 as a product of convergent series.

Keywords:

#### 1. Introduction

It is well known that the Taylor series expansion of  $(1 + z)^A$  is given by

$$(1+z)^A = \sum_{n=0}^{\infty} \binom{A}{n} z^n \tag{1}$$

for |z| < 1 where  $\binom{A}{n}$  is the binomial choose function. It is well known that the series expansion does not converge for |z| > 1 where A is a real number which is not equal to zero or a positive integer.

We could obtain a limited series expansion for |z| > 1 by writing the above expression as follows

$$(1+z)^A = (1+\frac{z}{2}+\frac{z}{2})^A = (1+\frac{z}{2})^A (1+\frac{z}{2})^A (1+\frac{z}{2})^A = (1+\frac{z}{2})^A (1+\frac{z}{z+2})^A$$
(2)

The second term in the above equation has a convergent series representation, given that  $\left|\frac{z}{z+2}\right| < 1$ . If  $\left|\frac{z}{2}\right| > 1$ , we can write

$$(1 + \frac{z}{2})^A = (1 + \frac{z}{4})^A (1 + \frac{z}{z+4})^A \tag{3}$$

Repeating this procedure iteratively, if  $m_0$  is the minimum value for which  $\left|\frac{z}{2^{m_0}}\right| < 1$ , we can write

$$(1+z)^A = (1+\frac{z}{2^{m_0}})^A \prod_{r=1}^{m_0} (1+\frac{z}{z+2^r})^A$$
 (4)

Each of the terms in the above product of terms has a convergent series representation. Given that we can write the convergent series expansion for each of the terms above as  $(1 + \frac{z}{2m_0})^A =$ 

Preprint submitted to Number Theory

 $\sum_{n=0}^{\infty} {A \choose n} (\frac{z}{2^{m_0}})^n$  and  $(1 + \frac{z}{(z+2^r)})^A = \sum_{m=0}^{\infty} {A \choose m} (\frac{z}{(z+2^r)})^m$ , where  ${A \choose n}$  represents the Choose function[1], we have the **series expansion for**  $(1+z)^A$  **expressed as a product of convergent series, which converges for** |z| > 1 as follows:

$$(1+z)^{A} = \left[\sum_{n=0}^{\infty} \binom{A}{n} (\frac{z}{2^{m_0}})^n\right] \left[\prod_{r=1}^{m_0} \sum_{m=0}^{\infty} \binom{A}{m} [\frac{z}{z+2^r}]^m\right]$$
 (5)

#### 2. Section 2

Let us take the case of  $m_0 = 1$  for 1 < |z| < 2. Expanding the term  $\frac{1}{(z+2^r)^m}$  in the above equation 5 as Taylor series around a point z = 0, we have for m > 0

$$\frac{1}{(z+2^r)^m} = \sum_{i=0}^{\infty} b(j,r,m)z^j$$
 (6)

where  $b(0, r, m) = \frac{1}{(2^r)^m}$  and b(j, r, m) is given as follows for j = 1, 2, 3...

$$b(j,r,m) = \binom{m+j-1}{j} \frac{(-1)^j}{(2^r)^{m+j}};$$
(7)

For m = 0,  $\frac{1}{(z+2^r)^m} = 1$ . Now we can write the series expansion of  $(1+z)^A$  which converges for 1 < |z| < 2, as a product of terms expanded in Taylor series as follows:

$$(1+z)^A = \left[\sum_{n=0}^{\infty} \binom{A}{n} (\frac{z}{2})^n\right] \left[\sum_{m=0}^{\infty} \binom{A}{m} z^m \sum_{j=0}^{\infty} b(j,1,m) z^j\right]$$
(8)

For the case of  $m_0 > 1$  for |z| > 2, we can write as follows:

$$(1+z)^{A} = \left[\sum_{n=0}^{\infty} \binom{A}{n} \left(\frac{z}{2^{m_0}}\right)^{n}\right] \left[\sum_{m=0}^{\infty} \binom{A}{m} z^{m} \sum_{i=0}^{\infty} b(j, m_0, m) z^{j}\right] \left[\prod_{r=1}^{m_0-1} \sum_{m=0}^{\infty} \binom{A}{m} z^{m} \left[\frac{1}{z+2^{r}}\right]^{m}\right]$$
(9)

The last term in the above equation  $(\frac{1}{7+2^r})^m$  can be expressed as follows:

$$\left(\frac{1}{z+2r}\right)^m = (z+2^r)^{-m} = 2^{-r*m}(1+\frac{z}{2r})^{-m} \tag{10}$$

The term $(1+\frac{z}{2^r})^{-m}$  can be recursively expanded using Eq.9 by substituting  $z\to \frac{z}{2^r}$  and  $A\to -m$  and  $m_0\to m_0-r$  to obtain the series expansion of  $(1+z)^A$  which converges for |z|>1 as a product of terms expanded in Taylor series.

## 3. Section 3

Let us consider the following binomial expression

$$(x+y)^A \tag{11}$$

where *A* is a real number which is not equal to zero or a positive integer and  $z = \frac{x}{y}$  and |z| > 1. Writing  $(x + y)^A = (1 + \frac{x}{y})^A y^A = (1 + z)^A y^A$ , we can write the series expansion of this expression using results obtained in equations 5 and 9 as follows:

$$(x+y)^A = y^A \sum_{n=0}^{\infty} {A \choose n} (\frac{z}{2^{m_0}})^n \prod_{r=1}^{m_0} \sum_{m=0}^{\infty} {A \choose m} [\frac{z}{z+2^r}]^m$$
 (12)

$$(x+y)^{A} = y^{A} \left[\sum_{n=0}^{\infty} \binom{A}{n} (\frac{z}{2^{m_0}})^{n}\right] \left[\sum_{m=0}^{\infty} \binom{A}{m} z^{m} \sum_{j=0}^{\infty} b(j, m_0, m) z^{j}\right] \left[\prod_{r=1}^{m_0-1} \sum_{m=0}^{\infty} \binom{A}{m} z^{m} \left[\frac{1}{z+2^r}\right]^{m}\right]$$
(13)

# 4. Conclusion

It has been shown that the Taylor series expansion of  $(1 + z)^A$  can be expanded as a product of convergent series, for |z| > 1 where A is a real number which is not equal to zero or a positive integer.

# 5. Acknowledgements

The author would like to thank Michael Schlosser(Universitat Wien) for his constructive comments on the paper.

#### 6. References

[1] Abramowitz, M. and Stegun, I. A. (Eds.). Handbook of Mathematical Functions with Formulas, Graphs, and Mathematical Tables, 9th printing. New York: Dover, 1972.