

On Taylor series expansion of $(1+z)^A$ for $|z| > 1$

Akhila Raman

University of California at Berkeley, CA-94720. Email: akhila.raman@berkeley.edu. Ph: 510-540-5544

Abstract

It is well known that the Taylor series expansion of $(1+z)^A$ does not converge for $|z| > 1$ where A is a real number which is not equal to zero or a positive integer. A limited series expansion of this expression is obtained in this paper for $|z| > 1$ as a product of convergent series.

Keywords:

1. Introduction

It is well known that the Taylor series expansion of $(1+z)^A$ is given by

$$(1+z)^A = \sum_{n=0}^{\infty} \binom{A}{n} z^n \quad (1)$$

for $|z| < 1$ where $\binom{A}{n}$ is the binomial choose function. It is well known that the series expansion does not converge for $|z| > 1$ where A is a real number which is not equal to zero or a positive integer.

We could obtain a limited series expansion for $|z| > 1$ by writing the above expression as follows

$$(1+z)^A = \left(1 + \frac{z}{2} + \frac{z}{2}\right)^A = \left(1 + \frac{z}{2}\right)^A \left(1 + \frac{\frac{z}{2}}{\left(1 + \frac{z}{2}\right)}\right)^A = \left(1 + \frac{z}{2}\right)^A \left(1 + \frac{z}{z+2}\right)^A \quad (2)$$

The second term in the above equation has a convergent series representation, given that $|\frac{z}{z+2}| < 1$. If $|\frac{z}{2}| > 1$, we can write

$$\left(1 + \frac{z}{2}\right)^A = \left(1 + \frac{z}{4}\right)^A \left(1 + \frac{z}{z+4}\right)^A \quad (3)$$

Repeating this procedure iteratively, if m_0 is the minimum value for which $|\frac{z}{2^{m_0}}| < 1$, we can write

$$(1+z)^A = \left(1 + \frac{z}{2^{m_0}}\right)^A \prod_{r=1}^{m_0} \left(1 + \frac{z}{z+2^r}\right)^A \quad (4)$$

Each of the terms in the above product of terms has a convergent series representation. Given that we can write the convergent series expansion for each of the terms above as $\left(1 + \frac{z}{2^{m_0}}\right)^A =$

$\sum_{n=0}^{\infty} \binom{A}{n} \left(\frac{z}{2^{m_0}}\right)^n$ and $(1 + \frac{z}{z+2^r})^A = \sum_{m=0}^{\infty} \binom{A}{m} \left(\frac{z}{z+2^r}\right)^m$, where $\binom{A}{n}$ represents the Choose function[1], we have the **series expansion for $(1 + z)^A$ expressed as a product of convergent series, which converges for $|z| > 1$** as follows:

$$(1 + z)^A = \left[\sum_{n=0}^{\infty} \binom{A}{n} \left(\frac{z}{2^{m_0}}\right)^n \right] \left[\prod_{r=1}^{m_0} \sum_{m=0}^{\infty} \binom{A}{m} \left[\frac{z}{z+2^r}\right]^m \right] \quad (5)$$

2. Section 2

Expanding the term $\frac{1}{(z+2^r)^m}$ in the above equation 5 as Taylor series around a point $z = 0$, we have

$$\frac{1}{(z + 2^r)^m} = \sum_{j=0}^{\infty} b_j z^j \quad (6)$$

where $b_0 = \frac{1}{(2^r)^m}$ and b_j is given as follows for $j = 1, 2, 3, \dots$

$$b_j = \binom{m+j-1}{j} \frac{(-1)^j}{(2^r)^{m+j}}; \quad (7)$$

Now we can write the the series expansion of $(1+z)^A$ which converges for $|z| > 1$, as a product of terms expanded in Taylor series as follows:

$$(1 + z)^A = \left[\sum_{n=0}^{\infty} \binom{A}{n} \left(\frac{z}{2^{m_0}}\right)^n \right] \left[\prod_{r=1}^{m_0} \left[\sum_{m=0}^{\infty} \binom{A}{m} z^m \sum_{j=0}^{\infty} b_j z^j \right] \right] \quad (8)$$

3. Section 3

Let us consider the following binomial expression

$$(x + y)^A \quad (9)$$

where A is a real number which is not equal to zero or a positive integer and $z = \frac{x}{y}$ and $|z| > 1$. Writing $(x + y)^A = (1 + \frac{x}{y})^A y^A = (1 + z)^A y^A$, we can write the series expansion of this expression using results obtained in equations 5 and 8 as follows:

$$(x + y)^A = y^A \sum_{n=0}^{\infty} \binom{A}{n} \left(\frac{z}{2^{m_0}}\right)^n \prod_{r=1}^{m_0} \sum_{m=0}^{\infty} \binom{A}{m} \left[\frac{z}{z+2^r}\right]^m \quad (10)$$

$$(x + y)^A = y^A \left[\sum_{n=0}^{\infty} \binom{A}{n} \left(\frac{z}{2^{m_0}}\right)^n \right] \left[\prod_{r=1}^{m_0} \left[\sum_{m=0}^{\infty} \binom{A}{m} z^m \sum_{j=0}^{\infty} b_j z^j \right] \right] \quad (11)$$

4. Conclusion

It has been shown that the Taylor series expansion of $(1 + z)^A$ can be expanded as a product of convergent series, for $|z| > 1$ where A is a real number which is not equal to zero or a positive integer.

5. References

[1] Abramowitz, M. and Stegun, I. A. (Eds.). Handbook of Mathematical Functions with Formulas, Graphs, and Mathematical Tables, 9th printing. New York: Dover, 1972.