Equivelar and d-Covered Triangulations of Surfaces. I Frank H. Lutz, Thom Sulanke, Anand K. Tiwari, Ashish K. Upadhyay

Abstract

We survey basic properties and bounds for q -equivelar and d -covered triangulations of closed surfaces. Included in the survey is a list of the known sources for q-equivelar and d-covered triangulations. We identify all orientable and non-orientable surfaces M of Euler characteristic $0 > \chi(M) \geq -230$ which admit non-neighborly q-equivelar triangulations with equality in the upper bound $q \leq \left(\frac{1}{2}(5+\sqrt{49-24\chi(M)})\right)$. These examples give rise to d-covered triangulations with equality in the upper bound $d \leq 2 \left| \frac{1}{2}(5 + \sqrt{49 - 24\chi(M)}) \right|$. A generalization of Ringel's cyclic 7mod12 series of neighborly orientable triangulations [53] to a two-parameter family of cyclic orientable triangulations $R_{k,n}$, $k \geq 0$, $n \geq 7 + 12k$, is the main result of this paper. In particular, the two infinite subseries $R_{k,7+12k+1}$ and $R_{k,7+12k+2}$, $k \ge 1$, provide non-neighborly examples with equality for the upper bound for q as well as derived examples with equality for the upper bound for d.

1 Introduction and Survey

A triangulation (as a finite simplicial complex) K of a (closed) surface M is equivelar if all vertices of the triangulation have the same vertex-degree q. The f -vector (or face vector) of K is the vector $f(K) = (f_0(K), f_1(K), f_2(K))$, where $f_0(K), f_1(K)$, and $f_2(K)$ are denoting the numbers of vertices, edges, and triangles of K, respectively. For brevity, we write $f = (n, f_1, f_2)$ for the f-vector of a triangulation, with $n = f_0$ denoting the numbers of vertices. If the vertices of an equivelar triangulation K of a surface M have degree q , then the triangulation is called q-equivelar or equivelar of type $\{3, q\}$. For q-equivelar triangulations it follows by double counting of incidences between vertices and edges as well as between edges and triangles that

$$
nq = 2f_1 = 3f_2,\tag{1}
$$

which yields $f = (n, \frac{nq}{2})$ $\frac{u}{2}$, $\frac{nq}{3}$ $\frac{u}{3}$). Moreover, by Euler's equation,

$$
\chi(M) = n - f_1 + f_2 = n - \frac{nq}{2} + \frac{nq}{3} = \frac{n(6-q)}{6},\tag{2}
$$

or equivalently,

$$
q = 6 - \frac{6\chi(M)}{n}.\tag{3}
$$

Figure 1: A vertex-star of degree six and its subdivision.

Since q is a positive integer, n has to be a divisor of $6|\chi(M)|$ if $\chi(M) \neq 0$. In particular, a surface M of Euler characteristic $\chi(M) \neq 0$ has only finitely many equivelar triangulations; see for example [61] for a list of the possible values of (n, q) for equivelar triangulations of surfaces M with $\chi(M) \geq -10$.

A triangulation K of a surface M is d-covered, if at least one vertex of each edge of K has degree d. The study of d-covered triangulations was initiated by Negami and Nakamoto in [50]. In particular, a d-covered triangulation K is covered by triangulated disks, which are vertex-stars of vertices of degree d. Any q-equivelar triangulation clearly is a q-covered triangulation. If we stack all the triangles of a q-equivelar triangulation, i.e., if we place for each triangle a new vertex inside the triangle and connect the new vertex with the three vertices of the triangle by an edge each, then the resulting triangulation is 2q-covered [50]; see Figure 1 for the respective subdivision of a vertex-star of degree six.

For an arbitrary triangulation K of a surface M of Euler characteristic $\chi(M)$, the f-vector of K can (by double counting, $2f_1 = 3f_2$, and by Euler's equation, $\chi(M) = n - f_1 + f_2$) be expressed as $f = (n, 3(n - \chi(M)), 2(n - \chi(M)))$. Since K has at most $f_1 \leq {n \choose 2}$ $n/2$ edges, it follows that

$$
n \ge \left\lceil \frac{1}{2}(7 + \sqrt{49 - 24\chi(M)}) \right\rceil,\tag{4}
$$

which is Heawood's bound [26] from 1890. A triangulation K is called *neighborly* if $f_1 = \binom{n}{2}$ $\binom{n}{2}$, i.e., if K has a complete 1-skeleton. Neighborly triangulations are equivelar with $q = n-1$. Moreover, we have equality $3(n - \chi(M)) = f_1 = \binom{n}{2}$ n_2) for neighborly triangulations, or, equivalently

$$
\chi(M) = \frac{n(7-n)}{6},\tag{5}
$$

which implies $n \equiv 0, 1, 3, 4 \mod 6$, or equivalently, $n \equiv 0, 1 \mod 3$, where $n \geq 4$.

Theorem 1 (Jungerman and Ringel [31], [52]) Let M be a surface different from the orientable surface of genus 2, the Klein bottle, and the non-orientable surface of genus 3, then there is a triangulation of M on n vertices if and only if

$$
n \ge \left\lceil \frac{1}{2}(7 + \sqrt{49 - 24\chi(M)}) \right\rceil,
$$

with equality if and only if the triangulation is neighborly. For the three exceptional cases, one extra vertex has to be added to the lower bound; cf. [28].

Let $G = Sk_1(K)$ be the 1-skeleton of K, $deg_G(v)$ the degree of a vertex v in the graph G, and $\delta(G) = \min_{v \in G} \deg_G(v)$ the minimum degree in G. For any triangulated surface K with *n* vertices it trivially holds that $3 \leq \delta(G) \leq n-1$. Moreover, $n\delta(G) \leq \sum_{v \in G} \deg_G(v) = 2f_1$, with the equality by double counting of vertex-edge incidences. By combining $\delta(G) \leq \frac{2f_1}{n}$ with $f_1 = 3(n - \chi(M))$, it follows that

$$
\delta(G) \le 6 - \frac{6\chi(M)}{n}.\tag{6}
$$

If $\chi(M) > 0$, then $\delta(G) \leq 5$. For $\chi(M) \leq 0$, we combine (6) with $\delta(G) \leq n-1$ and obtain $\delta(G) \leq 6 - \frac{6\chi(M)}{\delta(G)+1}$, which is equivalent to

$$
\delta(G) \le \left\lfloor \frac{1}{2}(5 + \sqrt{49 - 24\chi(M)}) \right\rfloor. \tag{7}
$$

In particular, we have $\delta(G) = q$ for any q-equivelar triangulation K of a surface M, and thus

$$
q \le \left\lfloor \frac{1}{2}(5 + \sqrt{49 - 24\chi(M)}) \right\rfloor. \tag{8}
$$

Let K be a d-covered triangulation of a surface $M, G = Sk_1(K)$, and H the subgraph of G induced by the vertices of degree d. If v is a vertex of H , then every edge of the link of v in G contains at least one vertex of H , since otherwise this edge would not be covered by a vertex of degree d. Hence, $\deg_H(v) \geq \frac{d}{2}$ $\frac{a}{2}$, from which it follows that

$$
\frac{d}{2} \le \delta(H). \tag{9}
$$

If K is a 3-covered triangulation and v a vertex of degree 3, then the link of v contains at least two vertices of degree 3, which immediately implies that the third vertex of the link also has degree 3, and thus K is the boundary of the tetrahedron with 4 vertices. If K is a 4-covered triangulation, it can be concluded that K is the suspension of a triangulated circle. For a characterization of 5- and 6-covered triangulations see [50].

Let K be a d-covered triangulation of a surface M and w a vertex of degree $\deg_G(w) \neq d$. Then all vertices of the link of w are of degree d and thus are vertices of H . Therefore, if we delete all vertices w of degree $\deg_G(w) \neq d$ and the edges these vertices are contained in from the triangulation K, we obtain a regular cellular decomposition C of M, with $H = Sk_1(C)$. (A cell decomposition of a surface is regular if there are no identifications on the boundary of any cell.) Let $f^C = (f_0^C, f_1^C, f_2^C)$ be the face vector of C. For a 2-face P of C let $p(P)$ be the number of vertices of the polygon P, where $p(P) \geq 3$. Then, by the regularity of the cell complex C, $2f_1^C = \sum_{P \in \{2\text{-faces of } C\}} p(P) \ge 3f_2^C$. Hence, $\chi(M) = f_0^C - f_1^C + f_2^C \le f_0^C - \frac{1}{3}$ $\frac{1}{3}f_1^C,$ or, equivalently, $f_1^C \leq 3(f_0^C - \chi(M))$. It then follows that

$$
\frac{d}{2} \le \delta(H) = \frac{f_0^C \delta(H)}{f_0^C} \le \frac{\sum_{v \in H} \deg_H(v)}{f_0^C} = \frac{2f_1^C}{f_0^C} \le 6 - \frac{6\chi(M)}{f_0^C},\tag{10}
$$

which, together with $\frac{d}{2} \leq \delta(H) \leq f_0^C - 1$, gives

$$
d \le 2\left\lfloor \frac{1}{2}(5+\sqrt{49-24\chi(M)}) \right\rfloor \tag{11}
$$

if $\chi(M) \leq 0$, whereas $\frac{d}{2} \leq 5$ for $\chi(M) > 0$.

Theorem 2 (Negami and Nakamoto [50]) Let K be a d-covered triangulation of a surface M.

- (a) If $\chi(M) > 0$, then $d \leq 10$.
- (b) If $\chi(M) \leq 0$, then $d \leq 2 \left| \frac{1}{2} \right|$ $\frac{1}{2}(5+\sqrt{49-24\chi (M)})\Big|$.

If $d \geq 13$, then $7 \leq \lceil \frac{d}{2} \rceil \leq \delta(H) \leq 6 - \frac{6\chi(M)}{f_0^C}$ $\frac{\zeta(M)}{f_0^C}$, or, equivalently $f_0^C \leq -6\chi(M)$. As there are only finitely many regular cell decompositions C of a surface M with $f_0^C \le -6\chi(M)$, we obtain:

Theorem 3 (Negami and Nakamoto [50]) Let $d \geq 13$ and M be a closed surface. Then M has only finitely many d-covered triangulations.

Remark: The proofs of the Theorems 2 and 3 given in [50] are somewhat vague and imprecise. Nevertheless, the statements of the two theorems are correct, as we verified above. In particular, we have for regular cell decompositions:

Theorem 4 Let C be a regular cell decomposition of a surface M and let $H = Sk_1(C)$.

- (a) If $\chi(M) > 0$, then $\delta(H) \leq 5$.
- (b) If $\chi(M) \leq 0$, then $\delta(H) \leq \left| \frac{1}{2} \right|$ $\frac{1}{2}(5+\sqrt{49-24\chi(M)})\,\Big|.$

Theorem 5 Let M be a closed surface. Then M has only finitely many regular cell decompositions C with $\delta(\text{Sk}_1(C)) \geq 7$.

Since any q-equivelar triangulation yields, by subdivision, a $2q$ -covered triangulation, we see by the bounds (8) and (11) , that q-equivelar triangulations with

$$
q = \left[\frac{1}{2} (5 + \sqrt{49 - 24\chi(M)}) \right]
$$
 (12)

give 2q-covered triangulation with

$$
d = 2q = 2\left[\frac{1}{2}(5 + \sqrt{49 - 24\chi(M)})\right],\tag{13}
$$

which means tightness in inequality (11) .

In the neighborly cases $n \equiv 0, 1, 3, 4 \mod 6$, with $n \geq 4$, there are, according to Theorem 1 and Equation (5) , neighborly triangulations of orientable surfaces M with

$$
\chi(M) = \begin{cases}\n-24k^2 + 14k & \text{for } n = 12k, k \ge 1, \\
-24k^2 + 2k + 2 & \text{for } n = 12k + 3, k \ge 1, \\
-24k^2 - 2k + 2 & \text{for } n = 12k + 4, k \ge 0, \\
-24k^2 - 14k & \text{for } n = 12k + 7, k \ge 0\n\end{cases}
$$
\n(14)

and neighborly triangulations of non-orientable surfaces M with

$$
\chi(M) = \begin{cases}\n-\frac{3}{2}m^2 + \frac{7}{2}m & \text{for } n = 3m, m \ge 2, \\
-\frac{3}{2}m^2 + \frac{5}{2}m + 1 & \text{for } n = 3m + 1, m \ge 3.\n\end{cases}
$$
\n(15)

The respective examples are equivelar with $q = n - 1$ and yield 2q-covered triangulations with tight bound (11).

Figure 2: The series of triangulated tori $T^2(n)$.

Apart from the series of neighborly triangulations of Ringel [52] and Jungerman and Ringel [31], infinite classes and series of equivelar triangulation are known for all $q \geq 6$. In addition, there are some sporadic examples as well as enumerational results for small values of n and $|\chi|$:

- The boundaries of the tetrahedron, the octahedron, and the icosahedron are the only equivelar triangulations of the 2-sphere S^2 with $\chi(S^2) = 2$. In these cases, $(n, q) = (4, 3), (6, 4)$, and (12, 5), respectively. The unique neighborly 6-vertex triangulation of the real projective plane $\mathbb{R}P^2$ with $\chi(\mathbb{R}P^2) = 1$ and $(n,q) = (6,5)$ is the only equivelar triangulation of $\mathbb{R}P^2$. Together, these four examples are the only equivelar triangulations with $q \leq 5$.
- There are infinitely many equivelar triangulations of the 2-torus T^2 with $\chi(T^2) = 0$ and $q = 6$. A complete classification of these examples is due to Brehm and Kühnel [11]. All of these examples have vertex-transitive cyclic symmetry [19].

Let for $n \geq 7$ the cyclic group \mathbb{Z}_n on the elements $0, 1, \ldots, n-1$ be generated by the permutation $(0, 1, \ldots, n-1)$ and let $T^2(n)$ be defined by the orbits of the triangles $[0, 1, 3]$ and [0, 2, 3]. Then $T^2(n)$, as depicted in Figure 2, is a cyclic series of triangulated tori [1], [37]. For $n \geq 7$, the example $T^2(n)$ is realizable geometrically in the 2-skeleton of the cyclic 3-polytope $C_3(n)$ [1].

- Series of *equivelar polyhedra* in 3-space (i.e., geometric realizations in \mathbb{R}^3 of orientable equivelar surfaces) were first described in [46]: There are equivelar polyhedra of the types $\{3, q; g\}$ for
	- (a) $\{3,7;g\}$ with $n = 12(g-1)$ for $g \ge 2$,
	- (b) $\{3, 8; g\}$ with $n = 6(g 1)$ for $g \ge 4$,
	- (b) $\{3, 9; g\}$ with $n = 4(g 1)$ for $g = 6, 9, 10$ and $g \ge 12$,
	- (d) $\{3, 12; g\}$ with $n = 2(g 1)$ for $g = 73 + 66k, k \ge 0$,

with $g = 1 - \frac{1}{2}$ $\frac{1}{2}\chi$ the genus of a respective equivelar triangulation of vertex-degree q and, as a consequence of (3), with $n = \frac{6(2-2g)}{6-g}$ $rac{z-2g}{6-q}$ vertices.

• McMullen, Schulz, and Wills [47] provided in 1983 a combinatorial construction method for equivelar triangulations, which, in a second inductive step, allows a large family of equivelar polyhedra to be obtained. In particular, for each $q \geq 7$ there are infinitely many combinatorially different equivelar polyhedra of type $\{3,q\}$. Since for $g \geq 2$ the orientable

surface M_g of genus g has (at most) finitely many equivelar triangulations, it follows that for each $q \ge 7$ there are infinitely many equivelar polyhedra of distinct genus in the family. For $r \geq 3$, the vertex-minimal equivelar examples of type $\{3, 2r\}$ in the constructed family have $n(r) = 9 \cdot 2^{r-3}$ vertices and are of genus $g(r) = 3(r-3)2^{r-4} + 1$. By showing that for each $r \geq 3$ and all $0 \leq g \leq g(r)$ there are equivelar polyhedra M_q of genus g with $f_0(M_q) \leq n(r)$ vertices, McMullen, Schulz, and Wills established the remarkable result that there are (equivelar) polyhedra of genus g with $f_0 = O(g/\log g)$ vertices. Asymptotically, this is the strongest known result for the realizability in \mathbb{R}^3 of triangulated (orientable) surfaces of genus g, for which, by Heawood's bound (4), $O(\sqrt{g})$ vertices are needed for a triangulation. A generalization of the McMullen-Schulz-Wills family of equivelar surfaces is given in [55].

- Equivelar triangulations of the Klein bottle are of type $\{3, 6\}$. The respective 1-skeleta are 6-regular graphs and were classified in [49]; see also [24] for additional comments and further references. Equivelar triangulations of the Klein bottle (and of the torus) with up to 15 vertices were classified in [19]. An enumeration for up to 100 vertices can be found in $[61]$. As observed in [19], there is an *n*-vertex equivelar triangulation of the Klein bottle if and only if $n \geq 9$ is not prime.
- Equivelar triangulations of the double torus have 12 vertices and are of type $\{3, 7\}$ by (3) and (4): there are six such examples [20].
- Equivelar triangulations with up to 11 vertices were classified in [18]: there are 27 such examples. By enumeration, there are 240 914 equivelar triangulations with 12 vertices [61].
- There are no equivelar triangulations of the non-orientable surface of genus 3 with $\chi = -1$. A complete enumeration of the equivelar triangulations of the non-orientable surface of genus 4 with $\chi = -2$ (28 examples of type $(3, q; n) = (3, 7; 12)$), of the non-orientable surface of genus 5 with $\chi = -3$ (two examples of type $(3, 8; 9)$ and 1401 examples of type $(3, 7; 18)$, of the non-orientable surface of genus 6 with $\chi = -4$ (6500 examples of type $(3, 8; 12)$ and 600 946 examples of type $(3, 7; 24)$, and of the orientable surface of genus 3 with $\chi = -4$ (24 examples of type (3, 8; 12) and 11 277 examples of type (3, 7; 24)) was given in [61].
- The neighborly triangulations with up to 13 vertices are: the boundary of the tetrahedron with 4 vertices as the unique neighborly triangulation of S^2 , the unique Möbius torus [48] with 7 vertices, 59 neighborly triangulations of the orientable surface of genus 6 with 12 vertices [2], the unique vertex-minimal 6-vertex triangulation of the real projective plane, two neighborly triangulations of the non-orientable surface of genus 5 with 9 vertices [3], 14 neighborly triangulations of the non-orientable surface of genus 7 with 10 vertices [3], 182 200 neighborly triangulations of the non-orientable surface of genus 12 with 12 vertices [23], [61], and 243 088 286 neighborly triangulations of the non-orientable surface of genus 15 with 13 vertices [23].

The 4-vertex triangulation of S^2 and Möbius' torus are the only neighborly triangulations for which geometric realizations are known in \mathbb{R}^3 (for the Császár torus see [16] and also [5], [39]). None of the 59 neighborly triangulations of the orientable surface of genus 6 with 12 vertices is realizable in \mathbb{R}^3 [6], [56], and it is assumed that neighborly triangulations of orientable surfaces of higher genus never are realizable. This leaves a wide gap to the realizability result of McMullen, Schulz, and Wills [47] that there are realizations of orientable surface of genus g with $O(g/\log g)$ vertices.

- A particular class of symmetric equivelar triangulations are vertex-transitive triangulations. An enumeration of vertex-transitive triangulations of surfaces with up to 15 vertices was given in [34], [38] and of vertex-transitive neighborly triangulations of surfaces with up to 22 vertices in [38].
- Regular simplicial maps with a flag-transitive automorphism group are equivelar with highest possible symmetry. The book by Coxeter and Moser [15] gives a classical treatment. A complete enumeration of all regular maps with Euler characteristic $\chi \geq -200$ was obtained by Conder [14]. Dyck's regular map [21], [22] and Klein's regular map [33] are well-known examples of regular simplicial maps for which geometric realization are known [4], [7], [8], [57], [58].
- There are two well-known infinite series of vertex-transitive triangulations of surfaces, the above series $T^2(n)$ of cyclic torus triangulations [1], [37] and Ringel's cyclic 7mod12 series of neighborly orientable triangulations [53]. In fact, it was shown by Heffter [27] that cyclic neighborly triangulations of orientable surfaces can only exist for $n \equiv 7 \mod 12$. The Altshuler series, the Ringel series, and the class of cyclic equivelar tori [11] are the only infinite series of vertex-transitive triangulations of surfaces that, so far, have been described explicitely in the literature, while additional series of cyclic triangulations of surfaces of higher genus are implicitly contained in $[9]$, $[10]$, $[17]$, and $[29]$.

Every equivelar triangulation of type $\{3,q\}$ gives rise to a 2q-covered triangulation. Other sources for d-covered triangulations are:

- The boundary of the tetrahedron is the only 3-covered triangulation. Any 4-covered triangulation is the suspension of a triangulated circle. A characterization of 5- and 6-covered triangulations is due to Negami and Nakamoto [50]. For given $5 \leq d \leq 12$, Katahira [32] proved that there are infinitely many distinct d-covered triangulations.
- A map on a surface M is a decomposition of M into a finite cell complex. A map is *equivelar* of type $\{p, q\}$ if M is decomposed into p-gons such that every vertex has degree q; cf. [46], [47]. A map is polyhedral if the intersection of any two of its polygons is either empty, a common vertex, or a common edge; see [12], [13]. An *equivelar polyhedral map* is a map, which is both equivelar and polyhedral.

If we place for each p-gon of a $\{p,q\}$ -equivelar polyhedral map a new vertex inside the pgon and connect the new vertex with the p vertices of the p -gon by an edge each, then the resulting triangulation is 2q-covered.

Equivelar quadrangulations with few vertices are enumerated in [44], [61]. McMullen, Schulz, and Wills [47] gave infinite families of equivelar polyhedral maps of the types $\{4, q\}$ and $\{p, 4\}$ along with geometric realizations in \mathbb{R}^3 of these examples. For geometric realizations of further equivelar polyhedral maps of these types see [7], [45], [46], [59], and [55].

It is not known whether there are geometric realizations of equivelar polyhedral maps of type $\{p, q\}$ for $p, q \geq 5$; cf. [13]. Examples of equivelar polyhedral maps of type $\{5, 5\}$ and of type $\{6, 6\}$ were first given by Brehm [9]. An infinite series of $\{k, k\}$ -equivelar polyhedral maps was constructed by Datta [17].

- A $\{p, q\}$ -equivelar map is regular if it has a flag-transitive automorphism group. Regular polyhedral maps therefore provide highly symmetric examples of equivelar polyhedral maps; see [14], [15].
- If a $\{p, q\}$ -equivelar map is not a polyhedral map, but at least gives a regular cell decomposition of a surface, then this suffices to obtain a 2q-covered triangulation of the surface by subdividing the p-gons. The Heffter surfaces [27] and the generalized Heffter surfaces by Pfeifle and Ziegler [51] provide such examples.

Infinite series of vertex-transitive triangulations of surfaces as well as of higher-dimensional manifolds are rare in the literature. The cyclic 7mod12 series of Ringel [53] from 1961, the cyclic series of triangulated tori of Altshuler [1] from 1971, and the class of all cyclic tori, as completely classified by Brehm and Kühnel [11] in 2008, are the only documented series of vertextransitive triangulations of surfaces. In higher dimensions, the boundaries of simplices, of crosspolytopes, and of cyclic polytopes are classical examples of vertex-transitive triangulations of spheres. A cyclic series of centrally symmetric 3-spheres is given in [40], series of wreath products are described in [30], vertex-transitive triangulations of higher-dimensional tori in [36], and series of cyclic triangulations of sphere bundles in [35], [37].

In the following section, we consider surfaces with Euler characteristic $0 > \chi \ge -230$. We show that in all non-neighborly cases with $q = \frac{1}{2}$ $\frac{1}{2}(5+\sqrt{49-24\chi(M)})\Big\vert$, that is, with equality in (8), there indeed are equivelar triangulations of the respective surfaces. By subdividing these examples, we obtain 2q-covered triangulations with equality in (11) .

In Section 3, we generalize Ringel's cyclic 7mod12 series to a two-parameter family $R_{k,n}$ of cyclic triangulations of the orientable surfaces of genus $g = kn + 1$ with $n \geq 7 + 12k$ vertices and $q = 6 + 12k$, $k \ge 0$. For $k = 0$, $R_{0,n}$ coincides with Altshuler's cyclic series of tori, whereas $R_{k,7+12k}$ is Ringel's cyclic series of neighborly triangulations with $n = 7 + 12k$ vertices. We show that, apart from the cyclic tori with $n \geq 8$ vertices, the two subseries $R_{k,7+12k+1}$ and $R_{k,7+12k+2}, k \ge 1$, provide infinitely many examples of non-neighborly cyclic triangulations with $q=6+12k=\left\lfloor \frac{1}{2}\right\rfloor$ $\frac{1}{2}(5+\sqrt{49-24\chi})\Big|$, i.e., with equality in (8). Again, by subdivision, these examples yield 2q-covered triangulations with equality in (11).

In addition to the generalized Ringel two-parameter family of cyclic triangulations, a generation scheme for a large class of series of cyclic triangulations of surfaces is presented in [41]. In particular, infinite series of cyclic q-equivelar triangulations of orientable and non-orientable surfaces are given in [41] for each

$$
q = \begin{cases} 3k & \text{for} \quad k \ge 2, \\ 3k+1 & \text{for} \quad k \ge 3. \end{cases} \tag{16}
$$

χ	q	$\,n$	Trans. or.	Or.	Trans. non-or.	Non-or.
-2	7	12	yes $[34]$	yes $[61]$	yes $[34]$	yes $[61]$
$-4\,$	8	12	yes [34]	yes $[61]$	\log [34]	yes $[61]$
$-6\,$	9	12	yes $[34]$	yes $[61]$	yes $[34]$	yes $[61]$
$-\mathbf{7}$	9	14			no [34]	yes, E1
$-8\,$	10	12	yes $[34]$	yes $[61]$	yes $[34]$	yes $[61]$
-14	12	14	yes $[34]$	yes $[34]$	yes $[34]$	yes $[34]$
-15	12	15			yes $[34]$	yes $[34]$
-16	12	16	yes, E2	yes, E2	yes, E3	yes, E3
$-27\,$	15	18			no	yes, E4
$-30\,$	16	18	yes, E5	yes, E5	yes, E6	yes, E6
-40	18	$20\,$	yes, E7	yes, E7	yes, E8	yes, E8
-42	18	21	yes, E9	yes, E9	yes, E10	yes, $E10$
-60	21	$24\,$	yes, E11	yes, E11	\cdot	yes, $E12$
-64	22	24	\mathcal{P}	yes, E13	yes, E14	yes, $E14$
					\cdot ?	
$-78\,$	24	$26\,$	yes, $E15$	yes, E15		yes, E16
-81	24	$27\,$			yes, E17	yes, E17
-105	27	$30\,$			yes, E18	yes, $E18$
-110	28	30	$\ddot{?}$	yes, E19	yes, E20	yes, E20

Table 1: Existence of non-neighborly examples with equality in (8) for $0 > \chi \ge -127$.

2 d-covered triangulations with $d=2\left|\frac{1}{2}\right|$ $\left. \frac{1}{2}(5+\sqrt{49-24\chi (M)})\right|$

In the orientable neighborly triangulated cases with $n \equiv 0, 3, 4, 7 \mod 12$ vertices, $n \geq 4$, and in the non-orientable neighborly triangulated cases with $n \equiv 0, 1 \mod 3$ vertices, $n \geq 8$, the Euler characteristic $\chi(M)$ of a respective orientable or non-orientable surface M is determined, as a consequence of (5), by (14) and (15), respectively. In all these cases of neighborly triangulations, $n=\frac{1}{2}$ $\frac{1}{2}(7+\sqrt{49-24\chi(M)})$, $q=n-1$, and therefore $d=2q=2n-2=2\cdot\frac{1}{2}$ $\frac{1}{2}(5+\sqrt{49-24\chi(M)})$ for a derived d-covered triangulation.

In contrast, if $-\chi(M) \ge 5$ is prime, then Equation (3), $q = 6 - \frac{6\chi(M)}{n}$ $\frac{M}{n}$, yields integer values for q if and only if $n = -\chi(M), -2\chi(M), -3\chi(M), -6\chi(M)$, with corresponding values $q = 12, 9, 8, 7$, respectively.

Proposition 6 Let M be a surface with $-\chi(M) \geq 5$ prime. If M has an equivelar triangulation of type $\{3,q\}$, then $q \in \{7,8,9,12\}$.

In [62], the fourth author raised the question, whether there are cases, other than the neighborly ones, where $d = 2\lfloor \frac{1}{2} \rfloor$ $\frac{1}{2}(5 + \sqrt{49 - 24\chi(M)})$ can be achieved by subdivisions of q-equivelar triangulations with equality in (8), that is, with $q = \lfloor \frac{1}{2} \rfloor$ $\frac{1}{2}(5+\sqrt{49-24\chi(M)})$].

Example	Group: (n,i)	Orbit generating triangles
E2	(16, 189)	$[1,3,5]_{16}, [1,3,7]_{48}$
E3	(16,28)	$[1,3,6]_{32}, [1,3,9]_{32}$
E ₅	(18,3)	$[1, 2, 5]_{18}$, $[1, 3, 8]_{18}$, $[1, 4, 9]_{18}$, $[1, 4, 17]_{6}$, $[1, 5, 12]_{18}$, $[1, 6, 11]_{18}$
E ₆	(18,3)	$[1, 2, 3]_{18}$, $[1, 4, 6]_{18}$, $[1, 5, 11]_{18}$, $[1, 5, 15]_{6}$, $[1, 6, 16]_{18}$, $[1, 7, 13]_{18}$
E7	(20,35)	$[1,3,5]_{120}$
E8	(20,2)	$[1,3,5]_{20}, [1,3,7]_{20}, [1,6,13]_{20}, [1,6,15]_{20}, [1,7,13]_{20}, [1,10,19]_{20}$
E9	(21,7)	$[1, 4, 10]_{21}$, $[1, 4, 11]_{63}$, $[1, 5, 12]_{21}$, $[1, 6, 16]_{21}$
E10	(21,7)	$[1,4,11]_{63}, [1,4,13]_{63}$
E11	(24, 84)	$[1,4,7]_{24}, [1,4,14]_{72}, [1,5,10]_{72}$
E14	(24,21)	$[1,3,8]_{48}, [1,3,23]_{16}, [1,4,13]_{48}, [1,4,15]_{48}, [1,5,16]_{16}$
E15	(26,5)	$[1,3,5]_{78}, [1,3,13]_{78}, [1,4,10]_{26}, [1,6,18]_{26}$
E17	(27, 23)	$[1, 4, 10]_{81}$, $[1, 4, 11]_{81}$, $[1, 8, 18]_{27}$, $[1, 9, 17]_{27}$
E18	(30,9)	$[1, 2, 21]_{30}$, $[1, 3, 8]_{60}$, $[1, 4, 9]_{60}$, $[1, 5, 14]_{60}$, $[1, 9, 28]_{60}$
E20	(30,9)	$[1, 3, 8]_{60}$, $[1, 4, 9]_{60}$, $[1, 5, 14]_{60}$, $[1, 9, 30]_{30}$, $[1, 10, 30]_{10}$, $[1, 15, 17]_{60}$

Table 2: Examples with vertex-transitive symmetry.

For $\chi = 0$, all equivelar triangulations of the torus and the Klein bottle are of type $\{3, 6\}$, and thus yield 12-covered triangulations. So let $\chi < 0$. Then for a q-equivelar triangulation with $q=\lfloor \frac{1}{2}$ $\frac{1}{2}(5 + \sqrt{49 - 24\chi(M)})$ we have by (3) that

$$
n = -\frac{6\chi(M)}{q-6} = \frac{-6\chi(M)}{\left[\frac{1}{2}(5+\sqrt{49-24\chi(M)})\right]-6},\tag{17}
$$

which needs to be an integer.

Table 1 lists for $0 > \chi \ge -127$ all those triples (χ, q, n) with values $n \ge \left\lceil \frac{1}{2} \right\rceil$ $\frac{1}{2}(7+\sqrt{49-24\chi})\Big],$ for which $q = \lfloor \frac{1}{2} \rfloor$ $\frac{1}{2}(5+\sqrt{49-24\chi})$ < n - 1 and n = $\frac{-6\chi}{\sqrt{16}+\sqrt{49-24\chi}}$ $\lfloor \frac{1}{2}$ $\frac{1}{2}(5+\sqrt{49-24\chi})$] – 6 is an integer. These are, apart from the neighborly cases, the potential cases where q -equivelar triangulations with equality $q = \lfloor \frac{1}{2} \rfloor$ $\frac{1}{2}(5+\sqrt{49-24\chi})$ can occur. Columns four and six of Table 1 indicate whether vertextransitive orientable and non-orientable examples are known, whereas columns five and seven provide information on orientable and non-orientable equivelar triangulations, respectively. The vertex-transitive examples E2–E3, E5–E11, E14–E15, E17–E18, and E20 are listed in Table 2, the non-transitive examples E1, E4, E12, E13, E16, and E19 are listed at the end of this section.

Theorem 7 There are exactly 42 (orientable and non-orientable) surfaces with $0 > \chi \ge -230$ for which there are q-equivelar non-neighborly triangulations with equality in (8) and derived 2qcovered triangulations with equality in (11).

Proof: A complete list of equivelar triangulations with up to 12 vertices is given in [61], whereas vertex-transitive triangulations with up to 15 vertices can be found in [34] and [38]. For $16 \le n \le 30$, we used the GAP program MANIFOLD VT [43] to search for vertex-transitive triangulations.

In most of the cases of Table 1, corresponding vertex-transitive examples were found. However, we know from [34], [38] that there are no vertex-transitive triangulations of the non-orientable surface of Euler characteristic $\chi = -4$ with $n = 12$ vertices and no vertex-transitive triangulations of the surface of Euler characteristic $\chi = -7$ with $n = 14$. It also turned out that there are no vertex-transitive triangulations of the surface of Euler characteristic $\chi = -27$ with $n = 18$. Nevertheless, there are equivelar triangulations in these three cases, 500 non-orientable equivelar examples in the case $(\chi, q, n) = (-4, 8, 12), 11300559$ equivelar examples in the case $(-7, 9, 14)$ [42], and at least 412 equivelar examples in the case (−27, 15, 18), as were found with the program **lextri** [60]. In the range $0 > \chi \ge -127$ there remain four undecided cases where we do not know whether vertex-transitive triangulations exist. For the search for vertex-transitive triangulations we made use of the classification of transitive permutations groups of small degree n . A complete list of these groups is available for $n \leq 30$ via the package GAP [25], imposing a restriction $\chi \ge -127$ for the search (since the next non-neighborly candidate case with $\chi = -128$ requires $n = 32$ vertices). For $n = 24, 26,$ and 30, we were only able to process all transitive permutations groups of order larger than n (as well as some of order n) with MANIFOLD VT. Thus, in the open cases, there might be vertex-transitive triangulations corresponding to transitive permutations groups of degree n and order n . In these cases also lextri was too slow to produce examples, but we were able to find respective examples by using bistellar flips (with pruning the search according to the fact that before the final flip of the search exactly four vertices have the wrong degree). The vertex-transitive examples we found are listed in Table 2, with (n, i) the *i*th transitive permutation group of degree n in the GAP catalog, and the size of an orbit indicated in subscript. The non-transitive examples E1, E4, E12, E13, E16, and E19 are listed at the end of this section.

In the range $31 \le n \le 40$, that is, $-128 \ge \chi \ge -230$, there are six further triples $(-128, 30, 32)$, $(-132, 30, 33)$, $(-162, 33, 36)$, $(-168, 34, 36)$, $(-190, 36, 38)$, and $(-195, 36, 39)$. We found orientable examples in the first five cases and non-orientable examples in all six cases. entable examples in the first five cases and non-orientable examples in all six cases.

The non-orientable surface of Euler characteristic $\chi = -3$ is the smallest case, where for an admissible triple, $(\chi, q, n) = (-3, 8, 9)$, there are no vertex-transitive triangulations, whereas two equivelar triangulations exist; see [34], [38], and [61]. Further cases without vertex-transitive, but with equivelar triangulations are the non-orientable cases $(-4, 8, 12)$, $(-7, 9, 14)$, and $(-27, 15, 18)$.

Conjecture 8 Let $\chi < 0, n \geq \left[\frac{1}{2}\right]$ $\frac{1}{2}(7+\sqrt{49-24\chi})\Big\}, \text{ and } q=6-\frac{6\chi(M)}{n}$ $\frac{d(u)}{n}$ an integer. For every admissible triple (χ, q, n) , the non-orientable surface of Euler characteristic χ and, if χ is even, the orientable surface of Euler characteristic χ have q-equivelar triangulations with n vertices.

Conjecture 8 holds for orientable surfaces in the cases $q = 6 + 12k, k \ge 1$; see Section 3. Moreover, we found respective examples for all admissible triples (χ, q, n) with $n \leq 40$ with the help of bistellar flips.

Theorem 9 For each admissible triple (χ, q, n) with $n \leq 40$ there is a q-equivelar triangulation with n vertices of the non-orientable surface of Euler characteristic χ and, if χ is even, then also of the orientable surface of Euler characteristic χ .

For $\chi > 0$, there is exactly one equivelar, vertex-transitive triangulation of S^2 for $(n, q) = (4, 3)$, $(6, 4)$, and $(12, 5)$ each, as well of $\mathbb{R}P^2$ for $(n, q) = (6, 5)$. The 2-torus has equivelar, vertextransitive triangulations (with $q = 6$) for all $n \ge 7$ [1], [11], whereas the Klein bottle has equivelar triangulations (with $q = 6$) for all non-prime $n \geq 9$ [19].

Example E1:

Example E16:

Example E19:

Figure 3: The directed graph $G_{k,7+12k}$.

3 A generalization of Ringel's cyclic 7mod12 series

Ringel's cyclic 7mod12 series [53] provides for every $k \geq 0$ a neighborly triangulation with $n = 7 + 12k$ vertices of the orientable surface of genus $g = 12k^2 + 7k + 1$. We generalize Ringel's series to a two-parameter series of cyclic triangulations.

Theorem 10 For every $k \geq 0$ and every $n \geq 7 + 12k$ the orientable surface of genus $g = kn + 1$ has a triangulation $R_{k,n}$ with n vertices and vertex-transitive cyclic symmetry. The examples $R_{k,n}$ are equivelar of type $\{3, 6 + 12k\}$ and, in terms of $q = 6 + 12k$, the examples have f-vector $f = (n, 3(1+2k)n, 2(1+2k)n) = (n, \frac{nq}{2})$ $\frac{iq}{2}, \frac{nq}{3}$ $\frac{aq}{3}$) and genus $g = 1 + \frac{(q-6)n}{12}$.

Proof: Step I: Let $k \ge 0$ and $n = 7+12k$. Then $f = (n, 3(1+2k)n, 2(1+2k)n) = (n, {n \choose 2})$ $\binom{n}{2}$, $\frac{2}{3}$ $\frac{2}{3}$ $\binom{n}{2}$ $\binom{n}{2}),$ which is the f-vector of a neighborly triangulation of a (orientable or non-orientable) surface of Euler characteristic $\chi = n - \frac{1}{3}$ $rac{1}{3}$ $\binom{n}{2}$ n_2). Ringel's cyclic 7mod12 series [53], [54, Ch. 2.3] provides examples $R_{k,7+12k}$ of triangulated orientable surfaces with the desired parameters.

We follow Ringel's description [54, pp. 25–28] of the series $R_{k,7+12k}$ and consider a directed graph $G_{k,7+12k}$ for any given $k \geq 0$ as displayed in Figure 3, where the leftmost arc and the rightmost arc are to be identified. The $6k + 3$ arcs of the digraph are labeled by $1, \ldots, 6k + 3$ in a way, such that Kirchhoff's rule holds at every vertex, i.e., at each vertex the sum of the values of the ingoing arcs equals the sum of the values of the outgoing arcs. The elements $1, \ldots, 6k + 3$ are regarded as elements of the cyclic group $\mathbb{Z}_n = \mathbb{Z}_{7+12k}$ with inverses $\overline{1}, \ldots, \overline{6k+3}$ and neutral element 0. We let the vertices of $R_{k,7+12k}$ be the elements of \mathbb{Z}_{7+12k} . Next, we read off the vertex-link of the vertex 0 in the triangulation $R_{k,7+12k}$ from Figure 3 by "walking along" the arcs of the digraph: Start with the arc with label, say, 1 and follow the arc in the indicated direction. At a vertex, turn left whenever the vertex is black and turn right whenever the vertex is white, respectively. Thus, at the end of arc 1 we turn left and follow the arc with label $5k + 3$ in the opposite direction of the arc. Then we turn left again and follow the arc $3k + 2$ in opposite direction, etc. Along our walk we record the labels of the arcs, respectively their inverses, if an arc is traversed in opposite direction. Thus, we record (for the link of the vertex 0):

$$
0: \quad 1, \overline{5k+3}, \overline{3k+2}, \overline{3k+3}, \ldots, \overline{2k+2}, \ 2k+1, \ 5k+3, \ 5k+4, \ \ldots \ 2, \ 3k+3.
$$

Every arc of the digraph is traversed twice, once into the indicated direction and once into the opposite direction. Hence, the link of the vertex 0 is a circle with $2 \cdot (6k + 3) = 6 + 12k$ vertices. For convenience, we write, from now on, $n - a$ for the inverse \overline{a} of an element $a \in \mathbb{Z}_n$.

The vertex links of the vertices $1, \ldots, n-1$ are obtained from the link of 0 by cyclic shifts:

0: 1, $n - 5k - 3$, $n - 3k - 2$, $n - 3k - 3$, ..., $n - 2k - 2$, $2k + 1$, $5k + 3$, $5k + 4$, ..., 2, $3k + 3$ 1: 2, $n - 5k - 2$, $n - 3k - 1$, $n - 3k - 2$, ..., $n - 2k - 1$, $2k + 2$, $5k + 4$, $5k + 5$, ..., 3, $3k + 4$ 2: 3, $n - 5k - 1$, $n - 3k$, $n - 3k - 1$, ..., $n - 2k$, $2k + 3$, $5k + 5$, $5k + 6$, ..., 4 , $3k + 5$

Step II: In order to describe a triangulated surface, we can, instead of the vertex links, give the triangles of the surface. In the case of the Ringel series $R_{k,7+12k}$ with cyclic symmetry it is sufficient to give one generating triangle from each of the $2+4k$ orbits of triangles, with each orbit of size $7 + 12k$. We can even choose (and this will be important in Step III) a set of generating triangles with vertices from the subset $\{0, 1, 2, \ldots, 6k + 3\}$; see Table 3.

In each orbit there are three triangles that contain the vertex 0 and thus contribute an edge to the vertex-link of 0. For the choice of generating triangles in Table 3, these $3 \cdot (2 + 4k)$ edges are as follows:

 $[1, 3k + 3],$ $[3k + 2, n - 1],$ $[n - 3k - 3, n - 3k - 2],$ $[2, 3k + 3],$ $[3k + 1, n - 2],$ $[n - 3k - 3, n - 3k - 1],$
 $[3, 3k + 4],$ $[3k + 1, n - 3],$ $[n - 3k - 4, n - 3k],$ $[4, 3k + 4],$ $[3k, n - 4],$ $[n - 3k - 4, n - 3k],$ $[3, 3k + 4],$ $[3k + 1, n - 3],$ $[n - 3k - 4, n - 3k]$, $[4, 3k + 4],$ $[3k, n - 4],$ $[n - 3k - 4, n - 3k]$,
 $[5, 3k + 5],$ $[3k, n - 5],$ $[n - 3k - 5, n - 3k]$, $[6, 3k + 5],$ $[3k - 1, n - 6],$ $[n - 3k - 5, n - 3k + 1]$ $[3k, n-5],$ $[n-3k-5, n-3k],$ $[2k-3, 4k+1], [2k+4, n-2k+3], [n-4k-1, n-2k-4], [2k-2, 4k+1], [2k+3, n-2k+2], [n-4k-1, n-2k-3],$
 $[2k-1, 4k+2], [2k+3, n-2k+1], [n-4k-2, n-2k-3], [2k, 4k+2], [2k+2, n-2k], [n-4k-2, n-2k-2],$ $[2k-1, 4k+2], [2k+3, n-2k+1], [n-4k-2, n-2k-3],$ $[4k+3, 6k+3], [2k, n-4k-3], [n-6k-3, n-2k], [2k+2, 4k+3], [2k+1, n-2k-2], [n-4k-3, n-2k-1],$ $[4k+4, 6k+2]$, $[2k-2, n-4k-4]$, $[n-6k-2, n-2k+2]$, $[4k+4, 6k+3]$, $[2k-1, n-4k-4]$, $[n-6k-3, n-2k+1]$, $[4k+5, 6k+1]$, $[2k-4, n-4k-5]$, $[n-6k-1, n-2k+4]$, $[4k+5, 6k+2]$, $[2k-3, n-4k-5]$, $[n-6k-2, n-2k+3]$, $[5k+1, 5k+5], \quad [4, n-5k-1], \quad [n-5k-5, n-4], \quad [5k+1, 5k+6], \quad [5, n-5k-1], \quad [n-5k-6, n-5],$ $[5k+2, 5k+4], \quad [2, n-5k-2], \quad [n-5k-4, n-2], \quad [5k+2, 5k+5], \quad [3, n-5k-2],$ $[2k+1, 5k+3], [3k+2, n-2k-1], [n-5k-3, n-3k-2], [5k+3, 5k+4], [1, n-5k-3], [n-5k-4, n-1].$

Figure 4: Rule R^* .

Of course, both link-descriptions yield the same triangulated circle as the link of the vertex 0. In Ringel's description, the edges of the link are read off from the digraph $G_{k,7+12k}$ in consecutive order, whereas the description via the set of generating triangles lists the edges of the link in correspondence to the orbits of triangles in which they occur.

Remark: The labeled graph $G_{k,7+12k}$ degenerates for $k = 0$ and $k = 1$, and so do the orbit descriptions of $R_{0,n}$ and $R_{1,n}$. For $k=1$, we have the orbit generating triangles

$$
\begin{array}{lll}\n[0, 1, 6], & [0, 2, 6], \\
[0, 7, 9], & [0, 4, 7], \\
\hline\n& - & - & - \\
[0, 3, 8], & [0, 8, 9],\n\end{array}
$$

whereas for $k = 0$, the orbit generating triangles are

$$
[0, 1, 3], \qquad [0, 2, 3],
$$

which precisely are the orbit generating triangles of Altshuler's series of cyclic tori $T^2(n)$ for $n \geq 7$; see Figure 2.

Step III: For $n \geq 7 + 12k$, let the vertices of $R_{k,n}$ be $\{0,1,2,\ldots,n-1\}$ and define $R_{k,n}$ via the same set of $2 + 4k$ generating triangles from Table 3 with respect to the action of the cyclic group \mathbb{Z}_n . Since all vertices of the generating triangles are elements of the set $\{0, 1, 2, \ldots, 6k+3\}$, the link of vertex 0 has the same description as before in the case $n = 7 + 12k$. In other words, the 'symbol' $n = 7 + 12k$ in the link of 0 in $R_{k,7+12k}$ is replaced by the 'symbol' $n \ge 7 + 12k$ in the link of 0 in $R_{k,n}$. This way, no unwanted identifications of vertices in the link take place, so the link is one circle of length $6 + 12k$. It follows, that $R_{k,n}$ is a triangulated surface, which is equivelar of type $\{3, 6 + 12k\}$, has f-vector $f = (n, 3(1 + 2k)n, 2(1 + 2k)n)$, and genus $g = kn + 1$.

It remains to show that $R_{k,n}$ is orientable for all $n \geq 7 + 12k$, $k \geq 0$. In Ringel's description of the link of 0 in the case $n = 7 + 12k, k \ge 0$, the edges of the link are listed in consecutive order, which induces an orientation of the triangles of the star of the vertex 0. Since the stars of the other vertices are obtained from the star of 0 by cyclic shifts, all vertex-stars inherit an orientation from the orientation of the star of 0. For $n = 7+12k$, the orientations of the individual vertex-stars are compatible, as is expressed by Ringel's rule R^* for the vertex-links [54, p. 28]:

Rule R^* : If in row *i* one has $i : \ldots jkl \ldots$, then row *k* appears as $k : \ldots lij \ldots$ (see Figure 4).

In particular, rule R^* holds for row 0, that is, if we have $0: \ldots$ *jkl* ..., then row k appears as $k: \ldots l0j \ldots$ If we apply a cyclic shift to row k by adding $-k$, we obtain $0: \ldots l-k, -k, j-k \ldots$

For $R_{k,7+12k}$, the link of 0 has the set $P \cup N$ as its set of vertices, with $P = \{1, 2, \ldots, 6k + 3\}$ and $N = \{n-1, n-2, \ldots, n-(6k+3)\}\.$ In the transition from $R_{k,7+12k}$ to $R_{k,n}$, that is, from $n = 7 + 12k$ to $n \ge 7 + 12k$, a vertex $a \in P$ of the link of 0 remains unchanged, whereas a vertex b in the link of 0 in $R_{k,7+12k}$ with $b \in N$ can be written as $b = (7+12k) - r$ with $r \in P$, and is replaced by $n-r$. As a consequence, if we have $0: \ldots jkl \ldots$ for the row 0 in $R_{k,n}$ for $n \geq 7+12k$, then also $0: \ldots l-k, -k, j-k, \ldots$, as originally in $R_{k,7+12k}$. If we now add k to this row, we obtain $k : \ldots l0j \ldots$, so the orientation of the star of 0 is compatible with the orientations of all the stars of the vertices in the link of 0. By the cyclic symmetry it follows that the orientations of any two neighboring vertex-stars are compatible. Therefore, $R_{k,n}$ is orientable. \Box

The generalized Ringel series $R_{k,n}$ interpolates between Ringel's cyclic neighborly series $R_{k,7+12k}$ and Altshuler's cyclic series of 6-equivelar tori $T^2(n) = R_{0,n}$. The examples $R_{k,n}$ settle Conjecture 8 for $q = 6 + 12k$, $k \ge 1$, for the orientable surfaces with $\chi = -2kn$.

Corollary 11 The two subseries $R_{k,7+12k+1}$ and $R_{k,7+12k+2}$ of the generalized Ringel series provide for $k \geq 1$ infinitely many examples of non-neighborly cyclic triangulations with $q = 6+12k =$ $\frac{1}{2}$ $\frac{1}{2}(5+\sqrt{49-24\chi})\Big\vert$, i.e., with equality in (8). By subdivision, these examples yield $2q$ -covered triangulations with equality in (11).

Proof: Let for $k \geq 1$, $n = 7 + 12k + m$ with $m \geq 0$. By Theorem 10, the examples $R_{k,n}$ of the generalized Ringel series have Euler characteristic $\chi = -2kn$.

For $m = 0, k \ge 1$, we have $n = 7 + 12k$. Therefore, the example $R_{k,7+12k}$ is neighborly with $q=\left\lfloor \frac{1}{2}\right\rfloor$ $\frac{1}{2}(5+\sqrt{49-24\chi})\Big|=\Big|\frac{1}{2}\Big|$ $\frac{1}{2}(5+\sqrt{49+24\cdot 2k(7+12k)})\bigg|=\bigg|\frac{1}{2}\bigg|$ $\frac{1}{2}(5+\sqrt{(7+24k)^2})\Big|=6+12k.$ We next determine those $m \geq 1$, such that $\frac{1}{2}$ $\frac{1}{2}(5+\sqrt{49+24\cdot 2k(7+12k+m)})\Big| = 6+12k,$ which is equivalent to find those $m \geq 1$, for which $\sqrt{(7 + 24k)^2 + 48km} < (7 + 24k) + 2$, that is, $m < 2 + \frac{2}{3k}$. Thus, $1 \le m \le 2$.

Corollary 12 For $k \geq 1$, the series $R_{k,7+12k+1}$ provides vertex-minimal triangulations of the orientable surfaces of Euler characteristic $\chi = -2kn$ with vertex-transitive cyclic symmetry on $n = 7 + 12k + 1$ vertices.

Proof: For
$$
k \ge 1
$$
, $\left[\frac{1}{2}(7 + \sqrt{49 + 24\chi})\right] = \left[\frac{1}{2}(7 + \sqrt{49 + 24 \cdot 2k(7 + 12k + 1)})\right] = 7 + 12k + 1.$

References

- [1] A. Altshuler. Polyhedral realization in \mathbb{R}^3 of triangulations of the torus and 2-manifolds in cyclic 4-polytopes. Discrete Math. 1, 211–238 (1971).
- [2] A. Altshuler, J. Bokowski, and P. Schuchert. Neighborly 2-manifolds with 12 vertices. J. Comb. Theory, Ser. A 75, 148–162 (1996).
- [3] A. Altshuler and U. Brehm. Neighborly maps with few vertices. Discrete Comput. Geom. 8, 93–104 (1992).
- [4] J. Bokowski. A geometric realization without self-intersections does exist for Dyck's regular map. Discrete Comput. Geom. 4, 583–589 (1989).
- [5] J. Bokowski and A. Eggert. Toutes les réalisations du tore de Möbius avec sept sommets/All realizations of Möbius' torus with 7 vertices. *Topologie Struct.* 17, 59–78 (1991).
- [6] J. Bokowski and A. Guedes de Oliveira. On the generation of oriented matroids. Discrete Comput. Geom. 24, 197–208 (2000).
- [7] J. Bokowski and J. M. Wills. Regular polyhedra with hidden symmetries. Math. Intell. 10, No. 1, 27–32 (1988).
- [8] U. Brehm. Maximally symmetric polyhedral realizations of Dyck's regular map. Mathematika 34, 229–236 (1987).
- [9] U. Brehm. Polyhedral maps with few edges. Topics in Combinatorics and Graph Theory: Essays in Honour of Gerhard Ringel (R. Bodendiek and R. Henn, eds.), 153–162. Physica-Verlag, Heidelberg, 1990.
- [10] U. Brehm, B. Datta, and N. Nilakantan. The edge-minimal polyhedral maps of Euler characteristic −8. Beitr. Algebra Geom. 43, 583–596 (2002).
- [11] U. Brehm and W. Kühnel. Equivelar maps on the torus. *Eur. J. Comb.* 29, 1843–1861 (2008).
- [12] U. Brehm and E. Schulte. Polyhedral maps. Handbook of Discrete and Computational Geometry (J. E. Goodman and J. O'Rourke, eds.), Chapter 18, 345–358. CRC Press, Boca Raton, FL, 1997.
- [13] U. Brehm and J. M. Wills. Polyhedral manifolds. Handbook of Convex Geometry, Volume A (P. M. Gruber and J. M. Wills, eds.), Chapter 2.4, 535–554. North-Holland, Amsterdam, 1993.
- [14] M. D. E. Conder. Regular maps and hypermaps of Euler characteristic −1 to −200. J. Comb. Theory, Ser. A 99, 455–459 (2009).
- [15] H. S. M. Coxeter and W. O. J. Moser. Generators and Relations for Discrete Groups. Ergebnisse der Mathematik und ihrer Grenzgebiete 14. Springer-Verlag, Berlin, 1957. Fourth edition, 1980.
- [16] A. Császár. A polyhedron without diagonals. Acta Sci. Math., Szeged 13, 140–142 (1949– 1950).
- [17] B. Datta. A note on the existence of $\{k, k\}$ -equivelar polyhedral maps. Beitr. Algebra Geom. 46, 537–544 (2005).
- [18] B. Datta and N. Nilakantan. Equivelar polyhedra with few vertices. Discrete Comput. Geom. 26, 429–461 (2001).
- [19] B. Datta and A. K. Upadhyay. Degree-regular triangulations of torus and Klein bottle. Proc. Indian Acad. Sci., Math. Sci. 115, 279–307 (2005).
- [20] B. Datta and A. K. Upadhyay. Degree-regular triangulations of the double-torus. Forum Math. 18, 1011–1025 (2006).
- [21] W. Dyck. Ueber Aufstellung und Untersuchung von Gruppe und Irrationalität regulärer Riemann'scher Flächen. Math. Ann. 17, 473–509 (1880).
- [22] W. Dyck. Notiz über eine reguläre Riemann'sche Fläche vom Geschlechte drei und die zugehörige "Normalcurve" vierter Ordnung. *Math. Ann.* **17**, 510–516 (1880).
- [23] M. N. Ellingham and C. Stephens. Triangular embeddings of complete graphs (neighborly maps) with 12 and 13 vertices. J. Comb. Des. 13, 336–344 (2005).
- [24] G. Fijavž. Minor-minimal 6-regular graphs in the Klein bottle. Eur. J. Comb. 25, 893–898 (2004).
- [25] The GAP Group. GAP Groups, Algorithms, and Programming, Version 4.4.9. http: //www.gap-system.org, 2006.
- [26] P. J. Heawood. Map-colour theorem. *Quart. J. Pure Appl. Math.* **24**, 332–338 (1890).
- [27] L. Heffter. Ueber das Problem der Nachbargebiete. Math. Ann. 38, 477–508 (1891).
- [28] J. P. Huneke. A minimum-vertex triangulation. J. Comb. Theory, Ser. B 24, 258–266 (1978).
- [29] M. Jamet. Polyhedral maps with cyclic symmetry. Diplomarbeit, Technische Universität Dresden, 2001, 73 pages.
- [30] M. Joswig and F. H. Lutz. One-point suspensions and wreath products of polytopes and spheres. J. Comb. Theory, Ser. A 110, 193–216 (2005).
- [31] M. Jungerman and G. Ringel. Minimal triangulations on orientable surfaces. Acta Math. 145, 121–154 (1980).
- [32] T. Katahira. Infinite series of triangulations on closed surfaces with vertex covers of given degree. Graphs Comb. 20, 73–78 (2004).
- [33] F. Klein. Ueber die Transformation siebenter Ordnung der elliptischen Functionen. Math. Ann. 14, 428–471 (1879).
- [34] E. G. Köhler and F. H. Lutz. Triangulated Manifolds with Few Vertices: Vertex-Transitive Triangulations I. arXiv:math.GT/0506520, 2005, 74 pages.
- [35] W. Kühnel. Higherdimensional analogues of Császár's torus. Result. Math. 9, 95–106 (1986).
- [36] W. Kühnel and G. Lassmann. Combinatorial d-tori with a large symmetry group. Discrete Comput. Geom. 3, 169–176 (1988).
- [37] W. Kühnel and G. Lassmann. Permuted difference cycles and triangulated sphere bundles. Discrete Math. **162**, 215–227 (1996).
- [38] F. H. Lutz. Triangulated Manifolds with Few Vertices and Vertex-Transitive Group Actions. Dissertation. Shaker Verlag, Aachen, 1999, 146 pages.
- [39] F. H. Lutz. Császár's torus. *Electronic Geometry Models* No. 2001.02.069 (2002). http: //www.eg-models.de/2001.02.069.
- [40] F. H. Lutz. Triangulated Manifolds with Few Vertices: Centrally Symmetric Spheres and Products of Spheres. arXiv:math.MG/0404465, 2004, 26 pages.
- [41] F. H. Lutz. Equivelar and d-covered triangulations of surfaces. II. Cyclic triangulations and tessellations. arXiv:1001.2779, 2010, 27 pages.
- [42] F. H. Lutz. The Manifold Page, 1999–2010. http://www.math.tu-berlin.de/diskregeom/ stellar/.
- [43] F. H. Lutz. MANIFOLD VT, Version Apr/2002. http://www.math.tu-berlin.de/ diskregeom/stellar/MANIFOLD_VT, 2002.
- [44] F. H. Lutz and T. Sulanke. Quadrangulations of surfaces. In preparation.
- [45] P. McMullen, E. Schulte, and J. M. Wills. Infinite series of combinatorially regular polyhedra in three-space. Geom. Dedicata 26, 299–307 (1988).
- [46] P. McMullen, Ch. Schulz, and J. M. Wills. Equivelar polyhedral manifolds in E^3 . Isr. J. Math. 41, 331–346 (1982).
- [47] P. McMullen, Ch. Schulz, and J. M. Wills. Polyhedral 2-manifolds in E^3 with unusually large genus. Isr. J. Math. 46, 127–144 (1983).
- [48] A. F. Möbius. Mittheilungen aus Möbius' Nachlass: I. Zur Theorie der Polyëder und der Elementarverwandtschaft. Gesammelte Werke II (F. Klein, ed.), 515–559. Verlag von S. Hirzel, Leipzig, 1886.
- [49] S. Negami. Classification of 6-regular Klein-bottlal graphs. Res. Rep. Inf. Sci. T.I.T. A-96 (1984).
- [50] S. Negami and A. Nakamoto. Triangulations on closed surfaces covered by vertices of given degree. Graphs Comb. 17, 529–537 (2001).
- [51] J. Pfeifle and G. M. Ziegler. Many triangulated 3-spheres. Math. Ann. 330, 829–837 (2004).
- [52] G. Ringel. Wie man die geschlossenen nichtorientierbaren Flächen in möglichst wenig Dreiecke zerlegen kann. Math. Ann. 130, 317–326 (1955).
- [53] G. Ringel. Über das Problem der Nachbargebiete auf orientierbaren Flächen. Abh. Math. Sem. Univ. Hamburg 25, 105–127 (1961).
- [54] G. Ringel. Map Color Theorem. Grundlehren der mathematischen Wissenschaften 209. Springer-Verlag, Berlin, 1974.
- [55] T. Rörig and G. M. Ziegler. Polyhedral surfaces in wedge products. $arXiv:0908.3159, 2009,$ 17 pages.
- [56] L. Schewe. Non-realizable minimal vertex triangulations of surfaces: Showing non-realizability using oriented matroids and satisfiability solvers. arXiv:0801.2582, 2008, 14 pages.
- [57] E. Schulte and J. M. Wills. A polyhedral realization of Felix Klein's map $\{3, 7\}$ on a Riemann surface of genus 3. J. Lond. Math. Soc., II. Ser. **32**, 539–547 (1985).
- [58] E. Schulte and J. M. Wills. Geometric realizations for Dyck's regular map on a surface of genus 3. Discrete Comput. Geom. 1, 141–153 (1986).
- [59] E. Schulte and J. M. Wills. On Coxeter's regular skew polyhedra. Discrete Math. 60, 253–262 (1986).
- [60] T. Sulanke. Source for lextri. http://hep.physics.indiana.edu/~tsulanke/graphs/ lex/downloads/, 2009. Version 0.16.
- [61] T. Sulanke and F. H. Lutz. Isomorphism free lexicographic enumeration of triangulated surfaces and 3-manifolds. Eur. J. Comb. **30**, 1965–1979 (2009).
- [62] A. K. Upadhyay. A note on upper bound for d-covered triangulations of closed surfaces. Int. J. Pure Appl. Math., to appear.

Frank H. Lutz Institut für Mathematik Technische Universität Berlin Straße des 17. Juni 136 10623 Berlin, Germany lutz@math.tu-berlin.de

Thom Sulanke Department of Physics Indiana University Bloomington, Indiana 47405, USA tsulanke@indiana.edu

Ashish K. Upadhyay, Anand K. Tiwari Department of Mathematics Indian Institute of Technology Patna Patliputra Colony, Patna – 800 013, India {upadhyay,anand}@iitp.ac.in